Study Protocol

Efficacy and Mechanisms of Cannabis Oil for Alleviating Side Effects of Breast Cancer Chemotherapy (CBC2): Protocol for Randomized Controlled Trial.

May Soe Thu[1,2], Krit Pongpirul[3,4,5*], Mawin Vongsaisuwon[6], Chanida Vinayanuwattikun[7,8], Kamonwan Banchuen[3], Thunnicha Ondee[3], Sunchai Payungporn[9], Phanupong Phutrakool[3,10], Preecha Nootim[11], Pajaree Chariyavilaskul[12], Sarocha Cherdchom[3], Kulthanit Wanaratna[11], Nattiya Hirankarn[1].

Department of Microbiology, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand.

Department of Infection Biology & Microbiomes, University of Liverpool, Liverpool, UK.

Department of Preventive and Social Medicine, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand.

Bumrungrad International Hospital, Bangkok, Thailand.

Department of International Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA

Department of Surgery, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand.

Division of Medical Oncology, Department of Medicine, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand.

King Chulalongkorn Memorial Hospital, Bangkok, Thailand.

Department of Biochemistry, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand.

Chula Data Management Center, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand.

Department of Thai Traditional and Alternative Medicine, Ministry of Public Health, Nonthaburi, Thailand.

Department of Pharmacology, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand.

*doctorkrit@gmail.com

Abstract

Background

In a pilot study using both cannabidiol (CBD) and tetrahydrocannabinol (THC) as single agents in advanced cancer patients undergoing palliative care in Thailand, the doses were generally well tolerated, and the outcome measure of total symptom distress scores showed overall symptom benefit. The current study aims to determine the intensity of the symptoms experienced by breast cancer patients receiving chemotherapy, to explore the microbiome profile, cytokines, and bacterial metabolites before and after the treatment with cannabis oil or no cannabis oil, and to study the pharmacokinetics parameters and pharmacogenetics profile of the doses.

Methods

A randomized, double-blinded, placebo-controlled trial will be conducted on the metastatic breast cancer cases currently receiving chemotherapy at King Chulalongkorn Memorial Hospital (KCMH), Bangkok, Thailand. Block randomization will be used to allocate the patients into three groups: Ganja Oil (THC 2 mg/ml; THC 0.08 mg/drop, and CBD 0.02 mg/drop) group), Metta Osot (THC 81 mg/ml; THC 3 mg/drop), and placebo oil. The Edmonton Symptom Assessment System (ESAS), microbiome profile, cytokines, and bacterial metabolites will be assessed before and after the interventions.

Thai Clinical Trial Registration: TCTR20220809001

Introduction

Breast cancer patients reported lower quality of life (QoL) globally (1-3) and chemotherapy was a factor in worsening QoL among the patients (4). Despite standard medical treatment, the impact of integrating complementary and alternative medicine was introduced into the current practice (5). Therefore, cannabinoids from *Cannabis sativa* have become an attractive choice due to their prominent impact in reducing pain associated with chemotherapy and tumor-associated symptoms (6-8).

In a pilot study using both cannabidiol (CBD) and tetrahydrocannabinol (THC) as single agents in advanced cancer patients undergoing palliative care, the doses of THC and CBD were generally well tolerated, and the outcome measure of total symptom distress scores (TSDSs) showed overall symptom benefit (9). The average daily amounts that were tolerated were 300 mg of CBD and 10 mg of THC, respectively (range 5–30 mg), and 43% met the response criteria of ≥6 point reduction in TSDSs (9).

Several clinically relevant mouse models of breast cancer have shown that naturally sourced cannabinoids possess anti-tumor activities, with CBD as the most potent inhibitor of cancer cell proliferation (10). In experimental models, psychoactive THC and non-psychoactive CBD disturbed the disease progression, and its effects on signaling pathways within cancer cells following activation of G-protein coupled cannabinoid-receptors (CB-Rs), other receptors such as GPR55 and TRPV1, and in a receptor-independent way. Both *in vitro* and *in vivo* studies have revealed that cannabinoids can activate these receptors, leading to anti-tumorigenic activity in most cases (11).

Recently, the human microbiota raised so much attention as a major risk modulator since they play a prominent role in the regulation of steroid hormone metabolism, through the stimulation of different enzymes, such as hydroxysteroid dehydrogenase (12, 13). Despite the lack of convincing evidence that breast cancer is caused by dysbiosis, the comparison of several breast samples reveals variations in the abundance and microbial diversity of some particular genera between healthy individuals and patients (14, 15). Additionally, it recognizes that various breast cancer subtypes are correlated to various intestinal microbial signatures (16).

Together with the dysbiosis of intestinal microbiota, it is also related to the dysfunction in the intestine leading to inflammatory and immune disorders. Subsequently, this indicates to occur diseases via microbial translocation. It was suggested that the intestinal microbiota is propagated to the skin, then reached the breast, and the dendritic cells might

transport the bacteria to the breast tissue (17). Therefore, it becomes critical to explore the bacterial translocation markers to understand this unclear underlying mechanism.

In addition, decreased fungal richness and diversity have been observed in patients with cancer (18) but whether certain species and their specific metabolic outputs define, and support carcinogenesis has not been fully explored. Fungal signatures are seen to be different in the tumor microenvironment of various hormonal subtypes of breast cancer, and higher fungal diversity was found in estrogen receptor-positive breast cancer patients in contrast to the triple-negative breast cancer group (16, 19).

Despite recent advances, the function of the breast skin microbiota is still uncertain because the skin microorganisms and the underlying mucosal immune function interact largely. According to a systematic analysis of the skin microbiota in breast cancer, metastases and breast cancer have been associated with large abundances of skin commensals, particularly several species of *Staphylococcus* (20). For microbial transfer to underlying tissue, several pathways have been hypothesized, including retrograde transfer through ductal networks, permeabilization of the epidermal barrier, and migration through the nipple-aspirate fluid (20).

Particularly, there are hints that breast cancer and the oral microbiome are related. Women with periodontal disease, which is brought on by particular bacteria like the red complex (*Porphyromonas gingivalis, Tannerella forsythia, and Treponema denticola*), and the orange complex (*Fusobacterium nucleatum, Prevotella intermedia, Prevotella nigrescens, Peptostreptococcus micros, Streptococcus constellatus, Eubacterium nodatum, Campylobacter showae, Campylobacter gracilis, and Campylobacter rectus*), have been found to have an increased risk of breast cancer in some studies (21-23).

Currently, the Herb and Thai Traditional Medicine Development Division, Department of Thai Traditional and Alternative Medicine (DTAM), Ministry of Public Health, Thailand, manufactures Metta Osot (THC 81 mg/ml; THC 3 mg/drop) and Ganja Oil (THC 2 mg/ml; THC 0.08 mg/drop and CBD 0.02 mg/drop), both of which are produced under license number 13/2562. The preliminary analysis of the Thai Cannabis Practice Patterns and Quality of Life (Thai Cannabis PQ) study revealed that cancer patients had significantly lower baseline EQ-5D-5L (0.80 ± 0.29 vs 0.85 ± 0.22) and ESAS pain (3.44 ± 3.21 vs 4.05 ± 3.30) and anxiety (2.03 ± 2.77 vs 2.37 ± 2.85) but higher tiredness (3.25 ± 3.08 vs 2.89 ± 2.83), nausea (1.00 ± 2.07 vs 0.69 ± 1.66), poor appetite (2.21 ± 2.94 vs 1.20 ± 2.20), and shortness of breath (1.74 ± 2.61 vs 1.22 ± 2.17) than non-cancer individuals (24). Both Ganja Oil and Metta Osot were added to the National List of Essential Medicines (NLEM) (25).

The pharmacokinetic profile of cannabinoids depends on routes of medication such as oral dosage and inhalation. A study revealed that the median value of the time taken to achieve the highest levels (T_{max}) of sublingual drops for CBD only and a combination of CBD and THC was 2.17 hours (range 1 – 4 hours) and 1.67 hours (range 1 – 3 hours) and the mean of the highest concentration of the drops (C_{max}) was 2.05 (SD = 0.92) and 2.58 (SD = 0.68) nanograms/ml, respectively (26).

Hence, the current randomized study aims to explore the efficacy of Thai cannabis oil on QoL of individuals diagnosed with advanced breast cancer and treated with chemotherapies and the adverse effects of Thai cannabis oil along with the study of microbial alteration and diversity by cannabis among those patients. We aim to determine the intensity of the symptoms experienced by cancer patients using the Edmonton Symptom Assessment System (ESAS), to explore the microbiome profile, cytokines, and bacterial metabolites in hormone-dependent and independent breast cancer patients at baseline, to study the alteration of microbiome profile, cytokines, and bacterial metabolites before and after the treatment with the cannabis oil or no cannabis, and to study the pharmacokinetics parameters and pharmacogenetics profile.

Materials and methods

The prospective randomized placebo-controlled study will recruit the participants at the King Chulalongkorn Memorial Hospital (KCMH) in Bangkok, Thailand. The study is scheduled to launch approximately in January 2023. The trial was approved by the Institutional Review Board of the Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand (COA No. 1713/2022; IRB No.0548/65) on 28[th] December 2022 in compliance with the international guidelines for human research protection as Declaration of Helsinki, The Belmont Report, CIOMS Guideline and International Conference on Harmonization in Good Clinical Practice (ICH-GCP). Only the patients providing written informed consent will proceed for the participation of the trial. All the data will be maintained confidentially.

Target population

We are including 90 adult breast cancer women patients currently receiving chemotherapy at King Chulalongkorn Memorial Hospital (KCMH) with an anticipated starting recruitment approximately in January 2023.

    Inclusion criteria

    1. Women aged 18-70 years old who were diagnosed with breast cancer

2. Metastatic breast cancer patients who are currently receiving different doses of chemotherapy regimen with high/low emetic risk such as AC (cyclophosphamide and doxorubicin), AC followed by T (cyclophosphamide, doxorubicin, and paclitaxel), DC (cyclophosphamide and docetaxel), CMF (cyclophosphamide, fluorouracil, and methotrexate), and Paclitaxel

3. Breast cancer patients at KCMH during the study period

4. Participants provided written informed consent

5. Individuals having at least ESAS score ≥3

Exclusion criteria

1. Women who have contraindications for chemotherapy such as anaphylaxis

2. Women who have serious chemotherapy complications such as anaphylactic shock, bone marrow suppression, liver failure, acute kidney injury, and cardiac arrest

3. Women who are scheduled for elective surgery or other procedures requiring general anesthesia during the study

4. Women who are pregnant or lactating, and who are planning for pregnancy during the study

5. Women who are terminally ill or inappropriate for placebo medication

6. Women with any other significant disease or disorder which, in the opinion of the investigator, may have put the patient at risk because of participation in the study, or may have influenced the result of the study, or the patient's ability to participate in the study

7. Women with a known history of substance abuse

8. Women with a history of cannabis ingredients allergy or suspected of having an adverse reaction to cannabis

9. Women using antibiotics within the past 3 months or using NSAIDs drugs

10. Women consuming probiotic or prebiotic supplements

Trial design and follow-up

Patient demographic information will be reviewed, and selection and recruitment will be performed using the inclusion and exclusion criteria. The medical record will be used to collect clinical and laboratory investigation data related to breast cancer staging, hormonal breast cancer subtypes, baseline symptoms, current medication including chemotherapy regime and other herbal use, emetic condition, history of surgery, history of other cancers,

sites of metastasis, history of smoking, history of alcohol use, history of drug abuse, and history of radiotherapy.

For recruitment, we will give the trial information to potential participants using surgeons and nurses in each surgical ward or outpatient clinic and a participant information sheet to confirm the participant's understanding of the trial objective, procedure, benefits, and risks. The recruitment will be performed in Thai language as native. If potential participants are willing to participate in the trial, they will be asked to sign an informed consent form with the understanding that they can quit the trial at any time without any consequence.

The recruited patients will be randomly allocated into 3 groups: Ganja Oil (THC 2 mg/ml; THC 0.08 mg/drop, and CBD 0.02 mg/drop) group), Metta Osot (THC 81 mg/ml; THC 3 mg/drop), and placebo oil groups. The allocation will be conducted with a 1:1:1 ratio using block randomization with a block of five by the table of random numbers for creating randomization ID which will be enveloped in an opaque envelope to conceal allocation, mask clinicians, data collectors, and patients. Following the cannabis practice patterns of the preliminary study (24), all the participants will be received cannabis or placebo oil once per day before bedtime for 90 days.

There will be four visits for the recruited participants during the trial. At the first visit, the enrollment will be performed by research investigators by checking the eligibility screening, informed consent form, and allocation plan. After setting the random assignment, further assessment will be conducted using the case report forms approved by the IRB. Then, the blood, saliva, stool, and skin samples will be collected as a baseline and started the treatment a day 1 trial. The second visit will be on another day to assess the adverse drug reaction by a medical officer. The first follow-up treatment as the third visit will be on day 30 and the participants will be checked. Day 90 will be the last visit and second follow-up of the daily treatment at which all the participants will be assessed, and the blood, saliva, stool, and skin samples will be collected post-treatment.

After the trial is finished, patients will be asked to stop taking cannabis oil from any sources. They will be monitored for another month for withdrawal symptoms.

Safety

It has been confirmed the effectiveness of the cannabis products in the preliminary analysis and monitored the side effect at the trials.

Statistical considerations

Sample size estimation

We target 30 patients in each arm considering the maximum follow-up loss of 50%. Since the target population is metastatic breast cancer cases, and both questionnaires and biological samples such as blood, stool, and breast skin scrapping will be collected before and after the treatment, it could increase the withdrawal rate at any time during the trial.

Statistical analysis

The statistical analysis will be performed on an intention-to-treat basis using Stata/MP software version 16.0 (StataCorp 2017, College Station, TX). Descriptive statistics will be carried out using mean with standard deviation for normally distributed continuous data, and median with interquartile range for non-normal distributed continuous data. Categorical data will be presented as counts and percentages and will be tested for significance with the Chi-square test. Continuous data will be assessed for normal distribution using a histogram and Shapiro–Wilk test. One-way ANOVA with Bonferroni correction will be used for normally distributed continuous data and the Kruskal-Wallis test will be used for non-normal distributed continuous data.

To identify the alpha diversity, Shannon's diversity index is used for evenness and Chao 1 index for the richness of operational taxonomic units (OTUs). For the detection of beta diversity, the Jaccard dissimilarity index is used, and it is statistically described by permutational multivariate analysis of variance using distance matrices (PERMANOVA) test. A statistically significant level is defined as $p < 0.05$. Correction for multiple analyses was not done but they will not be included in the conclusion of the study as strong findings. Sensitivity analysis using some subgroup to show robustness will be considered if the primary outcome was statistical significance.

Discussion

With a rising incidence of 22,158 new cases occupying a prevalence of 22.8% amongst all female breast cancer cases in Thailand in 2020 (27, 28), more emphasis becomes on the long-term quality of life and measurement of patients' symptoms using validated tools (29, 30). Thailand actively launched cannabis oil in the market in 2018, subsequently, the FDA (Food and Drug Administration) authorized all hospitals running under the Public Health Ministry for medical cannabis to be available on prescription to patients for approved conditions like cancer chemotherapy for relief of pain, to counter inflammation, and so on (31). Following the preliminary analysis of the Thai Cannabis PQ study, it is expected to improve the chemotherapy-related symptoms and their quality of life, the adverse effects as well.

The CBC2 study is a 1-year prospective study, and it is designed as a randomized, double-blinded, placebo-controlled trial that would be conducted on current female breast cancer cases attending a tertiary medical school. This is a phase 4 clinical trial to obtain additional information; the suggested therapy is an additional drug for treating breast cancer, and it has predictable and controllable side effects.

Since some studies found a particular pattern of bacterial and fungal profiles in different breast cancer subtypes (16, 19), we expected to identify the similar microbial profile in hormone-dependent and independent breast cancer patients before and after the treatment with cannabis oil and placebo oil and evaluate the statistical changes by the treatment.

Typically, dysbiosis is an imbalance of healthy and derogatory bacterial community structures leading to the development of chronic inflammatory conditions and cancers, particularly in the gastrointestinal tract (32, 33). Therefore, we will investigate proinflammatory cytokines such as interleukin-17 (IL-17), interferon-γ (IFN-γ), and inflammatory cytokines like interleukin-10 (IL-10), and tumor growth factor-β (TGF-β), then will explore the differences at the baseline and after the treatment.

Moreover, lipopolysaccharide-binding protein (LBP) and soluble CD14 (sCD14) levels as markers of translocation of bacteria from the gut to the blood will be measured to explore the possibility of a gut-breast cancer microbiota axis (34) and to evaluate the influence that cannabis oil treatment may have on the prevention of bloodstream transmission of oncogenic bacteria from the intestine to the breast. In addition, we will also examine levels of equol which is a potent estrogenic metabolite produced by intestinal bacteria (35).

Short-chain fatty acids (SCFAs), such as acetate, propionate, and butyrate, are present in the intestine exceeding the concentration of 100 mM, and they are mainly produced by 2 major gut bacterial phyla: *Bacteroidetes* and *Firmicutes* (36). And branched chain iso-SCFAs, which play a role in membrane permeability and fluidity (37), will be investigated as bioactive bacterial metabolites in breast cancer. The SCFAs (beneficial) and iso-SCFAs (potentially harmful) will be indicators for microbial dysbiosis and bowel health (38) in participants, pre- and post-cannabis oil treatment.

Primary and secondary bile acids will be diagnosed as bioactive bacterial metabolites in breast cancer. From a range of bile acid profile, deoxycholic acid (DCA) and chenodeoxycholic acid (CDCA) at the physiological level leads to significant reductions in cell invasion, migration, adhesion, and survival, but not to cytotoxicity or apoptosis (39). In addition, lithocholic acid (LCA) stimulates oxidative stress in breast cancer patients which can reduce cancer cell proliferation (40).

The pharmacokinetic profile of cannabis oil including THC and CBD on routes of medication through oral administration will be examined to achieve the T_{max} and C_{max} of the drugs (26). As is constantly expanding the cannabis treatment, it becomes important to predict the genes whose variations are responsible for the presence of the therapeutic and side effects of cannabis oil (41).

Conclusions

The clinical trial will provide the efficacy of Thai cannabis oil in metastatic breast cancer female patients in Thailand mainly in terms of alleviation of chemotherapy-related side effects. Furthermore, it will obtain secondary factors such as QoL indexes, adverse effects of the dosage, alteration of microbiome profile, bacterial metabolites, and cytokines before and after the use of the cannabis oil, pharmacokinetic parameters, and pharmacogenetic profiles.

Data Availability

All data produced in the present work are contained in the manuscript.

Funding

This study is financially supported by the Thai Traditional Medical Knowledge Fund, Department of Thai Traditional and Alternative Medicine, Ministry of Public Health, Thailand.

Acknowledgments

The authors are grateful to the Research Affairs, Faculty of Medicine, Chulalongkorn University for their support.

Author Contributions

Conceptualization: Krit Pongpirul, Mawin Vongsaisuwon, May Soe Thu, Nattiya Hirankarn

Data curation: Kamonwan Banchuen, May Soe Thu, Phanupong Phutrakool, Thunnicha Ondee, Sunchai Payungporn, Pajaree Chariyavilaskul

Formal analysis: Phanupong Phutrakool, Kamonwan Banchuen, Krit Pongpirul, May Soe Thu

Methodology: Krit Pongpirul, Kamonwan Banchuen, Kulthanit Wanaratna, Pajaree Chariyavilaskul, Preecha Nootim, May Soe Thu, Sarocha Cherdchom, Sunchai Payungporn, Thunnicha Ondee

Supervision: Chanida Vinayanuwattikun, Krit Pongpirul, Mawin Vongsaisuwon, Nattiya Hirankarn, Sunchai Payungporn

Writing – original draft: May Soe Thu, Krit Pongpirul

Writing – reviewing and editing: Krit Pongpirul, May Soe Thu

References

1. Hashemi SM, Balouchi A, Al-Mawali A, Rafiemanesh H, Rezaie-Keikhaie K, Bouya S, et al. Health-related quality of life of breast cancer patients in the Eastern Mediterranean region: a systematic review and meta-analysis. Breast Cancer Res Treat. 2019;174(3):585-96.

2. Yanez B, Thompson EH, Stanton AL. Quality of life among Latina breast cancer patients: a systematic review of the literature. J Cancer Surviv. 2011;5(2):191-207.

3. Paraskevi T. Quality of life outcomes in patients with breast cancer. Oncol Rev. 2012;6(1):e2.

4. Redeker NS, Lev EL, Ruggiero J. Insomnia, fatigue, anxiety, depression, and quality of life of cancer patients undergoing chemotherapy. Sch Inq Nurs Pract. 2000;14(4):275-90; discussion 91-8.

5. Mokhtari-Hessari P, Montazeri A. Health-related quality of life in breast cancer patients: review of reviews from 2008 to 2018. Health Qual Life Outcomes. 2020;18(1):338.

6. Tomko AM, Whynot EG, Ellis LD, Dupre DJ. Anti-Cancer Potential of Cannabinoids, Terpenes, and Flavonoids Present in Cannabis. Cancers (Basel). 2020;12(7).

7. Velasco G, Sanchez C, Guzman M. Anticancer mechanisms of cannabinoids. Curr Oncol. 2016;23(2):S23-32.

8. Kiskova T, Mungenast F, Suvakova M, Jager W, Thalhammer T. Future Aspects for Cannabinoids in Breast Cancer Therapy. Int J Mol Sci. 2019;20(7).

9. Good PD, Greer RM, Huggett GE, Hardy JR. An Open-Label Pilot Study Testing the Feasibility of Assessing Total Symptom Burden in Trials of Cannabinoid Medications in Palliative Care. J Palliat Med. 2020;23(5):650-5.

10. Grimaldi C, Pisanti S, Laezza C, Malfitano AM, Santoro A, Vitale M, et al. Anandamide inhibits adhesion and migration of breast cancer cells. Exp Cell Res. 2006;312(4):363-73.

11. Daris B, Tancer Verboten M, Knez Z, Ferk P. Cannabinoids in cancer treatment: Therapeutic potential and legislation. Bosn J Basic Med Sci. 2019;19(1):14-23.

12. Flores R, Shi J, Fuhrman B, Xu X, Veenstra TD, Gail MH, et al. Fecal microbial determinants of fecal and systemic estrogens and estrogen metabolites: a cross-sectional study. J Transl Med. 2012;10:253.

13. El-Talatini MR, Taylor AH, Konje JC. The relationship between plasma levels of the endocannabinoid, anandamide, sex steroids, and gonadotrophins during the menstrual cycle. Fertil Steril. 2010;93(6):1989-96.

14. Urbaniak C, Cummins J, Brackstone M, Macklaim JM, Gloor GB, Baban CK, et al. Microbiota of human breast tissue. Appl Environ Microbiol. 2014;80(10):3007-14.

15. Laborda-Illanes A, Sanchez-Alcoholado L, Dominguez-Recio ME, Jimenez-Rodriguez B, Lavado R, Comino-Mendez I, et al. Breast and Gut Microbiota Action Mechanisms in Breast Cancer Pathogenesis and Treatment. Cancers (Basel). 2020;12(9).

16. Banerjee S, Tian T, Wei Z, Shih N, Feldman MD, Peck KN, et al. Distinct Microbial Signatures Associated With Different Breast Cancer Types. Front Microbiol. 2018;9:951.

17. Zhang J, Xia Y, Sun J. Breast and gut microbiome in health and cancer. Genes Dis. 2021;8(5):581-9.

18. Vallianou N, Kounatidis D, Christodoulatos GS, Panagopoulos F, Karampela I, Dalamaga M. Mycobiome and Cancer: What Is the Evidence? Cancers (Basel). 2021;13(13).

19. Wang N, Sun T, Xu J. Tumor-related Microbiome in the Breast Microenvironment and Breast Cancer. J Cancer. 2021;12(16):4841-8.

20. Wang K, Nakano K, Naderi N, Bajaj-Elliott M, Mosahebi A. Is the skin microbiota a modifiable risk factor for breast disease?: A systematic review. Breast. 2021;59:279-85.

21. Socransky SS, Haffajee AD, Cugini MA, Smith C, Kent RL, Jr. Microbial complexes in subgingival plaque. J Clin Periodontol. 1998;25(2):134-44.

22. Freudenheim JL, Genco RJ, LaMonte MJ, Millen AE, Hovey KM, Mai X, et al. Periodontal Disease and Breast Cancer: Prospective Cohort Study of Postmenopausal Women. Cancer Epidemiol Biomarkers Prev. 2016;25(1):43-50.

23. Chung SD, Tsai MC, Huang CC, Kao LT, Chen CH. A population-based study on the associations between chronic periodontitis and the risk of cancer. Int J Clin Oncol. 2016;21(2):219-23.

24. Pramote Stienrut, Krit Pongpirul, Phanupong Phutrakool, Ornpapha Chanhom, Panthakan Jeamjumrus, Kanyarat Laopiyasakul, et al. Thai cannabis practice patterns and quality of life study (Thai Cannabis PQ): A preliminary analysis. APHA 2020 Annual Meeting and Expo; Oct 24-28; Virtual: American Public Health Association; 2020.

25. Guidelines for providing integrated medical cannabis clinic services [Internet]: Department of Medical Services, Ministry of Public Health; 2021 [cited 2022. Available from: http://www.mhso.dmh.go.th/fileupload/20210129308824218.pdf.

26. Millar SA, Stone NL, Yates AS, O'Sullivan SE. A Systematic Review on the Pharmacokinetics of Cannabidiol in Humans. Front Pharmacol. 2018;9:1365.

27. Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA Cancer J Clin. 2021.

28. International Agency for Research on Cancer: Thailand fact sheets; 2020 [

29. Pattanaphesaj J, Thavorncharoensap M, Ramos-Goni JM, Tongsiri S, Ingsrisawang L, Teerawattananon Y. The EQ-5D-5L Valuation study in Thailand. Expert Rev Pharmacoecon Outcomes Res. 2018;18(5):551-8.

30. Milton L, Behroozian T, Coburn N, Trudeau M, Razvi Y, McKenzie E, et al. Prediction of breast cancer-related outcomes with the Edmonton Symptom Assessment Scale: A literature review. Support Care Cancer. 2021;29(2):595-603.

31. Cannabis for Thailand [Internet]. 2021 [updated 2021; cited 2021. Available from: https://cannabisforthailand.com/.

32. Zhu J, Liao M, Yao Z, Liang W, Li Q, Liu J, et al. Breast cancer in postmenopausal women is associated with an altered gut metagenome. Microbiome. 2018;6(1):136.

33. Kwa M, Plottel CS, Blaser MJ, Adams S. The Intestinal Microbiome and Estrogen Receptor-Positive Female Breast Cancer. J Natl Cancer Inst. 2016;108(8).

34. Alpuim Costa D, Nobre JG, Batista MV, Ribeiro C, Calle C, Cortes A, et al. Human Microbiota and Breast Cancer-Is There Any Relevant Link?-A Literature Review and New Horizons Toward Personalised Medicine. Front Microbiol. 2021;12:584332.

35. Yuan JP, Wang JH, Liu X. Metabolism of dietary soy isoflavones to equol by human intestinal microflora--implications for health. Mol Nutr Food Res. 2007;51(7):765-81.

36. Mirzaei R, Afaghi A, Babakhani S, Sohrabi MR, Hosseini-Fard SR, Babolhavaeji K, et al. Role of microbiota-derived short-chain fatty acids in cancer development and prevention. Biomed Pharmacother. 2021;139:111619.

37. Taormina VM, Unger AL, Schiksnis MR, Torres-Gonzalez M, Kraft J. Branched-Chain Fatty Acids-An Underexplored Class of Dairy-Derived Fatty Acids. Nutrients. 2020;12(9).

38. Parada Venegas D, De la Fuente MK, Landskron G, Gonzalez MJ, Quera R, Dijkstra G, et al. Short Chain Fatty Acids (SCFAs)-Mediated Gut Epithelial and Immune Regulation and Its Relevance for Inflammatory Bowel Diseases. Front Immunol. 2019;10:277.

39. Phelan JP, Reen FJ, Dunphy N, O'Connor R, O'Gara F. Bile acids destabilise HIF-1alpha and promote anti-tumour phenotypes in cancer cell models. BMC Cancer. 2016;16:476.

40. Kovacs P, Csonka T, Kovacs T, Sari Z, Ujlaki G, Sipos A, et al. Lithocholic Acid, a Metabolite of the Microbiome, Increases Oxidative Stress in Breast Cancer. Cancers (Basel). 2019;11(9).

41. Hryhorowicz S, Walczak M, Zakerska-Banaszak O, Słomski R, Skrzypczak-Zielińska M. Pharmacogenetics of Cannabinoids. Eur J Drug Metab Pharmacokinet. 2018;43(1):1-12.