Study Protocol

Efficacy and Mechanisms of Cannabis Oil for Alleviating Side Effects of Breast Cancer Chemotherapy (CBC2): Protocol for Randomized Controlled Trial.

May Soe Thu[1,2], Krit Pongpirul[3,4,5*], Mawin Vongsaisuwon[6], Chanida Vinayanuwattikun[7,8], Kamonwan Banchuen[3], Thunnicha Ondee[3], Sunchai Payungporn[9], Phanupong Phutrakool[3,10], Preecha Nootim[11], Pajaree Chariyavilaskul[12], Sarocha Cherdchom[3], Kulthanit Wanaratna[11], Nattiya Hirankarn[1].

Department of Microbiology, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand.

Department of Infection Biology & Microbiomes, University of Liverpool, Liverpool, UK.

Department of Preventive and Social Medicine, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand.

Bumrungrad International Hospital, Bangkok, Thailand.

Department of International Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA

Department of Surgery, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand.

Division of Medical Oncology, Department of Medicine, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand.

King Chulalongkorn Memorial Hospital, Bangkok, Thailand.

Department of Biochemistry, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand.

Chula Data Management Center, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand.

Department of Thai Traditional and Alternative Medicine, Ministry of Public Health, Nonthaburi, Thailand.

Department of Pharmacology, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand.

*doctorkrit@gmail.com

Abstract

Background

In a pilot study using both cannabidiol (CBD) and tetrahydrocannabinol (THC) as single

agents in advanced cancer patients undergoing palliative care in Thailand, the doses were

generally well tolerated, and the outcome measure of total symptom distress scores showed

overall symptom benefit. The current study aims to determine the intensity of the symptoms

experienced by breast cancer patients receiving chemotherapy, to explore the microbiome

profile, cytokines, and bacterial metabolites before and after the treatment with cannabis oil

or no cannabis oil, and to study the pharmacokinetics parameters and pharmacogenetics

profile of the doses.

Methods

A randomized, double-blinded, placebo-controlled trial will be conducted on the metastatic

breast cancer cases receiving chemotherapy at King Chulalongkorn Memorial Hospital

(KCMH), Bangkok, Thailand. Block randomization will be used to allocate the patients into

three groups: Ganja Oil (THC 2 mg/ml; THC 0.08 mg/drop, and CBD 0.02 mg/drop), Metta

Osot (THC 81 mg/ml; THC 3 mg/drop), and placebo oil. The Edmonton Symptom

Assessment System (ESAS), microbiome profile, cytokines, and bacterial metabolites will be

assessed before and after the interventions.

Thai Clinical Trial Registration: TCTR20220809001

Introduction

Breast cancer patients reported lower quality of life (QoL) globally (1, 2, 3), and chemotherapy was a factor in worsening QoL among the patients (4). Despite standard medical treatment, the impact of integrating complementary and alternative medicine was introduced into the current practice (5). Therefore, cannabinoids from *Cannabis sativa* have become attractive due to their major impact in reducing pain associated with chemotherapy and tumor-associated symptoms (6, 7, 8).

In a pilot study using both cannabidiol (CBD) and tetrahydrocannabinol (THC) as single agents in advanced cancer patients undergoing palliative care, the doses of THC and CBD were generally well tolerated, and the outcome measure of total symptom distress scores (TSDSs) showed overall symptom benefit (9). The average daily amounts tolerated were 300 mg of CBD and 10 mg of THC, respectively (range 5–30 mg), and 43% met the response criteria of ≥6 point reduction in TSDSs (9).

Several clinically relevant mouse models of breast cancer have shown that naturally sourced cannabinoids possess anti-tumor activities, with CBD as the most potent inhibitor of cancer cell proliferation (10). In experimental models, psychoactive THC and non-psychoactive CBD disturbed the disease progression and its effects on signaling pathways within cancer cells following activation of G-protein coupled cannabinoid-receptors (CB-Rs), other receptors such as GPR55 and TRPV1, and in a receptor-independent way. Both *in vitro* and *in vivo* studies have revealed that cannabinoids can activate these receptors, leading to anti-tumorigenic activity in most cases (11).

The human microbiota has recently raised so much attention as a significant risk modulator since they play a prominent role in regulating steroid hormone metabolism through stimulating different enzymes, such as hydroxysteroid dehydrogenase (12, 13). Despite the lack of convincing evidence that breast cancer is caused by dysbiosis, the comparison of several breast samples reveals variations in the abundance and microbial diversity of some particular genera between healthy individuals and patients (14, 15). Additionally, it recognizes that various breast cancer subtypes are correlated to multiple intestinal microbial signatures (16).

It is linked to both the dysbiosis of the intestinal microbiota and the intestinal dysfunction that results in inflammatory and immunological ailments. This implies that diseases spread through microbial translocation. It was proposed that the gut microbiota spreads to the skin before making its way to the breast and that dendritic cells may carry the

bacteria there (17). Therefore, exploring the bacterial translocation markers to understand this
unclear underlying mechanism becomes critical.

Additionally, less fungal diversity and richness has been seen in cancer patients (18),
but it hasn't been thoroughly investigated if particular species and their particular metabolic
outputs characterize and support carcinogenesis. varied hormonal subtypes of breast cancer
exhibit varied fungal signatures in the tumor microenvironment, and individuals with
estrogen receptor-positive breast cancer showed higher levels of fungal variety than those
with triple-negative breast cancer (16, 19).

Because of the primary interactions between skin microorganisms and the underlying
mucosal immune function, the function of the breast skin microbiota is still unknown despite
recent developments. Metastases and breast cancer have been linked to high abundances of
skin commensals, particularly multiple species of *Staphylococcus*, in a thorough investigation
of the skin microbiota in breast cancer (20). Several pathways have been hypothesized for
microbial transfer to underlying tissue, including retrograde transfer through ductal networks,
permeabilization of the epidermal barrier, and migration through the nipple-aspirate fluid
(20).

There are hints that breast cancer and the oral microbiome are related. Women with
periodontal disease, which is brought on by particular bacteria like the red complex
(*Porphyromonas gingivalis, Tannerella forsythia, and Treponema denticola*), and the orange
complex (*Fusobacterium nucleatum, Prevotella intermedia, Prevotella nigrescens,*
Peptostreptococcus micros, Streptococcus constellatus, Eubacterium nodatum,
Campylobacter showae, Campylobacter gracilis, and Campylobacter rectus), have been
found to have an increased risk of breast cancer in some studies (21, 22, 23).

Currently, the Herb and Thai Traditional Medicine Development Division,
Department of Thai Traditional and Alternative Medicine (DTAM), Ministry of Public
Health, Thailand, manufactures Metta Osot (THC 81 mg/ml; THC 3 mg/drop) and Ganja Oil
(THC 2 mg/ml; THC 0.08 mg/drop and CBD 0.02 mg/drop), both of which are produced
under license number 13/2562. The preliminary analysis of the Thai Cannabis Practice
Patterns and QoL (Thai Cannabis PQ) study assessed the pain symptoms using the Edmonton
Symptom Assessment System (ESAS) and QoL using the EuroQoL Group's 5-dimension, 5-
level (EQ-5D-5L). It revealed that cancer patients had significantly lower baseline EQ-5D-5L
(0.80±0.29 vs. 0.85±0.22) and ESAS pain (3.44±3.21 vs. 4.05±3.30) and anxiety (2.03±2.77
vs. 2.37±2.85) but higher tiredness (3.25±3.08 vs. 2.89±2.83), nausea (1.00±2.07 vs.
0.69±1.66), poor appetite (2.21±2.94 vs. 1.20±2.20), and shortness of breath (1.74±2.61 vs.

1.22±2.17) than non-cancer individuals (24). Ganja Oil and Metta Osot were added to the

National List of Essential Medicines (NLEM) (25).

The pharmacokinetic profile of cannabinoids depends on routes of medication such as

oral dosage and inhalation. A study revealed that the median value of the time taken to

achieve the highest levels (T_{max}) of sublingual drops for CBD only and a combination of

CBD and THC was 2.17 hours (range 1 – 4 hours) and 1.67 hours (range 1 – 3 hours) and the

mean of the highest concentration of the drops (C_{max}) was 2.05 (SD = 0.92) and 2.58 (SD =

0.68) nanograms/ml, respectively (26).

Hence, the current randomized study aims to explore the efficacy of Thai cannabis oil

products on the QoL of individuals diagnosed with advanced breast cancer and treated with

chemotherapies and the adverse effects of Thai cannabis oil, along with the study of

microbial alteration and diversity by cannabis among those patients. We aim to determine the

intensity of the pain and QoL experienced by cancer patients using the ESAS and the EQ-5D-

5L forms, respectively, to explore the microbiome profile, cytokines, and bacterial

metabolites in hormone-dependent and independent breast cancer patients at baseline, to

study the alteration of microbiome profile, cytokines, and bacterial metabolites before and

after the treatment with the cannabis oil or no cannabis, and to study the pharmacokinetics

parameters and pharmacogenetics profile.

Materials and methods

Target population

We include 90 adult breast cancer women patients currently receiving chemotherapy at King

Chulalongkorn Memorial Hospital (KCMH) with an anticipated starting recruitment

approximately in June 2023. The sample size was calculated using the mean ESAS values of

the previous study (24).

Inclusion criteria

1. Women aged 18-70 years old who were diagnosed with metastatic breast cancer

2. Metastatic breast cancer patients who are currently receiving different doses of

chemotherapy regimen with high/low emetic risk such as AC (cyclophosphamide and

doxorubicin), AC followed by T (cyclophosphamide, doxorubicin, and paclitaxel), DC

(cyclophosphamide and docetaxel), CMF (cyclophosphamide, fluorouracil, and

methotrexate), and Paclitaxel

3. Breast cancer patients at KCMH during the study period

4. Participants provided written informed consent

5. Individuals having at least each score of ESAS parameters ≥3

Exclusion criteria

1. Women who have contraindications for chemotherapy, such as anaphylaxis

2. Women who have serious chemotherapy complications such as anaphylactic shock, bone

marrow suppression, liver failure, acute kidney injury, and cardiac arrest

3. Women who are scheduled for elective surgery or other procedures requiring general

anesthesia during the study

4. Women who are pregnant or lactating and who are planning for pregnancy during the

study

5. Women who are terminally ill or inappropriate for placebo medication

6. Women with any other significant disease or disorder which, in the opinion of the

investigator, may have put the patient at risk because of participation in the study, or may

have influenced the result of the study, or the patient's ability to participate in the study

7. Women with a known history of substance abuse

8. Women with a history of cannabis ingredients allergy or suspected of having an adverse

reaction to cannabis

9. Women using antibiotics within the past 3 months or using NSAID drugs

10. Women consuming probiotic or prebiotic supplements

Trial design and follow-up

Trial site

The prospective randomized placebo-controlled study will recruit the participants at the King

Chulalongkorn Memorial Hospital (KCMH) in Bangkok, Thailand. The study is scheduled to

launch approximately in June 2023.

Patient data confidentiality

Only the patients providing written informed consent will proceed to participate in the trial.

All the data will be maintained confidentially by the research team.

Patient enrollment and recruitment

Patient demographic information will be reviewed by physicians using KCMH's database.

Online (e.g., on Facebook) and onsite poster (e.g., at KCMH) will be used to advertise the

study. Patients will be recruited using the inclusion and exclusion criteria. The medical record

database will be used to collect clinical and laboratory investigation data related to breast

cancer staging, hormonal breast cancer subtypes, baseline symptoms, current medication

including chemotherapy regime and other herbal use, emetic condition, history of surgery,

history of other cancers, sites of metastasis, history of smoking, history of alcohol use, history

of drug abuse, and history of radiotherapy.

For the recruitment, we will give the trial information to potential participants using

surgeons and nurses in each surgical ward or outpatient clinic and a participant information

sheet to confirm the participant's understanding of the trial objective, procedure, benefits, and

risks. The recruitment will be performed in the Thai language as native. If potential

participants are willing to participate in the trial, they will be asked to sign an informed

consent form with the understanding that they can quit the test at any time without any

consequence.

Patient allocation and blinding

The recruited patients will be randomly allocated into 3 groups: Ganja Oil (THC 2

mg/mL; THC 0.08 mg/drop and CBD 0.02 mg/drop), Metta Osot (THC 81 mg/mL; THC 3

mg/drop), and placebo oil groups. The allocation will be conducted with a 1:1:1 ratio using

block randomization with a block of five by the table of random numbers for creating

randomization ID. The research team will provide the randomization code in an opaque

envelope to conceal allocation, mask clinicians, data collectors, and patients. Following the

cannabis practice patterns of the preliminary study (24), all the participants will be received

cannabis or placebo oil once per day before bedtime for 90 days.

Visits and Follow-ups

There will be four visits for the recruited participants during the trial. At the first visit,

research investigators will perform the enrollment by checking the eligibility screening,

informed consent form, and allocation plan. After setting the random assignment, further

assessment will be conducted using the case report forms approved by the IRB. Then, the

blood, saliva, stool, and skin samples will be collected as a baseline and start the treatment a

day 1 trial. The second visit will be on another day to assess the adverse drug reaction by a

medical officer. The first follow-up treatment, as the third visit, will be on day 30, and the

participants will be checked. Day 90 will be the last visit and second follow-up of the daily

treatment at which all the participants will be assessed, and the blood, saliva, stool, and skin

samples will be collected post-treatment.

After the trial, patients will be asked to stop taking cannabis oil from any sources.

They will be monitored for another month for withdrawal symptoms.

Trial timepoints

Assessment of adverse drug effects will be collected with 4 time points: 0^{th}, 1^{st}, 6^{th}, 12^{nd}

weeks of the trial day. The pain symptoms, QoL and diet pattern will be recorded with 3 time

points: 0^{th}, 6^{th}, 12^{nd} weeks. The blood collection (3 mL) for pharmacokinetic and
pharmacogenetic tests will be performed on 2 time points: 0^{th}, 6^{th} weeks. The blood (10 mL),
feces, skin scrapping, and saliva samples will be taken for biochemistry tests (plasma/
serum), microbiome (gut/ skin/ saliva), cytokines (plasma) and bacterial metabolites (serum/
feces) will be analyzed at 2 time points: 0^{th}, 12^{nd} weeks. The overview of the tests and visits
of the clinical trial was shown on the following Table 1.
Table 1. SPIRIT schedule of enrolment and assessments

#	Timepoint	Week-0 Baseline T1	Week-1 Follow-up T2	Week-6 Follow-up T3	Week-12 Follow-up T4
1	**Enrolment**				
	Eligibility screening	x			
	Informed consent	x			
	Allocation	x			
2	**Assessment**				
	CRF01 Case Record Form[a]	x			
	CRF02 FW & ADR[b]	x	x	x	x
	CRF03 ESAS	x		x	x
	CRF04 EQ-5D-5L	x		x	x
	CRF05 Food Frequency Questionnaire	x		x	x
3	**Blood Samples, Serum**				
	Blood Chemistry[c]	x			x
	Cytokines[d]	x			x
	Bacterial Metabolites[e]	x			x
	Pharmacokinetics parameters	x		x	
	Pharmacogenetics parameters	x		x	
4	**Feces Samples**				
	Fecal Microbiome Profile	x			x
	Fecal Bile Acid Profile[f]	x			x
	Fecal SCFAs[g] and BCFAs[h]	x			x
5	**Skin Samples**				
	Skin Microbiome Profile	x			x
6	**Saliva Samples**				

	Oral Microbiome Profile	x			x

[a.]Demographic and Epidemiological Factors: age, weight, height, waist circumference, hip
circumference, neck circumference, alcohol consumption, smoking status, occupation,
monthly income, method of health payment, and highest education.

[b.]Follow up and adverse drug reaction: vital signs, physical examination, lab results to assess
liver, kidney and heart function, pharmacokinetic record form, drug tolerance assessment
form, and adverse event assessment form.

[c.]Blood Chemistry: cholesterol, alkaline phosphatase, bilirubin, lactate dehydrogenase,
gamma-glutamyl transferase

[d.]Cytokines: interleukin-17, interferon-gamma, Interleukin-10, tumor growth factor-beta

[e.]Bacterial Metabolites: lipopolysaccharide binding protein, sCD14 levels, equol, cadaverine.

[f.]Fecal Bile Acid Profile: chenodeoxycholic acid, deoxycholic acid, lithocholic acid.

[g.]SCFAs: butyrate, propionate, acetate.

[h.]BCFAs: iso-butyric acid.

Primary outcome measures

The pain level of each patient will be assessed using the Edmonton Symptom Assessment
System (ESAS) form at the 4[th] visit as primary outcome. There are 10 symptoms that should
be assessed in the questionnaire and each scale is numbered 0-10, with 0 being asymptomatic
and 10 being the most symptomatic.

Secondary outcome measures

The ESAS at 0[th], 6[th] weeks visits will be taken as secondary outcome measures and will be
evaluated all the symptoms.

For QoL assessment, the Thai version of the EQ-5D-5L questionnaire will be applied to
evaluate at 0[th], 6[th], and 12[nd] weeks. At each dimension, it has the coefficients of the level of
severity utilized for calculating the utility score in accordance with the program of Health
Intervention and Technology Assessment Program (HITAP).

Adverse effects including headache, dizziness, increase nausea and vomiting, dry mouth,
diarrhea, muscle cramps, decreased appetite, sedation, somnolence, itching, or allergic
reactions will be examined using the approved case report form. The patients will be required
to evaluate the severity of the effects using 5-grade system, and the relationship to the study
drugs by 4 statuses.

Regarding feces, skin scrapping, and saliva microbiome, the comparison of the microbial
abundance at baseline and at the end of the trial for different arms will be performed to

explore the microbial changes by the cannabis oil treatment in terms of pain assessment. The

alteration of a variety of cytokines and different bacterial metabolites will be evaluated along

with the microbiome test.

Plasma concentration of THC and CBD will be measured quantitatively to determine the

pharmacokinetic and pharmacogenetic parameters.

Safety

It has been confirmed the effectiveness of the cannabis products in the preliminary analysis

and monitored the side effect at the trials.

Statistical considerations

Sample size estimation

We target 30 patients in each arm, considering the maximum follow-up loss of 50%. Since

the target population is metastatic breast cancer cases, questionnaires and biological samples

such as blood, stool, and breast skin scrapping will be collected before and after the

treatment, which could increase the withdrawal rate at any time during the trial.

Statistical analysis

The data collected from different questionnaires, including case report forms, will be

prepared in Microsoft Excel, followed by data cleaning and extraction. The statistical analysis

will be performed on an intention-to-treat basis using Stata/MP software version 16.0

(StataCorp 2017, College Station, TX). Descriptive statistics like pair t-tests for before and

after-treatment groups will be carried out using mean with standard deviation for normally

distributed continuous data and median with interquartile range for non-normal distributed

continuous data. Categorical data will be presented as counts and percentages using the Chi-

square test. Continuous data will be assessed for normal distribution using a histogram and

Shapiro–Wilk test. One-way ANOVA with Bonferroni correction will be used for normally

distributed continuous data, and the Kruskal-Wallis test will be used for non-normal

distributed continuous data among 3 arms.

To identify the alpha diversity, Shannon's diversity index is used for evenness and

Chao 1 index for the richness of operational taxonomic units (OTUs). The Jaccard

dissimilarity index is used to detect beta diversity, and it is statistically described by

permutational multivariate analysis of variance using distance matrices (PERMANOVA) test.

A statistically significant level is defined as $p < 0.05$. Correction for multiple analyses was

not done, but they will not be included in the conclusion of the study as factual findings.

Sensitivity analysis using some subgroup to show robustness will be considered if the
primary outcome is statistical significance.

Discussion

With a rising incidence of 22,158 new cases occupying a prevalence of 22.8% amongst all
female breast cancer cases in Thailand in 2020 (27, 28), more emphasis becomes on the long-
term quality of life and measurement of patients' symptoms using validated tools (29, 30).
Thailand actively launched cannabis oil in the market in 2018, subsequently, the FDA (Food
and Drug Administration) authorized all hospitals running under the Public Health Ministry
for medical cannabis to be available on prescription to patients for approved conditions like
cancer chemotherapy for relief of pain, to counter inflammation, and so on (31). Following
the preliminary analysis of the Thai Cannabis PQ study, it is expected to improve the
chemotherapy-related symptoms, their quality of life, and the adverse effects.

The CBC2 study is a 1-year prospective study, and it is designed as a randomized,
double-blinded, placebo-controlled trial that would be conducted on current female breast
cancer cases attending a tertiary medical school. This is a continual clinical trial to obtain
additional information; the suggested therapy is a different drug for treating breast cancer,
and it has predictable and controllable side effects.

Since some studies found a particular pattern of bacterial and fungal profiles in
different breast cancer subtypes (16, 19), we expected to identify the similar microbial profile
in hormone-dependent and independent breast cancer patients before and after the treatment
with cannabis oil and placebo oil and evaluate the statistical changes by the treatment.

Dysbiosis is an imbalance of healthy and derogatory bacterial community structures
leading to the development of chronic inflammatory conditions and cancers, particularly in
the gastrointestinal tract (32, 33). Therefore, we will investigate proinflammatory cytokines
such as interleukin-17 (IL-17), interferon-γ (IFN- γ), and inflammatory cytokines like
interleukin-10 (IL-10), and tumor growth factor-β (TGF-β), then will explore the differences
at the baseline and after the treatment.

Moreover, lipopolysaccharide-binding protein (LBP) and soluble CD14 (sCD14)
levels as markers of translocation of bacteria from the gut to the blood will be measured to
explore the possibility of a gut-breast cancer microbiota axis (34) and to evaluate the
influence that cannabis oil treatment may have on the prevention of bloodstream transmission
of oncogenic bacteria from the intestine to the breast. In addition, we will also examine levels
of equol, a potent estrogenic metabolite produced by intestinal bacteria (35).

Short-chain fatty acids (SCFAs), such as acetate, propionate, and butyrate, are present
in the intestine exceeding the concentration of 100 mM, and they are mainly produced by 2
major gut bacterial phyla: *Bacteroidetes* and *Firmicutes* (36). And branched chain iso-
SCFAs, which play a role in membrane permeability and fluidity (37), will be investigated as
bioactive bacterial metabolites in breast cancer. The SCFAs (beneficial) and iso-SCFAs
(potentially harmful) will be indicators for microbial dysbiosis and bowel health (38) in
participants, pre- and post-cannabis oil treatment.

Primary and secondary bile acids will be diagnosed as bioactive bacterial metabolites
in breast cancer. From a range of bile acid profile, deoxycholic acid (DCA) and
chenodeoxycholic acid (CDCA) at the physiological level leads to significant reductions in
cell invasion, migration, adhesion, and survival but not to cytotoxicity or apoptosis (39). In
addition, lithocholic acid (LCA) stimulates oxidative stress in breast cancer patients, which
can reduce cancer cell proliferation (40).

The pharmacokinetic profile of cannabis oil, including THC and CBD, on routes of
medication through oral administration will be examined to achieve the C_{max} of the drugs
(26). As is constantly expanding the cannabis treatment, it becomes essential to predict the
genes whose variations are responsible for the presence of the therapeutic and side effects of
cannabis oil (41).

Conclusions

The clinical trial will provide the efficacy of Thai cannabis oil in metastatic breast cancer
female patients in Thailand, mainly in alleviating chemotherapy-related side effects.
Furthermore, it will obtain secondary factors such as QoL indexes, adverse effects of the
dosage, alteration of microbiome profile, bacterial metabolites, and cytokines before and after
using the cannabis oil, pharmacokinetic parameters, and pharmacogenetic profiles.

Data Availability

All data produced in the present work are contained in the manuscript.

Funding

The Thai Traditional Medical Knowledge Fund, and the Department of Thai Traditional and

Alternative Medicine, Ministry of Public Health, Thailand, financially support this study.

Ethics

The trial was approved by the Institutional Review Board of the Faculty of Medicine,

Chulalongkorn University, Bangkok, Thailand (COA No. 1713/2022; IRB No.0548/65) on

20th December 2022 in compliance with the international guidelines for human research

protection as Declaration of Helsinki, The Belmont Report, CIOMS Guideline and

International Conference on Harmonization in Good Clinical Practice (ICH-GCP).

Acknowledgments

The authors are grateful to the Research Affairs, Faculty of Medicine, Chulalongkorn

University, for their support.

Author Contributions

Conceptualization: Krit Pongpirul, Mawin Vongsaisuwon, May Soe Thu, Nattiya Hirankarn

Data curation: Kamonwan Banchuen, May Soe Thu, Phanupong Phutrakool, Thunnicha

Ondee, Sunchai Payungporn, Pajaree Chariyavilaskul

Formal analysis: Phanupong Phutrakool, Kamonwan Banchuen, Krit Pongpirul, May Soe

Thu

Methodology: Krit Pongpirul, Kamonwan Banchuen, Kulthanit Wanaratna, Pajaree

Chariyavilaskul, Preecha Nootim, May Soe Thu, Sarocha Cherdchom, Sunchai Payungporn,

Thunnicha Ondee

Supervision: Chanida Vinayanuwattikun, Krit Pongpirul, Mawin Vongsaisuwon, Nattiya

Hirankarn, Sunchai Payungporn

Writing – original draft: May Soe Thu, Krit Pongpirul

Writing – reviewing and editing: Krit Pongpirul, May Soe Thu

References

1. Hashemi SM, Balouchi A, Al-Mawali A, Rafiemanesh H, Rezaie-Keikhaie K, Bouya S, et al.
Health-related quality of life of breast cancer patients in the Eastern Mediterranean region: a
systematic review and meta-analysis. Breast Cancer Res Treat. 2019;174(3):585-96.
2. Yanez B, Thompson EH, Stanton AL. Quality of life among Latina breast cancer patients: a
systematic review of the literature. J Cancer Surviv. 2011;5(2):191-207.
3. Paraskevi T. Quality of life outcomes in patients with breast cancer. Oncol Rev. 2012;6(1):e2.
4. Redeker NS, Lev EL, Ruggiero J. Insomnia, fatigue, anxiety, depression, and quality of life of
cancer patients undergoing chemotherapy. Sch Inq Nurs Pract. 2000;14(4):275-90; discussion 91-8.
5. Mokhtari-Hessari P, Montazeri A. Health-related quality of life in breast cancer patients:
review of reviews from 2008 to 2018. Health Qual Life Outcomes. 2020;18(1):338.
6. Tomko AM, Whynot EG, Ellis LD, Dupre DJ. Anti-Cancer Potential of Cannabinoids, Terpenes,
and Flavonoids Present in Cannabis. Cancers (Basel). 2020;12(7).
7. Velasco G, Sanchez C, Guzman M. Anticancer mechanisms of cannabinoids. Curr Oncol.
2016;23(2):S23-32.
8. Kiskova T, Mungenast F, Suvakova M, Jager W, Thalhammer T. Future Aspects for
Cannabinoids in Breast Cancer Therapy. Int J Mol Sci. 2019;20(7).
9. Good PD, Greer RM, Huggett GE, Hardy JR. An Open-Label Pilot Study Testing the Feasibility
of Assessing Total Symptom Burden in Trials of Cannabinoid Medications in Palliative Care. J Palliat
Med. 2020;23(5):650-5.
10. Grimaldi C, Pisanti S, Laezza C, Malfitano AM, Santoro A, Vitale M, et al. Anandamide inhibits
adhesion and migration of breast cancer cells. Exp Cell Res. 2006;312(4):363-73.
11. Daris B, Tancer Verboten M, Knez Z, Ferk P. Cannabinoids in cancer treatment: Therapeutic
potential and legislation. Bosn J Basic Med Sci. 2019;19(1):14-23.
12. Flores R, Shi J, Fuhrman B, Xu X, Veenstra TD, Gail MH, et al. Fecal microbial determinants of
fecal and systemic estrogens and estrogen metabolites: a cross-sectional study. J Transl Med.
2012;10:253.
13. El-Talatini MR, Taylor AH, Konje JC. The relationship between plasma levels of the
endocannabinoid, anandamide, sex steroids, and gonadotrophins during the menstrual cycle. Fertil
Steril. 2010;93(6):1989-96.
14. Urbaniak C, Cummins J, Brackstone M, Macklaim JM, Gloor GB, Baban CK, et al. Microbiota
of human breast tissue. Appl Environ Microbiol. 2014;80(10):3007-14.
15. Laborda-Illanes A, Sanchez-Alcoholado L, Dominguez-Recio ME, Jimenez-Rodriguez B, Lavado
R, Comino-Mendez I, et al. Breast and Gut Microbiota Action Mechanisms in Breast Cancer
Pathogenesis and Treatment. Cancers (Basel). 2020;12(9).
16. Banerjee S, Tian T, Wei Z, Shih N, Feldman MD, Peck KN, et al. Distinct Microbial Signatures
Associated With Different Breast Cancer Types. Front Microbiol. 2018;9:951.
17. Zhang J, Xia Y, Sun J. Breast and gut microbiome in health and cancer. Genes Dis.
2021;8(5):581-9.
18. Vallianou N, Kounatidis D, Christodoulatos GS, Panagopoulos F, Karampela I, Dalamaga M.
Mycobiome and Cancer: What Is the Evidence? Cancers (Basel). 2021;13(13).
19. Wang N, Sun T, Xu J. Tumor-related Microbiome in the Breast Microenvironment and Breast
Cancer. J Cancer. 2021;12(16):4841-8.
20. Wang K, Nakano K, Naderi N, Bajaj-Elliott M, Mosahebi A. Is the skin microbiota a modifiable
risk factor for breast disease?: A systematic review. Breast. 2021;59:279-85.
21. Socransky SS, Haffajee AD, Cugini MA, Smith C, Kent RL, Jr. Microbial complexes in
subgingival plaque. J Clin Periodontol. 1998;25(2):134-44.
22. Freudenheim JL, Genco RJ, LaMonte MJ, Millen AE, Hovey KM, Mai X, et al. Periodontal
Disease and Breast Cancer: Prospective Cohort Study of Postmenopausal Women. Cancer Epidemiol
Biomarkers Prev. 2016;25(1):43-50.

23.   Chung SD, Tsai MC, Huang CC, Kao LT, Chen CH. A population-based study on the
associations between chronic periodontitis and the risk of cancer. Int J Clin Oncol. 2016;21(2):219-
23.
24.   Pramote Stienrut, Krit Pongpirul, Phanupong Phutrakool, Ornpapha Chanhom, Panthakan
Jeamjumrus, Kanyarat Laopiyasakul, et al. Thai cannabis practice patterns and quality of life study
(Thai Cannabis PQ): A preliminary analysis. APHA 2020 Annual Meeting and Expo; Oct 24-28; Virtual:
American Public Health Association; 2020.
25.   Guidelines for providing integrated medical cannabis clinic services [Internet]: Department
of Medical Services, Ministry of Public Health; 2021 [cited 2022. Available from:
http://www.mhso.dmh.go.th/fileupload/20210129308824218.pdf.
26.   Millar SA, Stone NL, Yates AS, O'Sullivan SE. A Systematic Review on the Pharmacokinetics of
Cannabidiol in Humans. Front Pharmacol. 2018;9:1365.
27.   Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global cancer
statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185
countries. CA Cancer J Clin. 2021.
28.   International Agency for Research on Cancer: Thailand fact sheets; 2020 [
29.   Pattanaphesaj J, Thavorncharoensap M, Ramos-Goni JM, Tongsiri S, Ingsrisawang L,
Teerawattananon Y. The EQ-5D-5L Valuation study in Thailand. Expert Rev Pharmacoecon Outcomes
Res. 2018;18(5):551-8.
30.   Milton L, Behroozian T, Coburn N, Trudeau M, Razvi Y, McKenzie E, et al. Prediction of breast
cancer-related outcomes with the Edmonton Symptom Assessment Scale: A literature review.
Support Care Cancer. 2021;29(2):595-603.
31.   Cannabis for Thailand [Internet]. 2021 [updated 2021; cited 2021. Available from:
https://cannabisforthailand.com/.
32.   Zhu J, Liao M, Yao Z, Liang W, Li Q, Liu J, et al. Breast cancer in postmenopausal women is
associated with an altered gut metagenome. Microbiome. 2018;6(1):136.
33.   Kwa M, Plottel CS, Blaser MJ, Adams S. The Intestinal Microbiome and Estrogen Receptor-
Positive Female Breast Cancer. J Natl Cancer Inst. 2016;108(8).
34.   Alpuim Costa D, Nobre JG, Batista MV, Ribeiro C, Calle C, Cortes A, et al. Human Microbiota
and Breast Cancer-Is There Any Relevant Link?-A Literature Review and New Horizons Toward
Personalised Medicine. Front Microbiol. 2021;12:584332.
35.   Yuan JP, Wang JH, Liu X. Metabolism of dietary soy isoflavones to equol by human intestinal
microflora--implications for health. Mol Nutr Food Res. 2007;51(7):765-81.
36.   Mirzaei R, Afaghi A, Babakhani S, Sohrabi MR, Hosseini-Fard SR, Babolhavaeji K, et al. Role of
microbiota-derived short-chain fatty acids in cancer development and prevention. Biomed
Pharmacother. 2021;139:111619.
37.   Taormina VM, Unger AL, Schiksnis MR, Torres-Gonzalez M, Kraft J. Branched-Chain Fatty
Acids-An Underexplored Class of Dairy-Derived Fatty Acids. Nutrients. 2020;12(9).
38.   Parada Venegas D, De la Fuente MK, Landskron G, Gonzalez MJ, Quera R, Dijkstra G, et al.
Short Chain Fatty Acids (SCFAs)-Mediated Gut Epithelial and Immune Regulation and Its Relevance
for Inflammatory Bowel Diseases. Front Immunol. 2019;10:277.
39.   Phelan JP, Reen FJ, Dunphy N, O'Connor R, O'Gara F. Bile acids destabilise HIF-1alpha and
promote anti-tumour phenotypes in cancer cell models. BMC Cancer. 2016;16:476.
40.   Kovacs P, Csonka T, Kovacs T, Sari Z, Ujlaki G, Sipos A, et al. Lithocholic Acid, a Metabolite of
the Microbiome, Increases Oxidative Stress in Breast Cancer. Cancers (Basel). 2019;11(9).
41.   Hryhorowicz S, Walczak M, Zakerska-Banaszak O, Słomski R, Skrzypczak-Zielińska M.
Pharmacogenetics of Cannabinoids. Eur J Drug Metab Pharmacokinet. 2018;43(1):1-12.