The use of newborn foot length to identify low birth weight and preterm babies in Papua New Guinea: A diagnostic accuracy study

Alice Mengi[1] Lisa M Vallely[,2,3], Moses Laman[4], Eunice Jally[1], Janeth Kulimbao[1], Sharon Warel[1], Regina Enman[1], Jimmy Aipit[5], Nicola Low[6]*, Michaela Riddell[1,3].

1. Department of Infection and Immunity, Papua New Guinea Institute of Medical Research, Madang, Papua New Guinea
2. Sexual and Reproductive Health Unit, Papua New Guinea Institute of Medical Research, Madang, Papua New Guinea
3. Global Health Program, The Kirby Institute, University of New South Wales, Sydney, Australia
4. Department of Infection and Immunity, Papua New Guinea Institute of Medical Research, Goroka, Papua New Guinea
5. Madang Provincial Health Authority, Madang, Papua New Guinea
6. Institute of Social and Preventive Medicine, University of Bern, Bern, Switzerland

* Corresponding author

E-mail: nicola.low@ispm.unibe.ch (NL)

Abstract

Low birth weight (LBW, <2.50 kg) and preterm birth (PTB, <37 completed weeks of gestation) are important contributors to neonatal death. Newborn foot length has been assessed to identify LBW and PTB babies in Asia and Africa. The objectives of this observational study were to determine the diagnostic accuracy of foot length to identify LBW and PTB and to compare foot length measurements of a researcher with those of trained volunteers in Papua New Guinea. Newborn babies were enrolled prospectively with written informed consent from their mothers, who were participating in a clinical trial in Madang Province. The reference standards were birth weight, measured by electronic scales and gestational age at birth, based on ultrasound scan and last menstrual period at the first antenatal visit. Newborn foot length was measured within 72 hours of birth with a firm plastic ruler. Optimal foot length cut-off values for LBW and PTB were derived from receiver operating characteristic curve analysis. Bland-Altman analysis was used to assess inter-observer agreement between researcher and volunteer. From 12 October 2019 to 6 January 2021, we enrolled 342 newborns (80% of those eligible); 21.1% (72/342) were LBW and 7.3% (25/342) were PTB. The area under the curve for LBW was 87.0% (95% confidence intervals 82.8-90.2) and for PTB 85.6% (81.5-89.2). The optimal foot length cut-off was <7.7 cm for both LBW (sensitivity 84.7%, 74.7-91.2, specificity 69.6%, 63.9-74.8) and PTB (sensitivity 88.0% (70.0-95.8), specificity 61.8% (56.4-67.0). In 123 babies with paired measurements, the mean difference between the researcher and volunteer measurements was 0.07 cm (95% limits of agreement -0.55 to +0.70) and 7.3% (9/123) of the pairs were outside the 95% limits of agreement. Foot length measurement can identify LBW and PTB in newborns but needs appropriate training for community volunteers and evaluation of its impact on healthcare outcomes.

Introduction

Globally, complications of preterm birth (PTB), including low birth weight (LBW), are the leading cause of neonatal death in the first week of life [1]. Identifying the most vulnerable neonates so that they can receive timely essential care is challenging in resource-limited settings, particularly in rural areas or when births occur in the community [2]. Babies born LBW (<2.50 kg) [3] are either preterm (<37 completed weeks gestation) or small for gestational age (birth weight below the 10th percentile of the distribution for gestational age and sex) [3,4]. Many babies are, however, not weighed at birth [5] or do not have an accurate assessment of their gestational age at birth. These assessments require reliable equipment and skilled, trained health workers to conduct ultrasound scans during pregnancy, measure birth weight accurately, or apply clinical assessments, such as Dubowitz or New Ballard scores [6].

Foot length is one of a range of anthropometric measures that have been used as proxies for LBW and PTB in resource-limited settings [6-18]. In a systematic review and meta-analysis of studies published between 2007 and 2020, Folger et al. found 19 studies, all conducted in south and south-east Asia and sub-Saharan Africa [19]. The authors found that a newborn foot length of <7.7cm had reasonable accuracy for identification of LBW babies in 3 studies in Asia (pooled sensitivity 87.6%, 95% confidence interval, CI 55.7-97.5%, specificity 70.9%, 23.5-95.1%). They noted, however, that few studies used an accurate reference method for assessing gestational age and did not conduct a meta-analysis of foot length and PTB because of heterogeneity in study methods [19]. The optimal cut-off for newborn foot length may differ by country and region and performance characteristics could also vary, according to whether measurements are taken by clinically trained staff or community volunteers, field workers and care givers [11,20].

Papua New Guinea (PNG) is an island state in the South Pacific, which has one of the highest neonatal mortality rates in the world, with 22 newborn deaths per 1000 live births in 2020 [21,22].

Uptake of antenatal care in PNG is poor, with only 76% of women attending at least once and only

36-56% of women give birth in a health facility [21,23]. The objectives of this study were to 1)

determine the diagnostic accuracy of newborn foot length measurement to identify LBW or PTB

babies in health facilities and community settings and 2) to compare foot length measurements

taken by trained volunteers with those of a researcher to assess the feasibility of newborn foot

length measurement in a community setting.

Materials and methods

The Neofoot (neonatal foot length) study was a prospective single arm diagnostic test

accuracy study (S1 Protocol). We report the study using the Standards of Reporting of Diagnostic

accuracy 2015 (S1 Checklist). The study was conducted in three primary care clinics in Madang

Province, PNG, which serve a total population of approximately 35,000, according to the 2011 PNG

National Census [24]. The study population included both people living in urban shanty settlements

and rural subsistence farmers in areas with poor road conditions and difficult access to health

facilities. The Neofoot study was designed as a sub-study of the Women And Newborn Trial of

Antenatal Interventions and Management (WANTAIM) trial, a randomised controlled trial which

assessed the effect of point-of-care testing and treatment of curable sexually transmitted infections

and bacterial vaginosis during pregnancy on birth outcomes [25]. Women enrolled in the WANTAIM

study were aged 16 years or older at a gestational age below 26 completed weeks, based on the

date of the first day of the last menstrual period or an ultrasound scan performed by trained

midwives at the first antenatal visit [26]. Their newborn babies were eligible for inclusion if the

clinical researcher (AM) or senior study staff (EJ, JK, SW, RE) assessed them within 72 hours of birth,

their general condition was stable and the baby did not have abnormalities. There were no

additional exclusion criteria. At the first postnatal visit, the researcher or staff member used pictorial

diagrams to explain the Neofoot study in Tok-Pisin to consecutive women whom they attended,

assessed their understanding of the study and answered questions. The mother or guardian gave a

signature or thumbprint as informed consent for their baby to take part (S1 Protocol, page 20-21).

For women who were unable to read or write, an independent witness was present during the

consent procedures and also signed the consent form.

Reference standards

Birth weight was the average of two measurements taken at a 5 to 15 minute interval by

trained WANTAIM clinical staff within 72 hours of birth using an electronic infant scale (Charder

Cupid 1, Charder Medical, Taiwan) placed firmly on a hard surface and calibrated to zero [27]. Birth

weight was before the index test on a separate clinical record form. LBW was defined as below 2.50

kg [3]. Gestational age at birth was measured in completed weeks of pregnancy, calculated from the

best obstetric estimate of the due date [26], using the date of the first day of the last menstrual

period or an ultrasound scan (Logiq V2 portable ultrasound, GE Healthcare) performed by trained

midwives at the first antenatal visit. Gestational age at birth was calculated at the time of statistical

analysis. PTB was defined as birth before 37 completed weeks of pregnancy [4].

Index test

The index test was the length of the right foot, measured within 72 hours of birth. We

calculated the average of two measurements, taken 5 to 15 minutes apart by the clinical researcher.

Foot length was measured using a firm clear plastic 15cm ruler, bought from a local store and stuck

to an A4-sized cardboard box, which was cut open (Fig 1). The tip of the ruler was pushed into a slit

at the bottom of the box so that the zero level was at the base of the box. The researcher placed the

right heel at the zero level and held the plantar surface of the foot against the ruler. Foot length at

the tip of the hallux was measured to the nearest 1 mm, with a plastic setsquare perpendicular to

the ruler.

Fig 1. **Measurement of newborn foot length.** Measurements were made with a 15cm clear plastic

ruler stuck to the inside of a cardboard box. The zero mark of the ruler is at the base of the box. Foot

length is measured from the base of the heel to the tip of the hallux.

Training of volunteers

116 To assess the feasibility of community members or village health workers identifying at risk

117 babies, we enrolled nine volunteers, who were involved in the WANTAIM trial and were numerate

118 and literate. The clinical researcher gave a half-day training session at the postnatal ward of Madang

119 Provincial Hospital. The volunteers learned how to measure the baby's foot length, how to complete

120 the case record form and practised both tasks.

Measures to reduce the risk of bias

To reduce the risk of measurement bias in foot length, the researcher and the volunteer

recorded two measurements at a 5-to-15-minute interval. To ensure that the measurements were

independent, the researcher and volunteer recorded their measurements on separate pages of the

case record form and could not see the other person's record. Birth weight was measured before the

index test on a separate clinical record form. The clinician researcher had access to clinical

information about the newborn, but the best obstetric estimate of gestational age at birth [26] was

calculated at the time of statistical analysis.

Ethics approval

The Neofoot study protocol was approved by the PNG Institute of Medical Research

Institutional Review Board (IRB 1811) and the PNG Medical Research Advisory Committee (MRAC

18.18). The WANTAIM study (ISRCTN37134032) received ethical approval from the Institutional

Review Board of the PNG Institute of Medical Research (IRB number 1608); the Medical Research

Advisory Committee of the PNG National Department of Health (MRAC number 16.24); the Human

Research Ethics Committee of the University of New South Wales (HREC number 16708); and the

Research Ethics Committee of the London School of Hygiene and Tropical Medicine (REC number

12009) [25].

Data management

Data were recorded on printed case record forms designed using TELEform™ Elite version

10.5 (https://teleform.software.informer.com/10.5/). The case record forms were checked for

discrepancies or out of range values and corrected before scanning, after which they were machine-

read into a database (Microsoft). Variables collected from the mother at the first antenatal visit, as

part of the WANTAIM study, included demographic details (province of birth, age, and marital

status), gestational age at enrolment, estimated due date, haemoglobin, HIV test result

(Determine™ HIV 1/2 Test), syphilis (SD Bioline, Abbott Diagnostics) results and use of betel nut,

cigarettes, and alcohol during pregnancy. Data collected at the postnatal visit included date and

place of birth, reason for giving birth at that location, newborn birth weight and sex of the baby.

Data about foot length were the two measurements made by the researcher and the volunteers.

Statistical analysis

All statistical analyses used STATA (Stata 14.2 or 16, College Station, Texas, United States).

For each newborn we calculated the gestational age at birth using the estimated due date

established at the enrolment visit, and the average of the two measurements of birth weight and of

foot length. The distributions of continuous variables were examined using histograms and

described using the mean and standard deviation (SD) or median and interquartile range (IQR). For

the classification of LBW and PTB, birth weight and gestational age at birth were dichotomised in

accordance with the reference standard definitions.

We constructed receiver operating characteristics curves to display sensitivity against 1 minus

specificity for the classification of LBW and PTB at each 1 mm increment in foot length from the

lowest to the highest value. We calculated the overall area under the curve and test performance

characteristics sensitivity, specificity, positive and negative predictive values (as percentages with

95% CI). We selected the optimal foot length cut-off, separately for LBW and PTB, as the value with

the best balance of sensitivity and specificity, using the foot length cut-off with the highest area

under the curve. For each baby for whom paired foot length measurements were available from the

researcher and the volunteer, we assessed the level of agreement using Bland-Altman analysis [28].

Sample size calculation

166   The planned sample size was based on achieving an acceptable level of precision (95% CI +/-

167   4 to 5%) for values of sensitivity from 70 to 90%. Using projected WANTAIM trial enrolment at the

168   study clinics, we aimed to invite around 400 women to join the Neofoot study. Assuming 95% of

169   babies had a postnatal visit within the first 72 hours and that 90% of mothers agreed to foot length

170   measurement, the target was to have foot length measurements for 342 newborn babies.

Results

A total of 426 babies were born to 416 mothers enrolled in the WANTAIM trial from 12

October 2019 to 6 January 2021. Of these, 84 babies were excluded from the study for reasons

outlined in Fig 2. A total of 342 babies born (80.2% of those potentially eligible) to 335 mothers were

enrolled. The study population included 10 of 14 twins born to 7 mothers; 3 mothers had both twins

enrolled and four mothers had one twin enrolled (the other twin died before birth or before

neonatal assessment). The researcher recorded foot length measurements for all 342 babies. Owing

to disruptions during the COVID-19 pandemic and staff availability, a volunteer was available to

measure foot length at the time of the postnatal assessment for 123 (36.0%) babies.

Fig 2. Flow chart indicating number of mothers and newborn babies at each stage of the Neofoot

study.

Among the 335 mothers, their mean age was 25.5 (SD 5.9, range 16-44) years, 43.9% (147/335) were

primiparous and 89.3% (299/335) were married (Table 1). Sixty-seven women attended one of the

primary care clinics covering urban shanty settlement areas and 268 attended clinics in rural areas.

Most women reported that they currently chewed betel nut (281/335, 83.9%) and about a quarter

were current smokers (81/335, 24.2%), but few (6/335, 1.8%) reported that they consumed alcohol

during pregnancy. Forty-two women 12.5% (42/335) had a positive test result for syphilis and 2

women (0.6%) had a positive test result for HIV. The mean haemoglobin concentration was 8.6 g/dL

(SD 1.6), three quarters of women (76.7 %; 257/335) had mild anaemia (haemoglobin 7.0–9.9 g/dL)

and 13.7 % (46/335) had moderate to severe anaemia (haemoglobin <7.0 g/dL).

Table 1. Characteristics of mothers whose newborn babies were enrolled in the study (N=335)

Maternal characteristic		Total N = 335
Birthplace, region n (%)	Momase	295 (88.1)
	New Guinea Island	19 (5.7)
	Highlands	16 (4.8)
	Southern Region	5 (1.5)
Age, years	Mean (SD)	25.5 (5.9)
Marital status, n (%)	Married	299 (89.3)
	Single	29 (8.7)
	Divorced/separated/widowed	3 (0.9)
	Missing	4 (1.2)
Chew betelnut, n (%)	Currently chew	281 (83.9)
	Previously chew	5 (1.5)
	Never chew	49 (14.6)
Drink alcohol, n (%)	Currently drink	6 (1.8)
	Previously drink	31 (9.3)
	Never drink	298 (89.0)
Smoke cigarettes, n (%)	Currently smoke	81 (24.2)
	Previously Smoke	34 (10.2)
	Never smoke	220 (65.7)

Parity, n (%)	0	147 (43.9)
	1-3	141 (42.1)
	≥ 4	47 (14.0)
Current pregnancy, n (%)	Singleton	325 (97.0)
	Twin	10 (3.0)
Gestation in weeks, median (IQR)	USS	22 (19.0-24.0)
	Fundal height	22 (19.0-24.0)
	LMP	20 (17.0-23.0)
HIV test result, n (%)	Positive	2 (0.6)
	Negative	323 (96.4)
	Not tested	10 (3.0)
Syphilis test result, n (%)	Positive	42 (12.5)
	Negative	293 (87.5)
Haemoglobin, g/dL	Mean (SD)	8.6 (1.6)
Haemoglobin, category	≥ 10 g/dL	16 (4.8)
	7.0-9.9 g/dL	257 (76.7)
	< 7.0 g/dL	46 (13.7)
	Missing	16 (4.8)

Abbreviations: cm, centimetres; g/dL, grams per decilitre; HIV, human immunodeficiency virus; IQR,

interquartile range; LMP, last menstrual period; N, total number of participants; n, number of

participants in a category; SD, standard deviation; USS, ultrasound scan.

Table 2 describes the characteristics of the 342 babies. Most babies (330/342, 96.5%) were born by

normal vaginal birth at a health facility (72.8%; 249/342), with 27.2% (93/342) births taking place at

home or before arrival at a health facility. Birth weight was normally distributed (mean 2.8 kg, SD

0.4), with a high frequency of birth weights recorded as 2.50 kg (known as digit preference or

heaping, S1 Fig).

Table 2. Characteristics of newborn babies in the study (N=342)

Newborn characteristic		**Total** N = 342	
Place of birth, n (%)	Health facility	249	(72.8)
	Born before arrival at health facility	5	(1.5)
	Home/village	88	(25.7)
Type of birth, n (%)	Normal vaginal birth	330	(96.5)
	Caesarean section	3	(0.9)

	Assisted birth	9	(2.6)
Sex, n (%)	Male	177	(51.8)
	Female	165	(48.3)
Birth weight, kg	Mean (SD)	2.8	(0.40)
	Median (IQR)	2.8	(2.5-3.1)
	Range, lowest-highest		1.3-4.2
Birth weight category n (%)	LBW (< 2.50 kg)	72	(21.0)
	Normal birth weight (≥ 2.50 kg)	270	(79.0)
Gestational age at birth, weeks	Mean (SD)	38.8	(1.8)
	Median (IQR)	39	(38-40)
	Range, lowest-highest		30-42
Gestational age category n (%)	Preterm < 37 weeks	25	(7.3)
	Term ≥ 37 weeks	317	(91.7)
Foot length, cm	Mean (SD)	7.8	(0.5)
	Median (IQR)	7.8	(7.5-8.0)
	Range, lowest-highest		5.8-8.9

Abbreviations: IQR, interquartile range; LBW, low birth weight; N, total number of

participants; n, number of participants in a category; SD, standard deviation.

Fig 3 shows scatterplots of birth weight, gestational age at birth and the foot length measured by the

researcher. About one-fifth of all babies (21.1%, 72/342), were LBW (<2.50 kg). The mean

gestational at birth was 38.8 weeks (SD 1.8) and 7.3% (25/342) were preterm. Of the babies born

preterm (<37 completed weeks of gestation), 5/25 babies were of normal weight (≥2.50 kg) and

amongst 317 babies born at term, 52 babies had a birth weight <2.50 kg (Fig 3). Mean foot length

was 7.8 cm (SD 0.5) with no adverse events recorded as a result of measurement.

Fig.3. Scatter plots of newborn measurements recorded within 72 hours of birth for 342

babies. Panel A. Birth weight and foot length; B. gestational age at birth and foot length; C.

gestational age at birth and birth weight. r, Pearson correlation coefficient. Reference lines in

red indicate the cut-offs for low birth weight (2.5 kg) and preterm birth (37 completed weeks

of gestation).

Optimal foot length cut-offs for classifying LBW and PTB

The receiver operating curves summarise the overall accuracy of foot length measurement for the classification of newborn babies as LBW or PTB (Fig 4) at cut-off increments of 1 mm. The overall area under the curve for foot length was 87.0% (95% CI, 82.3-91.6) for LBW and 85.8% (77.1-94.6) for PTB.

Fig. 4. Receiver operating characteristics curves of the accuracy of foot length for the classification of outcomes in 342 newborn babies. Panel A, low birth weight, B preterm birth.

Optimal foot length cut-offs for the classification of LBW and PTB were determined as the value with the combination of the highest sensitivity and specificity (Table 3, S1 Table, S2 Table). The optimal foot length cut-off for identifying LBW was <7.7 cm. At this foot length, 61/72 babies were correctly classified as LBW (sensitivity 84.7%, 74.7-91.2). The birth weights of 11 babies with foot length ≥7.7 cm, who were incorrectly classified as being of normal birth weight (false negatives), ranged from 2.15-2.43 kg. Among all babies with foot length <7.7 cm, 42.7 (34.8-50.9) were LBW (positive predictive value). The negative predictive value was 94.5% (90.4-96.9).

Table 3. Diagnostic accuracy statistics at the optimal foot length cut-off for classification of babies as low birth weight or preterm

Outcome	<7.7 cm	≥ 7.7 cm	Sensitivity, % (95% CI)	Specificity, % (95% CI)	Positive predictive value, % (95% CI)	Negative predictive value, % (95% CI)
Birth weight						
< 2.50 kg	61	82	84.7	69.6	42.7	94.5
≥ 2.50 kg	11	188	(74.7-91.2)	(63.9-74.8)	(34.8-50.9)	(72.4-81.5)
Gestational age at birth						
< 37 weeks	22	121	88.0	61.8	15.4	98.5
≥ 37 weeks	3	196	(70.0 – 95.8)	(56.4 – 67.0)	(10.4 – 22.2)	(95.7 – 99.5)

The optimal foot length cut-off for PTB was <7.7 cm (Table 3), with 22/25 babies correctly classified

(sensitivity 88.0%, 70.0-95.8). Of the 3 preterm babies not identified as PTB at this cut-off (false

negatives), 2 were born at 36 weeks and 1 at 35 weeks gestation; none was very (28-32 weeks) or

extremely (<28 weeks) preterm. At this cut-off, 121 babies were false positives, being born at term

but with foot length <7.7 cm. The positive predictive value of a foot length of <7.7 cm was 15.4%

(10.4-22.2) and negative predictive value 98.5% (95.7-99.5).

Foot length measurements from both the researcher and a volunteer were available for 123 babies.

Values at full or half centimetre marks were over-represented for both sets of average

measurements (S2 Fig). Fig 5 shows the Bland-Altman plot, displaying the differences between the

123 paired measurements against the average of the measurements of the researcher and

volunteer. The mean difference between researcher and volunteer measurements (dashed line) was

0.07 cm. The 95% limits of agreement ranged from -0.55 cm to +0.70 cm. Differences between

measurements for 9/123 (7.3%) of the data pairs were outside the 95% limits of agreement.

Fig 5. Bland-Altman plot for researcher and volunteers' newborn foot length measurements

(N=123). Size of circle is proportional to number of measurements.

Discussion

Summary of main findings

This prospective study enrolled 342 newborn babies in Madang Province, PNG. Overall, 21.1%

(72/342) of newborns were LBW and 7.3% (25/342) were born preterm. The optimal foot length cut-

off to classify LBW babies was <7.7 cm (sensitivity 84.7%, 95% CI 74.7-91.2, specificity 69.6%, 63.9-

74.8) and <7.7 cm to classify PTB (sensitivity 88.0%, 70.0-95.8, specificity 61.8%, 56.4–67.0%). The

95% limits of agreement between researcher and volunteer measurements were from -0.55 to +

0.70 cm and 7.3% (9/123) of the pairs were outside these limits.

Strengths and limitations

Strengths of this study included the use of accurate methods for determination of the reference

standards for measuring birth weight and gestational age at birth. All babies were weighed twice

using electronic scales [27] and gestational age at birth was calculated using published

recommendations for the best obstetric estimate of the delivery date, which are based on the date

of the last menstrual period and an ultrasound scan at the first antenatal visit [26]. Foot length was

measured using items readily available in the community and the study involved trained volunteers

from communities, with babies assessed at local health centres or at home. Also, the target sample

size was reached, and outcome variables LBW and PTB were available for all babies in the study,

even though a COVID-19 lock down was in place for part of the study.

There were also limitations of the study methods. First, the measurement of birth weight was taken

within the first 72 hours of birth. Ideally, birth weight should be measured within a few hours of

birth, before any postnatal weight loss has occurred [27]. However, Marchant et al. found that foot

length increased by only 0.2 cm (SD 0.3) between the first and fifth day after birth in a study in

Tanzania [10], so the delay of up to 72 hours in our study should not have affected the results.

Second, the researcher and volunteers could not be blinded from the clinical status of the newborn,

which might have influenced their foot length measurements. Knowledge of birthweight could have

resulted in overestimation of the accuracy of classification of LBW. Since the overall areas under the

curve for LBW and PTB were similar, we think risk of bias was low. Third, whilst mothers were

enrolled at primary care clinics and most gave birth at a health centre or at home/village, the study

was done in only one coastal province of PNG. The 2020 PNG National Department of Health Sector

Review Performance found that Madang Province was the province with the highest percentage of

low birth weight babies (10 in 2019 and 13% in 2022) [21]. Optimal foot length cut-off values might

be different in other regions of PNG or other Pacific islands.

Comparison with other studies

Our study from PNG in the Pacific region adds to the evidence from previously published studies that

have explored foot length as a proxy to identify LBW and PTB and were conducted in countries in

Asia and Africa. The levels of LBW (around 20%) and PTB (around 7%) in the Neofoot study were

similar to those of several other published studies that have assessed the diagnostic accuracy of foot

length both in sub-Saharan Africa [10,15] and south Asia [29]. Our study in PNG, however, overcame

limitations of assessment of gestational age and of applicability of findings from tertiary care setting

that were identified in previous studies [19]. The Neofoot study shared several characteristics with a

study by Lee et al. among 710 babies in the community in Bangladesh, but with different findings

about the accuracy of foot length as a method to identify PTB [29]. In both studies, the investigators

assessed babies within 72 hours after birth, estimated gestational age based on ultrasound scan and

used a clear plastic ruler to measure foot length. Lee et al. found that foot length measured at home

visits by trained community health workers did not classify PTB well, with an overall area under the

curve of 51.9% and sensitivity at a cut-off of <7.7 cm of 28%. In contrast, we found an overall area

under the curve of 85.6% and sensitivity of 88.0% for the same cut-off. Lee et al. attributed the poor

performance of foot length (and other anthropometric measurements) to the high level of foetal

growth restriction (32.4% of babies were assessed as being small for gestational age). In our study,

however, levels of PTB and mean birth weight were similar to those of Lee et al. and 12.3% (42/342)

of term babies were LBW. One factor that might have contributed to the different findings was that

all babies in Lee et al.'s study were assessed at home by community health workers, whilst more

babies in our study were assessed at a health facility or community by a paediatrician researcher,

where the conditions for measuring foot length might have been easier.

Interpretation of the findings

In the Neofoot study, a foot length cut-off of <7.7 cm had similarly high overall accuracy for classification of LBW and PTB (Table 3). A common cut-off for identifying LBW and PTB was also found by Paulsen et al. (≤7.7 cm) [30] and Marchant et al. (<8 cm) in Tanzania [10]. The cut-off of <7.7 cm for LBW concurs with the pooled result from the meta-analysis of 15 studies by Folger et al., which included studies from a range of countries [19]. The lower prevalence of PTB, compared with LBW, results in a lower positive predictive value. At the optimal foot-length cut-off, around 15% of babies were preterm, according to the reference standard (compared with nearly half of babies classified as LBW). The referral of substantial numbers of babies for urgent care would increase the workload for health systems and should be assessed. The use of a plastic ruler was an appropriate technology for the measurement of foot length and taking the average of two measurements should have reduced error, although we still observed digit preference for measurements at the half or full centimetre (S2 Fig). A single training session for lay people and health facility staff to measure foot length resulted in average foot length measurements that were similar to those of the researcher, but paired measurements for individual babies differed by up to 1 cm. These findings were comparable to those of Marchant et al. who assessed 142 paired measurements, using a plastic ruler, between researchers and community volunteers in Tanzania [11]. As in our study, they found no overall bias but volunteers' measurements were shorter than those of researchers.

Implications for research and practice

The goal of universal health coverage to ensure that all sick and small newborns survive and thrive will take time to achieve [2].The findings of this study have implications for future research and practice in settings such as PNG, where it is difficult to identify high risk babies in populations with rural communities, high levels of home births and limited access to health facilities with skilled staff and equipment. Most women in PNG attend antenatal clinic late in pregnancy, often in the last trimester, so assessment of gestational age at birth is particularly challenging. Our study adds to the

published evidence [19], showing that both LBW and PTB can be classified accurately and could be

used by trained health workers and community health volunteers. Newborn foot length has not,

however, been widely implemented. Implementation research studies should develop appropriate

technologies to apply optimal cut-off measurements, training programmes that reduce

misclassification from digit preference (heaping) and examine the impacts on referrals to health

systems and health outcomes. In this study in PNG, newborn foot length was a simple and feasible

method for the identification of LBW and PTB babies and could help to improve neonatal health

outcomes.

Acknowledgments

This study was conducted during the Master of Medical Sciences, University of Papua New

Guinea of Alice Mengi. We sincerely thank all the parents of the 342 babies that participated, the

PNG Institute of Medical Research and WANTAIM staff, especially Kelly Masil for scanning and

verification of data and Andrew Vallely and William Pomat for acquisition of funding and oversight of

the WANTAIM trial. We thank Odile Stalder at the Clinical Trials Unit, University of Bern, who helped

with the statistical analyses. We thank Sister Hilda Marijembi, Sister Susie Yuangi, Naida Samson,

Sister Bridgette Nawali, Lina Puli, Josephine Fei, Christophilda Noubiri and Alvita Jimmy for

assistance with data collection.

References

1.   Lawn J, Blencowe H, Oza S, You D, Lee AC, Waiswa P, et al. Every Newborn: Progress,

Priorities, and Potential Beyond Survival. *Lancet*. 2014;384:189-205.

2.   UNICEF WHO. Survive and Thrive: Transforming Care for Every Small and Sick Newborn:

Geneva: World Health Organization; 2019 [Access Date; cited 2022 6/11/2022]. Available from:

https://www.who.int/publications/i/item/9789241515887.

3.   World Health Organization. Newborns with Low Birth Weight (%) 2006 [Access Date; cited

2021 15/5/2021]. Available from:

https://www.who.int/whosis/whostat2006NewbornsLowBirthWeight.pdf.

4.   World Health Organization. Preterm Birth 2018 [Access Date; cited 2021 15/5/2021].

Available from: https://www.who.int/news-room/fact-sheets/detail/preterm-birth.

5.   Blanc AK, Wardlaw T. Monitoring Low Birth Weight: An Evaluation of International Estimates

and an Updated Estimation Procedure. *Bulletin of the World Health Organization*. 2005;83(3):178-

85. PubMed Central PMCID: 15798841.

6.   Lee A, Panchal P, Folger LV, Whelan H, Whelan R, Rosner B, et al. Diagnostic Accuracy of

Neonatal Assessment for Gestational Age Determination: A Systematic Review. *Pediatrics*. 2017;140.

7.   Ashish KC, Nelin V, Vitrakoti R, Aryal S, Målqvist M. Validation of the Foot Length Measure as

an Alternative Tool to Identify Low Birth Weight and Preterm Babies in a Low-Resource Setting Like

Nepal: A Cross-Sectional Study. *BMC Pediatr*. 2015;15:43. Epub 2015/05/01. doi:

https://doi.org/10.1186/s12887-015-0361-4. PubMed PMID: 25928804; PubMed Central PMCID:

PMC4403672.

8.   Feresu SA, Harlow SD, Gillespie BW, Welch K, Johnson TR. Birthweight-Adjusted Dubowitz

Methods: Reducing Misclassification of Assessments of Gestational Age in a Zimbabwean

Population. *The Central African journal of medicine*. 2003;49(5-6):47-53. PubMed PMID: 15214282.

9.   Hadush MY, Berhe A, Medhanyie A. Foot Length, Chest and Head Circumference

Measurements in Detection of Low Birth Weight Neonates in Mekelle, Ethiopia: A Hospital Based

Cross Sectional Study. *BMC Pediatrics*. 2017;17.

10.   Marchant T, Jaribu J, Penfold S, Tanner M, Schellenberg JA. Measuring Newborn Foot Length

to Identify Small Babies in Need of Extra Care: A Cross Sectional Hospital Based Study with

Community Follow-up in Tanzania. *BMC Public Health*. 2010;10(1):624. doi: 10.1186/1471-2458-10-

624.

11.   Marchant T, Penfold S, Mkumbo E, Shamba D, Jaribu J, Manzi F, et al. The Reliability of a

Newborn Foot Length Measurement Tool Used by Community Volunteers to Identify Low Birth

Weight or Premature Babies Born at Home in Southern Tanzania. *BMC Public Health*.

2014;14(1):859. doi: 10.1186/1471-2458-14-859.

12.   Mukherjee S, Roy P, Mitra S, Samanta M, Chatterjee S. Measuring New Born Foot Length to

Identify Small Babies in Need of Extra Care: A Cross-Sectional Hospital Based Study. *Iranian journal of*

pediatrics. 2013;23(5):508-12. PubMed PMID: 24800008.

13.   Nabiwemba E, Marchant T, Namazzi G, Kadobera D, Waiswa P. Identifying High-Risk Babies

Born in the Community Using Foot Length Measurement at Birth in Uganda. *Child Care Health Dev*.

2013;39(1):20-6. doi: 10.1111/j.1365-2214.2012.01383.x. PubMed PMID: 22680250.

14.   Nabiwemba E, Orach C, Kolsteren P. Determining an Anthropometric Surrogate Measure for

Identifying Low Birth Weight Babies in Uganda: A Hospital-Based Cross Sectional Study. *BMC Pediatr*.

2013;13:54. Epub 2013/04/17. doi: 10.1186/1471-2431-13-54. PubMed PMID: 23587297; PubMed

Central PMCID: PMC3639798.

15.   Otupiri E, Wobil P, Nguah SB, Hindin MJ. Anthropometric Measurements: Options for

Identifying Low Birth Weight Newborns in Kumasi, Ghana. *PloS one*. 2014;9(9):e106712-e. doi:

10.1371/journal.pone.0106712. PubMed PMID: 25226505.

16.   Pratinidhi AK, Bagade AC, Kakade SV, Kale HP, Kshirsagar VY, Babar R, et al. Action-Oriented

Colour-Coded Foot Length Calliper for Primary Healthcare Workers as a Proxy for Birth Weight &

Gestational Period. *The Indian journal of medical research*. 2017;145(3):347-52. doi:

10.4103/ijmr.IJMR_36_14. PubMed PMID: 28749397.

17.   Singhal R, Jain S, Chawla D, Guglani V. Accuracy of New Ballard Score in Small-for-Gestational

Age Neonates. *J Trop Pediatr*. 2017;63(6):489-94. Epub 2017/10/05. doi: 10.1093/tropej/fmx055.

PubMed PMID: 28977559.

18.   Thi HN, Khanh DK, Thu Hle T, Thomas EG, Lee KJ, Russell FM. Foot Length, Chest

Circumference, and Mid Upper Arm Circumference Are Good Predictors of Low Birth Weight and

Prematurity in Ethnic Minority Newborns in Vietnam: A Hospital-Based Observational Study. *PLoS*

One. 2015;10(11):e0142420. doi: 10.1371/journal.pone.0142420. PubMed PMID: 26555356;

PubMed Central PMCID: PMC4640656.

19.   Folger LV, Panchal P, Eglovitch M, Whelan R, Lee AC. Diagnostic Accuracy of Neonatal Foot

Length to Identify Preterm and Low Birthweight Infants: A Systematic Review and Meta-Analysis.

BMJ Global Health. 2020;5(11):e002976. doi: 10.1136/bmjgh-2020-002976.

20.   Hirve SS, Ganatra BR. Foot Tape Measure for Identification of Low Birth Weight Newborns.

Indian Pediatr. 1993;30(1):25-9. Epub 1993/01/01. PubMed PMID: 8406702.

21.   Papua New Guinea National Department of Health. Sector Performance Annual Report

(Spar) for 2020 : Assessment for Sectors 2016-2022. 2021.

22.   The World Bank Data. Mortality Rate, Neonatal (Per 1,000 Live Births) - Papua New Guinea

2019 [Access Date; cited 2021 15/5/2021]. Available from:

https://data.worldbank.org/indicator/SH.DYN.NMRT?locations=PG.

23.   Emori R, Vallely AJ, Gouda H, Siba P, Phuanukoonnon S, Mola G, et al. Supervised and

Unsupervised Birth and Early Newborn Care Practices in Papua New Guinea: Findings of a Maternal

and Infant Health Survey Conducted in Three Provinces. *Papua New Guinea Medical Journal*.

2017;60(1/2):27–40. doi: 10.3316/informit.301803940285652.

24.   Papua New Guinea National Statistical Office. The 2011 National Population and Housing

Census , Ward Population: Momase Region: National Statistical office; 2014 [Access Date; cited

2022]. Available from: https://sdd.spc.int/collection/2011-population-and-housing-census-

collection-papua-new-guinea.

25.   Vallely AJ, Pomat WS, Homer C, Guy R, Luchters S, Mola GDL, et al. Point-of-Care Testing and

Treatment of Sexually Transmitted Infections to Improve Birth Outcomes in High-Burden, Low-

Income Settings: Study Protocol for a Cluster Randomized Crossover Trial (the Wantaim Trial, Papua

New Guinea). *Wellcome open research*. 2019;4:53-. doi: 10.12688/wellcomeopenres.15173.2.

PubMed PMID: 32030356.

26.   Committee Opinion No 700: Methods for Estimating the Due Date. *Obstet Gynecol*.

2017;129(5):e150-e4. Epub 2017/04/21. doi: 10.1097/aog.0000000000002046. PubMed PMID:

28426621.

27.   Cutland CL, Lackritz EM, Mallett-Moore T, Bardají A, Chandrasekaran R, Lahariya C, et al. Low

Birth Weight: Case Definition & Guidelines for Data Collection, Analysis, and Presentation of

Maternal Immunization Safety Data. *Vaccine*. 2017;35(48 Pt A):6492-500. Epub 2017/11/19. doi:

10.1016/j.vaccine.2017.01.049. PubMed PMID: 29150054; PubMed Central PMCID: PMC5710991.

28. Bland JM, Altman DG. Statistical Methods for Assessing Agreement between Two Methods

of Clinical Measurement. *Lancet*. 1986;1(8476):307-10. Epub 1986/02/08. PubMed PMID: 2868172.

29. Lee A, Mullany L, Ladhani K, Uddin J, Mitra D, Ahmed P, et al. Validity of Newborn Clinical

Assessment to Determine Gestational Age in Bangladesh. *Pediatrics*. 2016;138.

30. Paulsen C, Nielsen BB, Msemo O, Møller SL, Ekmann JR, Theander T, et al. Anthropometric

Measurements Can Identify Small for Gestational Age Newborns: A Cohort Study in Rural Tanzania.

BMC Pediatrics. 2019;19(1):120.

447 Supporting information

S1 Checklist. Standards of Reporting of Diagnostic accuracy studies 2015 checklist.

(https://www.equator-network.org/reporting-guidelines/stard/)

S1 Fig. Bar chart of birth weight, measured on electronic scales (N=342).

S2 Fig. Bar chart of foot length measurements, average of two measurements with a plastic ruler.

Panel A, researcher (N=342), B, volunteer (N=123).

S1 Table. Receiver operating curve statistics for accuracy of newborn foot length to classify low

birth weight, 1 mm increments (N=342). Abbreviations: CI, confidence interval; LHR, likelihood ratio;

PV, predictive value.

S2 Table. Receiver operating curve statistics for accuracy of newborn foot length to classify

preterm birth, 1 mm increments (N=342). Abbreviations: CI, confidence interval; LHR, likelihood

ratio; PV, predictive value.

S1 Protocol. Neofoot study protocol approved by the Papua New Guinea Institute of Medical

Research.

Cardboard box

15 cm plastic ruler

Clear plastic set square at tip of hallux

Hallux (big toe)

Zero cm mark of ruler at base of box

Heel at base of box

Figure1_footlength

Fig. 2. Flow chart indicating number of mothers and newborn babies at each stage of the Neofoot study.

Figure2_flowchart

Figure3_scatterplots

Figure4_roc_lbw_ptb

Figure5_blandaltman