Garetosmab, an inhibitor of activin A, reduces heterotopic

ossification and flare-ups in adults with fibrodysplasia

ossificans progressiva: a randomized, double-blind, placebo-

controlled phase 2 trial

Maja Di Rocco, MD, PhD,[1] Eduardo Forleo-Neto, MD,[2] Robert Pignolo, MD, PhD,[3]

Richard Keen, PhD, FRCP,[4] Philippe Orcel, MD, PhD,[5,6] Thomas Funck-Brentano,

MD, PhD,[5,6] Christian Roux, MD, PhD,[7] Sami Kolta, MD,[7] Annalisa Madeo, MD,[1]

Judith S Bubbear, MD, FRCP,[4] Jacek Tabarkiewicz, MD, PhD,[8] Małgorzata

Szczepanek, MD, PhD,[8] Javier Bachiller-Corral, MD,[9] Angela M Cheung, MD,

PhD,[10] Kathryn M Dahir, MD,[11] Esmée Botman, MD, PhD,[12] Pieter G Raijmakers,

MD, PhD,[13] Mona Al Mukaddam, MD,[14] Lianne Tile, MD,[10] Cynthia Portal-Celhay,

MD, PhD,[2] Neena Sarkar, PhD,[2] Peijie Hou, PhD,[2] Bret Musser, PhD,[2] Anita

Boyapati, PhD,[2] Kusha Mohammadi, PhD,[2] Scott Mellis, MD, PhD,[2] Andrew J.

Rankin, PhD,[2] Aris N. Economides, PhD,[2] Dinko Gonzalez Trotter, PhD,[2] Gary

Herman, MD,[2] Sarah J. O'Meara, PhD,[2] Richard DelGizzi, MS,[2] David M. Weinreich,

MD,[2] George D. Yancopolous, MD, PhD,[2] E. Marelise W. Eekhoff, MD, PhD,[12*]

Frederick S. Kaplan, MD,[14*] on behalf of the LUMINA-1 Investigators

*Co-senior authors: EMWE, Lead Clinical Principal Investigator; FSK, Global Principal

Investigator.

[1]Department of Pediatrics, Unit of Rare Diseases, IRCCS Istituto Giannina Gaslini,

Genoa, Italy; [2]Regeneron Pharmaceuticals, Tarrytown, NY, USA; [3]Department of

Medicine, Mayo Clinic, Rochester, MN, USA; [4]Centre for Metabolic Bone Disease Royal

National Orthopaedic Hospital NHS Trust, London, UK; [5]Department of Rheumatology-

DMU Locomotion, Assistance Publique - Hôpitaux de Paris, Paris, France; [6]INSERM

Université Paris Cité, Paris, France; [7]Department of Rheumatology, Cochin Hospital,

Assistance Publique - Hôpitaux de Paris, Paris, France; [8]Institute of Medical Sciences,

Medical College of Rzeszów University, Rzeszów University, Rzeszów, Podkarpackie,

Poland; [9]Department of Rheumatology, Hospital Universitario Ramón y Cajal, Madrid,

Spain; [10]University Health Network, University of Toronto, Toronto, ON, Canada;

[11]Vanderbilt University Medical Center, Program for Metabolic Bone Disorders,

Nashville, TN, USA; [12]Department of Internal Medicine Section of Endocrinology,

Amsterdam UMC, Vrije Universiteit, Amsterdam Bone Center, Amsterdam, The

Netherlands; [13]Department of Radiology and Nuclear Medicine , Amsterdam UMC, Vrije

Universiteit, Amsterdam, The Netherlands; [14]Departments of Orthopaedics, Medicine

and the Center for Research in FOP & Related Disorders, University of Pennsylvania

Perelman School of Medicine, Philadelphia, PA, USA.

ABSTRACT

Background

Fibrodysplasia ossificans progressiva (FOP), an ultra-rare disorder caused by mutations

in the gene encoding activin A receptor type 1 (*ACVR1)*, is characterized by painful

flare-ups and cumulative heterotopic ossification (HO). Garetosmab, a fully-human

monoclonal antibody blocking activin A, prevents HO in FOP mice.

Methods

LUMINA-1 (NCT03188666) was a phase 2 , multi-center, randomized, double-blind,

placebo-controlled study evaluating the safety, tolerability, and effects on HO of

intravenous (IV) garetosmab 10 mg/kg every 4 weeks (Q4W). Adult patients with FOP

were randomized to garetosmab or placebo for 28 weeks (Period_1), followed by an

open-label period (Period_2). After Period_2, patients were allowed to stay on

garetosmab in an open-label extension. For Period_1, primary endpoints were HO total

lesion activity (HO-TLA) by ^{18}F-sodium fluoride positron emission tomography (^{18}F-NaF

PET) and HO total lesion volume by computed tomography (CT). The Period_2 primary

endpoint compared the number of new lesions in Period_2 versus Period_1. The safety

primary endpoint was incidence and severity of TEAEs through the end of the Period 1

at week 28.

Findings

Patients (n=44) were randomized to garetosmab (n=20) or placebo (n=24). In Period_1,

there was a trend for garetosmab to decrease HO-TLA versus placebo (24.6%; *P*=0.07),

primarily driven by near complete prevention of new lesions (97% decrease by ^{18}F-NaF

PET, post-hoc P=0.009; 90% relative reduction by CT, post-hoc P=0.017); flare-ups

were significantly reduced (P=0.0005). For placebo patients transitioning to garetosmab

in Period_2, no patients developed new HO lesions (0% in Period_2 versus 40.9% in

Period_1; P=0.0027) by CT. All 44 patients met primary safety endpoint of at least one

TEAE during Period 1. Garetosmab was associated with more adverse events than

placebo: mild recurrent epistaxis, madarosis, and skin/soft tissue infections. Overall, the

AEs were predominantly mild in severity, with no effect on patients' ability to receive

garetosmab. Five deaths (5/44; 11.4%) occurred either in Period_2 or the open-label

extension. The deaths were associated with baseline disease severity in some,

preexisting comorbidities in others and occurred following 8-16 doses (median: 15) of

garetosmab in the open label/follow-up periods.

Interpretation

Garetosmab reduced flare-ups and prevented new HO lesions in FOP patients.

Although side effects were mild to moderate, there were a relatively high number of

deaths for a small study; the deaths were not related to epistaxis and considered

unlikely to be related to garetosmab.

Funding: Regeneron Pharmaceuticals, Inc.

Introduction

Fibrodysplasia ossificans progressiva (FOP; MIM#135100) is an ultra-rare, autosomal

dominant disorder[1] with an estimated prevalence of 0.36–1.36 per million.[2-5] The most

common and debilitating manifestation of FOP is the heterotopic ossification (HO) of

connective tissues, such as tendons, ligaments, and skeletal muscles excepting the

diaphragm, extraocular muscles and the tongue.[6] HO typically presents during the first

decade of life and may be accompanied by painful, soft tissue inflammatory swellings,

known as flare-ups, however, this feature of the disease is variable.[7] A diagnostic

hallmark of FOP at birth is bilateral malformation of the great toes, which is often missed

and can lead to a delay in diagnosis.[8]

HO in FOP is seemingly episodic, and may be precipitated by trauma.[9] The

resulting HO lesions accumulate, leading to joint immobility, skeletal deformity, severe

pain, disability, and early mortality.[6,10,11] Patients with FOP experience significant

disease burden including restriction of activity as well as hearing loss which worsens

social isolation leading to decreased quality of life. Surgical removal of bone is

ineffective as it causes new HO and is associated with high anesthetic challenges.[12]

The degree of immobility can be captured by the cumulative analog joint involvement

scale (CAJIS) score, measuring mobility across 15 anatomic locations, and ranging

from 0 to 30 (higher scores indicating more severe limitations in mobility and other

function).[13] On the basis of FOP features related to joint dysfunction (flare-up activity,

body regions affected, thoracic insufficiency, and other complications) as well as its

consequences (assistance in activities of daily living, ambulation, and CAJIS score), 5

clinical stages of FOP have been defined: early, moderate, severe, profound, and end

of life.[14] Estimated median age of survival for patients with FOP is 56 years.[15] Mortality

is correlated with disease severity and CAJIS score and is primarily due to

cardiorespiratory failure from thoracic insufficiency syndrome, pneumonia, and

complications of falls.[15] With no globally-approved disease-modifying therapies, clinical

management is mostly limited to symptomatic flare-up treatment with corticosteroids or

non-steroidal anti-inflammatory agents, evidence of which is anecdotal.[7]

FOP is caused by heterozygous missense mutations in *ACVR1*, the gene encoding

activin A receptor type 1, a bone morphogenetic protein (BMP) type I receptor.[16] A

significant breakthrough in understanding the molecular mechanism of HO in FOP was

the discovery that FOP-causing variants of ACVR1 recognize activin A as an agonist

(**Figure S1**).[17-20] In contrast, activin A normally inhibits BMP signalling via wild-type

ACVR1.[17,21] Garetosmab, a fully-human monoclonal antibody (generated using

VelocImmune™ technology),[1,2] binds activin A and blocks its ability to activate FOP-

mutant ACVR1. Inhibition of activin A with garetosmab blocked new HO lesions and

resulted in stasis or partial reduction in the volume of pre-existing lesions in a

genetically humanized murine FOP model in which FOP-mutant ACVR1 was

introduced.[17,18] As such, it was hypothesized that garetosmab may beneficially impact

FOP disease progression in humans. LUMINA-1 was designed to assess the safety and

efficacy of garetosmab as a potential disease-modifying treatment for FOP, and

consequently also validate the role of activin A as a key driver of HO in FOP.

Methods

Study design

LUMINA-1 (NCT03188666) was a phase 2, randomized, double-blind, placebo-controlled study evaluating the safety, tolerability, and effects on HO of intravenous (IV) garetosmab 10 mg/kg every 4 weeks (Q4W). The study was conducted at 11 sites in 8 countries across North America and Europe, between February 26, 2018 and September 16, 2021. [18]F-NaF PET and whole-body, low-dose X-ray computed tomography (CT)[22] were used to measure and track HO. The study consisted of a 28-day screening/baseline period followed by a 28-week randomized, double-blind, placebo-controlled period (Period_1), a 28-week open-label treatment period (Period_2), and a subsequent open-label extension from week 56 to end of study (**Figure S2**). The primary analysis was conducted when all patients completed Period_1. Further analyses were conducted at the end of Period_2. Amendments to the trial protocol are detailed in **Table S1**. Patient safety and welfare were monitored by an Independent Data Monitoring Committee. This study was conducted in accordance with the 2013 Declaration of Helsinki and the International Council for Harmonization guidelines for GCP. All patients provided written, informed consent. Detailed information and a full list of the Institutional Review Boards are in the **Supplementary Methods**.

Patients

Adult patients (aged 18–60 years) with a clinical diagnosis of FOP (based on congenital great toe malformation, episodic soft tissue swelling, and/or progressive HO) were included. Clinical diagnosis was confirmed with documentation of any FOP-causing

ACVR1 variant and disease activity (defined as pain, swelling, stiffness, and other

signs/symptoms associated with FOP flare-ups; or worsening of joint function, or

radiographic progression of HO) within 1 year of screening. Key inclusion/exclusion

criteria are in **Table S2** and **Supplementary Methods**.

Randomization and masking

Enrolled patients were randomized (1:1) to receive garetosmab 10 mg/kg Q4W as

previously assessed,[23] or placebo in Period_1, according to a central randomization

scheme. Block randomization was done using an interactive response technology (IRT)

provided to the designated study pharmacist or qualified designee. Randomization was

stratified by presence/absence of baseline active HO lesions, gender, and mutation

type. All PET/CT scans were reviewed by 2 independent readers and an adjudicator; all

3 were blinded to treatment assignment.

Procedures

Patients were assigned to receive garetosmab 10 mg/kg or placebo Q4W for 28 weeks

(Period_1). After week 28, all patients continued or were switched to receive

garetosmab 10 mg/kg Q4W (Period_2). At the conclusion of Period_2, patients were

given the option to continue on garetosmab in an open-label extension. Whole-body

PET/CT scans were acquired to identify pre-existing target lesions at baseline, identify

new HO lesions, and measure volume changes in pre-existing target and new HO

lesions. Baseline imaging with PET/CT was performed at most 7 days prior to study

drug administration and at Weeks 8, 28, 56, and 76 (**Figure S2**). Detailed information

on imaging acquisition and read procedures are in the **Supplementary Methods**.

Outcomes

Period_1: The pre-specified primary endpoint for efficacy was the effect of garetosmab

versus placebo on time-weighted average (TWA) of the percentage change from

baseline in total lesion activity (TLA) by PET; TLA is considered proportional to the

deposition rate of bone mineral into actively forming HO lesions (definition in

supplementary methods section). The next endpoint in the hierarchy was to assess

percentage change of the total volume of HO lesions by CT in Period_1 relative to

baseline. Other key secondary and exploratory endpoints are listed in **Table S3**.

Exploratory endpoints in Period_1 included the percentage of patients with flare-ups as

assessed by patient diary and investigator report.

Period_2: Based on the outcomes of Period_1 (see Results), the primary endpoint for

efficacy for Period_2 was prospectively changed to the number of new lesions in

patients transitioning from placebo to garetosmab as assessed by CT.

Safety: The primary safety endpoint for Period_1 and a primary endpoint for Period_2

was the incidence and severity of adverse events, which included those not present at

baseline or an exacerbation of a pre-existing condition.

Statistical analyses

Testing of primary and key secondary efficacy outcomes followed a pre-specified

hierarchical testing procedure to address multiplicity at an overall 2-sided alpha=0.05

significance for Period_1. Other secondary outcomes were tested at the nominal 2-

sided alpha=0.05 significance without multiplicity adjustment. In Period_2, a 2-sided

10% significant level was applied for the primary and key secondary endpoints. For the

Period_1 primary analysis, the TWA percent change from baseline in TLA ^{18}F-NaF PET

over 28 weeks was analyzed in AHO (baseline active HO analysis set) and AHOC

(baseline active HO classic ACVR1[R206H] mutation analysis set) populations using an

analysis of covariance model. The difference in least squares mean change from

baseline, 95% confidence interval, and *P*-value were provided from the model to

compare garetosmab against placebo group. For Period_2 primary analysis, the

Wilcoxon signed-rank test was used to compare the number of new HO lesions as

assessed by CT at week 56 (relative to week 28 scan) with the number of new HO

lesions at week 28 (relative to baseline scan) in patients who crossed over from placebo

to garetosmab. The estimate and 95% confidence interval of the rate of new HO lesions

at week 56 and that of the rate ratio (comparing Period_2 versus Period_1) were based

on a negative binomial model with repeated measures and using a generalized

estimating equation. Safety outcomes were analyzed using descriptive statistics.

Further details are provided in the **Supplementary Methods** and amendments to the

statistical methodology are summarized in **Table S1**.

Results

LUMINA-1 enrolled 44 patients with FOP, all of whom had active HO (AHO) at baseline

(see full definition in the **Supplementary Methods**). Overall, 42/44 (95.5%) of patients

had the "classic" FOP-causing variant of *ACVR1* (c.617G>A, p.R206H) (AHOC

population). Patients were randomized to garetosmab 10 mg/kg IV Q4W (n=20) or

placebo (n=24) (**Figure S2; Figure S3**). Baseline demographics and disease

characteristics were similar between groups (**Table S4**). The mean (SD) age was 27.6

(8.5) years. The mean CAJIS score at baseline was 15.7 (SD: 6.6; range, 6–30; Median

[Quartile1, Quartile3]: 15.0 [11, 19]). Data analyses presented are for the AHO

population and were similar to the AHOC population (**Table S5A**). The study was

initiated in February 2018; primary data cut-off was September 17, 2019 (week 28);

additional data cut-off dates for efficacy analyses were August 11, 2020 (week 56), and

October 30, 2020 (week 76); safety was reported until last patient last visit on

September 16, 2021.

Period_1 Results

Forty-three (98%) patients completed Period_1. During Period_1, garetosmab reduced

TLA (new and existing lesions) from baseline versus placebo by 24.6% (95%

confidence interval [CI]: -51.8%–2.5%; *P*=0.07; **Figure 1A**; **Supplementary Section:**

Results). Although this did not meet the pre-specified statistical threshold of *P*≤0.05,

patients treated with garetosmab had a ~97% relative reduction in new lesion activity, a

~90% relative reduction in new HO lesion volume compared with the placebo arm

(**Figure 1B,C; Table S6**). Additionally, a reduction in both the number of new HO

lesions per patient as assessed by PET (mean of 0.15 new lesions/patient for

garetosmab versus 1.19 for placebo; post-hoc analysis, nominal P=0.006) and CT (0.15

versus 1.13, respectively; post-hoc analysis, nominal P=0.009; **Table S6**) were

observed. The total number of new HO lesions in Period_1 for placebo was 27 by CT

and 29 by PET compared with 3 by CT and PET for garetosmab (**Figure 1B**). The

percentage of patients who developed new lesions in Period_1 was lower with

garetosmab (15% by CT; 15% by PET) versus placebo (45.8% by CT and PET; nominal

P=0.05; **Figure S4**).

Given that garetosmab appears to block the formation of new HO in patients with

FOP, its effects on normal bone were also examined as reference using PET.

Assessment of percent change from baseline of the mean [standard error] SUVmean of

selected normotopic bones (as defined in the protocol), showed no differences between

treatments at week 8 (garetosmab: 42.0% [9.9]; placebo: 20.3% [9.2]; nominal P=0.12)

or week 28 (garetosmab: 24.6% [8.4]; placebo: 28.9% [7.7]; nominal P=0.71; data not

shown), indicating that garetosmab does not affect bone turnover.

In addition to the reduction in the number of new HO lesions, an effect of

garetosmab was also observed on reported flare-ups. A marked reduction was

observed during Period_1 in patients reporting flare-ups on garetosmab versus placebo

based on patient diary entries (35.0% versus 70.8%, respectively; relative risk

[RR]=0.49; nominal P=0.03) and investigator adverse event reports (10.0% versus

41.7%, respectively; RR=0.24; post-hoc analysis, nominal P=0.039; **Table 1A; Figure**

S5A, B).

Since garetosmab's most pronounced effect on HO during Period_1 was the

suppression of new HO lesions formation, the study hypothesis for Period_2 was

redefined to be that garetosmab prevents formation of new HO lesions.

Period_2 Results

Of the 43 patients who completed Period_1, all entered Period_2 and 42 patients

completed Period_2. Patients randomized in Period_1 to placebo crossed-over to

receive garetosmab. In Period_2, the total number of new lesions measured by CT for

patients who crossed over from placebo to garetosmab was reduced 100% relative to

Period_1, (0 versus 22, respectively; P=0.0039) and 95% by PET (1 versus 23,

respectively; P=0.0039; **Figure 2A, B; Table S5B**). Mean new lesion volume and

activity were significantly lower for Period_2 versus Period_1 (volume: 0.05 cm^3 versus

9.29 cm^3, respectively, P=0.0039; activity: 13.20 versus 204.45, respectively; P=0.0273;

Figure 2C, D). This reduction was also reflected in the percentage of patients who

developed new lesions, which was significantly lower during Period_2 compared with

Period_1 by CT (0% versus 40.9%, respectively; P=0.0027) and PET (4.5% versus

40.9%, respectively; P=0.0047; **Figure 2E, F).**

For patients originally randomized to garetosmab and who remained on garetosmab

in Period_2, garetosmab's efficacy in preventing new HO lesions was maintained (0

new lesions by CT and PET in Period_2 versus 2 by CT and 1 by PET in Period_1;

Figure S6. Additionally, the percentage of patients with new lesions was 0% for

Period_2 by PET and CT (**Figure S6**) for patients remaining on garetosmab.

Open-Label Extension Results

During the subsequent open-label extension, 0 new lesions at week 76 were observed

in patients crossing to garetosmab after week 28 (n=17) while one new lesion was

observed by both PET and CT at 76 weeks among patients continuing garetosmab

since baseline (n=15). Overall, treatment with garetosmab resulted in a sustained and

pronounced effect in preventing new HO lesions from forming up to 76 weeks (the time

of last imaging scan assessments in the study). Thirty-four (77%) patients continued in

the study, after the open-label extension.

Flare-Ups and CAJIS

For patients switching to garetosmab after week 28, the proportion with new flare-ups

was significantly lower in Period_2 compared to Period_1 as reported by both patients

(13.6% versus 68.2%, respectively; P=0.0005) and investigators (13.6% versus 45.5%,

respectively; P=0.0196; **Table 1B; Figure S5C, D**). The total number of new flare-ups

by patient diary was 11 in Period_2 compared with 31 in Period_1 and by investigator

report was 4 and 22, respectively (**Figure S5E, F**). The relationship between baseline

age and CAJIS score is depicted in **Figure 3A.**

Safety

All patients in the LUMINA-1 trial reported ≥1 AE during Period_1 (**Table 2**). One patient

with a medical history of restrictive lung disease, pulmonary congestion, and mucus

plugging discontinued during Period_1 due to an TEAE of mild pyrexia that followed

recurrent episodes of pneumonia and hemoptysis. Most AEs were considered by

investigators to be mild to moderate in severity. In Period_1, notable imbalances in AEs

with garetosmab versus placebo included epistaxis (50.0% versus 16.7%), madarosis

(loss of eyebrows/eyelashes; 30.0% versus 0%), and a composite of skin and soft

tissue infections that included acne (65% versus 12.5%; **Table S7; Table S8**). Nine

serious adverse events (SAEs) in Period_1 occurred in 6 (13.6%) patients; of these, 4

(20.0%) patients received garetosmab versus 2 (8.3%) patients who received placebo

(**Table S9**). SAEs in the garetosmab group included 3 (15.0%) patients with infections

and infestations, 1 (5.0%) patient with epistaxis (5.0%), and 1 (5.0%) with intestinal

obstruction. The SAE of epistaxis resulted in hospitalization for nasal packing and was

assessed as related to garetosmab by the investigator, but patient required no

transfusions and had no drop in hemoglobin. The patient fully recovered and continued

in the study without recurrence. This SAE of epistaxis led to a protocol amendment to

include additional exclusion criteria (**Table S2**) and baseline (if applicable) and post-

treatment laboratory measures of coagulation parameters and platelet effector function

to exclude patients who may have existing propensity for bleeding or mitigate the

potential risk for epistaxis.

Coagulation tests and platelet functional assays including activated prothrombin

time (aPTT), prothrombin time (PTT), and prothrombin international normalized ratio

(INR), measured in a subset of patients at baseline and post treatment, were in the

normal range at baseline and fluctuations observed in placebo or in the garetosmab

treatment group remained within the normal ranges (**Figure S7**). In Period_1, 13

(65.0%) patients in the garetosmab treatment group experienced a bleeding event

compared to 9 (37.5%) patients in the placebo treatment group (**Table S10**).The only

bleeding event reported in more than two patients in the garetosmab group was

epistaxis. Except for the single epistaxis SAE, other events of epistaxis were non-

serious and the majority mild in severity. Other non-epistaxis bleeding events were

balanced between garetosmab and placebo recipients and non-serious. None of the

patients discontinued therapy due to a bleeding AE.

All patients reported ≥1 AE during Period_2 and subsequent open label extension,

which involved patients receiving garetosmab. The most frequently reported AEs were

consistent with those reported in Period_1, comprising of skin and soft tissue infections

(81.4%; acne (32.6%), madarosis (46.5%)) and epistaxis (34.9%), and were mostly

mild/moderate in severity (**Table S8; Table S11; Figure 3B and 3C**). During Period_2

and subsequent treatment until end of study, epistaxis was reported in 15 patients

(34.9%). Most events were mild, and no patients discontinued treatment with

garetosmab due to epistaxis. Twenty SAEs in 13 (30.2%) patients were reported during

Period_2 until end of study, including infections and infestations (16.3%) and

gastrointestinal disorders (4.7%; **Table S12**). Five (11.6%) patients experienced 6 SAEs

of abscess requiring a hospital admission or ER visit for incision and drainage. These

events resolved and patients continued garetosmab after a temporary interruption of

study drug; 5 of these 6 SAEs were deemed related to garetosmab by the investigator.

In order to determine if epistaxis events were related to changes in BMP/TGFß

family members other than activin A known to regulate angiogenesis and vascular

endothelium homeostasis, levels of BMP9 were measured.[24,25] Minor fluctuations in

serum BMP-9 were observed in both the placebo and in the garetosmab treatment

group. The onset of epistaxis occurred in patients with stable and fluctuating serum

BMP-9 concentrations. (**Figure S8**).

The percentage of patients with infusion reactions was balanced between

garetosmab (5/20 [25%]) and placebo (6/24 [25%]). Except for two infusion reactions of

moderate severity that occurred in patients who received garetosmab during Period_1,

the other infusion reactions were reported as mild. Some reactions required a temporary

infusion interruption or antihistamine (loratadine) pre-medication (n=2), but all infusions

were completed. No patient discontinued treatment due to infusion reactions and none

of the infusion reactions were associated with signs or symptoms of anaphylaxis or

development of antidrug antibodies.

There were five deaths (5/44; 11.4%) during the study, all of which occurred in

Period_2 (Open label) or in the open label extension between 8-16 (median-15) doses

of garetosmab; none occurred during the double-blind period (Period_1). The deaths

were reported by investigators as unrelated to garetosmab. The case summaries can be

found in Supplement **Table S13**. As can be seen in **Figure 3B and 3D**, there was no

relationship between subjects who died and the frequency or severity of epistaxis. The

causes of death were head and brain trauma due to a fall in the setting of severe motor

disability; hemorrhagic stroke in the setting of poorly controlled hypertension and

previous strokes; fatal intestinal obstruction in the setting of a previous episode of

intestinal obstruction; cardiac arrest during surgery procedure for traumatic spleen

rupture due to fall; sudden cardiac death in a patient with lung granulomatous

inflammation most likely attributable to chronic pulmonary aspiration.

Of the five deaths, three patients had a CAJIS score ≥24 and either profound or

end-of-life disease severity as measured by the clinical staging of FOP developed by

Pignolo & Kaplan.[14] Among the fatal events in study participants with lower CAJIS

scores, one occurred in a 26 to 30-year-old patient with a CAJIS score of 16 (moderate

disease staging) who died from a fall resulting in severe head and brain trauma. This

individual had significant rigidity and walking disability at baseline. Another was a 36 to

40-year-old patient with a CAJIS score of 19 (severe disease staging) who died from

apparent sudden cardiac arrest with extensive granulomatous formation in the lungs

consistent with chronic aspiration of foreign material.

Discussion

This phase 2 study was conducted to assess the safety and efficacy of garetosmab in

patients with FOP. The initial hypothesis for Period_1 was that garetosmab prevents

formation of new lesions and progression of pre-existing lesions. This was based on

preclinical data demonstrating that activin A is required for the formation of new

heterotopic bone lesions, as well as continued growth and expansion of pre-existing, yet

nascent lesions.[17,18] Data from Period_1 demonstrated that the most pronounced effect

of garetosmab was the suppression of the formation of new HO lesions. Garetosmab

did not significantly affect the continued growth and expansion of pre-exisiting HO

lesions. Contrary to our apriori hypothesis, these established lesions had already

achieved maturity prior to the administration of garetosmab as evidenced by no

significant growth in the placebo arm (**Table S14**). This finding led to prospective

redefining of the study hypothesis for Period_2 to be that garetosmab prevents

formation of new HO lesions in patients switching from placebo to garetosmab. The

hypothesis was agreed upon with the US Food and Drug Administration, along with

concordance that patients exposed to placebo during Period_1 could serve as the

control group when they transitioned to receive garetosmab during Period_2 given the

ultra-rare nature of FOP and striking results in Period_1. Notably, no new lesions were

detected in Period_2 by CT in patients transitioning to garetosmab, while efficacy

persisted in those who received garetosmab during Period_1 and continued

garetosmab in Period_2. This study also demonstrated that PET and CT scans were

similarly well correlated in the detection of new lesions.

In addition to the significant reduction in the number of new HO lesions, treatment

with garetosmab demonstrated a significant effect on the incidence and frequency of

soft-tissue inflammatory flare-ups based upon both patient diary and investigator

adverse event reporting. Flare-ups are a significant burden for patients with FOP, often

are painful and debilitating and require the use of corticosteroids in high doses or during

prolonged periods of time. The reduction in flare ups by garetosmab suggests that flare-

ups are also associated with the activin A pathway and highlights an additional potential

utility of garetosmab as a disease-modifying and steroid-sparing treatment for patients

with FOP.

Data from patients treated with garetosmab in both Period_1 and Period_2 highlight

the durability of garetosmab in preventing the formation of new HO lesions, as evident

from both CT and PET scan results. This effect by garetosmab was sustained through

week 76.

The clinical program for garetosmab included two previous phase 1 first-in-human

studies conducted in healthy volunteers (Clinicaltrials.gov: NCT02870400,

NCT02943239) that showed garetosmab in multiple doses up to 10 mg/kg (single dose)

and 10 mg/kg Q4W for 4 doses were well-tolerated. In one study of 30 healthy females

treated with garetosmab and 10 randomized to placebo, there were no occurrences of

epistaxis or skin and soft tissue infections, suggesting that these effects of garetosmab

maybe specific to FOP.[23] In another phase 1 study (NCT02943239) combining

garetosmab with an anti-myostatin, GDF8 mAb (n=82), there was a single epistaxis

event and 2 episodes of postmenopausal bleeding events. Overall there were 3 (3.75%)

bleeding event AEs (1 epistaxis and 2 episodes of postmenopausal bleeding) amongst

122 volunteers which were all non-serious and mild in severity. The review of the

healthy volunteer data from Phase I clinical trials did not suggest a signal between

garetosmab and epistaxis although the exposure to garetosmab was brief.

Overall, in this study, the AEs were predominantly mild in severity, with no effect on

patients' ability to receive garetosmab. Notable AE imbalances with garetosmab during

Period_1 included epistaxis, madarosis, and a composite of skin and soft tissue

infections (which included skin abscesses, folliculitis, cellulitis, furuncles and

carbuncles). Based on results from Period_1, higher incidences of epistaxis, madarosis,

and skin and soft tissue infections were determined to be associated with garetosmab.

Further investigation shows that while there was an apparent link between

garetosmab and epistaxis, there is no apparent link between garetosmab and bleeding

events in other organs or systems; these events were balanced during Period_1. The

great majority of epistaxis events throughout the study were mild in severity, with only 4

moderate events and one severe event. The biological mechanism behind inhibition of

activin A with garetosmab and the occurrence of epistaxis in a subset of FOP patients is

not understood. While activin A utilizes the same type II receptors as BMP9, which is

associated with angiogenesis and vascular endothelium homeostasis,[24,25] treatment-

related changes or epistaxis events did not correlate with fluctuations in serum BMP-9

levels. Furthermore, there was no difference in serum BMP9 levels among patients with

a fatal outcome. Coagulation tests and platelet functional assays were also in the

normal ranges. Based on available data, no abnormalities were reported suggesting

that the patients enrolled did not have coagulation defects at study entry or post

garetosmab treatment.

Five fatal events were observed during Period_2 and subsequent open-label

extension; in both periods patients received only garetosmab. Three of the 5 patients

who died experienced epistaxis and all events were mild to moderate in severity. Two of

these patients experienced one event (one of them had history of epistaxis prior to

study enrollment, and the other had medical history of brain cavernoma hemorrhage as

well as pigmented purpuric dermatosis, which is an inflammatory dermatitis with

vascular fragility); the third patient reported 2 epistaxis events (**Figure 3B**).

Of the five fatalites, two of the deaths were due to complications resulting from falls,

which is common in patients with FOP. One death was due to intestinal obstruction

which has been previously reported as a complication of FOP.[15] One death occurred as

a result of a hemorrhagic stroke to the deep structures of the brain leading to acute

respiratory failure. This patient presented with difficult-to-control arterial hypertension

and interpretation of the brain CT by two independent neuroradiologists concluded

evidence of chronic hypertensive effects and the findings suggested death was likely a

consequence of hypertension. One death was initially considered to be related to a

hemorrhagic event in the lung, which generated concern around the possible

association between bleeding and deaths, given that early in the course of the study

epistaxis had already been identified as a risk associated with garetosmab. As a result,

a temporary pause in patient dosing was instituted to allow the sponsor time to conduct

thorough investigations into the deaths, evaluate the autopsy reports performed in 2 of

the 5 patients with a fatal outcome, and further assess a potential association between

garetosmab and bleeding risk. The cause of death of the patient initially thought to have

experienced lung hemorrhage was ultimately determined to be sudden cardiac arrest

secondary to recurrent aspiration. There was no evidence of any bleeding which was

further confirmed by a second independent autopsy report from a pathologist at a

separate institution.

While there are published reports of mortality in FOP, there are no published

reports of annualized rates of death in this community.[13] With this in mind, post-hoc

assessment of safety data by all the authors concluded: 1) While epistaxis events were

related to garetosmab 2) there was no apparent association between epistaxis and

patient deaths; 3) causes of death appeared consistent with known causes of death and

life expectancy for patients with FOP of similar age and disease severity as measured

by clinical staging.[15] Four of the 5 patients who died on study had advanced CAJIS

scores. The one relatively young patient with a lower CAJIS score was known to have

significant gait difficulties due to FOP, and died from a fall down a flight of stairs not

believed to be precipitated by an antecedent bleed. Garetosmab was considered

unlikely to be the cause of death in these cases, which appeared more likely to have

been related to the underlying severity of their FOP disease. Although a definitive link to

garetosmab could not be established in this population with advanced disease, the

number of deaths was relatively high for this small study. As these deaths were in the

open label period, although a contribution by the drug can not be excluded, no common

pathogenetic mechanism of deaths related to garetosmab has been identified.

The major impact of garetosmab in reducing new HO suggests that the greatest

utility of garetosmab may be early in the disease course before substantial HO has

occurred and before permanent disability ensues. Together with previously published

preclinical data in murine models of FOP, these data indicate that activin A is also

required for the continued growth and progression of new lesions. The reduction in

flare-ups provides an additional and important benefit by significantly reducing disease

burden. Together, the data generated during LUMINA-1 demonstrate that activin A is a

required ligand for HO in FOP, and provides strong evidence that inhibition of activin A

using garetosmab is a promising disease-modifying therapy with the ability to not only

block HO but also reduce the number and severity of flare-ups. In the context of this

ultra-rare disease setting, the patient population was small with limited patient exposure.

A phase 3 adult study and a pediatric study are planned to continue to assess the

benefits and risks of garetosmab intervention in people with FOP, and further validate

the results from this small phase 2 study.

Conclusions

Garetosmab treatment has demonstrated substantial and persistent reduction of new

heterotopic bone lesion formation and flare-ups in adults with FOP. Although a definitive

link between garetosmab and the five deaths was not established in this population with

advanced disease, the number of deaths was relatively high for this small study,

therefore the benefit-risk of garetosmab will be further evaluated in Phase 3 clinical

studies. Garetosmab may provide a therapeutic option in this ultra-rare, severely

debilitating, life-threatening disease and further studies are planned.

Consortium Author Information

Authors and Affiliations

Department of Pediatrics, Unit of Rare Diseases, IRCCS Istituto Giannina Gaslini,

Genoa, Italy

Maja Di Rocco, Annalisa Madeo

Regeneron Pharmaceuticals, Tarrytown, NY, USA

Eduardo Forleo-Neto, Cynthia Portal-Celhay, Neena Sarkar, Peijie Hou, Bret Musser,

Anita Boyapati, Kusha Mohammadi, Scott Mellis, Andrew J. Rankin, Aris N.

Economides, Dinko Gonzalez Trotter, Gary Herman, Sarah J. O'Meara, Richard

DelGizzi, David M. Weinreich, George D. Yancopolous

Department of Medicine, Mayo Clinic, Rochester, MN, USA

Robert Pignolo

Centre for Metabolic Bone Disease Royal National Orthopaedic Hospital NHS

Trust, London, UK

Richard Keen, Judith S Bubbear

Department of Rheumatology- DMU Locomotion, Assistance Publique - Hôpitaux

de Paris, Paris, France

Philippe Orcel, Thomas Funck-Brentano

INSERM Université Paris Cité, Paris, France

Philippe Orcel, Thomas Funck-Brentano

Department of Rheumatology, Cochin Hospital, Assistance Publique - Hôpitaux

de Paris, Paris, France

Christian Roux, Sami Kolta

Institute of Medical Sciences, Medical College of Rzeszów University, Rzeszów

University, Rzeszów, Podkarpackie, Poland

Jacek Tabarkiewicz, Małgorzata Szczepanek

Department of Rheumatology, Hospital Universitario Ramón y Cajal, Madrid,

Spain

Javier Bachiller-Corral

University Health Network, University of Toronto, Toronto, ON, Canada

Angela M Cheung, Lianne Tile

Vanderbilt University Medical Center, Program for Metabolic Bone Disorders,

Nashville, TN, USA

Kathryn M Dahir

Department of Internal Medicine Section of Endocrinology, Amsterdam UMC, Vrije

Universiteit, Amsterdam Bone Center, Amsterdam, The Netherlands

Esmée Botman, E. Marelise W. Eekhoff

Department of Radiology and Nuclear Medicine , Amsterdam UMC, Vrije

Universiteit, Amsterdam, The Netherlands

Pieter G Raijmakers

Departments of Orthopaedics, Medicine and the Center for Research in FOP &

Related Disorders, University of Pennsylvania Perelman School of Medicine,

Philadelphia, PA, USA

Mona Al Mukaddam, Frederick S. Kaplan

Richard Keen: PI of clinical trials sponsored by Clementia/Ipsen and Regeneron

Pharmaceuticals Inc; non-paid member of the International Clinical Council on FOP and

IFOPA registry advisory board.

Philippe Orcel: PI of clinical trial sponsored by Regeneron Pharmaceuticals Inc.

Thomas Funck-Brentano: is a sub-investigator of the Regeneron Pharmaceuticals Inc.

LUMINA-1 trial and PI of IPSEN FALKON trial

Christian Roux: research grants, for the institution, from Regeneron Pharmaceuticals

Inc.

Sami Kolta: sub-investigator of the Regeneron Pharmaceuticals Inc. LUMINA-1 trial.

Annalisa Madeo: sub-Investigator in clinical trials sponsored by Regeneron

Pharmaceuticals Inc. and Clementia-Ipsen.

Judith S. Bubbear: sub-investigator of the Regeneron Pharmaceuticals Inc. LUMINA-1

and Clementia/Ipsen MOVE trials.

Jacek Tabarkiewicz: grants to institution (UR), speaker for Merck, Novartis; hired

scientific expert for SoftSystem.

Małgorzata Szczepanek: speaker for Roche.

Javier Bachiller-Corral: investigator of clinical trial sponsored by Regeneron

Pharmaceuticals Inc.

Angela M. Cheung: grant to institution (UHN) for clinical trial, no personal COI.

Kathryn M. Dahir: PI on the Regeneron Pharmaceuticals Inc. Trial. This project

described was supported by CTSA award No. UL1 TR002243 from the National Center

for Advancing Translational Sciences. Its contents are solely the responsibility of the

authors/sponsor and do not necessarily represent official views of the National Center

for Advancing Translational Sciences or the National Institutes of Health.

Esmée Botman: is a sub-investigator of the Regeneron Pharmaceuticals Inc. LUMINA-1

trial at the Amsterdam UMC, the Netherlands.

Pieter G Raijmakers: is a sub-investigator of the Regeneron Pharmaceuticals Inc.

LUMINA-1 trial at the Amsterdam UMC, the Netherlands.

Mona Al Mukaddam: PI of clinical trials sponsored by Clementia/Ipsen, Regeneron

Pharmaceuticals Inc. and Incyte; non-paid member of the International Clinical Council

on FOP and IFOPA registry advisory board.

Lianne Tile: grant to institution (UHN) for clinical trial (sub-investigator), no personal

COI.

Cynthia Portal-Celhay: employee of and holds stocks and shares in Regeneron

Pharmaceuticals Inc.

Melissa Simek-Lemos: employee of and holds stocks and shares in Regeneron

Pharmaceuticals Inc.

Neena Sarkar: employee of and holds stocks and shares in Regeneron

Pharmaceuticals Inc.

Peijie Hou: employee of and holds stocks and shares in Regeneron Pharmaceuticals

Inc.

Bret Musser: employee of and holds stocks and shares in Regeneron Pharmaceuticals

Inc.

Anita Boyapati: employee of and holds stocks and shares in Regeneron

Pharmaceuticals Inc.

Scott Mellis: employee of and holds stocks and shares in Regeneron Pharmaceuticals

Inc.

Andrew J. Rankin: employee of and holds stocks and shares in Regeneron

Pharmaceuticals Inc.

Aris N. Economides: employee of and holds stocks and shares in Regeneron

Pharmaceuticals Inc.

Dinko Gonzalez Trotter: employee of and holds stocks and shares in Regeneron

Pharmaceuticals Inc.

Gary Herman: employee of and holds stocks and shares in Regeneron Pharmaceuticals

Inc.

David M. Weinreich: employee of and holds stocks and shares in Regeneron

Pharmaceuticals Inc.

George D. Yancopolous: employee of and holds stocks and shares in Regeneron

Pharmaceuticals Inc.

Acknowledgements

We thank the patients and their families for their participation in this study and the healthcare professionals and investigators who treated these patients and made this study possible.

We acknowledge the contributions of Kevin Niswender, MD, PhD; Edward Sherwood. MD; Lana Howard, BSN,RN, CCRP; Margo Black, MSN, RN, CCRP, at Vanderbilt University Medical Center.

We acknowledge the contributions of Staci Kallish DO, Katherine S.Toder CCRC, and Renee Jurek CRC at the University of Pennsylvania.

We thank Donal O'Donovan Ph.D. for the safety figure concept and Richa Attre Ph.D. from Regeneron Pharmaceuticals, Inc. for assistance with development of the manuscript. Medical writing support under the direction of the authors was provided by Caroline Ridley, PhD, Prime Global (Knutsford, UK) according to Good Publication Practice guidelines (Link) and funded by Regeneron Pharmaceuticals, Inc.

The authors were involved in the study design and collection, analysis, and interpretation of data. All authors had full access to all the data in this study and take complete responsibility for the integrity of the data and the accuracy of the data analysis.

Data-sharing statement

Qualified researchers may request access to study documents that support the methods and findings reported in this manuscript. Individual anonymized patient data will be

638   considered for sharing once the product and indication has been approved by major

639   health authorities (eg, FDA, EMA, PMDA, etc), if there is legal authority to share the

640   data and there is not a reasonable likelihood of patient re-identification. Submit requests

641   to https://vivli.org/.

References

1. Pignolo, R.J., Shore, E.M. & Kaplan, F.S. Fibrodysplasia ossificans progressiva: clinical and genetic aspects. *Orphanet J Rare Dis* **6**, 80 (2011).

2. Baujat, G. et al. Prevalence of fibrodysplasia ossificans progressiva (FOP) in France: an estimate based on a record linkage of two national databases. *Orphanet J Rare Dis* **12**, 123 (2017).

3. Liljesthrom, M., Pignolo, R.J. & Kaplan, F.S. Epidemiology of the Global Fibrodysplasia Ossificans Progressiva (FOP) Community. *Journal of Rare Diseases Research & Treatment* **5**, 31-36 (2020).

4. Morales-Piga, A. et al. Fibrodysplasia ossificans progressiva in Spain: epidemiological, clinical, and genetic aspects. *Bone* **51**, 748-755 (2012).

5. Pignolo, R.J. et al. Prevalence of fibrodysplasia ossificans progressiva (FOP) in the United States: estimate from three treatment centers and a patient organization. *Orphanet Journal of Rare Diseases* **16**, 350 (2021).

6. Pignolo, R.J. et al. Natural history of fibrodysplasia ossificans progressiva: cross-sectional analysis of annotated baseline phenotypes. *Orphanet J Rare Dis* **14**, 98 (2019).

7. Pignolo, R.J. et al. The natural history of flare-ups in fibrodysplasia ossificans progressiva (FOP): a comprehensive global assessment. *J Bone Miner Res* **31**, 650-656 (2016).

8. Peng, K. et al. Longitudinal evaluation of pain, flare-up, and emotional health in fibrodysplasia ossificans progressiva: analyses of the International FOP Registry. *JBMR Plus* **3**, e10181-e10181 (2019).

9.   Kaplan, F.S. et al. Fibrodysplasia ossificans progressiva. *Best practice &*
research. Clinical rheumatology **22**, 191-205 (2008).

10.   Ortiz-Agapito, F. & Colmenares-Bonilla, D. Quality of life of patients with
fibrodysplasia ossificans progressiva. *J Child Orthop* **9**, 489-493 (2015).

11.   Cohen, R.B. et al. The natural history of heterotopic ossification in patients who
have fibrodysplasia ossificans progressiva. A study of forty-four patients. *J Bone*
Joint Surg Am **75**, 215-219 (1993).

12.   Liu, J.X., Hu, R., Sun, Y. & Jiang, H. General anesthesia in fibrodysplasia
ossificans progressive: a case report and clinical review. *Int J Clin Exp Med* **7**,
1474-1479 (2014).

13.   Kaplan, F.S., Al Mukaddam, M. & Pignolo, R.J. A cumulative analogue joint
involvement scale (CAJIS) for fibrodysplasia ossificans progressiva (FOP). *Bone*
101, 123-128 (2017).

14.   Pignolo, R.J. & Kaplan, F.S. Clinical staging of Fibrodysplasia Ossificans
Progressiva (FOP). *Bone* **109**, 111-114 (2018).

15.   Kaplan, F.S. et al. Early mortality and cardiorespiratory failure in patients with
fibrodysplasia ossificans progressiva. *J Bone Joint Surg Am* **92**, 686-691 (2010).

16.   Shore, E.M. et al. A recurrent mutation in the BMP type I receptor ACVR1 causes
inherited and sporadic fibrodysplasia ossificans progressiva. *Nat Genet* **38**, 525-
527 (2006).

17.   Hatsell, S.J. et al. ACVR1R206H receptor mutation causes fibrodysplasia
ossificans progressiva by imparting responsiveness to activin A. *Sci Transl Med*
7, 303ra137 (2015).

18. Upadhyay, J. et al. The expansion of heterotopic bone in fibrodysplasia
ossificans progressiva is Activin A-dependent. *J Bone Miner Res* **32**, 2489-2499
(2017).

19. Alessi Wolken, D.M. et al. The obligatory role of Activin A in the formation of
heterotopic bone in Fibrodysplasia Ossificans Progressiva. *Bone* **109**, 210-217
(2018).

20. Hino, K. et al. Neofunction of ACVR1 in fibrodysplasia ossificans progressiva.
Proc Natl Acad Sci U S A **112**, 15438-15443 (2015).

21. Aykul, S. et al. Activin A forms a non-signaling complex with ACVR1 and type II
Activin/BMP receptors via its finger 2 tip loop. *eLife* **9**, e54582 (2020).

22. Kenny, L.M. et al. Reproducibility of [11C]choline-positron emission tomography
and effect of trastuzumab. *Clinical cancer research : an official journal of the*
American Association for Cancer Research **16**, 4236-4245 (2010).

23. Vanhoutte, F. et al. Pharmacokinetics and Pharmacodynamics of Garetosmab
(Anti-Activin A): Results From a First-in-Human Phase 1 Study. *J Clin Pharmacol*
60, 1424-1431 (2020).

24. Robert, F. et al. Future treatments for hereditary hemorrhagic telangiectasia.
Orphanet J Rare Dis **15**, 4 (2020).

25. Wooderchak-Donahue, W.L. et al. BMP9 mutations cause a vascular-anomaly
syndrome with phenotypic overlap with hereditary hemorrhagic telangiectasia.
Am J Hum Genet **93**, 530-537 (2013).

Figures and Tables

Table 1. Percent of patients with flare-ups by patient diary and by investigator

adverse event (AE) verbatim term in (A) Period_1 and (B) Period_2 (mITT)

Incidence of flare-up events			
A	**Double-blind Period_1 (P1)*** **Inter-group comparison – n (%)**		
Patient diary	**Placebo (n=24)**	**Garetosmab (n=20)**	***P* value**
	17 (70.8)	7 (35.0)	0.03
Investigator AE verbatim term	10 (41.7)	2 (10.0)	0.04
B	**Period_2 (P2) compared to Period_1 (P1)**** **Intra-group comparison – n (%)**		
	Placebo (P1) vs Garetosmab (P2) (n=22)#		***P* value**
Patient diary	15 (68.2)	3 (13.6)	0.0005
Investigator AE verbatim term	10 (45.5)	3 (13.6)	0.019

715 *Active HO analysis set; ** mITT analysis set; #No comparison performed since the intent is to evaluate persistency of

716 garetosmab effect. mITT=modified intent-to-treat. *Of the 24 patients on placebo in Period_1, 22 went on to

717 garetosmab in Period_2. Shown here are only those 22 patients who transitioned from placebo to garetosmab.

 Table 2. Summary of TEAEs during the double-blind period of the study

 (Period_1)*

Patients, n (%)	Placebo (n=24)	Garetosmab 10 mg/kg Q4W (n=20)
≥1 TEAE	24 (100)	20 (100)
≥1 SAE	2 (8.3)	4 (20.0)
≥1 severe TEAE	3 (12.5)	3 (15.0)
≥1 drug-related TEAE	13 (54.2)	13 (65.0)
≥1 TEAE resulting in discontinuation from study	0	1 (5.0)
≥1 TEAE resulting in death	0	0
≥1 TEAE of AESI†	0	1 (5.0)
TEAEs occurring in ≥4 patients in any treatment group		
Headache	7 (29.2)	10 (50.0)
Epistaxis	4 (16.7)	10 (50.0)
Acne	3 (12.5)	6 (30.0)
Pain in extremity	9 (37.5)	5 (25.0)
Arthralgia	9 (37.5)	7 (35.0)
Diarrhea	6 (25.0)	5 (25.0)
Madarosis	0	6 (30)
Back pain	1 (4.2)	4 (20.0)
Neck pain	3 (12.5%)	4 (20.0%)
Toothache	0	4 (20.0)
Neck pain	3 (12.5)	4 (20.0)
Rhinitis	0	4 (20.0)
Dizziness	2 (8.3%)	4 (20.0%)
Nasopharyngitis	5 (20.8)	3 (15.0)
Rash	4 (16.7)	2 (10.0)

AESI=adverse event of special interest; Q4W=every 4 weeks; SAE=serious adverse event;
TEAE=treatment-emergent adverse event. *Safety analysis set. †AESI included epididymitis, orchitis,
hydrocele, scrotum pain, scrotum swelling, moderate to severe episodes of non-traumatic bleeding,

moderate epistaxis (≥30 minutes or requiring professional medical intervention), and severe epistaxis
(based on the definition of a severe TEAE). □Preferred term.

Figure 1. Effect of garetosmab on the (A) change from baseline in TLA compared

with placebo, (B) total number of new HO lesions assessed by quantitative

imaging, and (C) total volume of new HO lesions by CT per patient in Period_1

C.

(A) Time-weighted average of the percent change from baseline in total lesion activity as assessed by

[18]F-NaF PET over 28 weeks (active HO analysis set). **(B)** Total number of new lesions in all patients per

group (combined) by CT and [18]F-NaF PET during Period_1 relative to baseline (active HO analysis set).

(C) Total volume of new lesions per patient as assessed by CT in Period_1 (active HO analysis set). [18]F-

NaF PET=fluorine-18-labelled sodium fluoride positron emission tomography; HO=heterotopic

ossification; IV=intravenous; LS=least squares; Q4W=every 4 weeks; SE=standard error; TLA=total

lesion activity; TWA=time-weighted average.

 Figure 2. Effect of garetosmab in Period_2 relative to Period_1 on the (A, B) total

 number of new HO lesions, (C) mean total volume of new HO lesions (D) TLA in

 new HO lesions, and (E, F) percent of patients with new lesions assessed by

 quantitative imaging in patients originally randomized to placebo

(A) Total number of new lesions by CT during Period_2 relative to Period_1 (mITT analysis set). **(B)** Total

number of new lesions by ^{18}F-NaF PET during Period_2 relative to Period_1 (mITT analysis set). **(C)**

Mean total volume of new lesions as assessed by CT in Period_2 relative to Period_1 (mITT analysis

set). **(D)** TLA of new lesions by 18F-NaF PET in Period_2 relative to Period_1 (mITT analysis set). **(E)**

Percent of patients with new lesions by CT during Period_2 relative to Period_1 (mITT analysis set). **(F)**

Percent of patients with new lesions by [18]F-NaF PET during Period_2 relative to Period_1 (mITT analysis

set).[18]F-NaF PET=fluorine-18-labelled sodium fluoride positron emission tomography; CT=computed

tomography; HO=heterotopic ossification; mITT=modified intent-to-treat; TLA=total lesion activity.

A new flare-up was defined as a flare-up event starting in the corresponding period. Investigator-defined

flare-ups were defined as TEAEs with the verbatim term containing "flare". HO=heterotopic ossification;

mITT=modified intent-to-treat; TEAE=treatment-emergent adverse event.

Figure 3 (A) Relationship between patient baseline age and disease severity as measured by the clinical staging

of FOP (B) Lack of association of epistaxis events and deaths in LUMINA-1 study (C) madarosis events in the

LUMINA-1 study (D) number of epistaxis TEAEs in patients with or without fatal SAE.

A. Age vs Disease severity

*Cumulative Analog Joint Involvement Scale (CAJIS). CAJIS is an assessment of mobility limitation at 15 anatomic locations; scores are tabulated

for each site as normal unaffected (0), affected (1), or completely functionally ankylosed (2). The total score ranges from 0 to 30. On the basis of

FOP features (flare-up activity, body regions affected, thoracic insufficiency syndrome, other complications) and its consequences (impairments in

activities of daily living [ADLs] and ambulation, increasing CAJIS), five clinical stages of disease severity have been defined.[14] In early-stage FOP

the total CAJIS score is usually ≤ 4, in moderate stage 5-18, severe stage 19-24, profound ≥ 24, and end of life stage ≥ 28.

B. Epistaxis events and deaths

770 **C. Madarosis events**

D. Epistaxis events

Number of Epistaxis events	Death (N=5)	No Death (N=39)	Total (N=44)
0	2 (40.0%)	19 (48.7%)	21 (47.7%)
1	2 (40.0%)	1 (2.6%)	3 (6.8%)
2	1 (20.0%)	6 (15.4%)	7 (15.9%)
>=3	0	13 (33.3%)	13 (29.5%)

(A) Red triangles indicate patients with fatal SAEs. **(B)** In red are patients that received placebo during the 28-week Period_1, and in blue patients receiving garetosmab. Dots represent epistaxis events. Black dots are mild epistaxis events, yellow moderate (defined as any episode lasting longer than 30 minutes or requiring medical intervention) and red is severe. Red arrows indicate the patient experienced a fatal SAE. **(C)** In red are patients that received placebo during the 28-week Period_1, and in blue patients receiving garetosmab. Dots represent madarosis events. Black dots are mild madarosis events, yellow moderate. Red arrows indicate the patient experienced a fatal SAE. **(D)** The table shows the number of epistaxis TEAEs in patients with or without fatal SAE. No drug refers to period after dosing was discontinued following Clinical hold. CAJIS=Cumulative Analog Joint Involvement Scale; SAE=severe adverse event; TEAE=treatment-emergent adverse event.

a

b

c