Effectiveness of a chatbot in improving the mental wellbeing of health workers in Malawi during the COVID-19 pandemic: A randomized, controlled trial

Short title: Chatbot use and mental wellbeing of health workers

Eckhard F. Kleinau[1*], Tilinao Lamba[2], Wanda Jaskiewicz[3], Katy Gorentz[3], Ines Hungerbuehler[4], Donya Rahimi[3], Demoubly Kokota[2], Limbika Maliwichi[2], Edister S. Jamu[2], Alex Zumazuma[5], Mariana Negrão[4], Raphael Mota[4], Yasmine Khouri[4], Michael Kapps[4]

[1] Global-Latin America and Caribbean (LAC)-West Africa Region, University Research Co. (URC), Chevy Chase, Maryland, United States of America

[2] Dept. of Psychology, University of Malawi – Chancellor College, Zomba, Malawi

[3] Global Health Division, Chemonics International, Washington, District of Columbia, United States of America

[4] Clinical Division, Vitalk, São Paulo, Brazil

[5] Department of Mental Health, Kamuzu University of Health Sciences (KUHES), Blantyre, Malawi

* Corresponding author

E-mail: ekleinau@urc-chs.com (EK)

Abstract

After the deleterious effects of the COVID-19 pandemic on healthcare worker mental health, we tested

the effectiveness of an interactive chatbot, Vitalk, for improving wellbeing and resilience among

healthcare workers in Malawi, a country with few mental health professionals. We conducted a

randomized, controlled trial (RCT) to investigate our hypothesis that Vitalk is more effective in improving

mental health and resilience outcomes than passive Internet resources.

For our 2-arm, 8-week, parallel RCT (ISRCTN Registry: trial ID ISRCTN16378480), we recruited

participants from 8 professional cadres from public and private healthcare facilities. The treatment arm

used Vitalk; the control arm received links to Internet resources. Of 1,584 participants, 512 completed

baseline and endline assessments. Six assessments provided outcome measures for: anxiety (GAD-7);

depression (PHQ-9); burnout (OLBI); loneliness (ULCA); resilience (RS-14); and resilience-building

activities. We analyzed effectiveness using mixed-effects linear models, effect size estimates, and

reliable change in risk levels.

Results from mixed-model analyses support our hypothesis. Difference-in-differences estimators

showed that Vitalk reduced: depression (-0.68 [95% CI -1.15 to -0.21]); anxiety (-0.44 [95% CI -0.88 to

0.01]); and burnout (-0.58 [95% CI -1.32 to 0.15]). Changes in resilience (1.47 [95% CI 0.05 to 2.88]) and

resilience-building activities (1.22 [95% CI 0.56 to 1.87]) were significantly greater in the treatment

group. We observed no treatment effect on loneliness. Our RCT produced a medium effect size.

This is the first RCT of a mental health app for healthcare workers during the pandemic in Southern

Africa combining multiple mental wellbeing outcomes, and measuring resilience and resilience-building

activities. A significant number of participants could have benefited from mental health support (1 in 8

43   reported anxiety and depression; 3 in 4 suffered burnout; and 1 in 4 had low resilience). Such help is not

44   readily available in Malawi. Vitalk has the potential to fill this gap.

Introduction

Healthcare provision can be stressful even under normal circumstances, and maintaining the mental wellbeing of health workers is of utmost importance for optimal and safe patient care [1]. The COVID-19 pandemic has overwhelmed countries' health systems and increased care-related pressure to ensure patient and staff safety. Health workers confront life and death decisions, physical exhaustion, lack of protective equipment, and fear of infection as daily threats to their mental wellbeing. An increasing number of studies report a high proportion of health professionals globally suffering from depression, anxiety, and burnout. A systematic review of 59 studies found that a median of 24% of health workers suffered from anxiety, 21% from depression, and 37% from distress [2]. A meta-analysis from 36 countries reported slightly higher levels (8.0% for depression; 26.9% for anxiety; 24.1% for post-traumatic stress symptoms; 36.5% for stress; and 50.0% for distress) [3]. Recent studies from sub-Saharan Africa showed similar levels. A multi-center cross-sectional study from Ghana, found the following levels: depression (21.1%), anxiety (27.8%) and stress (8.2%) [4]. Research from Ethiopia found the following prevalence: depression (20.2%), anxiety (21.9%), and psychological distress (15.5%) [5]. Data from South Africa suggest high levels of mental disorders (around 50%) among all types of health workers due to work-related stress [6]. A study from Malawi using a small sample of nurses and the Coronavirus Anxiety Scale suggests that 26% (n=26) of respondents had COVID-19-related anxiety and 48% (n=49) functional impairment [7].

Mental disorders not only pose threats to patient safety and health workers' quality of life, but they also incur high economic costs. A World Health Organization (WHO)-led study, well before the COVID-19 pandemic, found that depression and anxiety disorders cost the global economy US$1 trillion each year [8]. This study estimated a return on investments of between 2.3 and 5.7 to 1 if treatment of these mental disorders was scaled up. Mental health interventions include basic psychosocial counseling for

milder cases, and more intensive psychosocial treatment plus antidepressant drugs for more severe

cases. Over the last decade, computer- or Internet-based cognitive behavioral therapy (c-CBT or i-CBT)

has been tested and implemented as alternatives to in-person treatment. Research has shown that c-

CBT-based self-administered interventions improve depression and anxiety in adults. A meta-analysis of

49 randomized controlled trials (RCT) revealed a significant medium to large effect size (Hedges' g=0.77,

95% CI 0.59 to 0.95) of c-CBT for depression and anxiety [9].

Another meta-analysis of 22 RCTs found an even greater effect size (g=0.88, 95% CI 0.76 to 0.99) [10].

However, a recent systematic review found small to medium post-treatment pooled effect sizes

regarding depressive symptoms (g=0.51, 95% CI 0.30 to 0.72) and anxiety symptoms (g=0.44, 95% CI

0.23 to0.65) of c-CBT for reducing these symptoms in adolescents and young adults compared to passive

controls [11]. Clinical trials have established that mobile applications can effectively deliver CBT

programs for the treatment of: depression [12]; self-management of chronic pain conditions [13]; and

social anxiety disorder [14]. C-CBT or i-CBT can lower barriers to seek help, which is especially important

in settings such as those in sub-Saharan Africa with low levels of access to psychologists and therapists

[15].

C-CBT and i-CBT have evolved into interactive, automated conversational agents (chatbots) that are

driven by artificial intelligence, including the recent entrant Vitalk (with versions available for the public

and adapted for health professionals). A panel study without control group in Brazil established the

effectiveness of the public version, finding a large post-intervention effect size of Cohen's d of -0.81 or

greater for anxiety, depression and stress [16]. The version for health workers has been pilot tested in

Malawi, a country in which access to mental health therapy is very limited, with only 0.02 psychologists,

0.01 psychiatrists and 0.04 occupational therapists per 100,000 population [17]. Only recently has the

country successfully trained 3 psychiatrists. The frequent deployment of mental health professionals,

especially psychiatric nurses, to other duties such as maternity services exacerbates this situation [18].

To date, there have been few studies, and even fewer RCTs, with the aim of establishing the

effectiveness of chatbots. Most of these trials were based on small samples of 70 participants or fewer,

had a short duration of 2-4 weeks, suffered from serious biases, and fell short of establishing that

chatbots lead to improved mental health outcomes. All trials were conducted in high-income countries,

and none in low-resource settings [19]. No RCTs have been conducted to assess the effectiveness of

chatbots in improving the mental health status of health workers specifically [15].

To fill these gaps in evidence, we conducted an RCT in Malawi (building on the earlier pilot test), to

investigate our working hypothesis that a virtual mental healthcare assistant chatbot, Vitalk, is an

acceptable source of psychosocial and mental wellbeing support for health workers to effectively

decrease work-related anxiety, depression, burnout, and loneliness (based on standard mental health

scales), and to increase resilience and resilience-building behaviors.

Methods

Study design

This study, a 2-arm, parallel RCT, with a pre-treatment assessment, 8-week intervention period, mid-

study assessments at 4-5 weeks, and an end-of-study assessment at 9 weeks, was conducted by 2 study

teams with distinct roles: a trial management team and a research team. The management team

reviewed registration data, removed exclusions, assigned groups and unique trial IDs, and provided

communications and technical support to participants. The research team developed the study protocol,

recruited participants, held participant workshops at the beginning and end of the study, held focus

group discussions (FGDs) during the final workshop, and analyzed the data (de-identified before the

analyses).

Participants were randomly assigned to either the treatment or control group after registration. Vitalk,

the developer of the app, created a web portal for participants to register, give consent, and enter their

demographic information. Participants received an email with their unique trial ID and a link to either

the treatment app, Vitalk, or to the website (control group). As a single-blinded design, only the

research team was blinded to the study arm assignments; participants in both arms were told they were

participating in a study on "online self-help for mental wellbeing" but they were aware of the specific

intervention they were using and may have found out that others used a different intervention. Neither

personally identifiable information nor the participants' facility affiliations were shared outside the trial

management team.

123   Study population and recruitment

124   The 8 professional cadres (Table 1) eligible for participation were recruited from all public and private

125   primary, secondary and tertiary care facilities within Blantyre and Lilongwe districts.

126   **Table 1.Type of health workers eligible.**

• Doctors	• Nurses
• Medical Assistants	• Clinical officers
• Laboratory technicians	• Physiotherapy technicians
• Pharmacists	• Physiotherapists

128   We recruited participants via posters placed in participating health facilities. All facilities were contacted

129   by phone and email; in larger facilities the research team made short presentations to leadership to seek

130   permission to recruit from their facilities and hang the posters.

Informed consent, screening and eligibility

Interested health workers were invited to a half-day workshop to provide general information and

answer questions about the study, and in which they could choose to join the study by signing an

informed consent form (kept under lock by the University of Malawi). The research team introduced the

trial as a "mobile-phone-based research study of online self-help for mental wellbeing" but provided no

details about the 2 study groups.

To be eligible, participants needed to meet the following inclusion criteria:

- Be currently employed as 1 of the 8 types of service providers listed above (a minimum

  educational qualification of a diploma adhering to the Ministry of Health employment criteria;

  medical assistants with a certificate and 2 years of college completed qualified)

- Possess some degree of English language proficiency

- Own a smartphone with an Android operating system

- Have no history – past or current – of counseling or therapy for severe mental health disorders

- Not have self-reported suicidal ideation (question 9 of the PHQ-9)

- Score below "very high risk" levels for depression, or below "very high risk" of anxiety combined

  with "high" or "very high risk" of depression on the initial and subsequent assessments

The management team emailed excluded participants Internet resources for self-help. For those

excluded on the basis of suicidal ideation or very high risk of depression and/or anxiety, the team also

sent the contacts of 2 local psychologists.

Study procedures and participant retention

The University Research Co. (URC) Institutional Review Board and the University of Malawi Research

Ethics Committee (UNIMAREC) approved the study protocol.

In Malawi, Internet access and the use of apps through mobile phones requires expensive data

packages. Participants received 3 data packages of 5 gigabytes (GB) valued at $10 each (distributed

during the startup workshop, halfway through the study, and at the final workshop). This data allowance

covered a minimum of at least 1 month of an hour of daily use of Vitalk or web resources At the startup

and final workshops, participants received a transport allowance ($10) and a food allowance ($5),

following government guidelines. These allowances were not tied to active participation in the trial to

avoid any undue influence, and participants received no additional incentives. As the principal means of

participant retention, both study groups received weekly email and WhatsApp messages encouraging

them to engage with their app or website resources. During the baseline period, the management team

provided the treatment group with detailed download and sign-in instructions for the app via

WhatsApp, after noticing low sign-in and receiving messages about sign-in difficulties. The research

team was blinded to this process.

Study timeline

The study spanned 56 days (October 25-December 19, 2021), with startup workshops the week of

October 18 and end-of-study workshops the week of December 20 (Fig 1). The research team held

several workshops each time to accommodate the large number of participants and their varied work

schedules, while complying with physical distancing requirements due to COVID-19. The length of the

study enabled treatment group participants to complete exercises for 2 "carelines" (thematic

conversations addressing anxiety, depression, stress, relationships, or the COVID-19 pandemic) over a

period of a month for each. The baseline, midline, and endline periods take into account that

174   participants completed their first mental health assessments at varying times after registration and

175   repeated assessments with varying frequencies before completing their selected carelines. Several

176   participants completed assessments in a post-endline period.

177   **Fig 1. 2021 Study timeline.**

Sample size and randomization

The required study population size for paired tests of correlated means was calculated using STATA 17

and is based on the following information from published literature:

● Pre-post-treatment difference of 1.5 points on standard test scales

● Standard deviation of the difference = 6

● Power 80%

● Significance level 5%

This resulted in a minimum of 128 participants per study arm. Given the continuity experience (the

proportion of participants completing the pre- and post-treatment assessments for depression, anxiety

and stress) of 20-45% reported by Daley et al. [16], we assumed that the dropout rate could be as high

as 75% for at least 1 of the standard mental health assessments. This required a sample of 512

participants per arm to yield an effective post-intervention sample of 128. Furthermore, if 20% of

potential participants did not meet inclusion criteria or dropped out for other reasons, about 640

enrollees were needed initially per arm.

The management team assigned randomized trial participant numbers (trial ID) using permuted block

randomization sequences, generated by the website Randomization.com [20], to maintain a balance

across the study arms. Block size varied between 4 and 10 randomly ordered treatment assignments. A

random sequence of trial IDs between 1001 and 3000 was generated using the Research Randomizer

[21]. Using randomized trial IDs and permuted block randomization ensured that only the management

team was able to identify participants' group assignment and that the research team remained blinded

throughout trial implementation.

199   Study interventions

200   Treatment group

Participants in the treatment group were asked to download and register with the mental health

chatbot app, Vitalk, and to use it daily over the 8-week study period. The aim of Vitalk is to improve

mental wellbeing by reducing stress, anxiety, depression, and burnout and increasing resilience using a

preventative approach to mental health. Vitalk is an automated chatbot, using an avatar named Viki (Fig

2) to deliver mental health content in an innovative conversational format.

Fig 2. Viki, Vitalk's virtual mental healthcare assistant.

The conversations are based on insights and strategies taken from Cognitive Behavioral Therapy (CBT)

and Positive Psychology [22, 23], which include (but are not limited to): psychoeducation; cognitive

restructuring; behavioral activation; gratitude; and practical exercises (breathing, relaxation, and

meditation). The goal of the conversations is to help users reflect on experiences and learn techniques

to manage stress, mood, and anxiety. A mood tracking tool aids reflection, "emojis" mimic a natural

human interaction, and gamification increases user engagement [24]. Vitalk users choose between

different thematic conversations ("carelines") and use standardized instruments to assess their mental

health and resilience status before and after completing a careline at monthly or more frequent

intervals. The app provides users with feedback and advice about their assessment scores. Carelines are

continuously refined to ensure the language and emotional expression matches the needs of different

users. If user input in the PHQ-9 questionnaire suggests suicidality (e.g., "life isn't worth living") or

suicidal ideation, the risk module is triggered and the chatbot delivers information about emergency

support, including the contact information of local psychologists.

The version used in Malawi consisted of 5 carelines: anxiety; depression; stress; relationships; and the

COVID-19 pandemic. Throughout the 56-day study period, users selected 2 carelines to complete (4

weeks for each). During the first careline period (the first 4 weeks of the study), the user was engaged in

daily conversations, each lasting 5-10 min. This changed to an engagement every other day during the

second careline period (4 weeks). After the study, users were free to continue using the app and

experience the other carelines, receiving a final data bundle at the last workshop to enable them to do

so.

The chatbot avatar Viki was developed through focus groups and pilot testing in Brazil. A team of clinical

psychologists and healthcare professionals from Vitalk wrote the conversations used by Viki and

adapted them to the Malawi context. Unlike that of Brazil, the Malawi version did not include natural

language understanding (NLU). The chatbot was styled and designed by an information technology,

product development and user experience (UX) team at Vitalk in São Paulo. The app is built on ruby and

JavaScript as a free-to-use service, hosted within an instant messenger platform, accessible from

Android devices.

Vitalk does not aim to replace a healthcare professional or to offer treatment. Users are made aware of

this limitation in the terms and conditions required for consent; they are advised to seek additional

support if they show a high risk of depression, anxiety or burnout in their mental health assessments.

Control group (wait-list)

Participants in the control group were provided access to a webpage in their Internet browser with links

to 4 mental health resources: Mental Health Foundation in the United Kingdom[25]; WHO/Geneva

mental well-being resources for the public [26]; WHO/Geneva #HealthyAtHome - Mental health [27];

and the Doing What Matters in Times of Stress: An illustrated Guide [28]. These resources had no

interactive features (passive) and relied on participants' initiative to access them. Control group

participants were considered to be wait-listed since they were given access to the Vitalk app after the

completion of the study.

Outcomes

Primary outcome measures

This study used the scores from several standardized mental health and resilience assessments as outcome measures. In addition, we introduced a new measure of resilience-building activities. In the analysis, we used total assessment scores as well as risk groupings to establish clinical relevance following classifications from published literature (based on the sensitivity and specificity of the assessments).

Patient Health Questionnaire (PHQ-9)

The PHQ-9 is a 9-item self-reported scale that evaluates symptoms of depression over the past 2 weeks (e.g., "how often have you been bothered by feeling down, depressed, or hopeless"). Item response options use a Likert scale ranging from 0 (not at all) to 3 (nearly every day), with a maximum score of 27. According to the literature, total scores are divided into 5 risk categories: none (0–4); mild (5–9); moderate (10–14); moderately severe (15–19); and severe (20+) symptoms. The PHQ-9 has been widely used and validated in Malawi [29].

Oldenburg Burnout Inventory (OLBI)

Burnout is linked to relatively high work requirements and limited resource availability for managing them. The discrepancy between resources and challenges creates a significant negative emotional state. The OLBI has 16 items rated 1 to 4, with a maximum score of 64. Eight items describe exhaustion (OLBI-E) and 8 describe disengagement (OLBI-D) [30]. The questionnaire includes both straight and reversely worded items in both dimensions. According to the literature, total scores are divided into 3 risk categories: low (16-35); moderate (36-43); and high (44+). OLBI has been validated and used in low- and middle-income countries (LMICs) [31].

Generalized Anxiety Disorder (GAD-7)

The GAD-7 is a 7-item self-reported scale to assess anxiety symptoms over the past 2 weeks (e.g., "how

often have you been bothered by feeling afraid something awful might happen"). Scores range from 0

(not at all) to 3 (nearly every day), with a maximum score of 21. The total scores are divided into 4 risk

categories: none (0–4); mild (5–9); moderate (10–14); and severe (15+) symptoms. GAD-7 has been

utilized effectively in Malawi and other LMICs with comparable demographics [32].

UCLA Short (three-item) Loneliness Scale (UCLA Loneliness)

UCLA Loneliness was added to Vitalk for this study. While there is no agreed upon definition of

"loneliness" in research, it is described as an unwelcome, painful and unpleasant feeling, and a fluid

experience that can come and go over a short time or persist in the longer term. The UCLA Loneliness

scale consists of 3 items rated 1 to 3, with a maximum score of 9. Total scores are divided into 2 risk

categories: non-lonely (3-5) and lonely (6+) [33, 34].

5-item Resilience-Building Behavior Scale

A novel addition to this study and Vitalk was an assessment of participants' resilience-building behaviors

related to stress management, self-awareness, self-care, purpose, and connection with others (adapted

from a toolkit for building health worker resilience [35]). While numerous scales for measuring resilience

exist, we did not find any publications on the validity and reliability of instruments measuring resilience-

building activities. The questionnaire had 5 items rated 1 to 4, with a maximum score of 20, related to

the frequency with which participants performed resilience-building activities over the past 2 weeks.

Scores were divided into 3 activity levels: low or none (5-10); moderate (11-15); and high (16-20).

14-item Resilience Scale (RS–14)

Resilience refers to the ability to withstand or adaptively recover from stressors, promoting

psychological and physical wellbeing. Resilience is negatively correlated with symptoms of generalized

anxiety and post-traumatic stress and positively correlated with gratitude, optimism, and positive affect.

The 5 characteristics of resilience are: meaningful and purposeful life; perseverance; equanimity; self-

reliance; and existential aloneness [36]. The RS-14 has 14 items rated 1 to 7, with a maximum score of

98. Total scores are divided into 6 resilience levels: very low (14–56); low (57–64); on the low end (65–

73); moderate (74–81); moderately high (82–90); and high (91+). RS-14 has been validated and used in

LMICs [37, 38]

While the app registers the actual scores, to simplify communication of risk to users, Vitalk reports 4

levels of mental health risks and resilience for its 4 standard assessments (GAD-7, PHQ-9, OLBI, and RS-

14): very low; low; high; and very high risk. The Vitalk risk classification differs from published levels as

follows: 1 additional level for OLBI; 4 instead of 5 levels for PHQ-9; and 4 instead of 6 levels for RS-14.

For the purposes of this trial, we used only published risk classifications based on actual test scores.

All 6 assessments together required about 20 minutes to complete. Participants in both arms were

allowed to take the assessments repeatedly and saw their results immediately after answering the 6

questionnaires. Participants were prompted either by the avatar Viki or What's App messages and

emails to take the assessments at specific intervals during the baseline, midline, and endline periods.

Secondary measures

Mood meter

Users of the Vitalk app had the option of assessing their mood using emojis representing 5 mood levels

(0-4): very bad; bad; okay; good; and very good. Emojis are successfully used in sentiment analysis and

show a stronger emotion compared to words [24]. The timing of mood assessments was matched to

mental health assessments prior to data analysis since participants took them independently. The mood

meter was part of the intervention to raise self-awareness, and only available to the treatment group.

Participant engagement

For the treatment group, we captured several measures related to participant engagement over the

study period: number of interactions with Vitalk; number of days with interactions with the app; number

of days between the first and last day of interaction with the app; and total number of hours spent on

the app. For the control group, we captured 2 measures related to participant engagement: number of

days they accessed web resources and number of web resource click-throughs. All raw engagement

measures were converted to z-scores for both groups to account for differences in metrics and

distributions in each group. Standardized hours spent on Vitalk, and standardized number of days with

web resources accessed, were retained as determinants of mental health and resilience scores.

Smartphone experience

During registration, we measured participants' smartphone experience by asking how often (never, less

often, weekly, several times a week, daily) they used the following features (each scored 0 to 4):

Internet; WhatsApp; Facebook, Instagram or twitter; chat app; mobile money; games; email; videos; and

music. Smartphone experience is a composite of the frequency of use for all 9 features, with a maximum

score of 36.

Current or past exposure to mental health counseling

During registration, participants were asked whether they had any current or past exposure to mental

health counseling. Possible responses were: currently receiving counseling; received counseling in the

last 6 months; received counseling in the last 2 years; received counseling more than 2 years ago; and

never received counseling. Participants with any counseling experience were then asked whether

counseling was for mild or moderate mental health issues, or for acute or severe mental health issues.

Participants who responded the latter were excluded from the study and given the contact for local

psychologists.

Current or past chat app and website use for mental health

At registration, participants were asked about their current use of a chat app for mental wellbeing (with

the response options: never; sometimes; and often). Past use was assessed through a binary yes or no

response. We created a composite "chat app for mental health" variable with values ranging from 0 (no

chat app use) to 5 (current often use and past use) by combining responses from both questions.

Current and prior use of Internet-based resources about mental wellbeing was assessed in the same

manner.

Impact of COVID-19

Four questions assessed how the COVID-19 pandemic impacted study participants. The first 3 questions

asked about the impact on workload, workhours, and stress levels over the past 12 months. Possible

response options for each question (scored from 0 to 4) were: greatly decreased; somewhat decreased;

stayed the same; somewhat increased; and greatly increased. The sum of the responses to these

questions created a composite COVID-19 effect variable. The fourth question inquired whether COVID-

19 prevented participants from going into work over the past 12 months. Response options (scored from

0 to 3) were: no disruptions at all; several days over the entire year; several days every month; several

days every week; and I did not work at all (coded as missing).

Anonymous online participant experience questionnaire

During the final workshops, participants from both arms had the opportunity to complete an

anonymous online questionnaire about their experience using either the Vitalk app or the web

resources. Questions focused on the frequency of use, reasons for infrequent use, relevance and

usefulness, ease of use, and specific difficulties encountered. Some key findings related to study

outcomes are highlighted in this paper; a separate comprehensive analysis will be published together

with the findings from focus group discussions.

358   Data analysis

Participant characteristics are presented as proportions or means (SD) for the treatment and control

groups.

Mixed linear model. We used multilevel mixed-effects generalized linear models (STATA *meglm*

command) to analyze primary participant-reported outcome measures for all 6 mental health and

resilience assessments over the baseline and endline periods. A full factorial model of study group and

assessment period was fitted to represent fixed effects. Participants' trial ID specified the random effect

in this model. The parameter estimate of interest was the coefficient of the interaction between study

group and assessment period, which is equivalent to a difference-in-differences (DiD) estimator. DiD

estimators were robust, because only valid assessments for the baseline and endline periods were

included in the analysis, avoiding the need for inter- or extrapolating missing values. Because of

randomization, covariates were not included in the mixed models. Data from the midline and any

assessments completed after the 8-week study period (post-endline) were not included in the analysis

because of a much lower number of observations in these compared to the other periods.

Effect size. To compare the different outcome measures, Cohen's d was calculated as the mean

difference between the 2 groups divided by their common SD and interpreted as small (d=0.2), medium

(d=0.5), and large (d=0.8).

Reliable change index. To assess whether the change in outcomes (the total score for each assessment,

between baseline and endline) is statistically reliable, and not due to measurement variability inherent

to the assessments alone, we calculated a reliable change index (RCI). The RCI is the standard error of

the difference between the 2 measurements, SE_{diff}, in equation (1). SD_{bas} is the standard deviation of the

baseline observations, and r is Cronbach's alpha representing the reliability of the measurement

instrument. Each assessment instrument has its own RCI.

$$RCI = 1.96 \times SE_{diff} = 1.96 \times SD_{bas} \sqrt{2} \sqrt{1-r} \qquad (1)$$

Change in risk levels. We used total scores from the 4 mental health and 2 resilience assessments for the

mixed linear model, effect size, and RCI estimation. To determine whether the changes between the

pre- and post-intervention periods matter from a clinical perspective, the total assessment scores – a

continuous variable – were converted to mental health risk and resilience levels –an ordinal variable –

using the published classifications described above for each type of assessment (e.g., depression scores

from the PHQ-9 are divided into 5 levels ranging from none to severe). A change in risk levels was

calculated as either a drop or increase of 1 or more levels for each mental health and resilience metric.

For participants registering a change in risk level, the proportion with reliable change was calculated.

Reliable change was established if the difference between baseline and endline scores were equal or

greater than the RCI.

Determinants of mental health and resilience. All mental health and resilience outcomes were regressed

against all participant characteristics using ordinary least squares models, controlling for study group

and assessment period. All observations with valid baseline and endline assessments for the control and

treatment groups were included in the analysis. A separate analysis was done for the treatment group

to assess which covariates unique to this group (mood meter scores and carelines) affected the

outcomes.

P ≤ 0.10 was considered to be indicative of statistical significance. Equally important as the p-value was

the consistency of patterns across all mental health and resilience assessments and their agreement

with behavioral theory. All data were analyzed with STATA 17.0 (StataCorp LLC, College Station, TX).

402 Results

The Consolidated Standards of Reporting Trials (CONSORT) diagram in Fig 3 shows that of 1,584

participants enrolled in the study, 836 completed one or more mental health and resilience assessments

(481 in the control and 355 in the treatment group). Based on participant feedback, the treatment group

saw a much greater attrition due to technical difficulties impacting the use of the Vitalk app. The

number of participants completing assessments varied substantially by study period, as seen in Table 2.

Fig 3. Study participant flow.

Table 2. Number of participants completing any mental health and resilience assessment by study
period.

Study period	Control Group	Treatment Group	Total
Baseline	469	318	787
Midline	77	181	258
Endline	304	260	564
Post-endline	13	86	99

Numbers include singular assessments that were completed in one study period only, which
were excluded from analysis.

For the DiD, effect size, and risk level analyses, Table 3 shows that the sample size differed by type of

assessment. Overall, 511 participants (296 in the control and 215 in the treatment group) had no

protocol violations and completed baseline and endline assessments for GAD-7. Only participant who

completed base- and midline assessments were included in the calculations of DiD estimators using

mixed-effects linear modeling, effect size estimation, and change in risk levels. Samples in the treatment

group for the loneliness and resilience-building activities assessments were markedly smaller than in the

control group. These 2 assessments were the last in the series of the 6 assessments suggesting that a

majority of Vitalk users stopped completing assessments after RS-14. The control group did not

experience a similar drop, suggesting that technical issues may have prevented app users from

completing all assessments.

Table 3. Effective sample size by type of mental health and resilience assessment with completed
baseline and endline.

Assessment	Control Group n	Treatment Group n	Total n
Anxiety (GAD-7)	296	215	511
Depression (PHQ-9)	286	201	487
Burnout (OLBI)	291	211	502
Resilience (RS-14)	280	183	463
Loneliness (UCLA)	280	70	350
Resilience building activities	278	85	363

427   Assessments shown in the order completed by participants.

428   These are the sample sizes used in DiD, effect size, and change in risk level analyses.

Participant characteristics

The distribution of characteristics for the 836 participants with valid mental health and resilience assessments was balanced between the control and treatment groups as shown in Fig 4. The difference between groups varied by 1 to 6 percentage points. Overall, participants were young (under 30 years) and female. Most were professional nurses working in public secondary or tertiary healthcare facilities in urban areas. Maternity care was the most common type of care provided. Fewer than 30% had ever used a chat app for mental health and more than 40% had ever accessed websites with mental health content. One in 6 participants had a history of mental health counseling. The level of experience in performing 9 different tasks with their smartphones was high (a mean of 29 out of a maximum score of 36) and nearly the same in both study groups (Fig 5), corresponding to several times per week to daily use. Both groups reported the same effect of the COVID-19 pandemic. More than 1 in 3 participants mentioned that the pandemic had disrupted their work. The average effect of COVID-19 on participants' workload, workhours, and stress levels was 8 out of a maximum score of 12 (Fig 5), corresponding to somewhat increased workload and stress levels.

Fig 4. Study participant characteristics. Only the most frequent categories are shown (n = 836 [481 control group, 355 treatment group] except Age)

Fig 5. Mean smartphone experience and COVID-19 impact scores (n = 836).

Participant adherence to treatment and control interventions

Treatment group participants had an average of 2,200 (SD 6,294, median 920, range 84-58,021) interactions with the Vitalk app over the 56-day study period, with a distribution skewed towards the lower end. The average duration of app use was 24 days (SD 14, median 23, range 1-67), and users spent an average of 10.4 hours (SD 7.7, median 9.2, range 0.4-51.7) on the app. Treatment group participants expressed their mood an average of 13 times (SD 15, median 8, range 0-81) over the study period. The mean mood score was 2.9 (SD 1.0, median 3.0, range 0-4).

Control group participants accessed the website resources an average of 1.9 days (SD 2.0, median 1, range 1-23) over the study period, with a distribution heavily skewed towards 1.0 days. The only other measure of adherence available in this group was an average number of 1.2 click-throughs (SD 0.7, median 1, range 1-8) of links to mental health resources on the website. We did not measure time spent on resource use and mood status for the control group.

Summary of outcomes at baseline and endline

Table 4 shows summary statistics for the 4 mental health and 2 resilience assessments at baseline and endline. Mean anxiety, depression, and loneliness scores from the GAD-7 (baseline mean 4.5 [SD 3.8], endline mean 2.9 [SD 3.5]), PHQ-9 (baseline mean 4.0 [SD 4.1], endline mean 2.9 [SD 3.8]) and UCLA loneliness (baseline mean 5.3 [SD 1.5], endline mean 4.7 [SD 1.6]) questionnaires, respectively, fell into the lowest risk level. Burnout scores were at moderate risk, with a mean of 38.4 (SD 6.0) and 36.6 (SD 6.7) at baseline and endline, respectively. Resilience was moderate to moderately high, with a mean of 78.6 (SD 10.4) at baseline and a mean of 81.5 (SD 11.4) at endline, and resilience-building activities were moderate, with a mean of 13.1 (SD 3.0) and a mean of 14.2 (SD 3.3) at endline. Both control and treatment groups saw an improvement in mental health and resilience scores between the 2 assessment periods. All summary statistics are based on participants with valid baseline and endline

469   data using all assessments completed during each assessment period. The average number of

470   assessments taken each period varied between 1 and 5, depending on the type of assessment and was

471   generally higher at endline.

Table 4. Mental health and resilience scores from completed assessments by study period and group.

Questionnaire	Control Group		Treatment Group		Total	
	Baseline	Endline	Baseline	Endline	Baseline	Endline
PHQ-9						
Frequency	510	634	429	496	939	1130
Mean score ***	3.67	2.99	4.29	2.72	3.95	2.87
Standard deviation	4.01	4.04	4.26	3.52	4.13	3.82
Median score	3	1	3	1.5	3	1
Avg. no. of assessments	2.9	3.3	3.2	4.6	3.0	3.9
OLBI						
Frequency	471	583	379	441	850	1024
Mean score ***	38.00	36.55	38.84	36.63	38.37	36.59
Standard deviation	6.18	6.83	5.74	6.41	6.00	6.65
Median score	39	37	39	37	39	37
Avg. no. of assessments	2.6	3.0	2.8	4.3	2.7	3.6
GAD-7						
Frequency	545	669	468	550	1013	1219
Mean score ***	4.24	2.89	4.73	2.93	4.46	2.91
Standard deviation	3.74	3.63	3.86	3.31	3.80	3.49
Median score	4	1	4	2	4	2
Avg. no. of assessments	3.1	3.6	3.3	4.9	3.2	4.2
UCLA Loneliness						
Frequency	433	531	84	106	517	637
Mean score ***	5.18	4.69	5.70	4.92	5.26	4.73
Standard deviation	1.52	1.59	1.45	1.74	1.52	1.61
Median score	5	4	6	5	6	4
Avg. no. of assessments	2.4	2.7	1.3	2.0	2.2	2.6
Resilience Building						
Frequency	422	521	128	117	550	638
Mean score ***	13.20	14.01	12.80	14.84	13.11	14.16
Standard deviation	3.11	3.40	2.62	2.65	3.00	3.29
Median score	13	14	13	15	13	14
Avg. no. of assessments	2.3	2.7	1.8	1.6	2.2	2.5
RS-14						
Frequency	438	540	327	394	765	934
Mean score ***	79.23	81.05	77.69	82.19	78.57	81.53
Standard deviation	10.02	11.36	10.81	10.57	10.39	11.04
Median score	80	83	79	84	80	83
Avg. no. of assessments	2.4	2.8	2.6	4.3	2.5	3.4

Mean score ***: A blue frame indicates statistically significant differences between mean scores at baseline and endline based on within group t-tests. P-values are < 0.01 for all assessments in the control and treatment groups.

Mean score ***: Shaded cells indicate statistically significant differences between mean scores comparing study periods for each study group based on between group t-tests. P-values are < 0.05 for differences at baseline for all assessments and not significant at endline except for Resilience Building where baseline scores did not differ significantly between groups but endline results did.

Using published reference levels and the average score from each assessment for each study period,

participants were classified according to "moderate to high" mental health risk and "very low to low"

resilience levels. As seen in Table 5, fewer participants in the control and treatment groups were

classified as "moderate to high" risk for mental wellbeing and "very low to low" resilience levels at

endline than at baseline. However, deceases in depression, burnout, and anxiety, and increases in

resilience- building activities and resilience were greater in the treatment than control group, consistent

with the changes in mean assessment scores shown in Table 4. Loneliness did not follow this pattern,

with a larger percentage of participants improving in the control than in the treatment group.

Table 5. Study participants with moderate to high risk levels for mental health or low levels of resilience.

| Assessment and risk level | Control Group | | | | | | Treatment Group | | | | | | Sample at base- and endline | | |
| | Baseline | | Endline | | Total | | Baseline | | Endline | | Total | | Control Group N | Treatment Group N | Total N |
	n	%	n	%	n	%	n	%	n	%	n	%			
Depression: moderate, moderately severe, severe	32	11%	25	9%	57	10%	30	14%	18	9%	48	11%	291	211	502
Burnout: moderate or high	211	74%	187	65%	398	70%	156	78%	121	60%	277	69%	286	201	487
Anxiety: moderate or severe	37	13%	24	8%	61	10%	29	13%	13	6%	42	10%	296	215	511
Loneliness: lonely	148	53%	105	38%	253	45%	44	63%	36	51%	80	57%	280	70	350
Resilience building: low or no activity	49	18%	44	16%	93	17%	14	16%	2	2%	16	9%	278	85	363
Resilience: very low, low, or on the low end	71	25%	54	19%	125	22%	56	31%	38	21%	94	26%	280	183	463

491   Risk levels are based on the average assessment score per participant for each study period.

Difference-in-differences (DiD) estimators

The DiD estimates shown in Fig 6 suggested a significant positive effect of Vitalk in reducing anxiety and depression, and in building resilience and increasing resilience-building activities. The coefficients for the interaction terms between study group and study period showed that Vitalk was more effective than the passive control intervention in reducing depression (DiD estimator -0.68 [95% CI -1.15 to -0.21]), anxiety (DiD estimator -0.44 [95% CI -0.88 to 0.01]), and burnout (DiD estimator -0.58 [95% CI -1.32 to 0.15]). Burnout was just short of statistical significance at the 90% level. Changes in resilience were positive and significantly greater in the treatment group (DiD estimator 1.47 [95% CI 0.05 to 2.88]). Similarly, this group showed significantly greater resilience-building activities (DiD estimator 1.22 [95% CI 0.56 to 1.87]). There seemed to exist no treatment effect on loneliness, which was omitted from the estimation of the RCI and risk level changes.

Fig 6. Difference-in-differences estimators with 95% confidence intervals from mixed effects linear models.

Effect size

Although the DiD estimates indicated that Vitalk use positively impacts mental health and resilience outcomes, they did not inform about the size of the effect since each outcome assessment uses a different scale. The effect size for each outcome was measured using standardized mean differences (SMD, Cohen's d). We also calculated Hedges' g – another SMD measure – to account for unequal sample sizes but do not show them separately since the values for 2 decimals were identical to Cohen's d. Fig 7 shows the results comparing the effect size for the control and treatment groups. Participants in the treatment group experienced a consistently larger effect for the mental health and resilience assessments (a medium effect), with the control group trending towards a small effect. Vitalk use had the largest impact on resilience-building activities with a large effect size (more than 3 times the effect in the control group). Effect size is calculated from statistical inference and therefore has confidence intervals (CI) shown in Fig 7. None of the 95% CI contain zero, indicating that both study groups experienced a positive effect – a decrease in depression, burnout, anxiety, and loneliness, and an increase in resilience-building activities and resilience. Depression and resilience-building activities showed the largest difference in effect size between the control and treatment groups, with no overlapping 95% CI.

522   **Fig 7. Effect size post-intervention by assessment type and study group.**

Reliable change index (RCI)

A comparison of the mean and median scores for mental health and resilience outcomes pre- and post-

intervention in Table 4 and Table 5 suggests that participants saw improved mental wellbeing and

resilience outcomes. However, this is the net result of a larger number of participants improving in

outcomes than participants deteriorating in outcomes. To assess whether this change may have arisen

by chance alone, we calculated the RCI and the proportion of participants with a change equal or greater

than the RCI. The results are shown in Fig 8A and Fig 8B. The RCI for each outcome was similar in the

control and treatment groups. Reliable improvement was greater in the treatment than in the control

group for depression and burnout, and was about the same for anxiety. The proportion that saw a

reliable increase in resilience-building activities was more than twice in the treatment than that of the

control group. Reliable change in resilience was the same in both groups. While overall, fewer

participants experienced a deterioration in mental health and resilience, reliable change was

consistently greater in the control than in the treatment group. Fig 8A also shows Cronbach's alpha as a

reliability measure for each assessment, which was used in the calculation of the RCI. Cronbach's alpha

is very good (greater and 0.80) for 4 assessment instruments and acceptable for resilience-building

activities.

Fig 8. Reliable change in mental wellbeing and resilience outcomes. [A] Reliable change index (RCI). [B]

Reliable improvements or deterioration in outcomes between baseline and endline.

Mental health and resilience risk level change

While any improvement in mean scores for mental health and resilience assessments between baseline

and endline is a positive outcome, a difference in means, although significant, did not impart whether

these were of clinical relevance. To assess whether a change in scores equated to a change in risk levels,

we categorized each participant's scores into risk levels using published cutoff points. We then

estimated the proportion of participants changing risk categories by either moving to a lower or higher

level (Fig 9). The improvement in risk levels was consistently greater in the treatment than in the control

group. The effect on anxiety was greater than on depression or burnout. The depression scale showed

the least amount of change, with about two-thirds of participants staying at the same risk level pre- and

post-intervention. The proportion of participants increasing resilience-building activities by at least 1

level was almost twice as high in the treatment group compared to the control group. The control group

consistently saw more cases with a deteriorated outcome than the treatment group. Not all the changes

of mental health risk or resilience levels were statistically significant. Therefore, Fig 9 also shows the

proportion of all those changing risk levels that were reliable based on the RCI. For burnout, the

proportion of reliable change was considerably higher in the treatment group (58%) than in the control

group (36%). For anxiety, 71% in the control group saw reliable change compared to 61% in the

treatment group. Participants in the control group whose mental wellbeing and resilience outcomes

deteriorated between study periods experienced a consistently greater reliable change than those in the

treatment group. The proportion of participants with reliable change in Fig 9 is equivalent to the

percentage of participants with reliable improvement or deterioration in Fig 8B.

Fig 9. Percent of participants with improved, worsened or same mental wellbeing and resilience risk

levels. White color text percent: percent without risk level change or with one or more risk levels

difference between baseline and endline. Black color text percent: percent of risk level change that is

reliable based on the RCI.

Determinants of mental health and resilience

Ordinary least squares models regressed participant characteristics on mental health and resilience

outcomes using data from all participants with valid mental health and resilience assessments,

controlling for study group and study period. Table 6 shows that several factors have a significant effect

on mental health outcomes. Anxiety increased by 0.02 points (95% CI 0.00 to 0.05) with each year of

age. Likewise, depression increased by 0.03 points (95% CI 0.01 to 0.06) and loneliness by 0.02 points

(95% CI 0.00 to 0.04). Burnout appeared to decrease with age by -0.04 points (95% CI -0.09 to -0.01).

Resilience-building activities increased with age by 0.04 points (95% CI 0.01 to 0.08). Participant gender

was negatively associated with anxiety and burnout, with women scoring higher than men by 0.43

points (95% CI 0.07 to 0.79) and 0.90 points (95% CI 0.19 to 1.61), respectively. In addition, women

showed less resilience-building activities of -0.40 points (95% CI -0.85 to 0.04) than men. While this was

consistent with women having -0.56-point lower resilience levels, this was not statistically significant.

Age and gender were also closely correlated with smartphone use experience. With each year increment

of age, smartphone experience decreased by -0.17 points [95% CI -0.17 to -0.15]; and females saw -0.99

point [95% CI -1.15 to -0.82] lower smartphone experience than men.

Table 6. Determinants of mental health and resilience outcomes for the control and treatment groups combined.

	Dependent variables (assessment scores)					
	Depression (PHQ-9)	Burnout (OLBI)	Anxiety (GAD-7)	UCLA LONELINESS	Resilience building activities	Resilience (RS-14)
Independent variables	Coefficient (SE)	Coefficient (SE)	Coefficient (SE)	Coefficient (SE)	Coefficient (SE)	Coefficient (SE)
z-score of time spent @	-0.43*** (0.10)	-0.44** (0.18)	-0.51*** (0.09)	-0.22*** (0.06)	0.09 (0.12)	-0.04 (0.33)
Cadre: clinical officer †	0.42 (0.36)	1.30** (0.63)	0.44 (0.32)	-0.17 (0.20)	-0.40 (0.40)	-2.54** (1.12)
Cadre: doctor †	-0.14 (1.01)	0.65 (1.74)	-0.42 (0.95)	1.37** (0.56)	-2.74** (1.17)	-0.94 (3.06)
Cadre: medical assistant †	-1.34*** (0.45)	1.62** (0.81)	-1.08*** (0.41)	-0.87*** (0.25)	0.57 (0.52)	-3.73*** (1.44)
Cadre: pharmacist †	0.87 (0.89)	2.45 (1.57)	1.14 (0.81)	0.43 (0.43)	0.78 (0.93)	-7.43*** (2.76)
Cadre: technician †	0.82 (0.85)	-1.15 (1.51)	0.93 (0.74)	0.36 (0.55)	-0.16 (1.12)	-1.77 (2.69)
Age (years)	0.03*** (0.01)	-0.04* (0.02)	0.02** (0.01)	0.02** (0.01)	0.04** (0.02)	-0.02 (0.04)
Sex ‡	0.17 (0.20)	0.90** (0.36)	0.43** (0.18)	0.11 (0.11)	-0.40* (0.23)	-0.56 (0.64)
Care level: secondary $	0.72** (0.33)	1.46** (0.59)	0.71** (0.30)	-0.06 (0.18)	0.37 (0.35)	-0.30 (1.05)
Care level: tertiary $	1.03*** (0.29)	2.01*** (0.52)	0.99*** (0.26)	0.05 (0.17)	0.47 (0.34)	-1.21 (0.93)
Facility type: private nfp !!	-0.62* (0.33)	-0.10 (0.59)	-0.60** (0.30)	0.25 (0.17)	-0.24 (0.34)	-2.08** (1.04)
Facility type: private fp !!	0.78* (0.44)	2.06*** (0.78)	0.48 (0.39)	0.96*** (0.25)	-0.63 (0.49)	-2.48* (1.37)
Location: peri-urban T̄	0.50 (0.34)	-0.18 (0.61)	0.48 (0.30)	0.01 (0.21)	0.04 (0.41)	-0.68 (1.08)
Location: rural T̄	0.52* (0.27)	-0.20 (0.49)	0.30 (0.25)	-0.08 (0.15)	0.04 (0.31)	1.50* (0.88)
Service: counseling §	-1.61*** (0.53)	1.15 (0.96)	-1.01** (0.48)	0.17 (0.33)	-1.18* (0.64)	0.33 (1.73)
Service: inpatient care §	-1.20*** (0.41)	-0.04 (0.74)	-1.04*** (0.37)	-0.47** (0.23)	0.24 (0.47)	-2.44* (1.32)
Service: intensive care §	-1.40*** (0.45)	0.67 (0.81)	-0.88** (0.40)	-0.05 (0.26)	-0.21 (0.52)	-0.91 (1.45)
Service: maternity care §	-0.93** (0.37)	1.46** (0.66)	-0.56* (0.33)	-0.42** (0.20)	0.10 (0.42)	-1.71 (1.18)
Service: support & diagnostics §	-1.69** (0.84)	-0.73 (1.52)	-1.36* (0.74)	-0.89* (0.53)	-0.29 (1.10)	3.69 (2.72)
Service: specialty care §	-2.05*** (0.45)	0.74 (0.81)	-1.59*** (0.41)	-0.76*** (0.25)	-0.53 (0.52)	0.24 (1.43)
Service: surgery care §	-1.55*** (0.48)	0.87 (0.87)	-1.49*** (0.44)	-1.00*** (0.27)	-0.90* (0.54)	-0.99 (1.55)
Smartphone experience score	0.06*** (0.02)	0.05* (0.03)	0.08*** (0.01)	0.03*** (0.01)	0.04** (0.02)	-0.09 (0.05)
History of MH counseling	0.63*** (0.11)	0.35* (0.20)	0.36*** (0.10)	-0.03 (0.07)	-0.07 (0.16)	0.27 (0.37)
COVID-19 impact score	0.02 (0.03)	0.20*** (0.06)	0.06** (0.03)	0.01 (0.02)	-0.13*** (0.04)	0.21** (0.10)
COVID-19 ability to work impact	0.62*** (0.10)	0.95*** (0.17)	0.39*** (0.09)	0.17*** (0.06)	-0.24** (0.12)	-0.11 (0.31)
History of chat app use for MH	-0.02 (0.08)	-0.24* (0.14)	-0.07 (0.07)	-0.06 (0.05)	0.28*** (0.09)	0.19 (0.26)

	Dependent variables (assessment scores)					
	Depression (PHQ-9)	Burnout (OLBI)	Anxiety (GAD-7)	UCLA LONELINESS	Resilience building activities	Resilience (RS-14)
Independent variables	Coefficient (SE)	Coefficient (SE)	Coefficient (SE)	Coefficient (SE)	Coefficient (SE)	Coefficient (SE)
History of website use for MH	0.20*** (0.08)	-0.07 (0.14)	0.17** (0.07)	0.05 (0.04)	0.07 (0.09)	0.30 (0.25)
Treatment group ¶	0.42** (0.18)	0.30 (0.32)	0.47*** (0.16)	0.52*** (0.13)	0.09 (0.24)	-0.45 (0.58)
Study period mid, end, post ≡	-0.60*** (0.09)	-0.95*** (0.15)	-0.80*** (0.08)	-0.27*** (0.05)	0.59*** (0.10)	1.63*** (0.27)

P-values: * <= 0.10, ** <= 0.05, *** <= 0.01

z-score of time spent @: represents standardized hours spent on Vitalk for the treatment group, and standardized number of days with
web resources accessed for the control group

Reference categories: † Cadre: nurse, ‡ Male, $ Care level: primary, ‼ Facility type: public, ⊤ Location: urban, § Service: outpatient care,
¶ Control group, ≡ Baseline

Abbreviations: nfp = not-for-profit, fp = for profit, MH = mental health, mid = Midline, end = Endline, post = Post-endline period

Secondary care providers had worse outcomes (higher scores) for anxiety, depression, and burnout

compared to primary care providers, ranging from 0.71 points (95% CI 0.12 to 1.30) for anxiety to 1.46

points (95% CI 0.30 to 2.62) for burnout. Tertiary care providers scored 2.01 points (95% CI 1.00 to 3.03)

higher for burnout than primary care providers. Outpatient care saw consistently higher anxiety,

depression, and loneliness than any of the other types of care, with surgery (depression -1.54 points,

95% CI -2.50 to -0.60) and specialty care (depression -2.06 points, 95% CI -2.93 to -1.18) having the

largest difference. Maternity care was the only type associated with a significantly 1.46-point (95% CI

0.16 to 2.76) higher burnout than outpatient care. Medical assistants scored significantly lower for

anxiety, depression, and loneliness compared to nurses, but they had 1.62-point (95% CI 0.03 to 3.20)

higher burnout, and -3.73-point (95% CI -6.56 to -0.90) lower resilience than the latter.

Anxiety, depression, burnout, and loneliness outcomes all decreased with the z-scores of time spent

with the treatment or control interventions, ranging from -0.22 points (95% CI -0.33 to -0.10) for

loneliness, to -0.51 points (95% CI -0.68 to -0.34) for anxiety. Conversely, greater use of smartphone

features was associated with significantly higher levels of anxiety, depression, burnout, and loneliness,

ranging from 0.03 points (95% CI 0.01 to 0.06) for loneliness to 0.08 points (95% CI 0.05 to 0.11) for

anxiety. Participants with greater smartphone experience had higher resilience-building activities by

0.043 points (95% CI 0.001 to 0.084). While smartphone use showed a -0.09-point lower level of

resilience, this was not statistically significant (p-value of 0.104).

Participants with a history of mental health counseling had higher levels of anxiety (0.36 points, 95% CI

0.16 to 0.56), depression (0.63 points, 95% CI 0.41 to 0.85), and burnout (0.35 points, 95% CI -0.05 to

0.76). The COVID-19 pandemic and its impact on work-related stress, workload, and working hours

increased participants' anxiety by 0.06 points (95% CI 0.00 to 0.11) and burnout by 0.20 points (95% CI

0.09 to 0.32). The coefficients for depression and loneliness also suggest a negative effect, but they were

smaller and not statistically significant. While resilience levels increased by 0.21 points (95% CI 0.02 to

0.41) with the impact of COVID-19, resilience-building activities were affected negatively by -0.13 points

(95% CI -0.20 to -0.06). We observed a similar but stronger effect on participants' ability to work during

the pandemic for all mental health outcomes and resilience-building activities, but not for resilience

levels, ranging from 0.17 points (95% CI 0.06 to 0.28) for loneliness, to 0.95 points (95% CI 0.61 to 1.29)

for burnout. Age and gender were significantly associated with both COVID-19 related variables, but the

effect was small (odds ratios close to 1). A history of using a chat app for mental health significantly

increased resilience-building activities by 0.28 points (95% CI 0.10 to 0.41). Participants who had

previously accessed Internet resources for mental health saw significantly greater anxiety (0.17 points,

95% CI 0.04 to 0.31) and depression (0.20 points, 95% CI 0.05 to 0.35).

While there were differences in outcomes between control and treatment groups, only higher anxiety

and loneliness were significantly associated with the treatment group. As expected from the DiD, effect

size, and reliable change analyses, there was a significant relationship between study period and all

mental wellbeing and resilience outcomes with strong gains post-intervention.

For the treatment group, Table 7 shows that the mood score was positively correlated with all mental

health outcomes and resilience levels but not with resilience-building activities. A 1-point increase in

mood status was associated with a -0.83 point (95% CI -1.04 to -0.61) decrease in anxiety, -0.71 point

(95% CI -0.95 to -0.47) decrease in depression, -1.44 point (95% C -1.85 to -1.03) decrease in burnout, -

0.38 point (95% CI -0.60 to -0.17) decrease in loneliness, and 2.37 point (95% CI 1.59 to 3.16) increase in

resilience. Specific carelines selected by participants had little impact on mental health or resilience

outcomes, except when comparing the stress and COVID-19 carelines to the anxiety careline. The stress

careline was associated with significantly greater depression but lower burnout than the anxiety

careline; there was no significant relationship between the stress careline and other outcomes. The

COVID-19 careline was associated with lower mental wellbeing scores for all 4 assessments and higher

resilience-building activities and resilience levels. However, these relationships were only marginally

significant at the 0.1 level of statistical significance for depression and loneliness.

Table 7. Additional determinants of mental health and resilience outcomes for the treatment group.

Independent variables	Dependent variables (assessment scores)					
	Depression (PHQ-9)	Burnout (OLBI)	Anxiety (GAD-7)	UCLA LONELINESS	Resilience building activities	Resilience (RS-14)
	Coefficient (SE)	Coefficient (SE)	Coefficient (SE)	Coefficient (SE)	Coefficient (SE)	Coefficient (SE)
Mood score	-0.71*** (0.12)	-1.44*** (0.21)	-0.83*** (0.11)	-0.38*** (0.11)	0.26 (0.19)	2.37*** (0.40)
Careline: depression †	0.20 (0.52)	-1.34 (0.89)	-0.26 (0.46)	-0.09 (0.46)	-1.18* (0.70)	-3.17* (1.67)
Careline: stress †	1.06** (0.43)	-1.79** (0.74)	0.62 (0.39)	-0.36 (0.39)	-0.02 (0.63)	-0.86 (1.44)
Careline: relationships †	-0.32 (0.43)	-1.03 (0.75)	0.36 (0.38)	-0.15 (0.38)	-0.14 (0.57)	-1.88 (1.44)
Careline: COVID-19 †	-0.60* (0.35)	-0.82 (0.60)	-0.21 (0.31)	-0.60* (0.31)	0.40 (0.56)	0.36 (1.15)

638   P-values: * <= 0.10, ** <= 0.05, *** <= 0.01

639   Reference category: † Careline: anxiety

Participant experience

At the end of the study, 820 participants completed an anonymous online participant experience

questionnaire (37% from the control group and 63% from the treatment group). The gender distribution

was the same (37% male and 63% female). Respondents were predominantly young (54% in the 18-to-

29-year age group). Nurses made up the majority (68%). These respondent characteristics are similar to

the attributes of the study population.

Fifty percent of participants from the treatment group reported using the app for more than 28 days out

of the total study period of 56 days (in line with an average of 24 days of use based on app data). When

asked why the app was not used more frequently, the top 3 responses were: "I did not have the time to

use the app" (45%); "The app took too long to work with" (30%); and "The app was too difficult to use"

(13%). Overall, 91% of users found the app easy to use; 87% found the app content relevant; 92%

derived benefits from the app; and 83% felt more resilient to everyday challenges. However, Vitalk users

encountered several difficulties: 32% found the trial welcome email confusing; 27% found it difficult to

download and install the app; 20% found it difficult to sign into the app; 19% had difficulties entering

the trial ID; 25% found getting to the mental health assessments confusing; and 33% noted that the app

took too long to work with. The top 3 likes of app use were: interactions with Viki (71%); the exercises

(58%); and taking the mental wellbeing and resilience assessments (55%). The top three dislikes or

challenges of app use were: exposure to new content every or every other day (36%); the mental health

exercises (28%); and completing the mental health and resilience assessments (25%). The net promotor

score (NPS) [39], was assessed by asking how likely users were on a scale from 1 (not likely) to 10 (very

likely) to recommend Vitalk to others. Sixty-six percent out of 515 respondents rated Vitalk a 9 or 10

(promoters), 23% a 7 or 8 (passives), and 11% a 6 or less (detractors). This resulted in a NPS – promoters

minus detractors – of 55.

Fifty-two percent of participants from the control group reported using the Internet resources for more

than 15 days out of the total study period of 56 days, a finding that is much higher than the one gleaned

through web analytics (an average of less than 2 days accessing web resources). When asked why the

Internet resources were not used more frequently the top 3 responses were: "I did not have the time to

use the resources" (46%); "The resources took too long to work with" (34%); and "The content was too

difficult to understand" (13%). Overall, 87% of users found the Internet resources easy to use; 92%

found the resource content relevant; 95% derived benefits from the resources; and 87% felt more

resilient to everyday challenges. However, Internet resource users encountered several difficulties: 17%

found the trial welcome email confusing; 22% found it difficult to access the resources; 11% found it

difficult to get to the website; 7% had difficulties entering the trial ID; 11% found getting to the mental

health assessments confusing; and 43% noted that the resources took too long to work with. The top

three likes of web resource use were: taking the mental wellbeing and resilience assessments (61%);

reading new content (55%); and the exercises (52%). The top three dislikes or challenges of resource use

were: reading the website content (44%); finding new content by following links (36%); and doing

mental health exercise (28%). Twenty-three percent found completing the mental health and resilience

assessments challenging. When asked how likely users were on a scale from 1 (not likely) to 10 (very

likely) to recommend the Internet resources to others, 64% out of 305 respondents rated Internet

resources a 9 or 10 (promoters), 23% a 7 or 8 (passives), and 13% a 6 or less (detractors). The resulting

NPS of 51 is similar to the treatment group.

Discussion

Based on our literature review, this study is the first RCT of a mental health chat app for health workers

combining multiple mental health and resilience outcomes. It was the first to include a measurement of

resilience and to explore the potential of a mental health chat app to stimulate resilience-building

activities among its users. The RCT took place in a low-resource setting in Southern Africa with very

limited access to mental health counseling and therapy. The challenges that Malawi faces meeting

population health needs have been further exacerbated by the COVID-19 pandemic.

The strengths of our RCT were the recruitment of more than 1,500 health professionals, and the

management team's continuous encouragement of participants over the 8-week study period.

Participants were successfully randomized into the 2 study arms while maintaining single blinding of the

research team. The overall enrollment rate was 94%, with group-specific retention rates of 28%

(treatment group) and 40% (control group) (see CONSORT diagram in Fig 3). Compared to other much

smaller RCTs of iCBT, retention rates were high due to weekly reminders through WhatsApp and email

messages, as well as instant support of users encountering issues during registration, sign in or app use.

A trial of adolescents with self-identified anxiety concern (with fewer than 50 participants per group)

reported 8-week retention rates of 28% (treatment group) and 58% (control group) [40]. An RCT of iCBT

of participants with depression and anxiety symptoms referred to care (substantially different from our

study population) reported much higher 8-week retention rates of 82% (treatment group) and 72%

(control group) [41].

The results from the DiD estimators obtained from mixed linear effect modeling support our working

hypothesis that use of a chatbot such as Vitalk can effectively improve mental health and resilience

outcomes among health professionals in Malawi. The coefficient for the interaction terms between

study group and assessment period varied between -0.68 for depression, -0.58 for burnout, -0.44 for

anxiety, 1.22 for resilience building activities, and 1.47 for resilience levels, indicating that active

engagement with the app was more effective in improving mental wellbeing (except loneliness) and

increasing health worker resilience compared to the passive offering of Internet resources. We are

assured that Vitalk app use causes greater improvements in mental wellbeing and resilience than the

alternative due to the experimental study design and the size of the DiD estimators and confidence

intervals (most do not include zero or absence of a treatment effect). It should be noted that such a

strong treatment effect was observed despite considerable technical issues faced by Vitalk users. These

challenges likely moderated the effect size as indicated by the disproportionate attrition in the

treatment group and the feedback from the anonymous user experience survey, suggesting that this

was a very realistic trial of a constantly evolving innovative mobile technology.

Based on DiD estimators, Vitalk use did not appear to be associated with changes in loneliness. Possible

reasons include: a small effective sample for the treatment group (n = 70); a coarse 3-question

instrument that may not be sufficiently sensitive for measuring the effect of Vitalk's relationship

careline; or app use in general. Although we explored behavioral theory about loneliness, we could not

identify any studies that examined whether iCBT or chatbot use influenced social isolation.

Our findings add substantial evidence to existing research. Unlike other published studies, we tested the

effectiveness of Vitalk by implementing a large RCT in a LMIC context where access to mental wellbeing

and resilience support is extremely limited. Compared to our sample sizes of 200 participants or more

per study arm for most of our outcome measures, the other existing RCTs tend to suffer from very small

samples of about 30 individuals or fewer. Our RCT is also the first of its kind to be applied to the health

workforce, which during normal times is at very high risk for work-related stress and burnout, but even

more so during the COVID-19 pandemic. In addition to mental wellbeing outcomes, our study not only

measured levels of resilience but also resilience-building activities by participants. We were not able to

identify studies that looked at resilience as an outcome to compare them to our findings.

While both study groups experienced improvements in mental wellbeing and resilience between

baseline and endline, the effect size was consistently larger for the treatment than for the control group,

further supporting our working hypothesis that Vitalk is more effective in improving mental health and

resilience outcomes than passive Internet resources. While we were not able to compare our DiD results

to the published literature, we can compare the effect size, which is commonly reported for RCTs. Our

findings suggest an approximately medium effect size, varying between d = -0.36 to -0.52 for mental

health, and d = 0.42 to 0.78 for resilience, which is similar to reported literature on the effectiveness of

c-CBT of depression and anxiety based on 24 RCTs with sample sizes between 19 and 257 adolescents

and young adults with symptoms of anxiety or depression [11]. However, it is smaller than an average

effect size of g = 0.77 (95 % CI 0.59–0.95) reported from a meta-analysis of 49 RCTs [9]. An RCT (n = 56)

on the use of Woebot (a chatbot) to deliver CBT to young adults with depression and anxiety reported a

significant reduction of both outcomes with an effect size of d = 0.44 and 0.37, respectively [42]. An RCT

of the use of the Vivibot chatbot (n = 45) by young people with cancer found a significant interaction

effect between time and study group of -0.41 for reducing anxiety, but no effect on depression [43]. A

review of 12 papers cited weak evidence of the ability of chatbots to reduce depression, with an effect

size of -0.55 (95% CI -0.87 to -0.23), which is very similar to our result, but reported outcomes for

anxiety contradicting our findings [19]. Effect size may be influenced by the characteristics of the study

population. While our study targeted healthy healthcare workers, other studies enrolled participants

with predetermined mental health issues or underlying illnesses. The fact that healthcare workers have

self-selected to cope with stressful situations compared to participants from the general population, or

those clinically diagnosed with symptoms, may explain some of the discrepancies in effect size.

Nonexperimental studies showed a greater effect size than RCTs. An earlier effectiveness study of self-

selected Vitalk users from the general population reported a large effect size of d = 0.81 and greater for

mental health outcomes [16]. Another uncontrolled trial of clients with severe symptoms of depression

and anxiety reported much higher scores for depression and anxiety (based on PHQ-9 and GAD-7

assessments) than our study, as well as effect sizes ranging from d = 0.61 to 0.83 [44].

The LMIC setting of our RCT may account for a smaller effect size than anticipated compared to results

from studies in middle- and high-income countries. Limited and expensive access to mobile phones in

Malawi provides less opportunity for self-help and other innovative app use among the population,

which may have constrained participants' ability to benefit from the full potential of a mental health

chatbot. This is supported by the fact that a high proportion of participants reported that they did not

have the time to use the app or Internet resources (1 in 5), or found that it took too long to work with

them (1 in 3). This could also imply that participants tried to conserve precious airtime. Large differences

in mean scores between our study and others suggest that culture may also play an important role in

shaping how participants respond to mental health and resilience questionnaires [45, 46]. For example,

our RCT in Malawi reported mean anxiety scores (4.5 at baseline and 2.9 at endline) and depression

scores (4.0/2.9) that were less than half of those found in Brazil (pre-/post-app use means of 12.2/8.1

for anxiety, and 15.9/10.4 for depression) [16].

Many of the improvements in mental health and resiliency scores documented in our study are of

statistical and clinical relevance and are welcome outcomes. Changes in risk levels were based on RCIs of

4 for the GAD-7 (anxiety) and PHQ-9 (depression) calculated from our sample. These are similar to or

lower than those reported by other studies (e.g., an RCI of 6 for PHQ-9 and an RCI of 4 for GAD-7 [44],

and an RCI of 6 for GAD-7 [47]). Based on our RCIs, we estimate that about one- to two-thirds of the

improvements in both study groups were reliable (see Fig 9). Applying the RCI, we saw reliable

improvements for depression of 14% (control group) and 17% (treatment group); and improvements in

anxiety of about 25% (both groups). This is considerably lower than the reliable change in clinical

caseness reported by other studies (e.g., 52% for anxiety and 45% for depression [16] and between 58%

and 70% across assessments [44]). These discrepancies are likely due to differences in study populations

(randomly selected health workers versus self-selected general population), with much lower mean

scores for our participants, and a small proportion classified at moderate or severe risk levels of about

13% at baseline and 8% at endline for depression and anxiety, respectively. A small proportion (0% to

7%) of participants saw a reliable worsening of mental health and resilience scores; the effect was

consistently higher in the control group. This decline is in line with about 2% to 5% reported by Daley et

al. [16].

While controlling for trial group affiliation and study period, our analysis explored which user or

engagement characteristics most significantly explained the outcomes of interest. Although these

factors may vary by country context, they must be considered when planning mental health support

strategies and helping health workers increase their resilience to meet everyday challenges.

Engagement with the app or the website, measured as z-scores of time spent, was strongly correlated

with reduced anxiety, depression, burnout, and loneliness. Gan et al. [48] also found a significant

association between engagement in digital mental health interventions and mental health

improvements. A systematic review by Molly and Anderson [49] identified several studies that showed

statistically significant associations between engagement and clinical improvement. Our data did not

indicate that time was a determinant of resilience or resilience-building activities.

Our study identified a need for user interface and app processes to be designed for people with less

experience using smartphones and online resources and with a higher risk of adverse mental health

outcomes. In our study, we found that older and female participants tended to have greater anxiety,

depression, loneliness (age only), and burnout (gender only), as well as less experience with smartphone

and Internet use. In a study of the health workforce in Jordan, age and gender were associated with

worse mental health outcomes for males [50]. A rapid scoping review of occupational stress, burnout,

and depression in women in healthcare during the COVID-19 pandemic showed that female health

workers were at increased risk [51].

Compared to those working in other types of medical and ancillary services, levels of anxiety and

depression appear to be highest among our participants working in outpatient care, a service in which

healthcare workers may be required to pay special attention to mental health support, but, according to

anonymous user feedback, they may also face the greatest time constraints for accessing online mental

health solutions. Participants working in secondary and tertiary care institutions were at higher risk for

anxiety, depression, and burnout compared to those in primary care, a finding similar to a study in Peru

[52]. In our study, we found that public facilities were associated with greater mental health risks than

private, not-for-profit facilities. Public facilities may be able to draw upon the strategies used by private

facilities to reduce anxiety and depression. However, the public sector in Malawi seems better equipped

to build health worker resilience compared to the private sector. Work location (urban versus rural), was

not significantly associated with the outcomes (at the 95% level).

Our findings that current or past exposure to mental health counseling was associated with greater

anxiety, depression and burnout confirmed those of Varma et al. [53]. Our participants saw a negative

impact of the COVID-19 pandemic across all outcomes (except for resilience scores), similarly to findings

from a study in Jordan [50]. A study of healthcare and allied workers involved in the COVID-19 pandemic

from India found high levels of severe and extremely severe anxiety (23.2%) and severe and extremely

severe depression (11.4%)[54]. The level of severe anxiety was almost double of that found in our study

(which did not focus on workers involved in the pandemic). The levels of severe depression were similar

to our findings. Although the Vitalk app demonstrated no effect on loneliness, the pandemic itself had a

negative impact on loneliness overall. This could be expected based on the results of a meta-analysis

reporting a small but significant effect size of g = 0.27 of the COVID-19 pandemic on loneliness [55].

Our study had several limitations. Due to technical issues with the app, the response rate for the

treatment group was substantially lower than for the control group. The attrition rate was greater for

the loneliness and resilience-building activity assessments, reducing the power of our statistical analysis.

Engagement with the app was an average of 11 minutes per day, including the completion of

assessments, and was within a range of 4 to 22 minutes (median) reported by a study of user

engagement with mental health apps [56]. The RCT provided participants with airtime to use Vitalk or to

peruse the web resources on their mobile phones, allowing us to test a mental wellbeing digital

intervention in a resource-poor environment in anticipation of a future rapid increase of affordable

cellular data access in sub-Saharan Africa. Our study was not representative of the entire health

workforce in Malawi, because the interventions required English language proficiency and smartphone

ownership, limiting the study to those with a diploma or at least 2 years of college education. Moreover,

participants were predominantly from 2 urban districts within major population centers and only 18%

from rural health facilities. Given that the classification of mental health and resilience relied entirely on

self-assessments, which can be influenced by sociocultural factors unique to a country or study

population, this phenomenon must be taken into account when comparing our findings to those of

other studies.

As indicated by NPS scores, user satisfaction with Vitalk and web resources was high in the treatment

(NPS of 55) and control (NPS of 51) groups. While we did not find NPS benchmarks for chat apps or iCBT,

a study by Marler [57] compared the NPS in an RCT of 2 smoking cessation apps and found values

ranging between 1 (control app) and 57 (intervention app). A study of community mental health services

found an NPS of 19, which varied considerably by respondent characteristics (from 10 for older, to 37 for

younger patients) [29]. Nonetheless, given the limitations of validity and reliability of single item

measurements such as the NPS [58], the magnitude of the NPS and difference between groups should

be viewed with caution. Moreover, apps such as Vitalk are constantly updated and enhanced as the

technology matures. During the RCT in Malawi, a newer Vitalk version with advanced artificial

intelligence and natural language processing features offering a potentially better user experience was

already under development in Brazil.

Conclusions

Given Vitalk's demonstrably positive effect on mental wellbeing and resilience, it has great potential to

be an effective tool in the arsenal of interventions to reduce work-related stress and burnout. Based on

our completion rate (1 in 3 study participants), there appears to be substantial demand for such an app

and willingness to engage with it for at least 10 minutes daily (a length of time that seems adequate for

most users to improve mental wellbeing and resilience, in addition to extra time for occasional

assessments). User feedback suggests that several factors encourage chatbot use including: natural

interaction with the app; frequently updated content and exercises; and carelines adapted to 2 diverse

types of users – those needing more general psychosocial support and those facing more pronounced

mental health issues. Users indicated that they enjoyed the interaction with the virtual assistant, Viki,

and appreciated the assessments on mental wellbeing and resilience as important benchmarks. Making

the assessments less tedious and more entertaining is an ongoing challenge for app developers. App

developers should weigh the advantages and disadvantages of changing the clinical risk classifications

based on standardized mental health and resilience assessments from their original to fixed groupings,

even if this makes sense from a user-friendliness perspective.

While apps such as Vitalk can help health professionals cope with daily stressors and improve mental

wellbeing, employers need to create less stressful work environments to maximize their effectiveness.

Proven interventions to strengthen health systems function and to increase healthcare provider

satisfaction include ensuring: essential supplies and equipment; an infrastructure that meets the needs

of patients and staff; and adequate and well-trained human resources for health. As Internet access

through mobile devices becomes more widespread and affordable in LMICs, employers in the public and

private sector should consider providing free app subscriptions to their employees, organizing group

sessions for mental wellbeing exercises, and encouraging app use.

Similarly to other studies, our RCT confirmed that mobile chat apps are one effective tool for improving

mental wellbeing and resilience among its target audience. For the first time, our findings show that

apps such as Vitalk can support the mental wellbeing and resilience of health workers to meet the

challenges unique to their profession, especially in a resource poor environment where access to mental

health services is very limited. Countries employ thousands of health workers, and many need support:

as many as 1 in 8 health workers in our RCT needed help with anxiety or depression; 3 in 4 faced some

degree of burnout; and 1 in 4 had low resilience. As countries struggle to increase access to mental

health professionals, apps such as Vitalk have potential to fill the gaps in support. In the aftermath of

the ongoing COVID-19 pandemic, healthcare providers in the public and private sectors should consider

offering access to mental wellbeing apps to all their employees to cope with everyday challenges and to

prepare for future public health emergencies.

883 Acknowledgments

884   The authors would like to thank the management and health professionals from health centers and

885   hospitals in Malawi for their support and participation in this study. We gratefully acknowledge

886   Elizabeth Leitman (Chemonics International) for editing the manuscript.

887 Disclaimer

Funding

Financial support for this RCT was provided by the United States Agency for International Development under the Cooperative Agreement No. AID-OAA-A-15-00046. The funders had no involvement in the analysis or in the decision to submit the paper for publication.

Role of the funding source

The study was funded by the U.S. Agency for International Development (USAID) through the Human Resources for Health 2030 Project, led by Chemonics International in partnership with University Research Company (URC) and other partners. USAID had no role in the data collection, data analysis, data interpretation, or writing of the paper. All authors had full access to all de-identified study data and had final responsibility for the decision to submit for publication.

Contributions

References

1.   Sovold LE, Naslund JA, Kousoulis AA, Saxena S, Qoronfleh MW, Grobler C, et al. Prioritizing the
Mental Health and Well-Being of Healthcare Workers: An Urgent Global Public Health Priority.
Front Public Health. 2021;9:679397.

2.   Muller AE, Hafstad EV, Himmels JPW, Smedslund G, Flottorp S, Stensland SO, et al. The mental
health impact of the covid-19 pandemic on healthcare workers, and interventions to help them:
A rapid systematic review. Psychiatry Res. 2020;293:113441.

3.   Nochaiwong S, Ruengorn C, Thavorn K, Hutton B, Awiphan R, Phosuya C, et al. Global prevalence
of mental health issues among the general population during the coronavirus disease-2019
pandemic: a systematic review and meta-analysis. Sci Rep. 2021;11(1):10173.

4.   Ofori AA, Osarfo J, Agbeno EK, Manu DO, Amoah E. Psychological impact of COVID-19 on health
workers in Ghana: A multicentre, cross-sectional study. SAGE Open Med.
2021;9:20503121211000919.

5.   Mulatu HA, Tesfaye M, Woldeyes E, Bayisa T, Fisseha H, Kassu RA. The prevalence of common
mental disorders among healthcare professionals during the COVID-19 pandemic at a tertiary
Hospital in Addis Ababa, Ethiopia. J Affect Disord Rep. 2021;6:100246.

6.   Msomi N. SA healthcare workers been experiencing burnout long before Covid-19 pandemic:
News24, health 24; 2021 [Available from: https://www.news24.com/health24/mental-
health/mental-health-in-sa/sa-healthcare-workers-been-experiencing-burnout-long-before-
covid-19-pandemic-20210318-2.

7.   Chorwe-Sungani G. Assessing COVID-19-related anxiety and functional impairment amongst
nurses in Malawi. Afr J Prim Health Care Fam Med. 2021;13(1):e1-e6.

8.   Chisholm D, Sweeny K, Sheehan P, Rasmussen B, Smit F, Cuijpers P, et al. Scaling-up treatment
of depression and anxiety: a global return on investment analysis. Lancet Psychiatry.
2016;3(5):415-24.

9.   Grist R, Cavanagh K. Computerised Cognitive Behavioural Therapy for Common Mental Health
Disorders, What Works, for Whom Under What Circumstances? A Systematic Review and Meta-
analysis. Journal of Contemporary Psychotherapy. 2013;43:243-51.

10.   Andrews G, Cuijpers P, Craske MG, McEvoy P, Titov N. Computer therapy for the anxiety and
depressive disorders is effective, acceptable and practical health care: a meta-analysis. PLoS
One. 2010;5(10):e13196.

11.   Christ C, Schouten MJ, Blankers M, van Schaik DJ, Beekman AT, Wisman MA, et al. Internet and
Computer-Based Cognitive Behavioral Therapy for Anxiety and Depression in Adolescents and
Young Adults: Systematic Review and Meta-Analysis. J Med Internet Res. 2020;22(9):e17831.

12.   Watts S, Mackenzie A, Thomas C, Griskaitis A, Mewton L, Williams A, et al. CBT for depression: a
pilot RCT comparing mobile phone vs. computer. BMC Psychiatry. 2013;13:49.

13.   Kristjansdottir OB, Fors EA, Eide E, Finset A, Stensrud TL, van Dulmen S, et al. A smartphone-
based intervention with diaries and therapist-feedback to reduce catastrophizing and increase
functioning in women with chronic widespread pain: randomized controlled trial. J Med Internet
Res. 2013;15(1):e5.

14.   Dagoo J, Asplund RP, Bsenko HA, Hjerling S, Holmberg A, Westh S, et al. Cognitive behavior
therapy versus interpersonal psychotherapy for social anxiety disorder delivered via smartphone
and computer: a randomized controlled trial. J Anxiety Disord. 2014;28(4):410-7.

15.   Bakker D, Kazantzis N, Rickwood D, Rickard N. Mental Health Smartphone Apps: Review and
Evidence-Based Recommendations for Future Developments. JMIR Ment Health. 2016;3(1):e7.

16.   Daley K, Hungerbuehler I, Cavanagh K, Claro HG, Swinton PA, Kapps M. Preliminary Evaluation of
the Engagement and Effectiveness of a Mental Health Chatbot. Front Digit Health.
2020;2:576361.
17.   Mental health atlas 2017. World Health Organization; 2018.
18.   Kauye F. Management of mental health services in Malawi. Int Psychiatry. 2008;5(2):29-30.
19.   Abd-Alrazaq AA, Rababeh A, Alajlani M, Bewick BM, Househ M. Effectiveness and Safety of Using
Chatbots to Improve Mental Health: Systematic Review and Meta-Analysis. J Med Internet Res.
2020;22(7):e16021.
20.   Dallal GE. Randomization plan generator 2020 [Available from:
http://www.jerrydallal.com/random/randomize.htm.
21.   Research Randomizer: Social Psychology Network; 2021 [Available from:
https://www.randomizer.org/.
22.   Seligman MEP, editor Building Human Strength: Psychology's Forgotten Mission. 2006.
23.   Beck JS, Beck AT. Cognitive Behavior Therapy, Second Edition: Basics and Beyond: Guilford
Publications; 2011.
24.   Yoo B-k, Rayz JT. Understanding Emojis for Sentiment Analysis. The International FLAIRS
Conference Proceedings. 2021.
25.   Looking after our mental health: Mental Health Foundation (MHF); 2021 [Available from:
https://www.mentalhealth.org.uk/publications/how-to-mental-health.
26.   Mental well-being: resources for the public: World Health Organization; 2021 [Available from:
https://www.who.int/news-room/feature-stories/mental-well-being-resources-for-the-public.
27.   #HealthyAtHome - Mental health: World Health Organization; 2020 [Available from:
https://www.who.int/campaigns/connecting-the-world-to-combat-
coronavirus/healthyathome/healthyathome---mental-health.
28.   Doing what matters in times of stress: an illustrated guide. Geneva: World Health Organization;
2020.
29.   Wilberforce M, Poll S, Langham H, Worden A, Challis D. Measuring the patient experience in
community mental health services for older people: A study of the Net Promoter Score using the
Friends and Family Test in England. Int J Geriatr Psychiatry. 2019;34(1):31-7.
30.   Demerouti E, Bakker AB. The Oldenburg Burnout Inventory: A good alternative to measure
burnout and engagement. Halbesleben JRB, editor. Hauppauge, NY: Nova Science Pub Inc.;
2008.
31.   Kaggwa MM, Kajjimu J, Sserunkuma J, Najjuka SM, Atim LM, Olum R, et al. Prevalence of
burnout among university students in low- and middle-income countries: A systematic review
and meta-analysis. PLoS One. 2021;16(8):e0256402.
32.   Mughal AY, Devadas J, Ardman E, Levis B, Go VF, Gaynes BN. A systematic review of validated
screening tools for anxiety disorders and PTSD in low to middle income countries. BMC
Psychiatry. 2020;20(1):338.
33.   Steptoe A, Shankar A, Demakakos P, Wardle J. Social isolation, loneliness, and all-cause mortality
in older men and women. Proc Natl Acad Sci U S A. 2013;110(15):5797-801.
34.   Hughes ME, Waite LJ, Hawkley LC, Cacioppo JT. A Short Scale for Measuring Loneliness in Large
Surveys: Results From Two Population-Based Studies. Res Aging. 2004;26(6):655-72.
35.   Building Health Worker Resilience: A Toolkit to Protect Against Burnout on the Front Lines. New
Brunswick, New Jersey, USA: Center for Health Worker Innovation; 2021.
36.   Wagnild G. The Resilience Scale user's guide for the US English version of the Resilience Scale
and the 14-item Resilience Scale (RS–14). Worden, MT: Resilience Center; 2009.
37.   Wagnild G. A review of the Resilience Scale. J Nurs Meas. 2009;17(2):105-13.

38. Siriwardhana C, Abas M, Siribaddana S, Sumathipala A, Stewart R. Dynamics of resilience in
forced migration: a 1-year follow-up study of longitudinal associations with mental health in a
conflict-affected, ethnic Muslim population. BMJ Open. 2015;5(2):e006000.
39. Reichheld FF. The one number you need to grow. Harv Bus Rev. 2003;81(12):46-54, 124.
40. O'Connor K, Bagnell A, McGrath P, Wozney L, Radomski A, Rosychuk RJ, et al. An Internet-Based
Cognitive Behavioral Program for Adolescents With Anxiety: Pilot Randomized Controlled Trial.
JMIR Ment Health. 2020;7(7):e13356.
41. Richards D, Enrique A, Eilert N, Franklin M, Palacios J, Duffy D, et al. A pragmatic randomized
waitlist-controlled effectiveness and cost-effectiveness trial of digital interventions for
depression and anxiety. NPJ Digit Med. 2020;3:85.
42. Fitzpatrick KK, Darcy A, Vierhile M. Delivering Cognitive Behavior Therapy to Young Adults With
Symptoms of Depression and Anxiety Using a Fully Automated Conversational Agent (Woebot):
A Randomized Controlled Trial. JMIR Ment Health. 2017;4(2):e19.
43. Greer S, Ramo D, Chang YJ, Fu M, Moskowitz J, Haritatos J. Use of the Chatbot "Vivibot" to
Deliver Positive Psychology Skills and Promote Well-Being Among Young People After Cancer
Treatment: Randomized Controlled Feasibility Trial. JMIR Mhealth Uhealth. 2019;7(10):e15018.
44. Duffy D, Enrique A, Connell S, Connolly C, Richards D. Internet-Delivered Cognitive Behavior
Therapy as a Prequel to Face-To-Face Therapy for Depression and Anxiety: A Naturalistic
Observation. Front Psychiatry. 2019;10:902.
45. Parkerson HA, Thibodeau MA, Brandt CP, Zvolensky MJ, Asmundson GJ. Cultural-based biases of
the GAD-7. J Anxiety Disord. 2015;31:38-42.
46. Harry ML, Coley RY, Waring SC, Simon GE. Evaluating the Cross-Cultural Measurement
Invariance of the PHQ-9 between American Indian/Alaska Native Adults and Diverse Racial and
Ethnic Groups. J Affect Disord Rep. 2021;4.
47. Bischoff T, Anderson SR, Heafner J, Tambling R. Establishment of a reliable change index for the
GAD-7. Psychology, community & health. 2020;8:176-87.
48. Gan DZQ, McGillivray L, Han J, Christensen H, Torok M. Effect of Engagement With Digital
Interventions on Mental Health Outcomes: A Systematic Review and Meta-Analysis. Front Digit
Health. 2021;3:764079.
49. Molloy A, Anderson PL. Engagement with mobile health interventions for depression: A
systematic review. Internet Interv. 2021;26:100454.
50. Alnazly E, Khraisat OM, Al-Bashaireh AM, Bryant CL. Anxiety, depression, stress, fear and social
support during COVID-19 pandemic among Jordanian healthcare workers. PLoS One.
2021;16(3):e0247679.
51. Sriharan A, Ratnapalan S, Tricco AC, Lupea D, Ayala AP, Pang H, et al. Occupational Stress,
Burnout, and Depression in Women in Healthcare During COVID-19 Pandemic: Rapid Scoping
Review. Front Glob Womens Health. 2020;1:596690.
52. Quispe-Sancho A, Chambi-Macedo KL, Laurel-Vargas V, Huamani-Merma E, Cuzcano-Gonzales
KV, Huaita-Rocha MA, et al. Depression, Anxiety, and Stress in Health Professionals Working
During the COVID-19 Pandemic in Peru: An Analytical Cross-Sectional Study. ELECTRON J GEN
MED. 2021;18(6).
53. Varma P, Junge M, Meaklim H, Jackson ML. Younger people are more vulnerable to stress,
anxiety and depression during COVID-19 pandemic: A global cross-sectional survey. Prog
Neuropsychopharmacol Biol Psychiatry. 2021;109:110236.
54. Shekhar S, Ahmad S, Ranjan A, Pandey S, Ayub A, Kumar P. Assessment of depression, anxiety
and stress experienced by health care and allied workers involved in SARS-CoV2 pandemic. J
Family Med Prim Care. 2022;11(2):466-71.

55.   Ernst M, Niederer D, Werner AM, Czaja SJ, Mikton C, Ong AD, et al. Loneliness before and during
1051   the COVID-19 pandemic: A systematic review with meta-analysis. Am Psychol. 2022;77(5):660-
1052   77.
1053   56.   Baumel A, Muench F, Edan S, Kane JM. Objective User Engagement With Mental Health Apps:
1054   Systematic Search and Panel-Based Usage Analysis. J Med Internet Res. 2019;21(9):e14567.
1055   57.   Marler JD, Fujii CA, Utley MT, Balbierz DJ, Galanko JA, Utley DS. Outcomes of a Comprehensive
1056   Mobile Smoking Cessation Program With Nicotine Replacement Therapy in Adult Smokers: Pilot
1057   Randomized Controlled Trial. JMIR Mhealth Uhealth. 2022;10(11):e41658.
1058   58.   Adams C, Walpola R, Schembri AM, Harrison R. The ultimate question? Evaluating the use of Net
1059   Promoter Score in healthcare: A systematic review. Health Expect. 2022;25(5):2328-39.

1060