SDF, SAP P$_{11}$-4, and GI Sealants for Managing Initial Caries Lesions Produce Clinic and Payor Savings in Financial Impact Model

Authors & affiliations

Savyasachi V. Shah, PhD*
  Senior Research Scientist & Senior Health Economist, National Opinion Research Center, Chicago, IL, USA
Laura J Kibbe, BS, RDH*
  Clinical Innovation Manager, CareQuest Innovation Partners, Boston, MA, USA
Lisa J. Heaton, PhD
  Science Writer, CareQuest Institute for Oral Health, Boston, MA, USA
Courtney Desrosiers, MPH
  Innovation Research Assistant, CareQuest Innovation Partners, Boston, MA, USA
John Wittenborn PhD
  Senior Research Scientist, National Opinion Research Center, Chicago, IL, USA
Mariya Filipova, MBA
  Chief Innovation Officer, CareQuest Institute for Oral Health, Boston, MA, USA
Kirill Zaydenman, MSFS
  Vice President, Innovation, CareQuest Innovation Partners, Boston, MA, USA
Jeremy Horst Keeper, DDS, PhD
  Director of Clinical Innovation, CareQuest Innovation Partners, Boston, MA, USA
  Affiliate Faculty, University of Washington Department of Oral Health Sciences, Seattle, WA, USA
  Instructor, dꞏxʷꞏx̌ayꞏbus Dental Therapy program at Skagit Valley College, Swinomish, WA, USA
  Corresponding author. Address:
  Email: jhorst@carequestinnovation.com
* These authors contributed equally to this work.

Conflict of interest

CareQuest Innovation Partners has a relationship with vVARDIS to assess and support the U.S. deployment of Curodont Repair Fluoride Plus (CRFP). The goal of the partnership is to ensure access for historically marginalized populations (e.g., Medicaid beneficiaries). Dr. Keeper has been paid honoraria for teaching continuing education courses on various noninvasive caries therapies by Elevate Oral Care and GC America, and practices pediatric dentistry.

Funding

All authors were funded by CareQuest Innovation Partners to conduct this evaluation.

Acknowledgements

Special thanks to the clinical innovation, operations, and data analytics teams of Advantage Dental+. Thanks to Dr. Leah Price for guiding the beginnings of this work. Thanks to our colleagues of CareQuest Institute for Oral Health for support and critical feedback, particularly Kelly Schroeder and Caroline McLeod. Thanks to Jean-Francois Blouvac of vVARDIS and to Steve Pardue of Elevate Oral Care for insightful conversations on the dynamics of costs and benefits in oral health product innovation.

Contributions

Conceptualization (JHK, LK, CD, KZ, JW, SS); design & investigation (JHK, LK, SS, JW); analysis (JHK, LK, LH, CD, SS, JW, KZ, MF); writing (JHK, LK, LH, SS); and editing (JHK, LK, LH, CD, SS, JW, KZ, MF).

Abstract

Introduction: Evidence-based noninvasive caries therapies for initial caries lesions recently became

available in the United States. Fundamental differences between noninvasive therapies and the traditional

surgical dental approach warrant study of the financial scalability.

Methods: The financial costs and benefits to fee-for-service clinics and payors were compared across

eleven scenarios simulating the treatment of 1,000 initial lesions over a three-year period. The scenarios

included varying combinations of noninvasive therapies (silver diamine fluoride (SDF), SAP P_{11}-4, and

glass ionomer sealants), no treatment, and various rates of one to three surface restorations to an estimated

current practice model. We used a decision tree microsimulation model for deterministic and probabilistic

sensitivity analyses. We derived assumptions from an initial lesion and noninvasive therapy-focused

cohort study with operations data from 16 sites accepting Medicaid in Alabama as a case study and

clinical data from all 92 sites.

Results: In comparison to the current practice model, scenarios that produce mutually beneficial results

for payors savings and clinics net profit and profit margin include: SAP P_{11}-4, SDF on non-cosmetic

surfaces, and a mix of three noninvasive therapies. When considering the limited resources of chair and

clinician time, the same scenarios as well as SDF with restorations emerged with substantially higher

clinic net profit.

Conclusion: Scenarios that include noninvasive therapies and minimize restorations achieve the balance

of improving outcomes for all parties.

Practical implications: Payors should appropriately reimburse and clinics should adopt noninvasive

caries therapies to improve oral health for all.

Key Words: Cost-benefit analysis, Nonrestorative dentistry, Initial caries lesions, Dental caries; Silver

Diamine Fluoride; Curodont Repair; glass ionomer cement

Introduction

Initial caries lesions are the first noticeable stage of dental caries, where a frank cavitation into dentin has not yet formed.[1] International consensus and United States (US) guidelines state consistently that drilling and restoring are not indicated for initial lesions.[2,3,4] There appears to be poor awareness that demineralization reaching as deep as the outer third of the dentin is unlikely to be cavitated and therefore infected.[1] Historically it has been ubiquitous for dental boards to have candidates drill and restore initial lesions to obtain a dental license. Further, 94% of U.S. dentists self-report performing restorations when radiographic demineralization is still within the range of initial lesions.[5] The same guidelines mentioned above indicate noninvasive treatments for initial lesions.[2,3] Tooth-specific noninvasive caries therapies include silver diamine fluoride (SDF), Self-assembling peptide (SAP P_{11}-4 such as Curodont Repair), and glass ionomer sealants (sealants), among others.[3,6,7] Noninvasive therapies have similar clinical outcomes to more invasive treatments without irreversibly removing tooth structures, are completed quickly and painlessly, and are often reimbursed by payors (e.g., Medicaid). However, to our knowledge, the clinic net profit and payor savings of noninvasive therapies compared to predominant current practice have not yet been studied.[4,5]

SDF is a brush-on therapy that prevents and arrests caries lesions at any stage excluding irreversible pulpitis. SDF-treated lesions turn black, while healthy enamel stays white.[8,9] There have not been sufficient clinical trials to assess the efficacy of SDF on initial lesions.[10] However, numerous clinical trials[3,10] support the American Dental Association (ADA)'s recommendation to arrest cavitated caries lesions on coronal surfaces using SDF. As initial lesions can be arrested with less-potent agents (e.g., fluoride varnish), it is presumed that SDF, a more potent agent, will readily arrest initial lesions.[3,10]

SAP P_{11}-4 is a biomimetic self-assembling peptide professionally applied as a brush-on liquid. SAP P_{11}-4 regenerates hydroxyapatite within the porous enamel of an initial lesion by absorbing into the lesion, self-assembling into scaffolds that span throughout the demineralized volume, and templating calcium and phosphate from the saliva into new hydroxyapatite.[11] SAP P_{11}-4 has clinically important

effects on arresting caries and shrinking initial lesions[6,7] while being safe.[12] In January 2019, SAP P_{11}-4

was added to the US Food and Drug Administration's National Drug Codes Directory under the Over-the-

Counter Anticaries Drug Monograph (NDC: 72247-101).

The ADA recommends the use of sealants in both preventing caries and arresting initial lesions in

pits and fissures.[13,3,14] Glass ionomer can be professionally applied to create a physical barrier and

strengthen the tooth with fluoride and metal ions (e.g., remineralization), which makes the tooth less

dissolvable by acid and less colonizable by bacteria.[15,16,17] Due to the incorporation of fluoride and metal

ions, caries prevention and arrest are maintained even if the bulk of the sealant falls out.[18]

SDF, SAP P_{11}-4, and sealants are evidence-based, noninvasive therapies that can be alternatives

to no-treatment or restorative approaches that could increase clinic net profits and reduce payor costs

while benefiting patients' dental treatment experience. This study analyzes the clinic costs, net profit, and

profit margin, as well as payor costs of using these three noninvasive therapies to treat initial lesions

compared to other restoration approaches and no treatment.

Methods

Study Overview

Clinic costs, gross revenue, net profit, and profit margin and payor costs were modeled for

eleven scenarios of different combinations for the three noninvasive therapies (SDF, SAP P_{11}-4, sealants),

restorations, and no treatment in 1,000 permanent teeth with initial caries lesions. For accuracy in larger

scale models, numbers were not rounded to the nearest whole tooth and therefore fractions of teeth are

included in the modeling. We used data from clinics in the state of Alabama that accepted Medicaid as

representative of a US region with a high disease burden and limited dental and financial resources. We

used a model of estimated *current practice* to which we compared other models in which 40% of teeth

with initial lesions received restorative treatment while 60% received no treatment (see current practice

model rationale below).

All scenarios were modeled for this purpose with assumptions derived from a cohort as described

below. The decision tree microsimulation model was built in Microsoft Excel (Redmond, WA) as per

guidelines from the International Society for Pharmacoepidemiology and Outcomes Research.[19,20] This

model was used for deterministic and probabilistic sensitivity analysis across the scenarios over three

years to compare the costs, net profit, and profit margin (all in US dollars) to fee-for-service clinics and

the fee-for-service payor savings. A three-year period was chosen to reflect the average duration members

remain with a dental insurance plan. We did not adjust for inflation as the primary objective was to

compare treatment approaches for managing initial caries lesions specifically, not the total cost of

managing the patient or population.

Data Source

The cohort consists of patients seen in ninety-two Advantage Dental+ clinics in eight states

(WCG IRB#: 1295572; clinicaltrials.gov ID: NCT04933331). Sixteen of these clinics are in Alabama.

These clinics focus on diagnosis and non-invasive management of initial lesions. The cohort includes

260,594 initial lesions diagnosed in 76,888 patients, with up to 6 years of follow-up. All data for the

current study come from the clinical management of this cohort, except where otherwise noted. Details of

the model variables involving current practice, clinician cost and time, material costs, clinic gross

revenue/payor costs, and treatment failures are presented below. Table 1 lists estimated labor salaries,

overhead costs, material costs, reimbursements, procedure times, and ranges thereof.

Current Practice Model

To determine a base case of current practice for comparison, we estimated the *expected* rate of

restoration for initial lesions. Dental health records typically include treatment codes rather than

diagnostic codes. Therefore, we consulted 20 experts with a national perspective on clinical care and

claims data with representation from payors, nonprofit institutes, academia, and DSOs and estimated that

40% of initial lesions are treated with 1-3 surface restorations. We considered that shallower (less severe)

lesions are most prevalent and least often drilled, while deeper initial lesions are nearly always drilled but

are less prevalent. We verified this expected initial lesions restoration rate by cross-referencing the

incidence of initial lesions and 1-3 surface restorations in the cohort data.

Clinician Costs and Time

We estimated the clinician and chair overhead time required to perform each treatment per patient

from consultation with individuals responsible for clinical management of this cohort. We then projected

this estimate to reflect the quantity of teeth treated and their distribution within the four quadrants of the

mouth, accounting for economies of scale when multiple lesions occur in the same patient. We modeled

hygienists as providers for noninvasive therapies and dentists with an assistant for restorations. Estimates

for provider time for each procedure per first and additional initial lesion(s) and quadrant(s) are shown in

Table 1. No additional time was calculated per additional tooth in the same quadrant for SDF, SAP P_{11}-4,

or sealants. Chair time for a procedure was estimated as either assistant (restoration) or hygienist

(noninvasive therapies) time. The median hourly wages for Alabama dentists, hygienists, and dental

assistants were obtained from Salary.com (accessed November 25[th], 2022) and 33% in indirect costs was

added for payroll taxes and benefits. These wage estimates were confirmed to be accurate with

individuals responsible for the clinical management of the cohort. The probability of the number of teeth

and distribution across four anatomical areas were factored into the model to account for economies of

scale in labor and material cost (Supplemental Table 1).

Material Costs

Material costs were estimated based on unit or dose cost and the number of teeth that can be

treated by each (Table 1). Cost for a planned more efficient SAP P_{11}-4 product for 4+ teeth was

estimated by the manufacturer. A 6.1% sales tax was added to material costs, as the average of those in

the Alabama clinic locations**.** The total treatment costs for each intervention were estimated by

considering the labor, overhead, and material costs, and treatment time, based on the number of teeth

treated and their anatomic distributions observed in the cohort.

Clinic Gross Revenue / Payor Costs

Fees were taken from the 2022 Alabama Medicaid fee schedule for each procedure code in the

existing Current Dental Terminology (CDT) and are listed in Table 1. SDF was modeled with annual

application with attrition, whereas SAP P_{11}-4 and sealants were modeled with a single application (Table

1). The weighted average reimbursement for restorations was calculated based on the relative incidence in

cohort patients for each of the nine CDT procedure codes for 1-3 surface restorations and similarly for re-

restoring any overlapping surface. For the initial placement of treatment, there are economies of scale

(reduced time and cost) represented when multiple lesions occur in the same patient to match clinical

practice (see Supplemental Table 2). The reimbursement for restoration within one year of any

noninvasive therapy was calculated by subtracting the reimbursement given for the prior treatment from

the reimbursement for the restoration. No compensation is given for re-restoration within one year in

Alabama. To account for multiple failures in the same tooth in sequential post-treatment years, separated

by more than 12 months, 50% of such repeat failures were reimbursed in the second and third years. To

establish a reimbursement rate and range for SAP P_{11}-4, a full deterministic analysis was run for the

current practice model and for SAP P_{11}-4 with a range from $20 to $50 per tooth. The break-even for

clinic net profit and payor cost with respect to the current practice model defined the minimum and

maximum of the range, and the midpoint was used in further analysis in attempt to balance payor and

clinic interests.

Treatment Failure / Patient Benefit

Treatment failure is defined as any tooth/surface receiving restoration after baseline (Table 2);

this affects costs. The greatest benefit to the patient is greater drill-free tooth survival (Figure 5;

Supplemental Table 3), preventing the need for additional restorative visits after baseline treatment.

Treatment failures were modeled at the end of each of 3 years. Due to the low treatment failure rate, we

anticipated failures to be independent of any economies of scale that would occur with multiple lesions

within one patient. Failure rates were derived from claims-level records from the cohort data for the

proportion of teeth with operative treatment (restoration, endodontics, extraction, or crown lengthening)

performed on an overlapping surface of an initial lesion previously treated with noninvasive therapies or

1-3 surface restorations. The break-even for clinic net profit and payor cost with respect to the current

practice model defined the minimum and maximum of the range, and the midpoint was used in further

analysis in attempt to balance payor and clinic interests.

Simulation Model of Initial Caries Management

We developed 11 scenarios to reflect several different approaches to initial lesions management,

and a formal model was built to reflect the 11 scenarios with 1000 teeth each (see Figure 1). In addition to

one condition that involved no treatment ("NoTx"), we included four scenarios in which differing

percentages of initial lesions were treated with traditional restorations while the remaining percentages

remained untreated. For example, in the scenario labeled "20%Restore," 20% of initial lesions received

traditional restorations while the remaining 80% were left untreated. We created two scenarios with SDF

applied on 75% of initial lesions, corresponding to the fraction of initial lesions occurring anywhere on

permanent teeth except the facial surfaces of anterior and premolar teeth in the cohort data: in the

"75%SDF" scenario, 75% of initial lesions were treated with SDF and 25% were left untreated; in the

"SDF+Restore" scenario, 75% of initial lesions were treated with SDF and 25% were treated using

traditional restorations. As sealants can only be applied to pits and fissures, similarly, we created two

scenarios with sealants applied to 24% of initial lesions, corresponding to the fraction of initial lesions of

permanent teeth occurring in pits and fissures in the cohort data: in the "24%Seal" scenario, 24% of initial

lesions were treated with sealants and 76% were left untreated; in the "Seal+Restore" scenario, 24% of

initial lesions were treated with sealants and 76% were treated using traditional restorations. There is no

contraindication to applying SAP P_{11}-4 to initial lesions on all surfaces, therefore one scenario estimated

treatment of all (100%) initial lesions using SAP P_{11}-4. Finally, we created one scenario ("NCTmix") in

which initial lesions were treated with the ideal noninvasive therapy with respect to the surface/location

of the initial lesions in the mouth. The 24% of initial lesions occurring in pit and fissure surfaces in cohort

data are treated with sealants; the 26% of initial lesions in facial surfaces of anterior teeth and premolars

(cosmetic surfaces), where the SDF stain may not be acceptable to dental teams and patients, are treated

with SAP P_{11}-4; and the remaining 50% of initial lesions are treated with SDF. In the model, initial lesion

diagnosis and management occur at baseline, and the treatment survival or failure is calculated at the end

of each year for three years. The payor cost and clinic cost, gross revenue, net profit, and profit margin for

each scenario, including treatment failure and repeat failure, are calculated using standard formulas

(Appendix A).

Sensitivity analysis

A Probabilistic Sensitivity Analysis (PSA) was conducted with ten thousand iterations to create

more accurate estimates and to assess the reliability of results. Hourly wage ranges of the 10^{th} to 90^{th}

percentile for dentists, hygienists, and dental assistants were used with an assumption of a triangle (linear)

distribution as the input parameters of salary for the three clinician types vary based on years of

experience and location. The failure ranges for each treatment or non-treatment were derived using an

assumed beta distribution around the arithmetic mean failure rate, by considering the number of actual

failures and observations. We modeled a uniform range of reimbursement for repeat application of SDF in

the 2^{nd} and 3^{rd} years, and for the reimbursement of SAP P_{11}-4. As the price of SAP P_{11}-4 may change

because it is a new product, we have used a uniform distribution with range of $27 to $33 for V2 (1-3

teeth) and $40 to $50 for V3 (4-6 teeth). In the PSA, the start sample of the scenarios was fixed to 1,000

ICLs at the beginning of the 3-year simulation, and random values were picked for each variable from a

range and distribution as previously described and in Table 1, resulting in different costs and gross

revenues for each scenario at the end of 3 years. This was repeated 10,000 times, and mean values with a

range between 2.5 and 97.5 percentile are reported.

Results

Reimbursement estimate

SAP P_{11}-4 reimbursement was estimated at $37 per tooth per the pricing analysis (Figure 2;

Supplemental Table 2c). There is a range ($32 - $43) in which shifting from the current practice model to

treating all teeth with SAP P_{11}-4 increases payor savings and increases clinic net profit. Reimbursement at

the midpoint increases clinic net profit by 41% and decreases payor costs by 10% compared to current

practice. The midpoint represents a balanced reimbursement to incentivize both stakeholders to switch.

Treatment Survival

Survival of treatments (i.e., without receiving restoration after baseline treatment) across three

years is shown in Table 2. Drill-free tooth survival across three years for 1,000 initial lesions managed

with each scenario are shown in Table 3.

Savings and net profits

Cumulative payor costs, clinic costs and clinic net revenue from the deterministic analysis are

shown broken out by year in Supplemental Figure 1 (numerical values in Supplemental Table 4), and in

summary for all three years in Figure 3 (numerical values in Supplemental Table 5). All scenarios

involving fewer restorations than the 40%Restore current practice model result in payor savings, in order

from greatest savings to least: NoTx, 24%Seal, 75%SDF, NCTmix, 20%Restore, and SAP P_{11}-4.

SDF+Restore is highly similar in payor costs. Scenarios involving more restorations than the current

practice model (60%Restore, Seal+Restore, and 80%Restore) result in higher payor costs.

Clinic costs follow the same pattern as payor savings, with all scenarios involving fewer

restorations than the 40% current practice model resulting in reduced clinic costs, and those involving

more restorations than the current practice model resulting in greater clinic costs. All scenarios that treat

more teeth than the current practice model (SDF+Restore, NCTmix, Seal+Restore, 75% SDF, SAP P_{11}-4,

80%Restore, and 60% Restore) result in higher clinic net profits. All deterministic outcomes fall within

the ranges observed in the probabilistic outcomes, as seen layered together in Figure 3.

All scenarios including SAP P_{11}-4 and/or SDF without restorations meet all three success

parameters of increasing payor savings, reducing clinic costs, and increasing clinic net profit: SAP P_{11}-4,

75%SDF, and NCTmix. SDF+Restore meets the clinic benefit criteria but does not reduce payor costs.

Immediate clinic net profit per patient visit

At a per-patient visit level, the increased net profit of noninvasive therapies is minimal or absent

for one tooth but becomes substantial with multiple teeth treated (Supplemental Figure 2). Net profits for

noninvasive therapies follow an economy of scale by number of teeth, due to the trivial additional time

and marginal increase in material costs for application to additional teeth. Distribution of restorations

across the mouth make a substantial impact on restorative dentistry due to the increased dentist and

assistant time (e.g., time for local anesthesia, use rubber dam isolation; Table 1). In comparison,

distribution across the mouth makes trivial impact on noninvasive therapy profitability.

Clinic net profit per hour

Clinic net profit per chair time is higher than the current practice model for all scenarios except

NoTx and 20%Restore (Figure 4). Treating the most teeth at baseline with the least invasive procedures

corresponds to the highest net profit per chair hour, with the top four being NCTmix ($328/hour), SAP

P_{11}-4 ($256/hour), 75%SDF ($232/hour), and SDF+Restore ($190/hour; Figure 4A). The use of time for

all personnel for each scenario is plotted against the corresponding clinic net profit in Figure 4B. Using

hygienist time for SDF and/or SAP P_{11}-4 scenarios saves considerable dentist and assistant time while

increasing clinic net profits compared to traditional restorations.

Financial and patient benefits

Payor savings, clinic profit margin, and drill-free tooth survival (patient benefits) are

simultaneously improved for all non-restorative NCT scenarios (Figure 5). Restorative-focused scenarios

have inverse relationships with payor savings and clinic profit margin. Restorative-focused scenarios with

less than 40% restorations increase payor savings and drill-free tooth survival at the expense of decreased

clinic profit margin. (Figure 5). The scenarios with the highest drill-free survival in order are NCTmix,

SAP P_{11}-4, 75%SDF, and 24%Seal (Supplemental Table 3).

When considering scenarios with greater treatment survival (preventing the need for additional

restorative treatments after baseline), scenarios including SAP P_{11}-4 and/or SDF without restorations lead

to reduced payor and clinic costs as the number of surviving treatments increase (Supplemental Figure 3;

NCTmix, 75%SDF, and SAP P_{11}-4). Non-SDF and non-SAP P_{11}-4 scenarios that treat fewer teeth at

baseline than the current practice model increase payor savings, but produce fewer surviving treatments

and have lower clinic net profit (24%Seal, 20% Restore, and NoTx). Scenarios involving more

restorations have higher payor costs and marginally higher clinic net profit (60%Restore, 80%Restore,

and Seal+Restore). The scenarios with the highest clinic net profit in order are, SAP P_{11}-4, SDF+Restore,

75%SDF, and NCTmix.

Discussion

Our study findings suggest that shifting from the current practice model (40%Restore) to non-

restoratively treating initial caries lesions with SDF or SAP P_{11}-4 (or a mix thereof) results in payor

savings, higher clinic net profit, and superior outcomes for patients. Additionally, all scenarios that treated

more teeth at baseline than the current practice model resulted in more treatments surviving after 3 years.

Longer-term payor savings may be even greater than our modeling shows, as costs accumulate over time

due to the progression of initial lesions to cavitation and restoration.

When considering the chair and clinician time needed to achieve increased net profits, the

noninvasive approaches NCTmix, SAP P_{11}-4, 75%SDF, and SDF+Restore demonstrate clear superiority.

The NCTmix net profit per chair time is six times that of the current practice model. SAP P_{11}-4 and

75%SDF resulted in a net profit per chair time over four times the current practice model. When

considering clinic costs and clinician time needed to achieve clinic net profit, four scenarios emerge as

superior: SDF+Restore, NCTmix, 75%SDF, and SAP P_{11}-4. By utilizing noninvasive therapies, dental

assistants and hygienists can work to the highest level of their scope of practice, freeing up dentists to

deliver care that requires advanced training. More complex procedures tend to have higher net profit per

time, creating opportunities for higher profits (not represented in this work). Shifting production from

managing initial lesions with restorations to performing higher net profit procedures amplifies the

financial benefits of managing initial lesions with SDF and SAP P_{11}-4.

Noninvasive therapies offer a new way to achieve improved treatment outcomes for patients while

financially incentivizing payors and clinics. They follow an economy of scale by teeth treated per patient

visit, so that two or more teeth treated per patient visit becomes increasingly more profitable. When

considering chair time, the mix of noninvasive therapies provides the highest clinic net profit, with SAP

P_{11}-4 and SDF emerging as much more profitable than other scenarios. The drivers of payor savings and

clinic net profit for SDF and/or SAP P_{11}-4 scenarios are minimal treatment times, combined lower cost of

material and clinician labor (compared to traditional restorations), and the applicability to most or all

tooth surfaces. Improvements in patient outcomes that are financially rewarding for clinics and payors are

hallmarks of value-based care. As the oral health care system makes meaningful, albeit gradual, progress

toward implementing value-based care, "there will be greater financial incentives [for oral health care

clinics]…to avoid invasive surgical interventions."[21] Noninvasive therapies are a critically important tool

in the value-based toolbox within oral health care.

Treatment of initial lesions with traditional restorations is extremely common, as 94% of US dentists

report restoring initial lesions[5] while only 11%-12% of control group initial lesions progress.[6,22] This

disconnect may be due to the ubiquitous misunderstanding that radiographic appearance of

demineralization reaching the dentin necessarily indicates cavitation.[5,23,24] However, international

guidelines recommend against restoring initial lesions as these lesions can be healed, or arrested, through

remineralization.[2,3,4] Although restorations have treatment failure rates similar to noninvasive therapies,

when noninvasive therapies fail the negative result is a restoration, whereas failed restorations require

deeper and more invasive procedures. Noninvasive therapies provide a treatment option that empowers

dental teams to heal the first signs of disease.

Sealants are a highly efficient and successful method of treating initial lesions and contribute to the

superiority of the NCTmix in the current analysis. The ADA recommends sealants for the management of

initial lesions.[3,13,14] Additionally, the World Health Organization (WHO) recently added glass ionomer,

modeled here as the sealant material, to their Lists of Essential Medicines for Children and for Adults in

recognition of its preventive and therapeutic value.[25,26,27] The optimal result of the NCTmix scenario

demonstrates that use of sealants to manage initial lesions is ideal when supplemented with other

noninvasive therapies (e.g., SDF or SAP P_{11}-4), rather than being used with other restorations or alone.

Since diagnosis is not widely implemented in American dentistry, payors and clinics may choose to use a

CDT code that indicates sealants placed over initial dental caries lesions for tracking purposes (e.g.

D1352 - preventive resin restoration, or D2940 - protective restoration).

In the short time since it has been used in the US for caries arrest, SDF has already been impactful,

for example by lowering the use of general anesthesia in young children by 70% in a large Medicaid-

focused network of community clinics.[28] SDF is the focus of the ADA evidence-based guideline on

NonRestorative Caries Treatment.[3] The WHO acknowledged SDF's potential to equitably decrease

suffering from caries by adding it to its Lists of Essential Medicines (for both children and adults).[3,24,26]

The simplicity of applying SDF was also recently recognized by the creation of a medical procedure

code.[26,29] However, current evidence primarily focuses on use for moderate to advanced caries lesions

rather than initial lesions.[3,10,30] The results of this study provide evidence of the cost-effectiveness of SDF

for treatment of initial lesions.

SAP P_{11}-4 is a relatively new drug that provides a therapeutic effect to rebuild enamel hydroxyapatite.

Unlike SDF, its lack of staining and neutral flavor positions it to be used throughout the mouth, and

particularly for permanent anterior teeth.[31] It is not used to seal in caries, and therefore may be more

accepted by dentists than sealants.[31,32] For dental clinics, the profit margin for SAP P_{11}-4 is superior,

because it takes less clinician time to perform the application than to place restorative fillings.

While not a direct focus of this analysis, noninvasive therapies are likely to be more beneficial for,

and favored by, patients over traditional dental treatments for initial lesions. Dental fear and anxiety are

well-documented barriers to receiving regular oral health care;[33,34,35,] and should be reduced given that

noninvasive therapies are pain, needle, and drill-free. Since noninvasive therapies can be easily delivered

by non-dentist professionals within and outside traditional dental care settings (e.g., schools,

communities), they should improve patient access to oral health care. Finally, since noninvasive therapies

stop and reverse disease progression, they help patients keep their teeth longer and avoid expensive

restorative care resulting from recurrent caries lesions.[36]

This study includes some limitations. The clinical outcome data on which the analysis depends cannot

be representative of all populations, and likely represent less disease progression than in the population.

This case study of Alabama Medicaid may not represent challenging dynamics in other geographic areas,

though any can be similarly modeled. Medicaid fee schedules usually represent lower fees than

commercial insurance reimbursements. The scenarios are meant to demonstrate practical options in

clinical practice but cannot represent all constellations of mixes. Real-life challenges will inevitably

deviate from the outcomes of this theoretical model, particularly during the change from a restorative to a

non-invasive paradigm; however, these data predict strong financial benefits after a transition time.

Conclusion

This work finds that the use of noninvasive therapies for initial lesions increases payor savings, is

efficient and profitable for clinics, and is beneficial for patients in the modeled scenarios. The key

component of the economic success of noninvasive therapies is to empower non-dentist clinicians to treat

dental caries. Accordingly, to replicate the success of this noninvasive paradigm cost model 1)

policymakers should support non-dentist clinicians to manage initial lesions; 2) payors should invest in

creating value-based reimbursements, and 3) clinics should adopt new procedures with consistent change

management. The time for these techniques has come.

Figure Legends

Figure 1. Initial caries lesion management model. The left side of the figure represents a table with
columns corresponding to the scenarios and rows corresponding to each management method. The 11
modeled scenarios (vertical bars) are simulated by applying the relevant fraction of teeth into each year 1
management category (green and black at left). The current practice scenario model is highlighted in
brown. Payor costs and clinic costs and net revenue are incorporated for the initial treatments (green) and
any progression of caries lesions reaching cavitation and therefore indicating a restoration (orange) or re-
restoration (crimson). Restoration from treatment failure within one year is reimbursed with the initial
treatment fee removed (-$); restoration thereafter is reimbursed fully. Teeth without restoration each year
are considered stable (grey) and have no costs.

Figure 2. Modeling SAP P_{11}-4 reimbursement to balance payor and clinic financial interests. Payor
savings (yellow) and clinic net profit (orange) with respect to the current practice model of 40%
restorations are shown for a range of per-tooth reimbursement amounts for SAP P_{11}-4 treatment.

Supplemental Figure 1: Deterministic clinic net profit, clinic costs, and payor costs. The clinic net
profits (green), clinic costs (orange), and payor costs (total) at the end of each year for 1,000 initial caries
lesions by each management scenario. Payor savings (yellow and brown dashed line) occurs when the
year 3 total bar height is less than that of the current practice model (brown outline). Increased clinic net
profit (dashed green and brown line) occurs when the year 3 green bar is higher than that of the current
practice model (brown outline). All options that include SDF and/or SAP P_{11}-4 increase clinic net profit
while also producing payor savings.

Figure 3. Increasing clinic net profit modified by clinic costs and payor costs. The clinic net profit
(green) and costs (orange) at the end of 3 years are shown for each scenario, from deterministic (bars) and
probabilistic (dots and 95% distributions) approaches. All deterministic data are within the range
observed in the corresponding simulations. Payor costs are shown as the combined bar height and is
written at bottom. From a payor financial perspective, all seven scenarios with lower payor costs are
improvements on the current practice scenario (shown as brown bar). From a clinical financial
perspective, scenarios ending in box "a" and not extending into box "b" are optimal. Basing clinical
decisions purely on net profit would lead to further overtreatment; considering clinic costs identifies three
scenarios as improvements (black outline) from current practice.

**Supplemental Figure 2. Per patient visit immediate clinic net profit by number of teeth and
anatomical areas,** with immediate costs only. Distribution of the teeth across anatomic areas (A) of the
mouth (e.g., quadrants) is much less impactful on noninvasive caries therapy net profit versus restoration
net profit because there is no need for local anesthesia or rubber dams. Treating more teeth (T) per patient
in one visit is more profitable for noninvasive caries therapies regardless of anatomic distribution; all
caries lesions can be managed in one visit.

Figure 4A and Figure 4B. Clinic net profit per hour. Left | A) Clinic net profit divided by the chair
time throughout 3 years is shown for each scenario. Higher clinic net profit per chair time occurs when
the corresponding bar height is higher than that of the current practice model (green box). All scenarios
with non-invasive caries therapies are more profitable per chair time than all restorative-only scenarios.
Right | B) Clinic net profit mapped against use of time from each clinician type: hygienist (purple),
dentist (lilac), and assistant (blue). Times for each scenario are connected by a grey line. Bolded scenarios
use the least amount of clinician time (often the most limited resource) while maximizing net profit.

Figure 5A and Figure 5B. Comparison of payor savings and clinic profit margin to drill-free teeth.
Left | A) Payor savings and **Right | B)** Clinic profit margin for each initial caries lesion management

scenario is plotted against teeth surviving without any caries progression to cavitation at the end of 3

years. Bolded scenarios improve patient outcomes, payor savings, and clinic profit margin. All

nonrestorative scenarios with SDF and/or SAP P_{11}-4 are superior for all stakeholders.

Supplemental Figure 3A and Figure 3B. Comparison of payor savings and clinic profit margin to

treatment survival. Left | A) Payor savings and **Right | B)** Clinic profit margin for each initial caries

lesion management scenario is plotted against treatments surviving without caries progression to

cavitation at the end of 3 years. Bolded scenarios improve patient outcomes, payor savings, and clinic

profit margin. All nonrestorative scenarios with SDF and/or SAP P_{11}-4 are superior for all stakeholders.

Appendix A: Key Terms Defined

Payor cost(s)	The amount of money that the payor pays the clinic for services provided to their covered members. In this model, this is the same as the clinic's gross revenue.
Payor savings	The financial benefit realized from reducing the payor's payments (payor costs). In this financial model payor savings occur anytime a scenario costs less than the current practice scenario – represented by the 40% restoration scenario, for services provided to manage 1000 initial lesions.
Clinic gross revenue	Total revenue generated from clinic operations or services provided.
Clinic cost(s)	The amount of money that the clinic pays to provide services to their patients. This includes expenses such as labor, overhead, and material costs.
Clinic net profit(s)	The amount of revenue remaining after all clinic costs are deducted from gross revenue. Clinic net profit = Clinic gross revenue – clinic costs
Clinic profit margin	A measure of clinic profitability that is expressed as a percentage. This measure displays the portion of gross revenue that the clinic keeps after accounting for clinic costs. Clinic profit margin = Clinic net profit / clinic gross revenue
Payor Financial benefit	In this work, a payor financial benefit is defined as payor costs (fees paid out to the clinic for services provided in each scenario) being less than the payor costs of current practice. This could also be reported as increased payor savings compared to current practice.
Clinic Financial benefit	In this work clinic financial benefits include increased clinic net profit and increased profit margin. These benefits are often realized through decreased clinic costs, increased time efficiency, and increased gross revenue.
Patient benefit	In this work, patient benefits include improved treatment survival rates which reduced the need for additional restorative visits after baseline treatment. Patient benefits can also include the number of teeth that survive without ever needing restorative treatment or drill-free teeth. Other indirect patient benefits include shorter appointment duration, and increased access to care.

References:

1. Young N, Newton J, Morris J, et al. Community water fluoridation and health outcomes in England: a cross-sectional study. *Community Dent Oral Epidemiol*. 2015;43(6):550-559.

2. Schwendicke F, Splieth C, Breschi L, et al. When to intervene in the caries process? An expert Delphi consensus statement. *Clin Oral Investig*. 2019;23(10):3691-3703.

3. Slayton RL, Urquhart O, Araujo MWB, et al. Evidence-based clinical practice guideline on nonrestorative treatments for carious lesions: A report from the American Dental Association. *J Am Dent Assoc*. 2018;149(10):837-849.e19.

4. Banerjee A, Splieth C, Breschi L, et al. When to intervene in the caries process? A Delphi consensus statement. *Br Dent J*. 2020;229(7):474-482.

5. Rechmann P, Doméjean S, Rechmann BMT, Kinsel R, Featherstone JDB. Approximal and occlusal carious lesions: Restorative treatment decisions by California dentists. *J Am Dent Assoc*. 2016;147(5):328-338.

6. Keeper JH, Skaret LJ, Thakkar-Samtani M, et al. Systematic Review and Meta-Analysis on the Effect of Self-Assembling Peptide P11-4 on Initial Caries Lesions. *J Am Dent Assoc. In press*.

7. Aparna BK, Yashoda R, Puranik MP. Remineralization of early enamel caries lesions using self-assembling peptides P11-4: Systematic review and meta-analysis. *J Oral Biol Craniofac Res*. 2022;12(3):324-331. doi:10.1016/j.jobcr.2022.03.012.

8. Espíndola-Castro LF, Rosenblatt A, Galembeck A, Monteiro G. Dentin Staining Caused by Nano-silver Fluoride: A Comparative Study. *Oper Dent*. 2020;45(4):435-441.

9. Horst JA, Heima M. Prevention of Dental Caries by Silver Diamine Fluoride. *Compend Contin Educ Dent*. 2019;40(3):158-163.

10. Horst JA. Silver Fluoride as a Treatment for Dental Caries. *Adv Dent Res*. 2018;29(1):135-140.

11. Alkilzy M, Santamaria RM, Schmoeckel J, Splieth CH. Treatment of Carious Lesions Using Self-Assembling Peptides. *Adv Dent Res*. 2018;29(1):42-47.

12. Brunton PA, Davies RPW, Burke JL, et al. Treatment of early caries lesions using biomimetic self-assembling peptides--a clinical safety trial. *Br Dent J*. 2013;215(4):E6.

13. Wright JT, Crall JJ, Fontana M, et al. Evidence-based clinical practice guideline for the use of pit-and-fissure sealants: A report of the American Dental Association and the American Academy of Pediatric Dentistry. *J Am Dent Assoc*. 2016;147(8):672-82.

14. Beauchamp J, Caufield PW, Crall JJ, et al. Evidence-based clinical recommendations for the use of pit-and-fissure sealants: a report of the American Dental Association Council on Scientific Affairs. *J Am Dent Assoc*. 2008;139(3):257-268.

15. Duque C, Negrini T de C, Hebling J, Spolidorio DMP. Inhibitory activity of glass-ionomer cements on cariogenic bacteria. *Oper Dent*. 2005;30(5):636-640.

16. Forss H, Seppä L. Prevention of enamel demineralization adjacent to glass ionomer filling materials. *Scand J Dent Res*. 1990;98(2):173-178.

17. Mount GJ, Hume WR, Ngo HC, Wolff MS. *Preservation and Restoration of Tooth Structure, 3rd*
Edition | Wiley.; 2016.

18. Mickenautsch S, Yengopal V. Caries-Preventive Effect of High-Viscosity Glass Ionomer and Resin-
Based Fissure Sealants on Permanent Teeth: A Systematic Review of Clinical Trials. *PLoS One*.
2016;11(1):e0146512.

19. Sullivan SD, Mauskopf JA, Augustovski F, et al. Budget impact analysis-principles of good practice:
report of the ISPOR 2012 Budget Impact Analysis Good Practice II Task Force. *Value Health*.
2014;17(1):5-14.

20. Mauskopf JA, Sullivan SD, Annemans L, et al. Principles of good practice for budget impact
analysis: report of the ISPOR Task Force on good research practices--budget impact analysis. *Value*
Health. 2007;10(5):336-347.

21. Gordon SC, Riedy CA, Stohler CS, Vujicic M. Trends in Scope of Practice for Oral Health Care:
Future Transformative Effects. *JDR Clin Trans Res*. 2022;7(1_suppl):31S-39S.

22. Griffin SO, Oong E, Kohn W, et al. The effectiveness of sealants in managing caries lesions. *J Dent*
Res. 2008;87(2):169-174.

23. Innes NPT, Schwendicke F. Restorative Thresholds for Carious Lesions: Systematic Review and
Meta-analysis. *J Dent Res*. 2017;96(5):501-508.

24. Doméjean S, Léger S, Maltrait M, Espelid I, Tveit AB, Tubert-Jeannin S. Changes in Occlusal
Caries Lesion Management in France from 2002 to 2012: A Persistent Gap between Evidence and
Clinical Practice. *Caries Res*. 2015;49(4):408-416.

25. Benzian H. Dental public health breakthrough. *Br Dent J*. 2022;232(7):421.

26. CareQuest Institute for Oral Health. New CPT code encourages health professionals to apply SDF to
treat cavities. Published November 21, 2022. Accessed January 11, 2023.
https://www.dentistryiq.com/dentistry/article/14286046/new-cpt-code-encourages-health-
professionals-to-apply-silver-diamine-fluoride-to-treat-cavities

27. World Health Organization. *World Health Organization Model List of Essential Medicines*.; 2021.

28. Raskin SE, Tranby EP, Ludwig S, Okunev I, Frantsve-Hawley J, Boynes S. Survival of silver
diamine fluoride among patients treated in community dental clinics: a naturalistic study. *BMC Oral*
Health. 2021;21(1):35.

29. American Medical Association. CPT® Editorial Summary of Panel Actions September 2022.
Published November 4, 2022. Accessed January 11, 2023. https://www.ama-
assn.org/system/files/september-2022-cpt-summary-panel-actions.pdf

30. Urquhart O, Tampi MP, Pilcher L, et al. Nonrestorative Treatments for Caries: Systematic Review
and Network Meta-analysis. *J Dent Res*. 2019;98(1):14-26.

31. Bagher SM, Sabbagh HJ, AlJohani SM, Alharbi G, Aldajani M, Elkhodary H. Parental acceptance of
the utilization of silver diamine fluoride on their child's primary and permanent teeth. *Patient Prefer*
Adherence. 2019;13:829-835.

32. Polk DE, Weyant RJ, Shah NH, Fellows JL, Pihlstrom DJ, Frantsve-Hawley J. Barriers to sealant

guideline implementation within a multi-site managed care dental practice. *BMC Oral Health*.
2018;18(1):17.

33. Dailey YM, Crawford AN, Humphris G, Lennon MA. Factors affecting dental attendance following
treatment for dental anxiety in primary dental care. *Prim Dent Care*. 2001;8(2):51-56.

34. Armfield JM, Stewart JF, Spencer AJ. The vicious cycle of dental fear: exploring the interplay
between oral health, service utilization and dental fear. *BMC Oral Health*. 2007;7:1.

35. Armfield JM. What goes around comes around: revisiting the hypothesized vicious cycle of dental
fear and avoidance. *Community Dent Oral Epidemiol*. 2013;41(3):279-287.

36. Rodriguez JL, Thakkar-Samtani M, Heaton LJ, Tranby EP, Tiwari T. Caries risk and social
determinants of health: A big data report. *J Am Dent Assoc*. Published online December 8, 2022.
doi:10.1016/j.adaj.2022.10.006

Table 1. Treatment cost factor estimates: estimated hourly wages, overhead cost, material costs and reimbursements, procedure times, applicable surfaces, and ranges thereof. These data are used as input in the probability sensitivity analysis. The type of distribution for each variable, where relevant, is listed at right.

Category	Sub-category	Amount	Range	Distribution
Median salary per hour[*]	Dentist	$ 85.00	$ 63 to $ 110	Triangle
	Dental Hygienist	$ 36.00	$ 27 to $ 45	Triangle
	Dental Assistant	$ 19.00	$ 15 to $ 23	Triangle
Overhead cost per hour	Without management	$ 11.65		
Material Costs[#]	SAP P_{11}-4 V2 for 1-3 teeth	$ 30	$ 27 to $ 33	Uniform
	SAP P_{11}-4 V3 for more than 3 teeth	$ 45	$ 40 to $ 50	Uniform
	SDF up to 4 teeth	$ 1.50		
	GI Sealant up to 2 teeth	$ 3.50		
	Composite Resin first tooth	$ 12		
	Composite Resin each additional tooth	$ 4		
Reimbursements per tooth[@]	SAP P_{11}-4	$ 37.31	$ 31.88 to $ 42.75	Uniform
	SDF	$ 20.00		
	GI Sealant	$ 26.00		
	Restoration	$ 76.60		
	Re-restoration	$ 93.72		
Procedure time	SDF – one anatomical area	3 min Hygienist		
	SDF – each additional area	1.5 min Hygienist		
	SAP P_{11}-4 – one anatomical area	5 min Hygienist		
	SAP P_{11}-4 – each	2.5 min		

	additional area	Hygienist
	GI Sealant – one anatomical area	5 min Hygienist
	GI Sealant – each additional area	2.5 min Hygienist
	Restoration – first tooth	25 min Assistant
		20 min Dentist
	Restoration – additional tooth in same area	5 min Assistant
		5 min Dentist
	Restoration – additional anatomic area	15 min Assistant
		15 min Dentist
	Re-restoration – first tooth	27.5 min Assistant
		22.5 min Dentist
	Re-restoration – additional tooth in same area	7.5 min Assistant
		7.5 min Dentist
	Re-restoration – additional anatomic area	15 min Assistant
		15 min Dentist
Applicable portion of initial caries lesions	SDF	75.1%
	SAP P_{11}-4	100%
	GI Sealants	24.3%
	Restoration	100%

SDF-treated teeth likely to receive additional treatments	2nd year	50%	35-70%	Triangle
	3rd year	25%	15-50%	Triangle

[*] Salary data for Alabama are obtained from salary.com, confirmed with clinical leadership from the cohort and the ranges are the 10th and 90th percentile values, with 33% fringe added to the salaries shown.

[#] Material costs for SAP P_{11}-4 and package size are sourced from the manufacturer while the material costs and package size for all other materials were sourced from cohort clinics in Alabama. 6.10% shipping and taxes is added to material costs shown.

[@] The reimbursement rates for SAP P_{11}-4 were hypothesized, and reimbursement rates for all other treatment approaches were obtained from the 2022 Alabama Medicaid fee schedule.

Table 2: Treatment survival rates for initial caries lesions management. Per-tooth treatment failure and survival from cohort data, unless otherwise noted. These data are used to define the beta distributions for treatment failure rates in the probability sensitivity analysis. †combination of control groups in Griffin et al., 2008 and Horst Keeper et al., *in press*. *Data from previous year used due to lack of robust data. In the model, survival rates were not rounded to the nearest whole tooth and fractions of teeth are included in the modeling for greater accuracy.

Materials	Failures	Observations	Survival proportion
Year 1 (from 6-12 month observations)			
SDF	400	15,648	97.4%
SAP P$_{11}$-4	61	1,717	96.4%
Sealant	4	398	99.0%
1-3 surface restoration	3,447	119,525	97.1%
Untreated†	57	473	88.0%
Year 2 (from 12-24 month observations)			
SDF	447	14,579	96.9%
SAP P$_{11}$-4	34	945	96.4%
Sealant	12	469	97.4%
1-3 surface restoration	6,682	177,601	96.2%
Untreated†*	57	473	88.0%
Year 3 (from 24-36 month observations)			
SDF	137	4,577	97.0%
SAP P$_{11}$-4*	34	945	96.4%
Sealant*	5	229	97.8%
1-3 surface restoration	4,784	120,240	96.0%
Untreated†*	57	473	88.0%