Subtyping Social Determinants of Health in *All of Us*:

Opportunities and Challenges in Integrating Multiple Datatypes for Precision Medicine

Suresh K. Bhavnani, Ph.D., M.Arch.[1,2]§ Weibin Zhang, Ph.D.,[1] Daniel Bao, B.S.,[1] Mukaila Raji, M.D., M.S., F.A.C.P.,[3] Veronica Ajewole, Pharm.D., BCOP,[4] Rodney Hunter, Pharm.D., BCOP,[4] Yong-Fang Kuo, Ph.D.[1], Susanne Schmidt, Ph.D.,[5] Monique R. Pappadis, Ph.D., MEd, FACRM,[1] Alex Bokov, Ph.D.,[6] Timothy Reistetter, Ph.D., OTR.,[6] Shyam Visweswaran*, M.D., Ph.D.,[7,8] Brian Downer*, Ph.D.[1]

[1]School of Public and Population Health, University of Texas Medical Branch, Galveston, TX, USA

[2]Institute for Translational Sciences, University of Texas Medical Branch, Galveston, TX, USA

[3]Division of Geriatric Medicine, Department of Internal Medicine, University of Texas Medical Branch, Galveston, TX, USA

[4]College of Pharmacy and Health Sciences, Texas Southern University, TX, USA

[5]Department of Population Health Sciences, Long School of Medicine, University of Texas Health San Antonio, San Antonio, TX, USA

[6]School of Health Professions, University of Texas Health San Antonio, San Antonio, TX, USA

[7]Department of Biomedical Informatics, University of Pittsburgh, Pittsburgh, PA, USA

[8]Intelligent Systems Program, University of Pittsburgh, Pittsburgh, PA, USA

§**Corresponding author**
Suresh K. Bhavnani, Ph.D., M.Arch., FAMIA
School of Public and Population Health
Department of Biostatistics and Data Science
Institute for Translational Sciences
University of Texas Medical Branch
301 University Blvd
Galveston, TX, USA
email: subhavna@utmb.edu

* *Shyam Visweswaran and Brian Downer share senior authorship*

A. ABSTRACT

Project Background. Social determinants of health (SDoH), such as financial resource strain and housing instability, account for between 30-55% of people's health outcomes. While many studies have identified strong associations among specific SDoH and health outcomes, most people experience multiple SDoH in their daily lives. Analysis of this complexity requires the integration of personal, clinical, social, and environmental information from a large cohort of individuals that have been traditionally underrepresented in research, which is only recently being made available through the *All of Us* research program. However, little is known about the range and response of SDoH in *All of Us*, and how they co-occur to form subtypes, which are critical for designing precision medicine interventions.

Research Questions. (1) What is the range and response to survey questions related to SDoH? (2) How do SDoH co-occur to form subtypes, and what are their risk for adverse health outcomes?

Methods. For Question-1, we analyzed the range of SDoH questions across the surveys with respect to the 5 domains in *Healthy People 2030* (HP-30), and analyzed their responses across the full *All of Us* cohort (n=372,397, V6). For Question-2, we used the following steps: (1) due to the missingness across the surveys, selected all participants with valid and complete SDoH data, and used inverse probability weighting to adjust their imbalance in demographics, compared to the full cohort; (2) asked three domain experts to group the SDoH questions into SDoH subdomains, for enabling a more consistent granularity; (3) used bipartite modularity maximization to identify SDoH biclusters, their significance, and their replicability; (4) measured the association of each bicluster to three outcomes (depression, delayed medical care, emergency room visits in the last year) using multiple data types (surveys, electronic health records, and zip codes mapped to Medicaid expansion states); and (5) asked three domain experts to infer the subtype labels, in addition to the potential mechanisms that precipitate their adverse health outcomes and interventions to prevent them.

Results. For Question-1, we identified 110 SDoH questions across 4 surveys, which were categorized into 18 SDoH subdomains, and covered all 5 domains in HP-30. However, the results also revealed a large degree of missingness in survey responses (1.76%-84.56%), with later surveys having significantly fewer responses compared to earlier ones, and significant differences in race, ethnicity, and age of participants when compared to the full cohort. For Question-2, the subtype analysis (n=12,913, d=18) identified 4 biclusters with significant biclusteredness (Q=0.13, random-Q=0.11, z=7.5, *P*<0.001), and significant replication (Real-RI=0.88, Random-RI=0.62, *P*<.001). Furthermore, there were significant associations between specific subtypes and the outcomes and with Medicaid expansion, each with meaningful interpretations and potential precision interventions. For example, the subtype *Socioeconomic Barriers* included the SDoH subdomains *employment, food security, housing, income, literacy*, and *education attainment*, and had a significantly higher odds ratio (OR=4.2, CI=3.5-5.1, *P*-corr<.001) for depression, when compared to the subtype *Sociocultural Barriers*. Individuals that match this subtype profile could be screened early for depression and referred to social services for addressing combinations of SDoH such as housing and income. Finally, the identified subtypes spanned one or more HP-30 domains revealing the difference between the current knowledge-based SDoH domains and the data-driven subtypes. These results reflect the complexity of how SDoH co-occur in the real world, and their potential use in the design of models to predict adverse health outcomes, and the design of interventions.

Community Impact. While several SDoH models including the Dahlgren-Whitehead conceptual model have identified SDoH domains, they have emphasized that real-world SDoH span multiple domains with complex interactions and feedback loops. However, this phenomenon has been difficult to analyze given the lack of large cohorts of individuals that have been traditionally underrepresented in biomedical research, and characterized by a wide range of SDoH and datatypes. The results from analyzing SDoH using the *All of Us* cohort provided direct evidence for this real-world phenomenon by showing that data-driven SDoH subtypes span one or more of the SDoH domains defined by HP-30. This result provides testable hypotheses for future studies that SDoH models based on data-driven subtypes will be more accurate and interpretable for predicting adverse health outcomes, when compared to models that do not use those subtypes. Furthermore, the characterization of the range and response to SDoH across the entire *All of Us* cohort using over one hundred SDoH, should enable researchers to use the approach for characterizing other cohorts to identify and address missingness. Finally, our workbench which focuses on subtyping SDoH, provides generalizable and scalable machine learning methods that can be used to periodically rerun the analysis as the *All of Us* cohort continues to evolve.

B. Introduction

Social determinants of health (SDoH), such as financial resource strain[1] and housing instability,[2] account for between 30-55% of people's health outcomes.[3] While many studies have identified strong associations among specific SDoH and health outcomes, most people experience multiple SDoH concurrently in their daily lives.[4-8] For example, limited access to education, unstable employment, and lack of access to healthcare tend to frequently co-occur across individuals leading to long-term stress and depression.[8] Such complex interactions among multiple SDoH make it critical to analyze combinations of SDoH versus single factors. However, analysis of such co-occurrences and their risks for adverse health outcomes requires the integration of personal, clinical, social, and environmental information, critical for designing cost-effective and precision interventions. Unfortunately, the lack of databases containing such multiple datatypes from the same individuals has resulted in a fragmented understanding of how SDoH co-occur and impact health, critical for designing precision interventions.

The *All of Us* program[9-11] provides an unprecedented opportunity to address this fragmented view of SDoH. This program aims to collect multiple data types from one million or more individuals with a focus on populations that have been traditionally underrepresented in biomedical research. These data types include electronic health records (EHRs), health surveys, whole sequence genome data, physical measurements, and personal digital information. Critically, *All of Us* provides several survey modules containing a wide range of SDoH, which in combination with other datatypes, could transform our understanding of high-risk combinations of SDoH.[9]

However, little is known about the range and response of SDoH in *All of Us*, and how they co-occur to form subtypes, which are critical for designing precision medicine interventions. To address these gaps, we characterized more than a hundred SDoH in *All of Us*, which guided the methods we used to analyze how they co-occur to form subtypes, and their risk for health outcomes. The results helped to highlight the opportunities and challenges for conducting subtype analysis in *All of Us*, which integrates multiple datatypes through the use of scalable and generalizable machine learning methods targeted to precision medicine.

C. Background

Social Determinants of Health

The World Health Organization (WHO) defines SDoH as the "non-medical factors that influence health outcomes."[3] Specifically, these include the conditions in which people are born, grow, work, live, and age. Furthermore, such conditions are shaped by a wider set of forces such as economic and social policies, and systems such as discriminatory laws and structural racism.

Several models have proposed the factors and mechanisms involved in SDoH.[4,12] These models were motivated by the concept of *social gradient*,[13] an empirical phenomenon observed within and across nations,[14,15] consistently showing that the lower an individual's social socioeconomic position, the worse their health. To help explain the factors underlying the social gradient, the Dahlgren-Whitehead model[4,16] proposed several inter-

Fig. 1. The Dahlgren-Whitehead conceptual model aimed to visually show the inter-related layers of SDoH domains that influence health.

connected layers of social determinants that influence health. As shown in Fig. 1, the innermost layer contains demographic and genetic factors which are largely unmodifiable. In contrast, the outer layers are modifiable to different degrees such as lifestyle (e.g., exercise and smoking), social and community networks (e.g., contact with supportive friends and family), living and working conditions (e.g., access to health care and employment), and broader socio-economic, cultural, and environmental conditions (e.g., crime in the neighborhood). While this model was not intended to provide explicit testable hypotheses,[4] the factors within each layer are expected to co-occur and impact each other, in addition to responding to external forces such as racism, and capitalism when it is focused on financial profits at the expense of societal benefits.

These early SDoH models motivated numerous studies[17] that analyzed associations among specific SDoH (e.g., immigration status and home density[7]), their association with health outcomes (e.g., education and mortality[18]), and how they manifest within subpopulations (e.g., patients with diabetes[19]). More recently, organizations such as Centers for Disease Control and Prevention (CDC) and *Healthy People 2030* have organized these empirical

results into SDoH domains that roughly map to the Dahlgren-Whitehead model. For example, *Healthy People*
2030 organizes SDoH empirical studies into five SDoH domains: (1) Economic Stability; (2) Education Access
and Quality; (3) Health Care Access and Quality; (4) Neighborhood and Built Environment; and (5) Social and
Community Context. Furthermore, the PhenX program (that provides well-established measurement protocols
for use in biomedical and translational research) has identified SDoH protocols to enable more systematic data
collection and analysis.[20-22]

While the above findings and categorizations have greatly improved our understanding of SDoH and their impact
on health, they have been mostly analyzed based on snapshots of associations among a few factors and health
outcomes. In contrast, SDoH models and recent empirical studies suggest that multiple SDoH tend to co-occur
and impact each other. For example, during the pandemic, Hispanic and Black or African American individuals
not only had a higher exposure to COVID-19 infections due to their front-line jobs and overcrowded living
conditions, but also had a higher risk for serious infections due to prior conditions not addressed due to lack of
healthcare access.[4] Similarly, undocumented immigrants with lower incomes living in neighborhoods with high
pollution, combined with the stress of deportation, have an increased risk of multiple chronic conditions such as
depression and lung cancer.[7] Such studies have resulted in the Centers for Medicare and Medicaid Services
(CMS) emphasizing that SDoH are a multi-level construct which includes both individual and contextual factors
that have complex interactions.[23]

The above co-occurrences of multiple SDoH and their impact on health directly reflect the interconnected layers
of the Dahlgren-Whitehead shown in Fig. 1. However, analysis of such co-occurrences and their health outcomes
requires large datasets with multiple datatypes that have only recently been made available through the *All of*
Us program.

154 *All of Us*: Multiple Datatypes Across a Large Cohort of Underrepresented Americans

The *All of Us* research program[9-11] (*All of Us*), sponsored by the Obama administration since 2015, aims to
accelerate biomedical research to enable discoveries leading to individualized and equitable prevention and
treatment. Such research is currently hampered due to the *limited range* of personal, clinical, social, and
environmental variables available for the same individuals, *limited representation* in research datasets of socially
marginalized populations, and *limited access* to individual-level data due to privacy laws.

To overcome these hurdles, *All of Us* provides three critical features: (1) a data repository that is projected to
contain one million or more participants, with multiple datatypes including electronic health records (EHRs),
health surveys, whole sequence genomic data, physical measurements, and personal digital information such
as from Fitbits; (2) a cohort targeted to include 75% participants from populations underrepresented in research
(race, ethnicity, gender, sex, sexual orientation, and disability) oversampled from the US population; and (3)
strictly-enforced rules to prevent reidentification of participants by disallowing the download of any participant
data, or reporting research results for subgroups less than 20. These rules allow analysis of the *All of Us* data to
be categorized as non-human subjects research, which combined with training and personal authentication by
researchers, has resulted in a substantial reduction in administrative hurdles.

As of 12/30/22 (Controlled Tier, version 6), *All of Us* contained 372,397 total participants, with 8.6% who had
attempted all 9 health surveys (7 related to demographics and general health, and 2 related to COVID-19), and
26.5% who had genomic data. Critical to the current study is the recent addition of a survey specifically targeted
to SDoH questions, which has been attempted by 15.5% in the *All of Us* cohort. A preliminary analysis revealed
that SDoH appear to be distributed across multiple health surveys and EHR codes, with participants providing
those data at different times on a rolling basis. However, little is known about the range and response of SDoH
in *All of Us*, and how they co-occur to form subtypes, a critical step for selecting the methods to identify and
interpret SDoH subtypes.

177 Computational Methods to Identify and Interpret Subtypes

A wide range of studies[24-32] on topics ranging from molecular to environmental determinants of health have
shown that most humans tend to share a subset of characteristics (e.g., comorbidities, symptoms, genetic
variants), forming distinct subtypes (also referred to as *subgroups* or *subphenotypes* depending on the condition
and variables analyzed). A primary goal of precision medicine is to identify such subtypes and infer their
underlying disease processes to design interventions targeted to those processes.[25,33] Methods to identify
subtypes include: (a) investigator-selected variables such as race for developing hierarchical regression

models,[34] or assigning patients to different arms of a clinical trial, (b) existing classification systems such as the
Medicare Severity-Diagnosis Related Group (MS-DRG)[35] to assign patients into a disease category for purposes
of billing, and (c) computational methods such as classification[36-38] and clustering[28,39] to discover subtypes.

Several studies have used computational methods to identify subtypes, each with critical trade-offs. Some
studies have used *combinatorial* approaches[40] (identify all pairs, all triples etc.), which are intuitive, but which
can lead to a combinatorial explosion (e.g., enumerating combinations of the 31 Elixhauser comorbidities would
lead to 2^{31} or 2147483648 combinations), with most combinations that do not incorporate the full range of
symptoms (e.g., the most frequent pair of symptoms ignores what other symptoms exist in the profile of patients
with that pair). Other studies have used *unipartite* clustering methods[38,39] (clustering patients or comorbidities,
but not both together) such as k-means, and hierarchical clustering; and dimensionality-reduction methods such
as principal component analysis (PCA) to help identify clusters of frequently co-occurring comorbidities.[40-46]
However, such methods have well-known limitations including the requirement of inputting user-selected
parameters (e.g., similarity measures, and the number of expected clusters), in addition to the lack of a
quantitative measure to describe the quality of the clustering (critical for measuring the statistical significance of
the clustering). Furthermore, because these methods are unipartite, there is no agreed-upon method to identify
the patient subgroup defined by a cluster of variables, and vice-versa.

More recently, bipartite network analysis[47] (see Appendix A for additional details) has been used to address the
above limitations by automatically identifying *biclusters,* consisting of patients and characteristics simultaneously.
This method takes as input any dataset such as *All of Us* participants and their SDoH, and outputs a quantitative
and visual description of biclusters (containing both participant subgroups, and their frequently co-occurring
SDoH). The quantitative output generates the number, size, and statistical significance of the biclusters,[48-50] and
the visual output displays the quantitative information of the biclusters through a network visualization.[51-53]
Bipartite network analysis therefore enables (1) the automatic identification of biclusters and their significance,
and (2) the visualization of the biclusters critical for their clinical interpretability. Furthermore, the attributes of
participants in a subgroup can be used to measure the subgroup risk for an adverse outcome, to develop
classifiers for classifying a new participant into one or more of the subgroups, and to develop a predictive model
that uses that subgroup membership for measuring the risk of an adverse outcome for the classified participant.

However, while several studies[50,54-61] have demonstrated the usefulness of bipartite networks for the identification
and clinical interpretation of subgroups, there has been no systematic attempt to identify SDoH subtypes mainly
because of the lack of large cohorts containing a wide coverage of SDoH. The *All of Us* program provides an
opportunity to use bipartite networks for the identification and interpretation of SDoH subtypes using a wide
range of variables in a large cohort, and for analyzing their risk for health outcomes, a critical step in advancing
precision medicine.

D. Method

Research Questions

219 Our analysis was guided by two research questions targeting the *All of Us* dataset:

   1. *What is the range and response to survey questions related to SDoH?*

   2. *How do SDoH co-occur to form subtypes, and what are their risk for adverse health outcomes?*

Data Description

Study Population. In Question-1, we analyzed the full *All of Us* cohort (n=372,397) and characterized their
responses to all the SDoH identified by the domain experts (described in the Variables subsection). For Queston-
2, we analyzed all participants (n=12,913) that had valid responses to the SDoH identified in Question-1, and
used them to identify subtypes, and their risks for specific outcomes.

Variables. For Question-1, 3 SDoH domain experts were asked to review all 1113 questions across 7 *All of Us*
non-COVID health surveys, each of which is attempted once per participant (*The Basics, Lifestyle, The Basics,*
Personal Medical History, Health Care Access & Utilization, Family Health History, and SDoH), and the 2843
Systematized Nomenclature of Medicine (SNOMED) codes related to SDoH.[62] The domain experts arrived at a
consensus for the SDoH across the surveys and the SNOMED codes. As the SDoH-related SNOMED codes in
the EHR had very low usage (see Appendix B for a characterization), they were not further characterized.

In Question-2, to identify SDoH subtypes, we used (a) the SDoH identified from Question-1, (b) covariates including 3-digit zip code (to determine if participants in each subtype came from a state that accepted Medicaid expansion providing greater access to health insurance), variants of the gene SLC6A4 (known to be highly prevalent in families with depression),[63,64] and demographics (age, sex, race). Outcomes included the following: (1) _Depression_ was selected as it is a common health outcome when individuals encounter SDoH in their daily lives such as long-term stress resulting from racism,[65] and dysregulation of the hypothalamic-pituitary-adrenal axis (HPA) axis.[66] Depression was defined as having a positive response to both of the following questions in the _The Basics_ survey ("_Are you still seeing a doctor or health care provider for depression?_" and "_Has a doctor or health care provider ever told you that you have Depression?_") or having SNOMED codes related to depression Codes in their EHR (35489007, 36923009, 370143000, 191616006, or 66344007), (2) _Delayed Medical Care_ was selected as it often results from the lack of medical insurance, which can impact the use of medical care when needed leading to poorer health outcomes.[67] Delayed medical care was defined as having one or more positive responses to 9 survey questions (delayed care due to: transportation, rural, nervousness, work, childcare, copay, elderly care, out of pocket, and deductible) from the _Health Care Access & Utilization_ survey. (3) _Emergency Room (ER) Visits in Last Year_ was selected because lack of medical insurance often results in individuals not seeking early medical care when needed, leading to an exacerbation of conditions precipitating one or more ER visits.[68] As the survey questions that we used for SDoH subtyping were based on outcomes in the past year, we defined ER visits for a participant as having one or more ER visits (CPT 99281-99285) one year preceding the date when the SDoH survey was completed.

Analytical Approach

Question-1: What is the range and response to survey questions related to SDoH?

To address this question, we characterized all SDoH in _All of Us_ at two levels of granularity: (1) SDoH questions based on the surveys used to collect the data, and (2) SDoH subdomains, which were groupings of the SDoH questions to form a coarser grained classification (see Table-1 which explains the difference between SDoH questions, domains, subdomains, and subtypes). These two levels of SDoH granularity in _All of Us_ were characterized as follows:

Term	Definition	Example
SDoH Domains	Five domains defined by health.gov	Economic Stability
SDoH Subdomains	19 subdomains present in All of Us	Food Security
SDoH Questions	SDoH related questions in All of Us surveys (categorized into SDoh domains and SDoH subdomains using domain knowledge)	How often do you have someone to help you read health-related materials?
SDoH Subtypes	Data-driven clusters of SDoH subdomains identified through bipartite network analysis	Cluster Members: Food Security, Educational Attainment, Employment, Housing, Income, and Literacy

Table 1. Definitions for SDoH domains, subdomains, questions, and subtypes.

Identification and Coding of SDoH (SDoH Questions and SDoH Subdomains)

A. Identification and Coding of SDoH Questions in All of Us. Three SDoH experts independently used their domain knowledge about SDoH to identify and code the SDoH questions, and to examine their range with respect to the five _Healthy People 2030_ domains using the following steps: (1) review all 1113 questions across 7 health surveys (excluding 2 related to COVID-19), and extract all SDoH questions that were relevant; (2) transform all positive or value-free questions into negative phrases and abbreviate them for interpretability in the graphs (e.g., "_How often do you have someone help you read health-related materials?_" was changed into "_No one to help read health materials_"); (3) reverse code, and dichotomize the abbreviated SDoH questions (e.g., Always/Often=1, and Never/Occasionally/Sometimes=0); and (4) categorize the SDoH questions into one of the five _Healthy People 2030_ SDoH domains (Economic Stability, Education Access and Quality, Health Care Access and Quality, Neighbourhood and Built Environment, and Social and Community Context). The three domain experts subsequently met and collaboratively resolved any differences between their coding schemes to arrive at a consensus.

B. Identification and Coding of SDoH Subdomains of Questions in All of Us. The same three SDoH experts arrived at a consensus to categorize one or more of the above SDoH questions in _All of Us,_ into SDoH subdomains, and to examine their range using the following steps: (1) review the subgrouping labels of questions in the _All of Us_ surveys, and integrate them to classify the SDoH into subdomains; (2) code a participant as having a "1" for a SDoH subdomain if they had one or more of the questions within that subdomain which had been answered with a "1"; and (3) categorize the SDoH subdomains into one of the five _Healthy People 2030_

SDoH domains (Economic Stability, Education Access and Quality, Health Care Access and Quality, Neighbourhood and Built Environment, and Social and Community Context).

Range and Responses to SDoH Questions and Subdomains

The above knowledge-based classification of SDoH questions and SDoH subdomains were analyzed to examine their range (with respect to the five *Healthy People 2030* domains), and their response (across all participants in *All of Us*), using the following methods. (1) Bar graph displaying the number of participants that had valid answers (all responses other than "skip" or "choose not to answer") to each of the SDoH questions, sorted by survey based on mean response, and then sorted by raw response within each survey. Additionally, to examine their range, each SDoH question/subdomain was colored by one of the five SDoH domains defined by *Healthy People 2030*. (2) Venn diagram showing how many participants had cross-sectionally valid responses to all identified SDoH questions/subdomains. (3) Table describing the number and proportion of race, ethnicity, sex, gender, and age between those that answered the SDoH questions/subdomains, versus those that did not have valid responses. (4) Frequency distribution of the number of SDoH questions/subdomains across participants that had valid responses for all the SDoH questions. The above plots are shown in the Results section.

Question-2: How do SDoH co-occur to form subtypes, and what are their associations with covariates and risks for adverse health outcomes?

Data. We used the cohort identified in Question-1 (participants who had valid answers to all the SDoH questions). However, *e*xamination of the SDoH questions revealed that some of them (e.g., cannot afford dental care, cannot afford prescriptions) had a finer level of granularity compared to others (e.g., single household). As the questions with a finer level of granularity tend to be more strongly co-related to each other in comparison to other coarser grained questions, they also tend to cluster together more strongly, confounding the interpretation of the subtypes. In contrast, as the SDoH subdomains had a more uniform granularity, and were at a level of abstraction that was appropriate to guide referral to the proper social services, we used them to identify the SDoH subtypes.

Analytical Model. To identify SDoH subtypes, their associations with outcomes and covariates, and their future translation into precision medicine, we used a three-part analytical framework called **H**eterogenization, **I**ntegration, and **T**ranslation (HIT). As shown in Fig. 2, the *heterogenization* step was used to identify the subtypes through the use of bipartite modularity maximization[48-50] (see Appendix A for more details), the *integration* step was used to measure the association of each subtype to multiple datatypes,[69] and the *translation* step was used to qualitatively interpret the subtypes,[69] with the goal of developing in the future a decision-support system to translate the subtypes into clinical practice. The following describes the specific methods used in each of the HIT steps:

Fig. 2. The three steps of the **HIT** framework to analyze SDoH. (1) **Heterogenization** of the data to identify subtypes. (2) **Integration** of multiple datatypes such as from EHRs (e.g., depression), genomic data (e.g., mutated genes), and state (e.g., to determine Medicaid expansion) to determine risk and enrichment of each subtype, and (3) **Translation** of subtypes through interpretation and predictive modeling, with the goal of designing of clinical decision-support systems for precision medicine.

1. Heterogenization: Identification of Subtypes. As there were many participants that did not have valid answers to the SDoH questions, dropping them resulted in differences in the proportion of demographic variables compared with the full *All of Us* cohort. The data therefore needed to be adjusted to better reflect the overall *All of Us* participants. To adjust the demographic distribution of the cohort to match the full *All of Us* cohort, we calculated the inverse probability weights (IPW)[70,71] for each participant in the cohort. IPW calculates weights to proportionally boost the values of participants that are underrepresented in the cohort, with respect to a comparison such as the full *All of Us* cohort, using the method similar to an earlier study on the *All of Us* cohort[72] (see Appendix E). Next, we multiplied the IPW generated weights with the original binary values for each participant in our cohort, and used *min-max* to range-normalize those weights within each SDoH subdomain. Finally, to test the replicability of the SDoH subdomain biclustering, we randomly divided the dataset into a training and a replication dataset.

We identified subtypes in the training dataset, and tested the degree to which the SDoH subdomain co-occurrences replicated in the test dataset using the following steps: (1) modelled participants and SDoH subdomains as a weighted bipartite network (see Step-1 in Fig. 2) where nodes were either participants (circles), or SDoH subdomains (triangles), and the associations between participant-SDoH subdomain pairs were weighted edges (lines) generated from IPW; (2) used a bipartite modularity maximization algorithm,[48-50] (which takes edge weights into consideration) to identify the number of biclusters, their members, and measure the degree of biclusteredness through bicluster modularity (Q, defined as the fraction of edges falling within a cluster, minus the expected fraction of such edges in a network of the same size with randomly assigned edges); (3) measured the significance of Q by comparing it to a distribution of the same quantity generated from 1000 random permutations of the network, while preserving the network size (number of nodes), and the distribution of weighted edges for each participant; (4) used the Rand Index (RI) to measure the degree to which SDoH occurred and did not co-occur in the same cluster in the training and test datasets,; and (5) measured the significance of RI by comparing it to the mean of a distribution of the same quantity generated by randomly permuting the training and replication datasets 1000 times, while preserving the size of the networks.

2. Integration: Risk and Enrichment of Subtypes. We used logistic regression to measure the odds ratio (OR) for each subtype compared pairwise to each of the other subtypes, for the three outcomes (Depression, Delayed Medical Care, and ER Visits in Last Year), and for living in a state with Medicaid expansion. To adjust for the difference in demographics due to the missingness, we used weights generated from IPW for each participant, and the comparisons were adjusted for demographics (age, sex, race) and corrected for multiple testing within each outcome using FDR. As 1688 (13.1%) participants did not have 3-digit zip code information, we used IPW to measure the weights of the cohort, and used them to account for potential sample selection bias.

3. Translation: Interpretation of Subtypes. The subtype interpretation was done using the following steps: (a) used the *Fruchterman-Reingold* [51] and *ExplodeLayout*[52,53] algorithms to visualize the bipartite network along with the risk for each of the outcomes; and (b) asked the domain experts to independently label the subtypes and infer their mechanisms with potential strategies to address modifiable SDoH with the goal of reducing the risk for each of the outcomes (Depression, Delayed Medical Care, and ER Visits in Last Year), and then collaboratively come to a consensus.

E. Results

Question-1: What is the range and response to survey questions related to SDoH?

Identification and Coding of SDoH Questions and Subdomains. The three domain experts identified 110 questions from 4 surveys (*The Basics*, *Overall Health*, *Healthcare Access & Utilization, and SDoH)*. Of these, 110 were abbreviated, and 48 were negatively-worded and coded (see Appendix C). The 110 SDoH questions were further categorized into 19 SDoH subdomains (one of these was *Delayed Medical Care* that was used as an outcome).

Response to SDoH Questions and Subdomains. As shown in Figure 3a, the number of valid responses for each of the 110 SDoH questions was largely dictated by the surveys in which they were solicited. SDoH from 2 surveys (*The Basics*, *Overall Health*) had the most valid responses (mean=349434, SD=23556), followed by *Healthcare Access & Utilization* (mean=149898, SD=6146), and finally the SDoH survey (mean=55960, SD=1083). This pattern of responses match how answers to each of the surveys were solicited: at enrollment,

all participants are required to do *The Basics*, and *Overall Health* surveys, and then on a rolling basis the other surveys responses are solicited. The *SDoH* survey is the latest survey that was solicited, which explained their lowest number of responses. As shown in Fig. 3b, this pattern of missingness held for the responses at the SDoH subdomain level, which was not unexpected as the SDoH subdomains were aggregations of the SDoH questions. However, as shown in Fig. 3A and 3B by the uneven number of valid responses within each survey block, there were several SDoH questions that had invalid responses ("skip" or "chose not to answer") at both levels of granularity: *The Basics*: 6%; *Health Access & Utilization* 6.1%; *Overall Health*: 4.39%; and *SDoH*: 2.61%. Furthermore, the proportion of valid to invalid responses between them was significantly different for the SDoH questions (χ^2 (2, N=365237)=57.489, *P*<.001), and for the SDoH subdomains (χ^2 (2, N=372063)=75.637, *P<.001*).

Range of SDoH Questions and Subdomains. As shown by the colored bars in Figure 3, the surveys spanned the full range of the five SDoH *Healthy People 2030* domains. The SDoH questions in *The Basics* and *Overall Health* surveys were predominantly related to economic stability (blue) and social and community context (purple), those in *Healthcare Access & Utilization* survey were all related to that topic (green), whereas those from the *SDoH* survey were a mix of all four domains. Overall, the five *Healthy People 2030* domains had 100% coverage across the four surveys: Social and Community Context=38; Neighborhood and Built Environment=19; Economic Stability=10; Education Access and Quality=2; Health care Access and Quality=42. This characterization suggests that while the SDoH in *All of Us* have broad domain coverage across the surveys, analysis of them requires access to all four surveys, which have different levels of completion and valid responses.

Cohort with Maximized Valid Responses. Given the large degree of missingness in 2 of the 4 surveys, we could not use multiple imputation to estimate the values. We therefore had to find a subset of participants that had valid responses to all the SDoH questions. An examination revealed that two SDoH questions had <10% responses (*English Verbal Frequency*=1.67%, and *Neighborhood has no recreation spaces*=8.4%), accounting for the largest loss in cohort size with valid responses. These questions were therefore dropped from further analysis. Furthermore, one question required a branched response (*Living Situation* branching to *Did not Live in a House*) which were merged. Finally, as we used *Delayed Medical Care* as an outcome, 9 questions related to that topic were removed, resulting in a total of 98 SDoH questions. As shown in Fig. 4, a Venn diagram of the overlap among the valid responses across the surveys revealed that 12,913 participants had valid responses to all 98 SDoH questions.

Fig. 3. The number of valid responses for (A) 110 SDoH questions, and (B) 19 SDoH subdomains. The colors denote how the SDoH in each were categorized based on the 5 *Healthy People 2030* (HP-30) domains.

Fig. 4. Venn diagram showing 12,913 participants (3.5% of the full cohort), who had valid responses to all 98 SDoH questions.

Fig. 5. Frequency distribution of (a) number of co-occurring responses to **SDoH questions** across the 12,913 participants with valid answers to the 98 SDoH questions, and (b) number of co-occurring **SDoH subdomains** across 19 SDoH subdomains.

Co-occurrence of the Number of SDoH across Responders. As shown in Fig. 5, participants had a median
of 15 SDoH question co-occurrences and a median of 9 SDoH subdomains co-occurrences. Furthermore,
participants of color or racial/ethnic minorities, who had valid responses to the 110 SDoH questions, had a
significantly higher median number of co-occurring SDoH compared to the equivalent White population (median
participants of color or racial/ethnic minorities=20, median White=14, P<.001). These results show the high co-
occurrences of SDoH at both levels of granularity, with a significant difference in median co-occurrences between
the White and the participants of color or racial/ethnic minority populations, with valid responses.

Participant Demographics with Valid Responses to SDoH Questions. As the cohort size dropped to 3.5%,
we analyzed how that impacted the demographic distribution compared with the overall *All of Us* cohort. As
shown in Table 2, there were statistically significant differences in race (χ^2(5, N=372,397)=2073.1, P<.001), and
ethnicity (χ^2(9, N=372,397)=6292.2, P<.001) between the two cohorts, after multiple testing correction, with a
higher proportion of White participants having valid answers compared to participants of color or racial/ethnic
minorities. Furthermore, there was a statistically significant difference in age between the participants who had
valid answers, versus those that did not (H(1)=148.08, P<.001). These results show the demographic differences
between the cohort with complete and valid answers to the SDoH questions, in comparison to the full *All of Us*
cohort.

Demographics		All AoU Participants: 372,397 (100%)	All AoU Participants with valid^a SDoH anwers: 12,913 (3.5%)
Race	**White**	201149 (54.01%)	11279 (87.35%)
	Black or African American	73383 (19.71%)	482 (3.73%)
	Asian	12459 (3.35%)	324 (2.51%)
	Other or >1 population	26890 (7.22%)	343 (2.66%)
	None Indicated	58516 (15.71%)	485 (3.76%)
Ethnicity	**Not Hispanic or Latino**	288227 (77.4%)	12095 (93.67%)
	Hispanic or Latino	66704 (17.91%)	751 (5.82%)
	Additional Options	17466 (4.69%)	67 (0.52%)
Sex at birth	**Female**	222495 (59.75%)	8236 (63.6%)
	Male	138831 (37.28%)	4674 (36.09%)
	Intersex	80 (0.02%)	20 (0.15%)
	Additional Options	10991 (2.95%)	20 (0.15%)
Gender	**Female**	220833 (59.3%)	8113 (62.82%)
	Male	138140 (37.09%)	4642 (35.95%)
	Non Binary	920 (0.25%)	60 (0.46%)
	Transgender	464 (0.12%)	20 (0.15%)
	Additional Options	12040 (3.23%)	79 (0.61%)
Age		Median=56 (19-122^b)	Median=58(19-93)

Table 2. The demographic differences between the total *All of Us* participants, and those that had valid answers to all 110 SDoH questions. ^aParticipants that completed all questions, and did not skip, or choose not to answer a question; ^bAge 122 = a participant chose the least birth year (1900).

Fig. 6. Four biclusters in the training dataset consisting of subgroups of participants (n=6492), and their most frequently co-occurring SDoH subdomains (d=18). The biclustering was significant (Q=0.13, random-Q=0.11, z=7.5, *P*<0.001) and the SDoH subdomains significantly replicated in the replication dataset (Real-RI=0.88, Random-RI=0.62, *P*<.001). Across all three outcomes, Cluster-1 had a significantly higher OR compared to Cluster-4. The labels in bold text represent the consensus interpretion of the clusters by three domain experts.

Question-2: How do SDoH subdomains co-occur to form subtypes, and what are their risk for adverse health outcomes?

The cohort used to identify the subtypes consisted of the 12,913 participants, of which 12,886 had valid IPW weights. The latter cohort were split randomly into the training and replication datasets, each with complete data for 18 SDoH subdomains (identified in Question-1), in addition to the three outcomes (depression, delayed medical care and ER visits in last year), and covariates (demographics).

1. Heterogenization: Identification of Subtypes

The weighted (using IPW generated weights) bipartite network analysis of the training dataset (n=6492) and the 19 SDoH subdomains revealed 4 biclusters with statistically significant bicluster modularity (Q=0.13, random-Q=0.11, z=7.5, *P*<0.001). As shown in Fig. 6, there were four clusters with participant subgroups and their most frequently co-occurring SDoH subdomains (*Cluster-1 (pink)*: education attainment, literacy, income, employment, food security, and housing; *Cluster-2 (green)*: afford medical care, discrimination in everyday life, discrimination in medical settings, interactions with providers; *Cluster-3 (blue)*: neighborhood cohesion, and relationships with others; and *Cluster-4 (gray)*: demographics, language, healthcare coverage, provider characteristics, neighborhood characteristics, and supportive relationships). These co-occurrences of SDoH subdomains, significantly replicated in the replication data set (Real-RI=0.88, Random-RI=0.62, *P*<.001). As shown in Fig. 7, while the 18 SDoH subdomains have a hierarchical relationship with the five *knowledge-driven Healthy People 2030* domains (shown on the left), those same SDoH subdomains have a more complex relationship with the four *data-driven* biclusters (shown on the right).

2. Integration: Risk and Enrichment of Subtypes

Table 3 shows the association of each subtype to the three outcomes. As shown by the dark orange row, Cluster-1 (education attainment, literacy, income, employment, food security, and housing) had a significantly higher OR for each of the three outcomes compared to Cluster-4 (provider characteristics, neighborhood characteristics, healthcare coverage, demographics, supportive relationships, language). Furthermore, within the *Depression* outcome, each of the clusters had a significantly higher OR compared to one other cluster forming a ranking of risk among all the four clusters (1>3>2>4). In contrast, *Delayed Medical Care* had two other significant

associations (2>1, 3>4), with *ER Visit in the Last Year* having only the one significant pair-wise association that fit into the overall trend.

As shown in Table 4, this trend continued in the enrichment analysis of association with living in a state with *No Medicaid Expansion*. As shown, Cluster-1 had a significantly higher OR compared to Cluster-4, in addition to the other clusters. The overall results suggests that Cluster-1 and Cluster-4 form "book ends" representing the high and low ends of risk among the clusters.

3. Translation: Interpretation of SDoH Subtypes and Design of Potential Interventions

The three domain experts examined the co-occurrences of SDoH subdomains within each bicluster shown in the network visualization (Fig. 6), and integrated them with the quantitative ORs in Table 3 and 4. The consistent "book ends" result where Cluster-1 had significantly higher ORs compared with Cluster-4 across all four variables was of strong interest, and interpreted as follows: (1) **Cluster-1** was labeled *Socioeconomic Barriers* as it contained multiple high risk SDoH. These co-occurring SDoH

Fig. 7. 18 SDoH subdomains (center) have a hierarchical relationship with the 5 SDoH domains define by HP-30 (left), both of which are knowledge driven. In contrast, the SDoH subdomains have a complex relationship with the subtypes identified through machine learning (ML), reflecting how they co-occur in the real-world, and suggested by models such as the Dahlgren-Whitehead model (shown in Fig. 1).

could have resulted from cascades over time such as low education leading to unstable employment, which in turn led to low income, and food and housing insecurity. Such cascading factors could be perceived by participants to be unmodifiable, leading to chronic stress and depression. Furthermore, the strong association of this subtype with the outcomes *Delayed Medical Care* and *ER Visits in Past Year*, and that participants in this subtype were more likely to be from a US state with *No Medicaid Expansion*, began to provide a more comprehensive understanding of this high-risk SDoH subtype. For example, individuals that match this subtype profile could be screened early for depression and simultaneously triaged to social services prioritized to address specific SDoH triggers such as food security, housing, and income if feasible. (2) **Cluster-4** was labeled *Sociocultural Barriers* as it contained a combination of SDoH related to neighborhood and supportive relations, in addition to cultural barriers related to language, provider interactions, and language. In contrast to the more

Cluster Comparison		Outcomes		
Cluster-A vs. Cluster-B		Depression	Delayed Medical Care	ER Visit in Last Year
1	2	OR=1.7, CI=1.5-2, P-corr=2.5e-10 <.001	OR=0.78, CI=0.67-0.92, P-corr=0.0038 <.01	OR=1.2, CI=0.91-1.6, P-corr=0.24
1	3	OR=1.3, CI=1.1-1.6, P-corr=0.019 <.05	OR=0.88, CI=0.72-1.1, P-corr=0.23	OR=1.4, CI=0.96-1.9, P-corr=0.13
1	4	OR=4.2, CI=3.5-5.1, P-corr=3.5e-52 <.001	OR=3.5, CI=3-4.1, P-corr=1.8e-53 <.001	OR=1.8, CI=1.4-2.3, P-corr=0.00016 <.001
2	3	OR=0.79, CI=0.64-0.97, P-corr=0.022 <.05	OR=1.2, CI=0.98-1.4, P-corr=0.094	OR=1, CI=0.75-1.5, P-corr=0.8
2	4	OR=2.3, CI=1.9-2.7, P-corr=5.2e-21 <.001	OR=4.3, CI=3.7-5, P-corr=3.4e-85 <.001	OR=1.3, CI=1-1.7, P-corr=0.12
3	4	OR=2.9, CI=2.3-3.5, P-corr=1.5e-21 <.001	OR=3.6, CI=3-4.4, P-corr=1.6e-38 <.001	OR=1.4, CI=0.95-1.9, P-corr=0.13

Table 3. Across all three outcomes, Cluster-1 had a significantly higher risk compared to Cluster-4 (dark orange row). The Depression outcome had a distinct ranking of risks (light orange), whereas the other two outcomes had a subset of them.

Cluster Comparison		Enrichment
Cluster-A vs. Cluster-B		No Medicaid Expansion
1	2	OR=1.5, CI=1.3-1.8, P-corr=1.7e-05 <.001
1	3	OR=1.3, CI=1-1.6, P-corr=0.048 <.05
1	4	OR=1.3, CI=1.1-1.5, P-corr=0.0057 <.01
2	3	OR=0.99, CI=0.8-1.2, P-corr=0.97
2	4	OR=0.99, CI=0.86-1.2, P-corr=0.97
3	4	OR=1, CI=0.82-1.2, P-corr=0.97

Table 4. Cluster-1 had a significantly higher OR compared to Cluster-4 (dark orange) for no Medicaid expansion, in addition to Cluster-2 and Cluster-3 (light orange).

devastating socioeconomic barriers in Cluster-1, the sociocultural barriers could be perceived as potentially modifiable, resulting in a lower risk for depression, delayed medical care, and ER visits. Participants that match this profile could be screened for language and communication barriers, useful for providing culturally-competent care, identifying providers that better match the profile of the individuals, and for providing resources to facilitate contact with matching nationality or cultural groups online or in the vicinity.

While Cluster-1 and Cluster-4 formed the "book ends" of risk across the three outcomes, **Cluster-2** was flagged as critical and labeled *Lived Experience Barriers*. The SDoH in this cluster included discrimination in everyday life, medical settings, in addition to problems in interactions with providers and affording medical care. These frequently co-occurring SDoH could explain why this subtype had a significantly higher OR for *Delayed Medical Care* compared to Cluster-1. Finally, **Cluster-3** was labeled *Social Context Barriers* as the SDoH related to neighborhood cohesion and interactions with others. While not as critical as Cluster-1 and Cluster-2, this cluster still had significantly higher OR for depression compared to Cluster-4.

The clusters were therefore meaningful, and additionally did not have a one-to-one mapping to the 5 SDoH domains defined by *Healthy People 2030*. As shown in Fig. 7, these data-driven clusters have a complex relationship with the SDoH domains and subdomains. While one subtype belonged to a single domain (subtype *Social Context* belonged to the domain *Social and Community Context*), three of the four subtypess belonged to two or more domains (e.g., the subtype *Socioeconomic Barriers* belonged to the domains *Economic Stability*, and *Education Access and Quality*). Such interdomain relationships reflect how SDoH co-occur in the real world reflecting the complex cross-domain interactions described in the Dahlgren-Whitehead model (Fig. 1). These relationships could be useful for refining conceptual models to explain the complex association beween SDoH and adverse health outcomes, and to build more accurate SDoH models for predicting adverse health outcomes.

F. Discussion

The mechanisms through which SDoH precipitate adverse health outcomes are complex consisting of many interacting factors and feedback loops among individual and environmental/contextual factors. While this phenomenon has been studied for more than three decades, the *limited range* of data types, *limited representation* of populations that have been socially marginalized, and *limited access* to individual-level data at scale due to privacy laws have been critical hurdles for researchers. Recognizing that *All of Us* has well-articulated plans and resources to overcome these limitations, but is still in a rapidly evolving stage, we conducted a systematic characterization of more than a hundred SDoH available in *All of Us,* and used them to identify SDoH subtypes with the future goal of designing precision medicine interventions. This attempt led to the following opporunities and challenges related to data, methods, and theory.

Data: Missingness and Granularity

All of Us data contained 110 SDoH across 4 surveys, and 93 SDoH-related SNOMED codes in the EMRs. While these provided a comprehensive coverage of SDoH with respect to domains and subdomains identified by *Healthy People 2030*, our analysis uncovered the following patterns of missingness and SDoH granularity.

Missingness. The analysis revealed three types of missingness: (1) *Rollout Missingness:* This type of missingness was largely dictated by how the surveys were rolled out to participants. As all participants at enrollment are required to do *The Basics*, and *Overall Health* surveys, they had the highest responses, followed by the later solicited survey *Healthcare Access & Utilization*, and the *SDoH* survey rolled out more recently in 2022. This order of rollout was the main source of missingness resulting in a precipitous reduction in cohort size for those that had answers to all the SDoH questions. (2) *Valid Answer Missingness.* As participants can choose not to answer any survey questions, the data contained "PMIs" related to "skip" and "choose not to answer". These accounted for a much smaller reduction in cohort size for complete data. (3) *Low Usage Missingness*. Although there were 259 SDoH SNOMED codes, only 93 (3.3%) had such information for >20 participants that are allowed to be reported. This could be because most clinicians currently do not screen for SDoH, as it is typically done by the social worker. Furthermore, we also attempted to use 3-digit zip codes to determine which subtypes had a significant association to living in a state that did not offer Medicaid expansion, and a mutation on the gene. However, 13.1% (1688) of the participants did not have zip code information (adjusted by using IPW), and 3248 (25.2%) did not have genomic data (which we are in the process of being analyzed).

Together, the above three types of missingness impacted the size of the resulting cohort that had valid answers, in the following two ways: (1) a drastic reduction in cohort size by 93.5%. However, because of the size of the overall cohort (n=372,397), we were still left with a large subcohort (n=12,886), which to the best of our knowledge is the largest subset of individuals to be analyzed for such a wide range of SDoH; and (2) significant differences in the proportion of race, ethnicity, and age in the above cohort when compared to the overall *All of Us* population. Specifically, the cohort with valid answers had significantly more White, or non-Hispanic, or older participants, when compared to the overall cohort. This could potentially be because once a participant has been

enrolled, there is a 90-day delay in sending subsequent solicitations to complete surveys, a policy that is currently being re-assessed due to its impact on missingness. We therefore had to correct this imbalance in demographics by using IPW for identifying subtypes that were representative of the overall *All of Us* cohort.

Granularity. Because our goal was to use machine learning methods to identify SDoH subtypes, we encountered uneven granularity in the SDoH questions. Some questions were fine-grained and highly correlated and therefore would cluster more strongly because of the nature of the granularity of the questions, not because of the SDoH mechanisms. To address this uneven granularity, and to make the results more interpretable, we used SDoH subdomains which had a coarser but more consistent level of granularity. We chose this approach because SDoH subdomains had already been defined, were understood by domain experts, and appeared to be at the right level of abstraction useful for clinical applications such as referring a patient to the appropriate social services. However, because the use of coarse-grained variables loses information, future research could explore aggregating only those SDoH questions that are highly correlated, and preserving the rest at the finer level of granularity.

Method: Scalability, Generalizability, and Extensibility

We designed the HIT analytical framework to be scalable enabling its use for the growing size of the cohort in *All of Us*, to be generalizable across cohorts and conditions, and to be extensible for including additional methods as needed in the future. Testing the HIT framework on the *All of Us* data provided insights for the strengths and limitations of the framework, and for the *All of Us* workbench where the analysis was conducted.

Scalability. We used three types of code to conduct the analysis for both research questions. (1) Automatically generated code to extract the cohort, produced by *All of Us* once a cohort was selected using the point and click interface. This code was adequately scalable and generalizable and so will not be discussed further. (2) Customized code to extract specific parts of the data. For example, the analysis of co-occurrences required customized code in R to plot the diagrams in Fig. 3. As expected, these tasks required strong programming skills, but fortunately we did not encounter any coding or execution problems using the R or Python programming languages. However, there were significant server issues which hampered our analysis. Although the workbench instructions state that code running on the workbench for more than 2 weeks would be terminated and all intermediate results deleted, we frequently encountered our work disappearing at shorter intervals. These disruptions resulted in a higher consumption of the free server time credits, and fewer analyses that we could conduct due to the computation time. (3) Machine learning code we had previously developed and disseminated on CRAN[73-75] to conduct the bipartite network analysis and the significance testing, and to visualize the network. As this code was designed to be generalizable and scalable, we did not encounter any issues in the execution of our code (besides the same server issues mentioned above). Finally, the visualization of our networks worked as expected, and we used them to help interpret the patterns in the data.

Generalizability. Our code for the first two steps of the HIT framework is in Jupyter notebooks and have been used to analyze other cohorts that were filtered for age and prior conditions. For example, we extracted a cohort (n=4090) of participants with diabetes aged >=65 with complete data on 18 SDoH variables selected through consensus by 2 experienced health services researchers, and guided by Andersen's behavioral model. The analysis[76,77] revealed 7 SDoH subtypes with statistically significant modularity compared with 100 random permutations of the data (*All of Us*=.51, Random Mean=.38, z=20, *P*<.001), and which were not only clinically meaningful, but also significant in different degrees for the outcome. Our subsequent attempt at increasing the number of SDoH variables from 18 to 110 for participants with diabetes that had valid answers, led to an extremely small cohort size (n=926) (see Appendix D) due to the missingness that we described above. While this reduction resulted in our current strategy of analyzing all particpants regardless of condition or age, these experiments demonstrate that our approach is generalizable to other subsets of the data.

Extensibility. The HIT model is designed to be extensible to include other methods. For example, the model could use other biclustering (e.g., Non-negative Matrix Factorization[78]) and causal modeling methods, and use different types of classification (e.g., deep learning[79]), and prediction methods (e.g., subgroup-specific modeling [38]) to build the decision-support system in the Translational Step (Fig. 2). Furthermore, the model can integrate a wide range of data types to enable analysis of how each subtype is associated with them, resulting in a layered interpretation of the SDoH subtypes as we have demonstrated. Finally, the integration of different subtypes required a diverse team consisting of experts in machine learning, biostatistics, programming, clinical care, health services research, and gerontology to enable a 360 analysis and interpretation of the subtypes. The use

of the workbench to share results operationalized *team-centered informatics*[80] designed to facilitate multidisciplinary translational teams[81] to work more effectively across disciplinary boundaries, with the goal of analyzing and designing interventions for precision medicine.

Theory: Model Building, and Translational Implications

The identification of SDoH subtypes have strong implications for model building in addition to translational applications. As shown in Fig. 7, while the current classification of five SDoH domains has a hierarchical relationship with the SDoH subdomains, the data-driven clusters have a more complex association with the same SDoH subdomains. This reflects the complexity of how SDoH occur in the real-world.

Future models should develop predictive models using the data-driven subtypes to determine whether they improve the accuracy of predicting adverse health outcomes when compared to models that do not use those subtypes. Because the subtypes were clinically interpretable, they could be used to build classification and predictive models, and used with an interface to develop a clinical decision-support system that help to triage patients to critical services. For example, the St. Vincent House (https://www.stvhope.org/) in Galveston, Texas provides several services to address SDoH including free walk-in clinical care, nurse practitioner with small co-pay requested, English and Spanish-speaking free mental health counseling, free dental health clinic, utility and rental assistance, case management, financial literacy, expanded food pantry, weekly free home delivery of pantry groceries, snack pack for people experiencing homeless, free transportation for doctor's appointment, immigration legal services, and spiritual counseling. Given the availability of this wide range of services in many communities across the US, a decision-support system could help to classify an individual based on their SDoH profile into one or more of the subtypes, measure their risk for an adverse health outcome, and proactively recommend one or more of such local services (Fig. 2, Step-3).

Notebooks for *All of Us* Community Use

Because the missingness of SDoH should reduce in the near future, their characterization and subtyping will need to be repeated and verified for different cohorts. Therefore, we have made the following two sets of code available for general use by *All of Us* researcher community:

1. *SDoH Valid Answer Tracker.* This set of notebooks generate four plots which can be used by other researchers on *All of Us* to characterize any cohort: (1) valid responses plot to show how many participants have data with valid responses, and colored by SDoH domains; (2) Venn diagram showing how many participants have valid responses for all questions within each survey; (3) frequency distribution plot showing co-occurrence of SDoH across the selected cohort. This set of tools should enable researchers to characterize SDoH across different cohorts, to help determine methods that are appropriate to adjust for missingness in those cohorts.

2. *SDoH Subtyper.* This set of notebooks can be used to conduct the following analyses: (1) bicluster modularity of a cohort with the 18 SDoH subdomains to identify the number and members of biclusters, and the measure Q representing the quality of the biclustering; (2) visualization of the bipartite network; and (3) significance of the network with respect to null models.

Limitations

This study has two main limitations. The first emerges from the temporary limitations of the large amount of missingness in the survey data, precluding the use of imputation methods. We could therefore use only complete data, which led to a large drop in cohort size, and which also introduced a bias in the demographics requiring a rebalancing through IPW. While such rebalancing is typically done for large datasets, the IPW method requires judgement to decide which variables to include in the model, and therefore could have introduced additional unknown biases. Therefore, the model should be refined to determine which variables to include in the regression models that estimate the IPWs. However, because the clustering was similar between the unweighted and IPW weighted networks, we believe that the current subtypes are stable, meaningful, and represent the demographic composition of the full *All of Us* cohort. This limitation is expected to be addressed as *All of Us* has recently removed the requirement of waiting for 90 days before a subsequent survey is given to an enrollee in the program, potentially reducing the degree of missingness. The second limitation is due to the high computational cost of empirically determining the significance of the biclustering. As such analysis is computationally expensive and time-consuming, it limited the experiments we could do to test different cohorts and models. We therefore look forward to the *All of Us* workbench providing the ability to run batch processes more efficiently, and which will

657 be uninterrupted for extended periods of time (exceeding the current time window), which together could help
658 alleviate this computational hurdle in the future.

G. Conclusion

How SDoH impact health is a complex phenomenon involving many interconnected factors which have yet to be
fully elucidated. While this phenomenon has been studied for more than 30 years, the analyses have been
hampered by the lack of large cohorts representing diverse populations with multiple datatypes, and with easy
access by researchers. *All of Us* provides an unprecedented opportunity to directly address these limitations with
the goal of doing justice to early conceptual models such as the social gradient and the Dahlgren-Whitehead
model, both of which drew international attention to the complex ways in which individual and contextual SDoH
factors impact health. The *All of Us* dataset is also timely because of the visceral health disparities that were
revealed during the pandemic, which pushed the abstract phenomenon of SDoH into the public and policy
realms. However, because *All of Us* is still rapidly evolving to meet its target of one million participants or more,
we conducted a systematic characterization of SDoH variables in *All of Us,* and used the results to guide the
analysis of SDoH subtypes, with the future aim of designing precision medicine interventions.

Our first goal of characterizing the data revealed the nature of the missingness in SDoH, and the uneven
granularity in the SDoH questions. Both these results led us to select the IPW method to address the
missingness, and analysis of subtypes at the SDoH subdomain level of granularity. Our second goal of identifying
SDoH subtypes led not only to statistically significant biclusters, but also to their statistically significant
replication, and meaningful domain interpretations. These results set the stage for further investigations to build
and evaluate prediction models, and decision-support systems. The results also led to the design, use, and
dissemination of general-purpose tools currently available on *All of Us* for other researchers, which will be useful
to reanalyze the *All of Us* data as it grows over the next few years. These collaborative advances should position
All of Us to revolutionize research for analyzing complex phenomena such as how SDoH impact health and
beyond, with the goal of enabling a more equitable future that all of us deserve.

H. References

1. Weida EB, Phojanakong P, Patel F, Chilton M. Financial health as a measurable social determinant of health.
PloS one. 2020;15(5):e0233359.
2. Kushel MB, Gupta R, Gee L, Haas JS. Housing instability and food insecurity as barriers to health care
among low-income americans. *Journal of general internal medicine.* 2006;21(1):71-77.
3. WHO. Social determinants of health. https://www.who.int/health-topics/social-determinants-of-
health#tab=tab_1.
4. Dahlgren G, Whitehead M. The Dahlgren-Whitehead model of health determinants: 30 years on and still
chasing rainbows. *Public health.* 2021;199:20-24.
5. Cook WK. Paid sick days and health care use: An analysis of the 2007 national health interview survey data.
American Journal of Industrial Medicine. 2011;54(10):771-779.
6. Kaplan GA, Keil JE. Socioeconomic factors and cardiovascular disease: a review of the literature. *Circulation.*
1993;88(4 Pt 1):1973-1998.
7. CMS. Center for Migration Studies. Mapping Key Determinants of Immigrants' Health in Brooklyn and
Queens. 2021; https://cmsny.org/wp-content/uploads/2021/02/Mapping-Key-Health-Determinants-for-
Immigrants-Report-Center-for-Migration-Studies.pdf.
8. NEJM. Social Determinants of Health (SDOH). *NEJM Catalyst.*
9. Ramirez AH, Gebo KA, Harris PA. Progress With the All of Us Research Program: Opening Access for
Researchers. *Jama.* 2021;325(24):2441-2442.
10. Denny JC, Rutter JL, Goldstein DB, et al. The "All of Us" Research Program. *The New England journal of
medicine.* 2019;381(7):668-676.
11. Ramirez AH, Sulieman L, Schlueter DJ, et al. The All of Us Research Program: Data quality, utility, and
diversity. *Patterns (New York, NY).* 2022;3(8):100570.
12. Marmot MG. Status Syndrome: A Challenge to Medicine. *Jama.* 2006;295(11):1304-1307.
13. Adler NE, Ostrove JM. Socioeconomic status and health: what we know and what we don't. *Annals of the
New York Academy of Sciences.* 1999;896:3-15.
14. Mackenbach JP, Bos V, Andersen O, et al. Widening socioeconomic inequalities in mortality in six Western
European countries. *Int J Epidemiol.* 2003;32(5):830-837.

15. McDonough P, Duncan GJ, Williams D, House J. Income dynamics and adult mortality in the United States, 1972 through 1989. *American journal of public health.* 1997;87(9):1476-1483.

16. Goran D, Whitehead M. Policies and strategies to promote social equity in health. Background document to WHO - Strategy paper for Europe, 1991. *Arbetsrapport, Institute for Futures Studies.* 2007;14.

17. Lucyk K, McLaren L. Taking stock of the social determinants of health: A scoping review. *PloS one.* 2017;12(5):e0177306.

18. Muller A. Education, income inequality, and mortality: a multiple regression analysis. *BMJ (Clinical research ed).* 2002;324(7328):23-25.

19. Hill-Briggs F, Adler NE, Berkowitz SA, et al. Social Determinants of Health and Diabetes: A Scientific Review. *Diabetes care.* 2020;44(1):258-279.

20. NIMHD. PhenX Social Determinants of Health Assessments Collection. 2022; https://www.nimhd.nih.gov/resources/phenx/. Accessed Januray, 2023.

21. Hamilton CM, Strader LC, Pratt JG, et al. The PhenX Toolkit: get the most from your measures. *American journal of epidemiology.* 2011;174(3):253-260.

22. PhenX. Social Determinants of Health Collections. 2017; https://www.phenxtoolkit.org/collections/view/6.

23. CMS. CMS Framework for Health Equity 2022–2032. 2022; https://www.cms.gov/files/document/cms-framework-health-equity.pdf.

24. McClellan J, King M-C. Genetic Heterogeneity in Human Disease. *Cell.*141(2):210-217.

25. Waldman SA, Terzic A. Therapeutic targeting: a crucible for individualized medicine. *Clinical Pharmacology & Therapeutics.* 2008;83(5):651–654.

26. Rouzier R, Perou CM, Symmans WF, et al. Breast cancer molecular subtypes respond differently to preoperative chemotherapy. *Clinical cancer research: an official journal of the American Association for Cancer Research.* 2005;11(16):5678-5685.

27. Sorlie T, Perou CM, Tibshirani R, et al. Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. *Proceedings of the National Academy of Sciences of the United States of America.* 2001;98(19):10869-10874.

28. Fitzpatrick AM, Teague WG, Meyers DA, et al. Heterogeneity of severe asthma in childhood: confirmation by cluster analysis of children in the National Institutes of Health/National Heart, Lung, and Blood Institute Severe Asthma Research Program. *The Journal of allergy and clinical immunology.* 2011;127(2):382-389.e381-313.

29. Haldar P, Pavord ID, Shaw DE, et al. Cluster analysis and clinical asthma phenotypes. *American journal of respiratory and critical care medicine.* 2008;178(3):218-224.

30. Lotvall J, Akdis CA, Bacharier LB, et al. Asthma endotypes: a new approach to classification of disease entities within the asthma syndrome. *The Journal of allergy and clinical immunology.* 2011;127(2):355-360.

31. Nair P, Pizzichini MMM, Kjarsgaard M, et al. Mepolizumab for Prednisone-Dependent Asthma with Sputum Eosinophilia. *New England Journal of Medicine.* 2009;360(10):985-993.

32. Ortega HG, Liu MC, Pavord ID, et al. Mepolizumab Treatment in Patients with Severe Eosinophilic Asthma. *New England Journal of Medicine.* 2014;371(13):1198-1207.

33. Collins FS, Varmus H. A new initiative on precision medicine. *The New England journal of medicine.* 2015;372(9):793-795.

34. Lacy ME, Wellenius GA, Carnethon MR, et al. Racial Differences in the Performance of Existing Risk Prediction Models for Incident Type 2 Diabetes: The CARDIA Study. *Diabetes care.* 2015.

35. Baker JJ. Medicare payment system for hospital inpatients: diagnosis-related groups. *Journal of health care finance.* 2002;28(3):1-13.

36. Lipkovich I, Dmitrienko A, Denne J, Enas G. Subgroup identification based on differential effect search--a recursive partitioning method for establishing response to treatment in patient subpopulations. *Statistics in medicine.* 2011;30(21):2601-2621.

37. Kehl V, Ulm K. Responder identification in clinical trials with censored data. *Comput Stat Data Anal.* 2006;50(5):1338-1355.

38. Hastie T, Tibshirani R, Friedman J. *The Elements of Statistical Learning.* New York, NY, USA: Springer New York Inc.; 2001.

39. Abu-jamous B, Fa R, Nandi AK. *Integrative Cluster Analysis in Bioinformatics.* Chichester, West Sussex, United Kingdom: John Wiley & Sons, Ltd.; 2015.

40. Lochner KA, Cox CS. Prevalence of multiple chronic conditions among Medicare beneficiaries, United States, 2010. *Preventing chronic disease.* 2013;10:E61.

41. Aryal S, Diaz-Guzman E, Mannino DM. Prevalence of COPD and comorbidity. *European Respiratory*
Monograph. 2013;59:1-12.
42. Baty F, Putora PM, Isenring B, Blum T, Brutsche M. Comorbidities and burden of COPD: a population based
case-control study. *PloS one.* 2013;8(5):e63285.
43. Moni MA, Lio P. Network-based analysis of comorbidities risk during an infection: SARS and HIV case studies.
BMC bioinformatics. 2014;15:333.
44. Cramer AO, Waldorp LJ, van der Maas HL, Borsboom D. Comorbidity: a network perspective. *The Behavioral*
and brain sciences. 2010;33(2-3):137-150; discussion 150-193.
45. Islam MM, Valderas JM, Yen L, Dawda P, Jowsey T, McRae IS. Multimorbidity and comorbidity of chronic
diseases among the senior Australians: prevalence and patterns. *PloS one.* 2014;9(1):e83783.
46. Folino F, Pizzuti C, Ventura M. A comorbidity network approach to predict disease risk. Proceedings of the
First international conference on Information technology in bio- and medical informatics; 2010; Bilbao, Spain.
47. Newman MEJ. *Networks: An Introduction.* Oxford, United Kingdom: Oxford University Press; 2010.
48. Treviño S, Nyberg A, Del Genio CI, Bassler KE. Fast and accurate determination of modularity and its effect
size. *Journal of Statistical Mechanics: Theory and Experiment.* 2015;2015(2):P02003.
49. Chauhan R, Ravi J, Datta P, et al. Reconstruction and topological characterization of the sigma factor
regulatory network of Mycobacterium tuberculosis. *Nature communications.* 2016;7:11062.
50. Bhavnani SK, Dang B, Penton R, et al. How High-Risk Comorbidities Co-Occur in Readmitted Patients With
Hip Fracture: Big Data Visual Analytical Approach. *JMIR Med Inform.* 2020;8(10):e13567.
51. Fruchterman T, Reingold E. Graph Drawing by Force-Directed Placement. *Software – Practice & Experience.*
1991;21(11):1129–1164.
52. Dang B, Chen T, Bassler KE, Bhavnani SK. ExplodeLayout: Enhancing the Comprehension of Large and
Dense Networks. AMIA Jt Summits Transl Sci Proc. ; 2016.
53. Bhavnani SK, Chen T, Ayyaswamy A, et al. Enabling Comprehension of Patient Subgroups and
Characteristics in Large Bipartite Networks: Implications for Precision Medicine. *Proceedings of AMIA Joint*
Summits on Translational Science. 2017:21-29.
54. Bhavnani SK, Eichinger F, Martini S, Saxman P, Jagadish HV, Kretzler M. Network analysis of genes
regulated in renal diseases: implications for a molecular-based classification. *BMC bioinformatics.* 2009;10
Suppl 9:S3.
55. Bhavnani SK, Bellala G, Ganesan A, et al. The nested structure of cancer symptoms. Implications for
analyzing co-occurrence and managing symptoms. *Methods of information in medicine.* 2010;49(6):581-591.
56. Bhavnani SK, Ganesan A, Hall T, et al. Discovering hidden relationships between renal diseases and
regulated genes through 3D network visualizations. *BMC research notes.* 2010;3:296.
57. Bhavnani SK, Victor S, Calhoun WJ, et al. How cytokines co-occur across asthma patients: from bipartite
network analysis to a molecular-based classification. *Journal of biomedical informatics.* 2011;44 Suppl
1:S24-30.
58. Bhavnani SK, Bellala G, Victor S, Bassler KE, Visweswaran S. The role of complementary bipartite visual
analytical representations in the analysis of SNPs: a case study in ancestral informative markers. *Journal of*
the American Medical Informatics Association: JAMIA. 2012;19(e1):e5-e12.
59. Bhavnani SK, Dang B, Bellala G, et al. Unlocking proteomic heterogeneity in complex diseases through
visual analytics. *Proteomics.* 2015;15(8):1405-1418.
60. Bhavnani SK, Dang B, Kilaru V, et al. Methylation differences reveal heterogeneity in preterm
pathophysiology: results from bipartite network analyses. *Journal of perinatal medicine.* 2018;46(5):509-521.
61. Bhavnani SK, Kummerfeld E, Zhang W, et al. Heterogeneity in COVID-19 Patients at Multiple Levels of
Granularity: From Biclusters to Clinical Interventions. *Proceedings of the American Medical Informatics*
Association Summits. 2021:112-121.
62. Patel SB, Nguyen NT. Creation of a Mapped, Machine-Readable Taxonomy to Facilitate Extraction of Social
Determinants of Health Data from Electronic Health Records. *Proceedings of AMIA Annual Symposium.*
2021;2021:959-968.
63. Fratelli C, Siqueira J, Silva C, Ferreira E, Silva I. 5HTTLPR Genetic Variant and Major Depressive Disorder:
A Review. *Genes (Basel).* 2020;11(11):1260.
64. Wurtman RJ. Genes, stress, and depression. *Metabolism: clinical and experimental.* 2005;54(5 Suppl 1):16-
19.
65. Allen J, Balfour R, Bell R, Marmot M. Social determinants of mental health. *International Review of Psychiatry.*
2014;26(4):392-407.

66. Busse D, Yim IS, Campos B, Marshburn CK. Discrimination and the HPA axis: current evidence and future
directions. *Journal of behavioral medicine.* 2017;40(4):539-552.
67. Decker A, Weaver R. Health and Social Determinants Associated With Delay of Health Care Among Rural
Older Adults. *Innovation in Aging.* 2021;5(Supplement_1):210-211.
68. Newton MF, Keirns CC, Cunningham R, Hayward RA, Stanley R. Uninsured adults presenting to US
emergency departments: assumptions vs data. *Jama.* 2008;300(16):1914-1924.
69. Bhavnani SK, Zhang W, Visweswaran S, Raji M, Kuo YF. A Framework for Modeling and Interpreting Patient
Subgroups Applied to Hospital Readmission: Visual Analytical Approach. *JMIR Med Inform.*
2022;10(12):e37239.
70. van der Wal WM, Geskus RB. ipw: An R Package for Inverse Probability Weighting. *Journal of Statistical*
Software. 2011;43(13):1 - 23.
71. Thoemmes F, Ong AD. A Primer on Inverse Probability of Treatment Weighting and Marginal Structural
Models. *Emerging Adulthood.* 2016;4(1):40-59.
72. Lee YH, Liu Z, Fatori D, et al. Association of Everyday Discrimination With Depressive Symptoms and
Suicidal Ideation During the COVID-19 Pandemic in the All of Us Research Program. *JAMA psychiatry.*
2022;79(9):898-906.
73. Chen T, Zhang W, Bhavnani S. BipartiteModularityMaximization: CRAN R Package. 2022; https://cran.r-
project.org/web/packages/BipartiteModularityMaximization/index.html, 2023.
74. Bhavnani   SK,   Zhang   W.   ExplodeLayout: CRAN   R   Package.   2022;   https://cran.r-
project.org/web/packages/ExplodeLayout/index.html, 2023.
75. DataScienceMeta.   CRAN   R   Packages   by   Number   of   Downloads.
http://www.datasciencemeta.com/rpackages, 2023.
76. Bhavnani S, Zhang W, Bao D, et al. The Impact of Critical Social Determinants of Health on Personal Medical
Decisions: Analysis of Older Americans in All of Us. (in press).
77. Bhavnani S, Zhang W, Bao D, Hatch S, Reistetter T, Downer B. Generalizable Machine Learning Methods
for Subtyping Individuals on National Health Databases: Case Studies Using Data from HRS, N3C, and All
of Us. (in press).
78. Dhillon IS, Sra S. Generalized nonnegative matrix approximations with Bregman divergences. Proceedings
of the 18th International Conference on Neural Information Processing Systems; 2005; Vancouver, British
Columbia, Canada.
79. Dilsizian ME, Siegel EL. Machine Meets Biology: a Primer on Artificial Intelligence in Cardiology and Cardiac
Imaging. *Current cardiology reports.* 2018;20(12):139.
80. Bhavnani SK, Visweswaran S, Divekar R, Brasier AR. Towards Team-Centered Informatics: Accelerating
Innovation in Multidisciplinary Scientific Teams Through Visual Analytics. *The Journal of Applied Behavioral*
Science. 2018:0021886318794606.
81. Wooten KC, Calhoun WJ, Bhavnani S, Rose RM, Ameredes B, Brasier AR. Evolution of Multidisciplinary
Translational Teams (MTTs): Insights for Accelerating Translational Innovations. *Clinical and translational*
science. 2015;8(5):542-552.

Appendix A: Description of Bipartite Network Analysis

Fig. 1. The distinction between a unipartite network (A), a bipartite network (B), and how the latter can be used to identify biclusters of participants and SDoH (C).

A network consists of nodes and edges; nodes represent one or more types of entities (e.g., participants or
SDoH), and edges between the nodes represent a specific relationship between the entities. Figure 1A shows a
unipartite network where nodes are the same type (typically used to analyze co-occurrence of comorbidities[46]).
In contrast, Figure 1B shows a bipartite network where nodes are of two types, and edges exist only between
different types such as between participants (circles) and SDoH (triangles).

Bipartite network analysis takes as input any dataset such as *All of Us* participants and their SDoH, and
automatically outputs a quantitative and visual description of biclusters (containing both participant subgroups,
and their frequently co-occurring SDoH). The quantitative output provides the number, size, and statistical
significance of the biclusters,[48-50] and the visual output displays the quantitative information of the biclusters
through a network visualization.[51-53] Bipartite network analysis therefore enables (1) the automatic identification
of biclusters and their significance, and (2) the visualization of the biclusters critical for their clinical interpretability
including labeling the subtypes, inferring potential mechanisms that precipitate adverse outcomes in each
subtype, and designing targeted interventions to prevent them. Furthermore, the characteristics (e.g., outcomes
and covariates) of participants in a subtype can be used to measure the risk of a subtype for an adverse outcome
when compared to a reference group (e.g., a control group or another subtype), and therefore enables the
integration of multiple data types. Finally, the biclusters can be used to develop classifiers for classifying a new
participant into one or more of the subtypes, and developing a predictive model that uses those subtype
membership for measuring the risk of an adverse outcome for that new participant.[69]

Appendix B: SNOMED Codes Related to SDoH, and their Use in the Electronic Health Records of
Participants in *All of Us.*
879

880
881

Appendix C: All SDoH questions identified from four *All of Us* surveys (Column-1), abbreviated (Column-2), negatively labeled and reverse coded (red text in Column-2), categorized into the five *Healthy People 2030* domains (shown by the five colors), and further categorized into subdomains (Column-5). *Delayed Medical Care* was used as an outcome.

Survey Name	Question/Field	Abbreviated and Negatively Labeled	SDoH Domain (healthy.gov)	Sub Domain (Domain Knowledge)
Social Determinants of Health	People around here are willing to help their neighbors	**Neighborhood people unwilling to help**	Social & community context	Neighborhood cohesion
Social Determinants of Health	People in my neighborhood generally get along with each other	**Neighborhood people do not get along**	Social & community context	Neighborhood cohesion
Social Determinants of Health	People in my neighborhood can be trusted	**Neighborhood people cannot be trusted**	Social & community context	Neighborhood cohesion
Social Determinants of Health	People in my neighborhood share the same values	**Neighborhood people do not share values**	Social & community context	Neighborhood cohesion
Social Determinants of Health	I'm always having trouble with my neighbors	Neighborhood people troublesome	Social & community context	Neighborhood cohesion
Social Determinants of Health	In my neighborhood, people watch out for each other	**Neighborhood people do not watch out**	Social & community context	Neighborhood cohesion
Social Determinants of Health	Someone to help you if you were confined to bed	**No one to help out of bed**	Social & community context	Supportive relationships
Social Determinants of Health	Someone to take you to the doctor if you need it	**No help for doctor visit**	Social & community context	Supportive relationships
Social Determinants of Health	Someone to prepare your meals if you were unable to do it yourself	**No one help with meal prep**	Social & community context	Supportive relationships
Social Determinants of Health	Someone to help with daily chores if you were sick	**No help when sick**	Social & community context	Supportive relationships
Social Determinants of Health	Someone to have a good time with	**No one to have good time with**	Social & community context	Supportive relationships
Social Determinants of Health	Someone to turn to for suggestions about how to deal with a personal problem	**No one to suggest problem solutions**	Social & community context	Supportive relationships
Social Determinants of Health	Someone who understands your problems	**No one who understands problems**	Social & community context	Supportive relationships
Overall Health	How often do you have someone help you read health-related materials?	**No one to help read health materials**	Social & community context	Supportive relationships
Social Determinants of Health	Someone to love and make you feel wanted	**No one to make you feel wanted**	Social & community context	Supportive relationships
Social Determinants of Health	I lack companionship	Lack companionship frequenncy	Social & community context	Relationships with others
Social Determinants of Health	There is no one I can turn to	No one to turn to frequency	Social & community context	Relationships with others
Social Determinants of Health	You are treated with less courtesy than other people are	Day to day less courtesy frequency	Social & community context	Discrimination in everyday life
Social Determinants of Health	you are treated with less respect than other people are	Day to day less respect frequency	Social & community context	Discrimination in everyday life
Social Determinants of Health	You receive poorer service than other people at restaurants or stores	Day to day poorer service frequency	Social & community context	Discrimination in everyday life
Social Determinants of Health	People act as if they think you are not smart	Others think you as less smart	Social & community context	Discrimination in everyday life
Social Determinants of Health	People act as if they are afraid of you	Others are afraid of you	Social & community context	Discrimination in everyday life
Social Determinants of Health	People act as if they think you are dishonest	Others think you are dishonest	Social & community context	Discrimination in everyday life
Social Determinants of Health	People act as if they're better than you are	Others think they're better than you	Social & community context	Discrimination in everyday life
Social Determinants of Health	You are called names or insulted	Called names or insulted frequency	Social & community context	Discrimination in everyday life
Social Determinants of Health	You are threatened or harassed	Threatened or harrassed frequency	Social & community context	Discrimination in everyday life
Social Determinants of Health	Do you speak a language other than English at home?	**Not bilingual**	Social & community context	Language
Social Determinants of Health	Since you speak a language other than English at home, how well would you say you sp	English verbal proficiency	Social & community context	Language
The Basics	In what country were you born?	**US Born**	Social & community context	Demographics
The Basics	Which categories describe you? Select all that apply. Note, you may select more than	**Non-White Race**	Social & community context	Demographics
The Basics	What was your biological sex assigned at birth?	**Nonbinary Sex**	Social & community context	Demographics
The Basics	What terms best express how you describe your gender identity (check all that apply)	**Nonbinary Gender Identity**	Social & community context	Demographics
The Basics	Which of th following best represents how you think of yourself?	**Non-heterosexual**	Social & community context	Demographics
The Basics	What is your current marital status?	**Unmarried**	Social & community context	Demographics
The Basics	Not including yourself, how many other people live at home with you?	**Single Household**	Social & community context	Demographics
The Basics	Think of other people who live with you. How many are under the age of 18 years?	**Housing dependents**	Social & community context	Demographics
The Basics	Have you ever served on active duty in the United States Armed forces?	**Active Duty Status**	Social & community context	Demographics
Social Determinants of Health	There is alot of graffiti in my neighborhood	Neighborhood has a lot of graffiti	Neighborhood & built environment	Neighborhood characteristics
Social Determinants of Health	My neighborhood is noisy	Neighborhood is noisy	Neighborhood & built environment	Neighborhood characteristics
Social Determinants of Health	Vandalism is common in my neighborhood	Neighborhood vandalism is common	Neighborhood & built environment	Neighborhood characteristics
Social Determinants of Health	There are lot of abandoned building in my neighborhood	Neighborhood has many abandoned buil	Neighborhood & built environment	Neighborhood characteristics
Social Determinants of Health	My neighborhood is clean	**Neighborhood not clean**	Neighborhood & built environment	Neighborhood characteristics
Social Determinants of Health	People in my neighborhood take good care of their houses and apartments	**Neighborhood is not well maintained**	Neighborhood & built environment	Neighborhood characteristics
Social Determinants of Health	There are too many people hanging around on the streets near my home	Neighborhood has too many loiters	Neighborhood & built environment	Neighborhood characteristics
Social Determinants of Health	There is a lot of crime in my neighborhood	Neighborhood has a lot of crime	Neighborhood & built environment	Neighborhood characteristics
Social Determinants of Health	There is too much drug use in my neighborhood	Neighborhood has too much drug use	Neighborhood & built environment	Neighborhood characteristics
Social Determinants of Health	There is too much alcohol use in my neighborhood	Neighborhood has too much alcohol	Neighborhood & built environment	Neighborhood characteristics
Social Determinants of Health	My neighborhood is safe	**Neighborhood is not safe**	Neighborhood & built environment	Neighborhood characteristics
Social Determinants of Health	What is the main type of housing in your neighborhood?	**Neighborhood mostly apartments**	Neighborhood & built environment	Neighborhood characteristics

Section	Question	Label	Domain	Subdomain
Social Determinants of Health	Many shops, stores, markets or other places to buy things I need are within easy walkin...	No shopping resource in walking proxim...	Neighborhood & built environment	Neighborhood characteristics
Social Determinants of Health	It is within a 10-15 minutes walk to a transit stop from home	No transportation in walking proximity	Neighborhood & built environment	Neighborhood characteristics
Social Determinants of Health	There are sidewalks on most of the streets in my neighborhood	Neighborhood has no sidewalks	Neighborhood & built environment	Neighborhood characteristics
Social Determinants of Health	There are facilities to bicycle in or near my neighborhood (e.g., special lanes, trails, pat...	Neighborhood is not bike friendly	Neighborhood & built environment	Neighborhood characteristics
Social Determinants of Health	My neighborhood has several free or low-cost recreation facilities (e.g., parks, pools, p...	Neighborhood has no recreation spaces	Neighborhood & built environment	Neighborhood characteristics
Social Determinants of Health	The crime rate in my neighborhood makes it unsafe to go on walks at night	Neighborhood is unsafe at night	Neighborhood & built environment	Neighborhood characteristics
Social Determinants of Health	The crime rate in my neighborhood makes it unsafe to go on walks during the day	Neighborhood is unsafe in the day	Neighborhood & built environment	Neighborhood characteristics
Social Determinants of Health	Think about the place you live. Do you have problems with any of the following (check...	Home has problems	Neighborhood & built environment	Housing
Social Determinants of Health	Within the past 12 months, we worried whether our food would run out before we got...	Food insecurity	Economic stability	Food security
Social Determinants of Health	Within the past 12 months, the food we bought just didn't last and we didn't have mon...	Food inadequate and insecure	Economic stability	Food security
Social Determinants of Health	In the last 12 months, how many times have you or your family moved from one home...	Moved home	Economic stability	Housing
The Basics	Do you own or rent the place where you live?	Lack of Home Ownership/Rent	Economic stability	Housing
The Basics	Where are you currently living?	Current Living Situation	Economic stability	Housing
The Basics	How many years have you lived at your current address?	Lived Less than 1 Year	Economic stability	Housing
The Basics	In the past 6 months, have you been worried or concerned about NOT having a place t...	Unstable Housing	Economic stability	Housing
The Basics	What is your current employment status? Please select 1 or more of these categories.	Unemployed	Economic stability	Employment
The Basics	What is your annual household income from all sources?	Poverty Income	Economic stability	Income
Overall Health	How often do you have problems learning about your medical condition because of dif...	Health Illiteracy	Education access and quality	Literacy
The Basics	What is the highest grade or year of school you completed?	Lack college education	Education access and quality	Education attainment
Health Care Access and Utilization	During the past 12 months, were you told by a health care provider or doctor's office t...	Health insurance not accepted	Health care access and quality	Healthcare coverage
Health Care Access and Utilization	In regard to your health insurance care coverage, how does it compare to a year ago?	Health insurance coverage worse	Health care access and quality	Healthcare coverage
Health Care Access and Utilization	Is there a place that you USUALLY go to when you are sick or need advice about your h...	No place for health advice	Health care access and quality	Healthcare coverage
Health Care Access and Utilization	If yes, what kind of place do you go to most often?	No kind of doctor place	Health care access and quality	Healthcare coverage
Health Care Access and Utilization	About how long has it been since you last saw or talked to a doctor or other health car...	More than 1 year since last spoken to h...	Health care access and quality	Healthcare coverage
Health Care Access and Utilization	How often were you treated with respect by your doctors or health care providers?	Healthcare discrimination less respect	Health care access and quality	Interactions with providers
Health Care Access and Utilization	How ofen did you doctor or health care providers ask for your opinions or beliefs abou...	Not asked for opinion by provider	Health care access and quality	Interactions with providers
Health Care Access and Utilization	How often did you doctors or health care providers tell or give you information about t...	Advice not easy to understand	Health care access and quality	Interactions with providers
Health Care Access and Utilization	Have you delayed getting care in the past 12 months because you didn't have transport...	Delayed care d/t Transportation	Health care access and quality	Delayed medical care
Health Care Access and Utilization	Have you delayed getting care in the past 12 months because you live in a rural area w...	Delayed care d/t rural area	Health care access and quality	Delayed medical care
Health Care Access and Utilization	Have you delayed getting care in the past 12 months because you were nervous about...	Delayed care d/t nervousness	Health care access and quality	Delayed medical care
Health Care Access and Utilization	Have you delayed getting care in the past 12 months because you couldn't get time off...	Delayed care d/t work	Health care access and quality	Delayed medical care
Health Care Access and Utilization	Have you delayed getting care in the past 12 months because you couldn't get child car...	Delayed care d/t childcare	Health care access and quality	Delayed medical care
Health Care Access and Utilization	Have you delayed getting care in the past 12 months because you couldn't afford the c...	Delayed care d/t copay	Health care access and quality	Delayed medical care
Health Care Access and Utilization	Have you delayed getting care in the past 12 months because you provide care to an a...	Delayed care d/t elderly care	Health care access and quality	Delayed medical care
Health Care Access and Utilization	You had to pay out of pocket for some or all of the procedure?	Delayed care d/t out of pocket	Health care access and quality	Delayed medical care
Health Care Access and Utilization	Have you delayed getting care in the past 12 months because your deductible was too...	Delayed care d/t deductible	Health care access and quality	Delayed medical care
Health Care Access and Utilization	During the past 12 months, was there any time when you needed prescription medicine...	Can't afford Rx	Health care access and quality	Afford medical care
Health Care Access and Utilization	During the past 12 months, was there any time when you needed mental health care o...	Can't afford mental health	Health care access and quality	Afford medical care
Health Care Access and Utilization	During the past 12 months, was there any time when you needed emergency care but c...	Can't afford emergency care	Health care access and quality	Afford medical care
Health Care Access and Utilization	During the past 12 months, was there any time when you needed dental care but didn't...	Can't afford dental care	Health care access and quality	Afford medical care
Health Care Access and Utilization	During the past 12 months, was there any time when you needed eyeglasses but didn't...	Can't afford eyeglasses	Health care access and quality	Afford medical care
Health Care Access and Utilization	During the past 12 months, was there any time when you needed to see a regular docto...	Can't afford healthcare provider	Health care access and quality	Afford medical care
Health Care Access and Utilization	During the past 12 months, was there any time when you needed to see a specialist but...	Can't afford specialist	Health care access and quality	Afford medical care
Health Care Access and Utilization	During the past 12 months, was there any time when you needed follow-up care but di...	Can't afford followup care	Health care access and quality	Afford medical care
Health Care Access and Utilization	If you get sick or hav an accident, how worried are you that you will be able to pay you...	Worried about not paying for care	Health care access and quality	Afford medical care
Health Care Access and Utilization	During the past 12 months, you skipped medication doses to save money?	Skipped meds to save money	Health care access and quality	Afford medical care
Health Care Access and Utilization	During the past 12 months, you took less medicine to save money?	Took less meds to save money	Health care access and quality	Afford medical care
Health Care Access and Utilization	During the past 12 months, you delayed filling a prescription to save money?	Delayed filling Rx to save money	Health care access and quality	Afford medical care
Health Care Access and Utilization	During th past 12 months, you asked your doctor for a lower cost medication to save m...	Lower cost Rx to save money	Health care access and quality	Afford medical care
Health Care Access and Utilization	During the past 12 months, you bought prescription drugs from aother country to save...	Bought Rx from another country to save	Health care access and quality	Afford medical care
Health Care Access and Utilization	During the past 12 months, you used alternative therapies to save money?	Alternative therapy to save money	Health care access and quality	Afford medical care
Health Care Access and Utilization	How important is it to you that your doctors or health care providers understand or ar...	Health provider race religion similar imp...	Health care access and quality	Provider characteristics
Health Care Access and Utilization	How often were you able to see doctors or health care providers who were similar to ...	Health provider race religion dissimilar f...	Health care access and quality	Provider characteristics
Health Care Access and Utilization	How often have you either delayed or not gone to see doctors or health care providers...	Health provider race religion delayed car...	Health care access and quality	Provider characteristics
Social Determinants of Health	When you go to a doctor's office or other health care provider, how often are you trea...	Healthcare discrimination less courtesy	Health care access and quality	Descrimination in medical settings
Social Determinants of Health	When you go to a doctor's office or other health care provider, how often are you trea...	Healthcare discrimination less respect	Health care access and quality	Descrimination in medical settings
Social Determinants of Health	When you go to a doctor's office or other health care provider, how often do you rece...	Healthcare discrimination poorer service	Health care access and quality	Descrimination in medical settings
Social Determinants of Health	When you go to a doctor's office or other health care provider, how often does a doct...	Healthcare discrimination intellect	Health care access and quality	Descrimination in medical settings
Social Determinants of Health	When you go to a doctor's office or other health care provider, how does a doctor or r...	Healthcare discrimination fear	Health care access and quality	Descrimination in medical settings
Social Determinants of Health	When you go to a doctor's office or other health care provider, how often does a doct...	Healthcare discrimination inferiority	Health care access and quality	Descrimination in medical settings
Social Determinants of Health	When you go to a doctor's office or other health care provider, how often do you feel...	Healthcare discrimination not listened to	Health care access and quality	Descrimination in medical settings

Appendix D: Condition-Specific Cohort Extraction for type II diabetes (T2DM), breast cancer, and coronary
artery disease (CAD).

Figure 1. Inclusion and exclusion criteria for selecting three condition-specific cohorts.

Appendix E: Inverse Probability Weighting

We found significant differences in the demographic characteristics between the total *All of Us* sample, and the
12,913 participants with valid answers for all SDoH questions. To account for potential sample selection bias,
we calculated inverse probability weights (IPW) using the ipwpoint function in the R package ipw.[66] This function
uses a logistic regression model to estimate the predicted probability of having valid responses on all SDoH
variables based on age, sex, race, ethnicity, being born in the United States, currently employed, having a college
degree or higher, health insurance, owning a home, and being married. We stabilized the weights according to
the observed probability of being in our cohort.