Wildfire smoke PM$_{2.5}$ and mortality in the contiguous United States

Yiqun Ma[1,2], Emma Zang[3], Yang Liu[4], Yuan Lu[5,6], Harlan M. Krumholz[5,6], Michelle L. Bell[7], Kai Chen[1,2,*]

[1]Department of Environmental Health Sciences, Yale School of Public Health, New Haven, CT, USA
[2]Yale Center on Climate Change and Health, Yale School of Public Health, New Haven, CT, USA
[3]Department of Sociology, Yale University, New Haven, CT, USA
[4]Department of Environmental Health, Rollins School of Public Health, Emory University, Atlanta, GA, USA
[5]Center for Outcomes Research and Evaluation, Yale New Haven Hospital, New Haven, CT, USA
[6]Section of Cardiovascular Medicine, Department of Medicine, Yale School of Medicine, New Haven, CT, USA
[7]School of the Environment, Yale University, New Haven, CT, USA

*Corresponding author: Kai Chen
Email: kai.chen@yale.edu

Author Contributions: Y.M. conducted formal analyses and drafted the manuscript. K.C. conceived of and supervised the conduct of this study and edited the manuscript. E.Z. applied and managed the mortality dataset, supervised the statistical analysis, and revised the manuscript. Y.L., Y.L., H.M.K., and M.L.B. contributed to the interpretation of results and manuscript revision. All authors reviewed and approved the final version of this manuscript.

Competing Interest Statement: The authors declare no conflict of interests.

Classification: Biological Sciences, Environmental Sciences

Keywords: wildfire, mortality, fine particulate matter, United States

Abstract

Despite the growing evidence on the health effects of wildfire smoke in the western U.S., the nationwide mortality risk and burden attributable to wildfire smoke fine particles ($PM_{2.5}$) remain unclear. This study aims to investigate the association between wildfire smoke $PM_{2.5}$ and mortality from all causes, cardiovascular diseases, respiratory diseases, and mental disorders, and calculate the corresponding attributable mortality burden in all 3,108 counties in the contiguous U.S., 2006–2016. Monthly county-level mortality counts were collected from National Center for Health Statistics. Wildfire smoke $PM_{2.5}$ concentration was derived from a 10×10 km^2 resolution spatiotemporal model. Controlling for non-smoke $PM_{2.5}$, air temperature, and unmeasured spatial and temporal confounders, we found that a 1 µg/m^3 increase in smoke $PM_{2.5}$ was significantly associated with an increase of 0.14% (95% confidence interval [CI]: 0.11%, 0.17%) in all-cause mortality, 0.13% (95% CI: 0.08%, 0.18%) in cardiovascular mortality, 0.16% (95% CI: 0.07%, 0.25%) in respiratory mortality, and 1.08% (95% CI: 0.93%, 1.23%) in mental disorder mortality. Smoke $PM_{2.5}$ contributed to approximately 1,141 all-cause deaths/year (95% CI: 893, 1,388) in the contiguous U.S., of which over three-fourths were from cardiovascular, respiratory, and mental causes. We found a higher vulnerability among males than females, people aged 0 to 64 years than those ≥ 65 years, and racial/ethnic minorities than non-Hispanic White people. Mild droughts were found to enhance the association between smoke $PM_{2.5}$ and mortality. Our results indicate that wildfire smoke $PM_{2.5}$ harms both physical and mental health, which suggests the need for more effective wildfire mitigation strategies and public health responses in the U.S.

Significance Statement

Wildfires activities have significantly increased in the United States in recent decades. Smoke

pollutants emitted by wildfires, particularly $PM_{2.5}$, can bring adverse health effects. However, the

nationwide wildfire smoke $PM_{2.5}$-related mortality risk and burden remain unclear. Utilizing wildfire

smoke $PM_{2.5}$ and mortality data in the contiguous U.S. from 2006 to 2016, we found significant

associations between smoke $PM_{2.5}$ and increased mortality risks from all causes, cardiovascular

diseases, respiratory diseases, and mental disorders. Each year, smoke $PM_{2.5}$ contributed to an

estimated over one thousand deaths in the U.S., indicating a great number of indirect deaths

brought by wildfires that official tolls could not capture. This study demonstrates the detrimental

effects of wildfire smoke $PM_{2.5}$ on both physical and mental health and calls for more effective

wildfire prevention and mitigation policies in the U.S.

Main Text

Introduction

Wildfire is a growing public health concern in the United States. The country has witnessed a

marked increase in the area affected by wildfires over the past few decades, with the burned area

roughly quadrupling (1). In recent years, wildfire contributed to up to 25% of fine particulate

matter ($PM_{2.5}$) across the U.S., and up to half in some Western regions (1). Driven by the

warming climate, the prevalence, frequency, and intensity of wildfire activities are expected to

increase in the future (2). Among the various air pollutants emitted by wildfires, $PM_{2.5}$ has

received great attention due to its ability to deeply penetrate the respiratory system and

demonstrated links to public health (3). In addition, because of different chemical composition and

smaller particle size, studies suggested that the toxicity of wildfire smoke $PM_{2.5}$ could be different

from urban background $PM_{2.5}$, potentially more lethal (4, 5). Previous studies on the health effects

of wildfire smoke mostly focused on the western U.S., where the majority of large fires occurred

(5-7). However, the pollutants from wildfire smoke can travel long distances from the source,

potentially affecting human health thousands of kilometers away outside the West (8).

In recent years, studies consistently reported a positive relationship between wildfire

smoke exposure and all-cause mortality (7, 9, 10). Among specific causes of mortality, respiratory

and cardiovascular diseases received most attention, with growing studies linking human

exposure to wildfire smoke with increased risks of respiratory and cardiovascular mortality (9-11).

However, in addition to the physical health harms, mental health can also be potentially

threatened by wildfire smoke. Some studies have linked wildfires to mental outcomes, such as

anxiety, depression, poor sleep quality, and posttraumatic stress disorder (7, 12, 13), but the

relationship between wildfire smoke and mental disorder mortality was rarely analyzed in previous

studies.

Furthermore, the effects of wildfire smoke $PM_{2.5}$ can be heterogenous among population

subgroups due to physiological, behavioral, and socioeconomic factors (14). Previous studies

indicated that demographic factors such as sex, age and race/ethnicity may modify the

association between smoke PM$_{2.5}$ and health outcomes (9, 14, 15). However, existing research

generated contradictory findings and is insufficient to identify specific subpopulations that are

more susceptible to wildfire smoke exposure (7). For example, a study in Australia reported that

people ≥ 65 years old had higher bushfire smoke-related rates of respiratory hospitalizations

compared to their younger counterparts (16), but another study in North Carolina found that the

association between peat bog wildfire smoke and emergency department visits for chronic

obstructive pulmonary disease, pneumonia, and acute bronchitis were higher among people < 65

years old than those older (17). Greater understanding of susceptibility to wildfire exposure

among specific subgroups is needed to help inform wildfire mitigation policies and address

environmental justice issues.

Besides demographic factors, drought, which often co-occurs with wildfires, may also

serve as an effect modifier. Drought is a complex phenomenon resulting from precipitation

deficiency, high temperature, low relative humidity, and other meteorological and hydrological

factors (18). Studies have shown stronger mortality effects of air pollution at higher air

temperatures and lower humidity levels (19-21), but little is known about whether drought, the

manifestation of the complicated interactions among multiple environmental factors, can enhance

the association between wildfire smoke PM$_{2.5}$ and mortality.

Constrained by a lack of nationwide validated data on pollutant concentrations

attributable to wildfire smoke, many previous studies on the impact of wildfire smoke on health

outcomes have been focusing on episodes with high wildfire smoke exposure (or smoke wave)

using binary measures of smoke concentrations (22). Recently, a machine learning model was

developed to estimate daily wildfire smoke PM$_{2.5}$ concentrations for the contiguous U.S., using a

combination of meteorological factors, fire variables, aerosol measurements, and land use and

elevation data (23). This new, high-resolution wildfire smoke PM$_{2.5}$ dataset (10×10 km^2) enabled

us to further examine the impact of wildfire smoke over a full range – from the more common,

low-level smoke concentrations to the increasingly frequent extremely high concentrations.

Utilizing the nationwide wildfire smoke PM$_{2.5}$ and mortality data, this study aimed to (a)

estimate the associations of smoke $PM_{2.5}$ exposure with county-level all-cause mortality and

mortality of cardiovascular and respiratory diseases and mental disorders, (b) calculate the

attributable cause-specific mortality burden from 2006 to 2016, and (c) examine whether the

smoke $PM_{2.5}$ related mortality risk varies among people of different sex, age, and racial/ethnicity

groups and different levels of drought.

Results

Description of wildfire smoke PM$_{2.5}$ and mortality in the contiguous U.S.

From 2006 to 2016, all 3,108 counties in the contiguous U.S. had experienced some amount of wildfire smoke PM$_{2.5}$, with the western, north central, and southeastern counties being exposed to higher monthly average concentrations than other regions (**Fig. 1A**). Although wildfire smoke PM$_{2.5}$ concentrations were generally higher in summer months than in winter months, the average concentration remained above zero for the majority of the study period (**Fig. S1**). The average concentration of smoke PM$_{2.5}$ across all county-months was 0.39 µg/m^3, but the highest smoke PM$_{2.5}$ concentration could exceed 70 µg/m^3 (e.g., 70.95 µg/m^3 levels observed in September 2012 in Lemhi County, Idaho). A total of 27,812,837 deaths were included in this study, including 8,826,110 deaths from cardiovascular diseases, 2,713,137 deaths from respiratory diseases, and 1,364,807 deaths from mental and behavioral disorders (hereafter referred to as "mental disorders"; **Table S1**).

Wildfire smoke PM$_{2.5}$ related cause-specific mortality risk and mortality burden

The association between smoke PM$_{2.5}$ and all-cause mortality was found to be near-linear in the linearity test (**Fig. S2**), so we used a linear term to model this relationship in the main model. Using a two-way fixed effects (TWFE) model to control for air temperature, non-smoke PM$_{2.5}$, and unmeasured spatial and temporal confounders, we found that a 1 µg/m^3 increase in smoke PM$_{2.5}$ concentration was significantly associated with a 0.14% (95% confidence interval [CI]: 0.11%, 0.17%) increase in all-cause mortality (**Fig. 2A**). On average, approximately 1,141 all-cause deaths (95% CI: 893, 1,388) were attributable to smoke PM$_{2.5}$ in the contiguous U.S. per year. The spatial distribution of this attributable burden was generally consistent with the distribution of smoke PM$_{2.5}$ concentration (**Fig. 1B**).

As shown in **Fig. 2A**, for cause-specific mortality, a 1 µg/m^3 increase in smoke PM$_{2.5}$ was significantly associated with an increase of 0.13% (95% CI: 0.08%, 0.18%) in cardiovascular disease mortality, 0.16% (95% CI: 0.07%, 0.25%) in respiratory disease mortality, and 1.08%

(95% CI: 0.93%, 1.23%) in mental disorder mortality. Each year, smoke $PM_{2.5}$ contributed

approximately 333 deaths from cardiovascular diseases (95% CI: 207, 459), 118 deaths from

respiratory diseases (95% CI: 48, 187), and 417 deaths from mental disorders (95% CI: 360,

473). Among the total wildfire smoke $PM_{2.5}$-attributable deaths, approximately 76.07% were from

these three specific causes (**Fig. 2B**).

Wildfire smoke $PM_{2.5}$ related all-cause mortality risk in different subgroups

The association between smoke $PM_{2.5}$ and all-cause mortality varies across different sex,

age, and race/ethnicity groups (**Fig. 3**). The estimated percent change in all-cause mortality per 1

$\mu g/m^3$ increase in smoke $PM_{2.5}$ was significantly higher in males (0.18%; 95% CI: 0.15%, 0.22%)

compared to females (0.10%; 95% CI: 0.06%, 0.14%; *p* value of difference: 0.002). Compared to

people aged 65 years and above (0.17%; 95% CI: 0.14%, 0.21%), people aged 0 to 64 years

(0.31%; 95% CI: 0.26%, 0.36%) had significantly higher smoke $PM_{2.5}$ related all-cause mortality

risk (*p* value of difference: <0.001). In addition, the between-group difference was particularly

evident across race/ethnicity, with non-Hispanic Black people (0.78%; 95% CI: 0.72%, 0.84%)

and Hispanic people (0.69%; 95% CI: 0.64%, 0.74%) significantly more vulnerable than non-

Hispanic White people (0.12%; 95% CI: 0.08%, 0.16%; *p* values of difference: < 0.001).

Effect modification by drought

From 2006 to 2016, 23.28% of the county-months had mild droughts and 8.68% had

severe droughts, with western and southern counties being exposed most. **Fig. 4A & B** display

the overlaps of the distribution of mild and severe droughts and wildfire smoke $PM_{2.5}$ in

contiguous U.S. counties, indicating the prevalent co-occurrence of these two conditions.

Specifically, the western and southern counties experienced the most months with both mild

drought and smoke $PM_{2.5}$, while severe drought co-occurred with smoke $PM_{2.5}$ mostly in southern

U.S. counties in Texas, Louisiana, and Florida.

We observed a significant effect modification by drought for the association between

smoke PM$_{2.5}$ and all-cause mortality (**Fig. 4C**). Compared to county-months with no drought

(0.10%; 95% CI: 0.07%, 0.14%), the estimated percent change in all-cause mortality per 1 µg/m^3

increase in smoke PM$_{2.5}$ became significantly higher during mild droughts (0.50%; 95% CI:

0.42%, 0.57%; *p* value of difference: < 0.001). No significant association was observed when

there were severe droughts.

Sensitivity analyses

Sensitivity analyses showed that our results generally remained robust when additionally

adjusting for dew point temperature and nitrogen dioxide (NO$_2$) and ozone (O$_3$), not adjusting for

non-smoke PM$_{2.5}$, using alternative numbers of degree of freedom for air temperature, and

restricting the analysis to counties that have high smoke PM$_{2.5}$ exposure (≥ 20 µg/m^3). The

association between smoke PM$_{2.5}$ and all-cause mortality tend to be the highest during the

current month and decreasing over the later one and two months.

Discussion

To the best of our knowledge, this is the first study evaluating the wildfire smoke $PM_{2.5}$

related cause-specific mortality risk and attributable mortality burden for all ages in the whole

contiguous U.S. Using 11-year full spatial and temporal coverage data, we found that wildfire

smoke $PM_{2.5}$ exposure was significantly associated with increased mortality risks of all causes,

cardiovascular diseases, respiratory diseases, and mental disorders. Each year, smoke $PM_{2.5}$

contributed to an estimated more than one thousand all-cause deaths in the contiguous U.S., of

which over three-fourths were from cardiovascular, respiratory, and mental causes. In addition,

males, people aged 0 to 64, racial/ethnic minorities, and those living in mild drought conditions

appeared to be more sensitive to smoke $PM_{2.5}$ exposure.

Most previous studies have explored the relationship between wildfire and health

outcomes by comparing periods with no fire and periods during or after fire events, or comparing

regions affected by wildfire smoke and unaffected regions (22), which generated effect estimates

that are difficult to directly compare to our estimates. Only a few studies have isolated wildfire-

specific $PM_{2.5}$ from other sources, and used a continuous variable for smoke $PM_{2.5}$ concentrations

instead of a binary variable to estimate the health effects of smoke $PM_{2.5}$ (9, 10). Our findings on

the associations between smoke $PM_{2.5}$ and all-cause, cardiovascular, and respiratory mortality

are consistent with these studies in terms of both magnitude and direction of the estimates. For

example, using a two-stage time-series analysis, a recent multi-country study based on 749 cities

reported that a 1 $\mu g/m^3$ increase in wildfire-related $PM_{2.5}$ associated with an increase of 0.19%,

0.17%, and 0.19% in all-cause, cardiovascular, and respiratory mortality, respectively (10).

Despite the different statistical methods and data sources used in our study and this global study,

the estimated effect estimates were roughly similar.

In addition to cardiovascular and respiratory mortality, we found a strong association

between smoke $PM_{2.5}$ and mortality from mental disorders. In a scoping review, the authors

proposed a multi-level conceptual framework for understanding the pathways connecting wildfire

smoke with mental health and well-being, which includes the interaction among individual, social

and community networks, living and working conditions, and ecological levels (12). However,

empirical evidence on this topic is still limited. More high-quality studies assessing smoke

exposure independently of other wildfire-related experiences and exploring the specific sub-

categories of mental disorders are needed to further explain the observed high smoke $PM_{2.5}$

related mortality risk from mental disorders.

We reported a higher vulnerability to smoke $PM_{2.5}$ among males than females, people

aged 0 to 64 than older adults, and non-Hispanic Black and Hispanic people than non-Hispanic

White people. Previous wildfire studies on the most vulnerable subpopulations have been

inconclusive (7, 14). The observed high risks among males and younger population may be

related to occupational and behavioral factors. For example, wildland firefighters, mostly non-

elderly males, are at increased risk of lung cancer and cardiovascular disease mortality

regardless of career durations or fire days per year (24). Compared to females and older adults,

males and younger people also tend to spend more time outdoors, resulting in higher exposure to

wildfire smoke (25, 26). Consistent with our finding, racial/ethnic minorities were found to be

vulnerable subgroups in several previous studies (14, 15), which could be explained by pre-

existing health conditions and limited access to high-quality healthcare due to structural racism in

the U.S. (27, 28).

A higher association between smoke $PM_{2.5}$ and mortality was found in counties and

months with mild drought compared to those with no drought. We did not observe a statistically

significant association during severe droughts, possibly due to the relatively small sample size for

this category. This effect modification can be potentially explained by both increased risk of

exposure and higher human susceptibility brought by droughts. First, droughts can produce

conditions that are favorable to dust storms, which increase the distribution and concentration of

particulate matter in the air (13). In addition, drought has been linked to a variety of adverse

health outcomes, such as cardiovascular disease, respiratory diseases, and psychosocial stress

(29-31), leading to higher susceptibility to smoke pollutants. This effect modification by drought

may be of great importance to public health benefits because both drought and wildfire smoke are

projected to increase in many regions in the coming century (32, 33).

Our study found that about 1,141 deaths per year were contributed by wildfire smoke

$PM_{2.5}$ in the contiguous U.S., which is over 100 times higher than the recorded wildfire deaths in

the U.S. Billion-dollar Weather and Climate Disasters report by the National Oceanic and

Atmospheric Administration's National Centers for Environmental Information (10 deaths/year)

(34). This indicates a great number of indirect deaths brought by wildfires that could not be

captured by official tolls. In addition, the U.S. Billion-dollar Weather and Climate Disasters report

estimated that wildfire events cost about 3.1 billion dollars per year in the U.S. (34), but this

estimate does not take into account health care related losses or values associated with loss of

life (35). A recent study reported that the economic value of the health impacts of wildfires could

be in the tens to hundreds of billions of U.S. dollars, but the exposure-response function for the

$PM_{2.5}$-mortality relationship they used was for all-source $PM_{2.5}$, not wildfire-specific $PM_{2.5}$ (36).

Our study suggests a tremendous wildfire smoke-related mortality burden, and our effect

estimates for the relationship between wildfire $PM_{2.5}$ and mortality could be applied in future

estimations of the wildfire costs.

Some limitations of this study should be noted. First, the wildfire smoke $PM_{2.5}$

concentrations were modelled and subject to uncertainty. Because direct measurements of the

smoke contribution to $PM_{2.5}$ pollution are not available, the smoke $PM_{2.5}$ prediction was based on

$PM_{2.5}$ anomalies at monitoring stations, which may be an imprecise estimate of the

concentrations from smoke (23). A more accurate exposure assessment of wildfire smoke $PM_{2.5}$,

especially different chemical components of smoke $PM_{2.5}$, is needed in the future. Second, we

assumed that the non-smoke $PM_{2.5}$ was the difference between the all-source $PM_{2.5}$ and the

smoke $PM_{2.5}$ concentrations, but they were generated using different methods and may introduce

measurement error. Third, lacking detailed location information, this county-level ecological study

is susceptible to ecological fallacy. We were also unable to analyze the influence of wildfire

evacuation in this study. In addition, although our model specification can account for

unmeasured confounders that vary only across time or space, the association between smoke

$PM_{2.5}$ and mortality may be confounded by unmeasured time-varying individual county effects,

such as other wildfire-related pollutants including carbon monoxide. Furthermore, the monthly

resolution of mortality data is not fine enough for us to capture more acute effects of wildfire

smoke within a few days, especially on respiratory outcomes. Finally, there is no standard way to

define and categorize drought. Our findings on the effect modification by drought may be

influenced by different drought definitions and categorizations.

In conclusion, our study estimated the detrimental effects of wildfire smoke $PM_{2.5}$ on both

physical and mental health in the U.S. With wildfire intensity and frequency anticipated to

increase in the future driven by climate change (2), more effective wildfire preparedness and

mitigation strategies are urgently needed. Public health responses that protect people against

wildfire smoke pollution is crucial both in and outside the areas where the wildfires occur.

Materials and Methods

Mortality and population data

We obtained mortality data for all 3,108 counties or county equivalents in the contiguous U.S. from 2006 to 2016 from the National Center for Health Statistics. The mortality dataset includes the year and month of death, the cause of death (International Statistical Classification of Diseases and Related Health Problems, 10th Revision [ICD-10] codes), and the sex, age, race, and ethnicity of each deceased person. Based on the findings from previous studies (7, 14, 22), this study focused on all-cause mortality (ICD-10 code: A00-Z99) and mortality from three major specific causes: cardiovascular diseases (I00-I99), respiratory diseases (J00-J99), and mental and behavioral disorders (F00-F99). For each county, we summarized the monthly mortality counts for each cause (all-cause, cardiovascular, respiratory, and mental), sex (male and female), age group (0 to 64, 65 and above), and race/ethnicity (non-Hispanic White, non-Hispanic Black, and Hispanic).

County-level population data was collected from the Surveillance, Epidemiology, and End Results Program, National Cancer Institute (37). The total population and population estimate by sex, age, race, and Hispanic origin were extracted for each county, 2006–2016. Using anonymized monthly county-level mortality data, this study was determined as a Not Human Subject research by the Yale Institutional Review Boards (protocol ID: 2000026808).

Wildfire smoke PM$_{2.5}$ and other air pollutants

Ambient wildfire smoke PM$_{2.5}$ estimates for the contiguous U.S. was provided by a recent study by Childs et al. (23). In brief, station-based ground smoke PM$_{2.5}$ was defined as anomalies above the median on days in which wildfire smoke was overhead, then a model was trained to predict the station-based smoke PM$_{2.5}$ using meteorological factors, fire variables, aerosol measurements, and land use and elevation data. Finally, the trained model was applied to produce daily estimates of smoke PM$_{2.5}$ over the contiguous U.S. at a resolution of 10×10 km^2 (23). This model performed well over the entire range of observed smoke PM$_{2.5}$, without

saturation at very high daily $PM_{2.5}$ levels (23). We additionally validated this model against a

recently published wildfire-specific $PM_{2.5}$ model in California, which applied a novel ensemble-

based statistical approach to isolate wildfire-specific $PM_{2.5}$ from other sources of emissions (38).

This external validation showed a great consistency between the monthly county-level predictions

from these two models in California, 2006–2016, with an R-squared (R^2) value of 0.92 and a root-

mean-square error (RMSE) of 0.57 µg/m^3 (**Fig. S3**).

Data of daily NO_2, O_3, and total all-source $PM_{2.5}$ concentrations at 1×1 km^2 resolution

were obtained from the NASA Socioeconomic Data and Applications Center (39-44). Non-smoke

$PM_{2.5}$ concentrations were calculated by subtracting the smoke $PM_{2.5}$ from the all-source $PM_{2.5}$

concentrations. For negative values produced by this subtraction (0.4% of the total observations),

the non-smoke $PM_{2.5}$ concentrations were recoded as 0. The daily smoke $PM_{2.5}$, non-smoke

$PM_{2.5}$, NO_2, and O_3 concentrations were aggregated into monthly county-level average using

population-weighted averaging to match with the mortality data. The cartographic boundary for

counties in the contiguous U.S. was downloaded from the U.S. Census Bureau's TIGER/Line

geodatabase (45).

Meteorological factors and drought

Monthly mean air temperature and mean dew point temperature data at 4×4 km^2 were

obtained from the PRISM Climate Group (46). Similar to the air pollution data, we generated

monthly averages for these two variables for each county.

Drought was measured by standardized precipitation evapotranspiration index (SPEI), a

recently developed index considering both precipitation and temperature variables (47).

Compared with other drought indices, SPEI has the advantages of being multi-scalar, reflecting

the responses of different systems based on several water deficit accumulation periods, and

being sensitive to changes in evaporation demand caused by temperature (48). This index has

been increasingly applied in epidemiological studies in recent years to determine the onset,

duration and magnitude of drought conditions (49, 50).

We extracted SPEI data at one-month timescale with a 0.5 degrees spatial resolution for

each county from the Global SPEI database (51). Based on the severity of drought, we classified

each month of each county into no drought (SPEI ≥ -0.4), mild drought (-1.2 ≤ SPEI ≤ -0.5), and

severe drought (SPEI ≤ -1.3) (50, 52).

Statistical analysis

We applied a TWFE model with a quasi-Poisson regression to estimate the association

between wildfire smoke PM$_{2.5}$ exposure and mortality. TWFE model has been increasingly

applied in environmental epidemiology in recent years (53, 54). By introducing the indicators for

each county and each month, TWFE model can potentially control for all spatial confounders that

only vary across counties (e.g., urbanicity) and all temporal confounders that only vary by time

(e.g., seasonality), either measured or unmeasured (55). In our study, the main TWFE model can

be expressed as

$\ln[E(Y_{i,t})] = \mu + \alpha_i + \theta_t + \beta_1 SmokePM_{2.5_{i,t}} + \beta_2 NonSmokePM_{2.5_{i,t}} + ns(Temperature_{i,t}, df =$
$5) + \varepsilon_{i,t} + offset[\ln(Population_{i,t})],$

where $Y_{i,t}$ represents the number of all-cause or cause-specific deaths in county i, month t.

$SmokePM_{2.5_{i,t}}$ and $NonSmokePM_{2.5_{i,t}}$ are the mean smoke PM$_{2.5}$ and non-smoke PM$_{2.5}$

concentrations in county i, month t. α_i refers to time-invariant county effects and θ_t refers to time-

varying effects that are common in all counties. Air temperature was controlled by a flexible

natural cubic spline with five degrees of freedom (df). $\ln(Population_{i,t})$ is an offset term which

represents the natural log of the population in county i, month t. μ is the intercept term and $\varepsilon_{i,t}$ is

the error term. We weighted models using the population size in each county to improve the

precision of our estimates (56). The TWFE model requires no unmeasured confounders that

display different temporal variations across counties (i.e., time-varying individual county effects)

(55). Here, we assumed that non-smoke PM$_{2.5}$ and air temperature are the only candidates of

such confounders.

Based on the estimated coefficient of $SmokePM_{2.5_{i,t}}$ (β_1) for each cause, we calculated

the number of deaths attributable to smoke $PM_{2.5}$ (AN) using the attributable fraction (AF)

method. Specifically, AF was defined as follows: $AF = 1 - e^{-\beta_1 SmokePM_{2.5_{i,t}}}$. AF was then

multiplied by the monthly cause-specific number of deaths to estimate the smoke $PM_{2.5}$

attributable deaths in each county in each month.

In subgroup analyses, we further estimated the association between smoke $PM_{2.5}$ and all-

cause mortality by sex, age, and race/ethnicity. We also conducted stratified analysis by drought

severity (no drought, mild drought, and severe drought) to investigate its potential modification

effect. We tested the statistical difference in effect estimates between different subgroups and

drought levels by calculating the z score as $(\hat{Q}_1 - \hat{Q}_2)/\sqrt{(S\hat{E}_1)^2 + (S\hat{E}_2)^2}$, where $\hat{Q}_1$ and $\hat{Q}_2$ are the

estimates, and $S\hat{E}_1$ and $S\hat{E}_2$ are their respective standard errors (57).

Several sensitivity analyses were performed to test the robustness of our results: (a) we

additionally adjusted for NO_2, O_3, or dew point temperature in the model; (b) we removed non-

smoke $PM_{2.5}$ from the model; (c) an alternative four or six dfs was used in the natural cubic spline

of air temperature; (d) we tested the delayed effects of smoke $PM_{2.5}$ in the previous one or two

months; and (e) we restricted the analysis to counties that had at least one day with smoke $PM_{2.5}$

higher than 20 µg/m³. To test the linearity of the relationship between smoke $PM_{2.5}$ and mortality,

we replaced the linear term of smoke $PM_{2.5}$ by a natural cubic spline with three dfs in the model

and visualized the estimated curve.

Acknowledgments

Dr. E. Zang received support from the National Institute on Aging (R21AG074238-01), the

National Institute on Minority Health and Health Disparities (1R01MD017298-01), the Research

Education Core of the Claude D. Pepper Older Americans Independence Center at Yale School

of Medicine (P30AG021342), and the Institution for Social and Policy Studies at Yale University.

Dr. Y. Liu received support from the National Institute of Environmental Health Sciences

(1R01ES034175).

References

1. Burke M*, et al.*, The changing risk and burden of wildfire in the United States. *Proc Natl Acad Sci U S A* 118(2):e2011048118 (2021).
2. Spracklen DV*, et al.*, Impacts of climate change from 2000 to 2050 on wildfire activity and carbonaceous aerosol concentrations in the western United States. *J Geophys Res Atmos* 114(D20)2009).
3. Kim K-H, Kabir E, & Kabir S, A review on the human health impact of airborne particulate matter. *Environ Int* 74:136-143 (2015).
4. Xu R*, et al.*, Wildfires, global climate change, and human health. *N Engl J Med* 383(22):2173-2181 (2020).
5. Liu JC*, et al.*, Wildfire-specific fine particulate matter and risk of hospital admissions in urban and rural counties. *Epidemiology* 28(1):77 (2017).
6. Heft-Neal S, Driscoll A, Yang W, Shaw G, & Burke M, Associations between wildfire smoke exposure during pregnancy and risk of preterm birth in California. *Environ Res* 203:111872 (2022).
7. Reid CE*, et al.*, Critical review of health impacts of wildfire smoke exposure. *Environ Health Perspect* 124(9):1334-1343 (2016).
8. O'Dell K*, et al.*, Estimated mortality and morbidity attributable to smoke plumes in the United States: Not just a western US problem. *GeoHealth* 5(9):e2021GH000457 (2021).
9. Ye T*, et al.*, Short-term exposure to wildfire-related PM2.5 increases mortality risks and burdens in Brazil. *Nat Commun* 13(1):1-9 (2022).
10. Chen G*, et al.*, Mortality risk attributable to wildfire-related PM2.5 pollution: a global time series study in 749 locations. *Lancet Planet Health* 5(9):e579-e587 (2021).
11. Chen H, Samet JM, Bromberg PA, & Tong H, Cardiovascular health impacts of wildfire smoke exposure. *Part Fibre Toxicol* 18(1):1-22 (2021).
12. Eisenman DP & Galway LP, The mental health and well-being effects of wildfire smoke: a scoping review. *BMC Public Health* 22(1):1-17 (2022).
13. Bell JE*, et al.*, Changes in extreme events and the potential impacts on human health. *J Air Waste Manag Assoc* 68(4):265-287 (2018).
14. Kondo MC*, et al.*, Meta-analysis of heterogeneity in the effects of wildfire smoke exposure on respiratory health in North America. *Int J Environ Res Public Health* 16(6):960 (2019).
15. Liu JC*, et al.*, Who among the elderly is most vulnerable to exposure to and health risks of fine particulate matter from wildfire smoke? *Am J Epidemiol* 186(6):730-735 (2017).
16. Morgan G*, et al.*, Effects of bushfire smoke on daily mortality and hospital admissions in Sydney, Australia. *Epidemiology* 21:47-55 (2010).
17. Rappold AG*, et al.*, Peat Bog Wildfire Smoke Exposure in Rural North Carolina Is Associated with Cardiopulmonary Emergency Department Visits Assessed through Syndromic Surveillance. *Environ Health Perspect* 119(10):1415-1420 (2011).
18. National Drought Mitigation Center, Drought In-depth. (2022).
19. Chen K*, et al.*, Two-way effect modifications of air pollution and air temperature on total natural and cardiovascular mortality in eight European urban areas. *Environ Int* 116:186-196 (2018).
20. Li J*, et al.*, Modification of the effects of air pollutants on mortality by temperature: a systematic review and meta-analysis. *Sci Total Environ* 575:1556-1570 (2017).
21. Leitte AM*, et al.*, Respiratory health, effects of ambient air pollution and its modification by air humidity in Drobeta-Turnu Severin, Romania. *Sci Total Environ* 407(13):4004-4011 (2009).
22. Liu JC, Pereira G, Uhl SA, Bravo MA, & Bell ML, A systematic review of the physical health impacts from non-occupational exposure to wildfire smoke. *Environ Res* 136:120-132 (2015).

23. Childs ML, *et al.*, Daily local-level estimates of ambient wildfire smoke PM2.5 for the
contiguous US. *Environ Sci Technol* 56(19):13607-13621 (2022).
24. Navarro KM, *et al.*, Wildland firefighter smoke exposure and risk of lung cancer and
cardiovascular disease mortality. *Environ Res* 173:462-468 (2019).
25. Matz CJ, *et al.*, Effects of age, season, gender and urban-rural status on time-activity:
CanadianHuman Activity Pattern Survey 2 (CHAPS 2). *Int J Environ Res Public Health*
11(2):2108-2124 (2014).
26. Kerr J, *et al.*, The relationship between outdoor activity and health in older adults using
GPS. *Int J Environ Res Public Health* 9(12):4615-4625 (2012).
27. Van Horne YO, *et al.*, An applied environmental justice framework for exposure science.
J Expo Sci Environ Epidemiol 2022).
28. Seltenrich N, The one-two-three punch: exposure, susceptibility, and disease burden
among U.S. populations of color. *Environ Health Perspect* 130(3):034001 (2022).
29. Sugg M, *et al.*, A scoping review of drought impacts on health and society in North
America. *Clim Change* 162(3):1177-1195 (2020).
30. Berman JD, Ebisu K, Peng RD, Dominici F, & Bell ML, Drought and the risk of hospital
admissions and mortality in older adults in western USA from 2000 to 2013: a
retrospective study. *Lancet Planet Health* 1(1):e17-e25 (2017).
31. Berman JD, *et al.*, The association between drought conditions and increased
occupational psychosocial stress among US farmers: An occupational cohort study. *Sci*
Total Environ 798:149245 (2021).
32. Ford B, *et al.*, Future fire impacts on smoke concentrations, visibility, and health in the
contiguous United States. *GeoHealth* 2(8):229-247 (2018).
33. IPCC (2021) *Climate Change 2021: The Physical Science Basis. Contribution of Working*
Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate
Change (Cambridge University Press, Cambridge, United Kingdom and New York, NY,
USA).
34. NOAA National Centers for Environmental Information (NCEI), U.S. Billion-Dollar
Weather and Climate Disasters.   (2023).
35. Smith AB & Katz RW, US billion-dollar weather and climate disasters: data sources,
trends, accuracy and biases. *Nat Hazards* 67(2):387-410 (2013).
36. Fann N, *et al.*, The health impacts and economic value of wildland fire episodes in the
U.S.: 2008–2012. *Sci Total Environ* 610-611:802-809 (2018).
37. The SEER Program, U.S. County Population Data - 1969-2020.   (2022).
38. Aguilera R, *et al.*, A novel ensemble-based statistical approach to estimate daily wildfire-
specific PM2.5 in California (2006-2020). *Environ Int* 171:107719 (2023).
39. Di Q, *et al.*, Daily and Annual PM2.5 Concentrations for the Contiguous United States, 1-
km Grids, v1 (2000 - 2016). (NASA Socioeconomic Data and Applications Center
(SEDAC), Palisades, New York) (2021).
40. Di Q, *et al.*, Assessing NO2 concentration and model uncertainty with high
spatiotemporal resolution across the contiguous United States. *Environ Sci Technol*
54(3):1372-1384 (2019).
41. Di Q, *et al.*, Daily and Annual NO2 Concentrations for the Contiguous United States, 1-
km Grids, v1 (2000 - 2016). (NASA Socioeconomic Data and Applications Center
(SEDAC), Palisades, New York) (2022).
42. Requia WJ, *et al.*, An ensemble learning approach for estimating high spatiotemporal
resolution of ground-level ozone in the contiguous U.S. *Environ Sci Technol*
54(18):11037-11047 (2020).
43. Requia WJ, *et al.*, Daily 8-Hour Maximum and Annual O3 Concentrations for the
Contiguous United States, 1-km Grids, v1 (2000 - 2016). (NASA Socioeconomic Data
and Applications Center (SEDAC), Palisades, New York) (2021).
44. Di Q, *et al.*, An ensemble-based model of PM2.5 concentration across the contiguous
United States with high spatiotemporal resolution. *Environ Int* 130:104909 (2019).
45. United States Census Bureau, Cartographic Boundary Files.   (2019).

46. PRISM Climate Group, PRISM Climate Data Recent Years (Jan 1981 - Jan 2022). (Oregon State University,) (2016).
47. Vicente-Serrano SM, Beguería S, & López-Moreno JI, A multiscalar drought index sensitive to global warming: the standardized precipitation evapotranspiration index. *J Clim* 23(7):1696-1718 (2010).
48. Vicente-Serrano SM*, et al.*, Performance of drought indices for ecological, agricultural, and hydrological applications. *Earth Interact* 16(10):1-27 (2012).
49. Salvador C*, et al.*, Drought effects on specific-cause mortality in Lisbon from 1983 to 2016: Risks assessment by gender and age groups. *Sci Total Environ* 751:142332 (2021).
50. Wang P, Asare E, Pitzer VE, Dubrow R, & Chen K, Associations between long-term drought and diarrhea among children under five in low-and middle-income countries. *Nat Commun* 13(1):1-10 (2022).
51. Beguería S, Vicente-Serrano SM, Reig F, & Latorre B, Standardized precipitation evapotranspiration index (SPEI) revisited: parameter fitting, evapotranspiration models, tools, datasets and drought monitoring. *Int J Climatol* 34(10):3001-3023 (2014).
52. Federal Office of Meteorology and Climatology MeteoSwiss, Calculation of SPI and SPEI. (2018).
53. Wang Y*, et al.*, Estimating causal effects of long-term PM2.5 exposure on mortality in New Jersey. *Environ Health Perspect* 124(8):1182-1188 (2016).
54. Renzi M*, et al.*, Long-term PM10 exposure and cause-specific mortality in the Latium Region (Italy): a difference-in-differences approach. *Environ Health Perspect* 127(6):067004 (2019).
55. Goodman-Bacon A, Difference-in-differences with variation in treatment timing. *J Econom* 225(2):254-277 (2021).
56. Zang E & Kim N, Intergenerational upward mobility and racial differences in mortality among young adults: Evidence from county-level analyses. *Health Place* 70:102628 (2021).
57. Chen K*, et al.*, Triggering of myocardial infarction by heat exposure is modified by medication intake. *Nat Cardiovasc Res* 1(8):727-731 (2022).

Figures
520

521
522 **Fig. 1. Map of monthly county-level mean wildfire smoke PM$_{2.5}$ concentration and annual**
523 **average attributable all-cause mortality burden in the contiguous U.S., 2006–2016. A**: The
524 distribution of monthly mean smoke PM$_{2.5}$ in contiguous U.S. counties (μg/m^3). **B**: The distribution
525 of annual average all-cause mortality burden attributable to wildfire smoke PM$_{2.5}$ in contiguous
526 U.S. counties (deaths per 1,000,000 people).
527
528
529

531
532
Fig. 2. Association between wildfire smoke PM$_{2.5}$ and cause-specific mortality risk and
annual average attributable cause-specific mortality burden. A: estimated percentage
changes in monthly mortality per 1 μg/m^3 increase in smoke PM$_{2.5}$ concentration, for all-cause
mortality, cardiovascular disease mortality, respiratory disease mortality, and mental disorder
mortality. The error bars indicate 95% confidence intervals. **B**: estimated annual average mortality
burden attributable to smoke PM$_{2.5}$, for all causes, cardiovascular diseases, respiratory diseases,
and mental and behavioral disorders. The numbers in parentheses are 95% confidence intervals.
540
541

Fig. 3. Percent change in all-cause mortality associated with a 1 µg/m³ increase in smoke PM₂.₅ for each subgroup. This figure shows the estimated associations between each 1-µg/m³ increase in smoke PM₂.₅ concentration and all-cause mortality risks for different sex, age, and race/ethnicity groups. Based on the z score calculated using the coefficients and standard errors for different subgroup, we tested the statistical difference in effect estimates among different sex, age, and race/ethnicity groups, using males, people aged 0 to 64 years, and non-Hispanic White people as the reference group.

553 **Fig. 4. Distribution of smoke PM$_{2.5}$ and drought and effect modification by drought. A, B**:
554 This bivariate choropleth map shows the percentage of months with non-zero smoke PM$_{2.5}$
555 concentrations and the percentage of months with mild (A) or severe (B) drought in each
556 contiguous U.S. county, from 2006 to 2016. **C**: The estimated percent changes in all-cause
557 mortality per 1 µg/m³ increase in smoke PM$_{2.5}$ concentration when there was no drought (86.03%
558 county-months), mild drought (23.28% county-months), or severe drought (8.68% county-
559 months). Based on the z score calculated using the coefficients and standard errors for different
560 levels of drought, we tested the statistical difference in effect estimates between different drought
561 levels, using no drought as the reference.
562