

1 Word count: 4545
2 Abstract word count: 221
3 Key words: 5
4 Number of references: 48
5 Number of tables: 05
6 Number of figures: 00
7

8 **Factors associated with suffering domestic violence in women during the last**
9 **semester of the pandemic in Honduras- Central America**

10

11 Eleonora Espinoza-Turcios^{1,2}

12 Lysien Ivania Zambrano¹

13 Carlos Sosa-Mendoza^{2,3}

14 Henry Noel Castro Ramos¹

15 Dennis Arias-Chávez⁴

16 Christian R. Mejia⁵

17

18 **Affiliations:**

19 ¹ Institute for Research in Medical Sciences and Right to Health (ICIMEDES)/ Faculty
20 of Medical Sciences (FCM), National Autonomous University of Honduras (UNAH).
21 Tegucigalpa, Honduras.

22 ²Mental health research group, FCM-UNAH

23 ³Honduran Institute for the Prevention of Alcoholism, Drug Addiction and Drug
24 Dependence (IHADFA) Tegucigalpa, Honduras.

25 ⁴Universidad Continental, Arequipa, Perú

26

27 ⁵Translational Medicine Research Centre. Universidad Norbert Wiener. Lima, Perú.

28

29 **Corresponding authors:** Eleonora Espinoza Turcios eleonora.espinoza@unah.edu.hn,
30 Instituto de Investigación en Ciencias Médicas y Derecho a la Salud
31 (ICIMEDES)/Unidad de Investigación Científica (UIC), Edificio CM1, FCM, Calle La

32 Salud, contiguo al Hospital Escuela, Tegucigalpa, M. D.C., Honduras, C.A. Código
33 Postal: 11101.

34 **ABSTRACT**

35 **Introduction:** Mental health deterioration had already been reported prior to the
36 pandemic, resulting in domestic violence in women, but this has not yet been reported
37 in the pandemic in Central America. **AIM:** To determine the Factors associated with
38 suffering domestic violence in women in the last half year of the pandemic in Honduras.
39 **Methodology:** Analytical and retrospective cross-sectional study, carried out through a
40 survey in 17 departments of Honduras, in hospitals and first level health care centers.
41 The main variable was obtained through the question "if you suffered domestic violence
42 in the last 6 months", being the possible answer verbal, physical, psychological.
43 Descriptive and analytical results were obtained. **Results:** Of the 8442 Hondurans
44 surveyed, 4.2% (352) perceived verbal violence, 1.9% (165) physical violence and 1.3%
45 (113) psychological violence. In the multivariate analysis, it was found that women
46 ($p=0.001$), those with a history of alcohol ($p=0.002$) or drug use ($p=0.015$), previous
47 mental illness ($p<0.001$), mild ($p<0.001$), moderate ($p<0.001$) or severe ($p=0.025$)
48 depression ($p<0.001$) had experienced more domestic violence; On the contrary, there
49 was less perception of domestic violence at higher economic income (all p-values were
50 <0.029), among single ($p=0.003$) and married people ($p<0.001$). **Discussion:** Important
51 Factors associated with suffering domestic violence in women, especially social factors
52 and a history of mental illness, as well as alcohol and drug use in the home.

53 **Key words:** COVID-19, Honduras, Central America, Women, Domestic Violence.

54

55 **INTRODUCTION**

56 Domestic violence (DV) is defined as "a pattern of behavior that is used to gain
57 or maintain power and control over an intimate partner in a relationship, may include a
58 child or other relative, or any other member of the household" [1]. Globally, the lifetime
59 prevalence of intimate partner violence is 26% among married or intimate partner
60 women over the age of 15; estimating that 6% of women over the age of 15 have been
61 subjected to non-partner sexual violence during some point in their lifetime [2].

62 At the beginning of the COVID-19 pandemic, in the first months of 2020,
63 domestic violence increased worldwide, Spain recorded a 20% increase in calls to
64 women's helplines; [3]. United Kingdom reported an additional 25% one week after the
65 announcement of the social distancing measures; [4] in France the increase of 36% was
66 reported; [5] in Jingzhou-China the number of reported domestic violence cases was
67 three times higher in February 2020, this compared to the same period last year [6]. This
68 is in addition to the global and public health problems that were known prior to this
69 pandemic that was generated worldwide, where one in three women had suffered in her
70 life some type of physical and/or sexual violence by an intimate partner [7].

71 In Honduras, the problem of domestic violence behaves similarly to the rest of
72 the world, and worsened as a result of the COVID-19 pandemic. According to data from
73 the National Demographic and Health Survey (ENDESA /MICS 2019), 20% of
74 Honduran women aged 15-49 years have ever been beaten or physically abused by
75 someone after the age of 15, psychological violence occurs more frequently in urban
76 areas (with 17% compared to 12% of cases in rural areas) and 4% of women aged 15-49
77 have suffered sexual abuse since the age of 12, being in 12.5% of cases produced by the
78 current boyfriend/ex-boyfriend [8]. And as if this were not enough, according to the
79 University Institute for Democracy, Peace and Security UNAH, there were 636 and 330
80 deaths of women due to violence in 2013 and 2021, respectively. And from January to
81 October 2022 there were 252 violent deaths of women and/or femicide, that is, every 28
82 hours and 57 minutes a woman dies in Honduras [9]. The domestic violence women in
83 Honduras involves multiple factors, such as socioeconomic, cultural, religious, and
84 represents a serious social problem with serious consequences [10]. Study conducted by
85 the National Autonomous University of Honduras through the Faculty of Medical
86 Sciences ICIMEDES, that women suffer some type of violence 45% in five very low-
87 income communities [11], also the national survey of the country ENDESA in 2019
88 shows that women between 15-49 years have suffered 16% some type of violence [8].
89 For all these reasons, the objective was to determine the Factors associated with
90 suffering domestic violence in women in the last half year of the pandemic in Honduras.

91

92 **METHODOLOGY**

93 An analytical, cross-sectional and retrospective research was generated. It was
94 based on the information collected during the pandemic stage, which included the last
95 third of the year 2021 and the first third of 2022. For the collection of information, a
96 baseline instrument was elaborated, which measured the baseline mental health situation
97 of the Honduran population in 17 of the departments [12]. This was carried out in health
98 care centers, both hospitals and first level health care facilities.

99 In order for a respondent to participate in the research, it was taken into account
100 that he/she was 18 years of age or older, that he/she agreed to participate in the research
101 and that he/she had answered the main question, ("if you suffered domestic violence in
102 the last 6 months", being the possible answer verbal, physical, psychological). There
103 were no exclusions because all respondents had answered the main question and met the
104 other inclusion criteria. To determine whether the number of respondents was adequate
105 for the main crosstabs, statistical power was calculated, where there was adequate
106 power for the crosstabs of general perception of violence versus sex (power = 100%),
107 according to the amount of family income (power = 100% for all categories), marital
108 status (power = 100% for all categories), alcohol consumption in the last 6 months
109 (power = 98%), drug consumption in the last 6 months (power = 100%), history of
110 mental illness (power = 100%) and versus depression (power = 100% for all categories).

111 The main variable was obtained through the question "if you have suffered
112 domestic violence in the last 6 months", which was a semi-open question with one of
113 four possible alternatives: no violence, or verbal, physical, psychological violence. For
114 the individual analysis of each of the types of violence, they were crossed with each of
115 the independent variables. But for the final analysis, the perception of violence was
116 considered if one or more of the three categories (verbal, physical or psychological) was
117 perceived, which was also crossed against the other variables. Secondary variables were
118 age (in years completed), sex (male or female), amount of family income (<\$400, \$400-
119 800, \$801-1199, \$1200-1600 and >\$1600), marital status (common-law/cohabiting,
120 single, married or widowed/divorced), alcohol use in the past 6 months (no or yes), drug
121 use in the past 6 months (no or yes), history of mental illness (no or yes) and versus
122 depression (have not, mild, moderate, severe or very severe).

123 To proceed with the data analysis, the information was first extracted from the
124 database, where an author did the general filtering of the data to be analyzed. Then a
125 second quality control was carried out by another of the authors, where the variables
126 were categorized, labeled and a glossary was constructed. Then all this information was
127 passed to the statistical program Stata (license obtained by the authors who carried out
128 the analysis). Then a table was created showing the frequencies and percentages of the
129 suffering of the three types of domestic violence, then three tables were generated
130 where these types of violence were crossed according to the socio-pathological
131 variables, where the p-values were obtained with the Wilcoxon test (for the age
132 variable) and the chi-square (for the other variables). Then a final table was generated,
133 where the dependent variable was having suffered at least one type of violence, also
134 being crossed against the socio-pathological variables; in this step the prevalence ratios,
135 95% confidence intervals and p-values were obtained, this with the use of generalized
136 linear models: Poisson family, log link function and adjusted models for obtaining
137 robust variances). The p-value <0.05 was the criterion for a variable to pass from the
138 bivariate to the multivariate model, as well as to determine statistical significance.

139 Ethics were always respected, the initial study was approved by a local ethics
140 committee (2019-62), The survey was anonymous, and participants were informed of
141 the objective of the study.

142

143 **RESULTS**

144 Of the 8442 Hondurans surveyed, 4.2% (352) perceived verbal violence, 1.9%
145 (165) perceived physical violence and 1.3% (113) perceived psychological violence.
146 When the 3 types of perceptions were added together, 97.9% (8265) did not perceive
147 any violence, while 0.1% of respondents perceived all 3 types of violence in their
148 homes during the last six months. Table 1

149

150 **Table 1. Frequencies and percentages of experiencing domestic violence in women**
151 **during the last semester of the pandemic in Honduras, n=8442.**

Type of violence	Frequency	Percentage
Verbal abuse		
No	8090	95.8
Yes	352	4.2
Physical violence*		
No	8277	98.1
Yes	165	1.9
Psychological violence		
No	8329	98.7
Yes	113	1.3
Sum of perceptions		
No violence of any kind	8265	97.9
1 type of violence	126	1.5
2 type of violence	45	0.5
3 type of violence	6	0.1

152 *don't include violence sexual.

153 Verbal domestic violence was associated with gender ($p=0.009$), family income
154 ($p<0.001$), marital status ($p=0.013$), alcohol consumption at home ($p<0.001$) or drug use
155 ($p<0.001$), a history of mental illness ($p<0.001$) and depression ($p<0.001$). Table 2

156

157

158 **Table 2. Household verbal abuse associated with socio-pathological variables in the last**
 159 **half year of the pandemic in Honduras, n=8442.**

Variables	Verbal abuse		P value
	No n (%)	Yes n (%)	
Age*	35(26-49)	35 (26-48)	0.740
Gender			
Male	2915 (96.6)	103 (3.4)	0.009
Female	5175 (95.4)	249 (4.6)	
Family income (monthly)			
<400.00 (\$)	2709 (94.6)	154 (5.4)	<0.001
400.00-800.00 (\$)	2134 (95.9)	92 (4.1)	
801.00-1199.00 (\$)	999 (97.2)	29 (2.8)	
1200.00-1600.00 (\$)	446 (97.6)	11(2.4)	
>1600.00 (\$)	570 (97.4)	15 (2.6)	
Marital status			
Unmarried/cohabitating partner	2821 (94.9)	151 (5.1)	0.013
Singles	2635 (96.2)	105 (3.8)	
Married	2269 (96.6)	79 (3.4)	
Widowed or divorced	365 (95.6)	17 (4.4)	
Consumption of alcohol**			
No	5894 (96.5)	212 (3.5)	<0.001
Yes	2196 (94.0)	140 (6.0)	
Consumption of drugs**			
No	7906 (96.1)	322 (3.9)	<0.001
Yes	184 (86.0)	30(14.0)	
Antecedent of mental illness			
No	7547 (96.2)	298 (3.8)	<0.001
Yes	543 (91.0)	54 (9.0)	
Depression***			
None	5972 (97.2)	174 (2.8)	<0.001

Mild	1668 (92.5)	136 (7.5)
Moderate	310 (91.7)	28 (8.3)
Severe	87 (89.7)	10 (10.3)
Highly severe	53 (93.0)	4 (7.0)

160 *Quantitative variable (median and interquartile ranges are shown). **Household
 161 consumption in the last six months. ***Measured with Hamilton's test. P values were
 162 obtained with Wilcoxon test (age) and chi-square.

163 Associated with physical violence at home were home use of alcohol (p=0.004) or drugs
 164 (p<0.001), history of mental illness (p<0.001) and depression (p<0.001). Table 3

165

166 **Table 3. Physical violence at home associated with socio-pathological variables in the last**
 167 **half year of the pandemic in Honduras, n=8442.**

Variables	Physical violence		P value
	No n (%)	Yes n (%)	
Age*	35 (26-49)	36 (27-46)	0.741
Gender			
Male	2968 (98.3)	50 (1.7)	0.140
Female	5309 (97.9)	115(2.1)	
Family income (monthly)			
<400.00 (\$)	2794 (97.6)	69 (2.4)	0.256
400.00-800.00 (\$)	2177 (97.8)	49 (2.2)	
801.00-1199.00 (\$)	1008 (98.1)	20 (1.9)	
1200.00-1600.00 (\$)	452 (98.9)	5 (1.1)	
>1600.00 (\$)	577 (98.6)	8 (1.4)	
Marital status			
Unmarried/cohabitating partner	2910 (97.9)	62 (2.1)	0.465
Singles	2691 (98.2)	49 (1.8)	
Married	2305 (98.2)	43 (1.8)	
Widows or divorced	371 (97.1)	11 (2.9)	
Consumption of alcohol**			
No	6003 (98.3)	103 (1.7)	0.004

Yes	2274 (97.4)	62 (2.6)	
Consumption of drugs**			
No	8078 (98.2)	150 (1.8)	<0.001
Yes	199 (93.0)	15 (7.0)	
Antecedent of mental illness			
No	7713 (98.3)	132 (1.7)	<0.001
Yes	564 (94.5)	33 (5.5)	
Depression ***			
None	6061 (98.6)	85 (1.4)	<0.001
Mild	1751 (97.1)	53 (2.9)	
Moderate	322(95.3)	16 (4.7)	
Severe	92 (94.9)	5 (5.1)	
Highly severe	51 (89.5)	6 (10.5)	

168 *Quantitative variable (median and interquartile ranges are shown). **Household
 169 consumption in the last six months. ***Measured with Hamilton's test. P values were
 170 obtained with Wilcoxon test (age) and chi-square.

171

172 Family income (p=0.031), home consumption of alcohol (p<0.001) or drugs
 173 (p<0.001), history of mental illness (p<0.001) and depression (p<0.001) were associated
 174 with verbal violence at home. Table 4

175

176 **Table 4. Domiciliary sexual violence associated with socio-pathological variables in the**
 177 **last half year of the pandemic in Honduras, n=8442.**

Variables	Sexual violence		P value
	No n (%)	Si n (%)	
Age	35 (26-49)	36 (29-47)	0.808
Gender			
Male	2985 (98.9)	33 (1.1)	0.144
Female	5344 (98.5)	80 (1.5)	
Family income (monthly)			
<400.00 (\$)	2810 (98.2)	53 (1.8)	0.031

400.00-800.00 (\$)	2200 (98.8)	26 (1.2)	
801.00-1199.00 (\$)	1017 (98.9)	11 (1.1)	
1200.00-1600.00 (\$)	456 (99.8)	1 (0.2)	
>1600.00 (\$)	576 (98.5)	9 (1.5)	
Marital status			
Unmarried/cohabitating partner	2927 (98.5)	45 (1.5)	0.554
Singles	2710 (98.9)	30 (1.1)	
Married	2316 (98.6)	32 (1.4)	
Widows or divorced	376 (98.4)	6 (1.6)	
Consumption of alcohol**			
No	6042 (99.0)	64 (1.0)	<0.001
Yes	2287 (97.9)	49 (2.1)	
Consumption of drugs**			
No	8124 (98.7)	104 (1.3)	<0.001
Yes	205 (95.8)	9 (4.2)	
Antecedent of mental illness			
No	7760 (98.9)	85 (1.1)	<0.001
Yes	569 (95.3)	28 (4.7)	
Depression***			
None	6098 (99.2)	48 (0.8)	<0.001
Mild	1753 (97.2)	51 (2.8)	
Moderate	328 (97.0)	10 (3.0)	
Severe	95 (97.9)	2 (2.1)	
Highly severe	55 (96.5)	2 (3.5)	

178 *Quantitative variable (median and interquartile ranges are shown). **Household
 179 consumption in the last six months. ***Measured with Hamilton's test. P values were
 180 obtained with Wilcoxon test (age) and chi-square.

181

182 In the multivariate analysis, it was found that women (PRa: 1.92; 95%CI: 1.33-
 183 2.77; p-value=0.001), those with a history of alcohol (PRa: 1.77; 95%CI: 1.24-2.52; p-
 184 value=0.002) or drug use (PRa: 2.08; 95%CI: 1.15-3.75; p-value=0.015), previous

185 mental illness (PRa: 2.94; 95%CI: 2.00-4.31; p-value<0.001), mild (RPa: 2.48; 95%CI:
 186 1.75-3.53; p-value<0.001), moderate (RPa: 3.24; 95%CI: 1.85-5.70; p-value<0.001) or
 187 severe depression (RPa: 3.11; 95%CI: 1.15-8.39; p-value=0.025); on the contrary, there
 188 was less perception of domestic violence at higher economic income (all p-values were
 189 <0.029), among single (p=0.003) and married (p<0.001). **Table 5**

190 **Factors associated with suffering domestic violence in women during the last**
 191 **semester of the pandemic in Honduras- Central America**

192

193 **Table 5. Bivariate and multivariate analysis of factors associated domestic violence in**
 194 **women during the last semester of the pandemic in Honduras, n=8442.**

Variable	Bivariate analysis	Multivariate analysis
Age*	0.998 (0.988-1.008) 0.685	Not entered in the final model
Gender female	1.68 (1.20-2.36) 0.003	1.92 (1.33-2.77) 0.001
Family income (monthly)		
<400.00 (\$)	Comparison category	Comparison category
400.00-800.00 (\$)	0.63 (0.44-0.90) 0.012	0.65 (0.45-0.93) 0.018
801.00-1199.00 (\$)	0.39 (0.21-0.71) 0.002	0.40 (0.21-0.75) 0.004
1200.00-1600.00 (\$)	0.36 (0.15-0.89) 0.027	0.36 (0.14-0.89) 0.028
>1600.00 (\$)	0.34 (0.15-0.78) 0.010	0.30 (0.13-0.68) 0.004
Marital status		
Unmarried/cohabitating partner	Comparison category	Comparison category
Singles	0.51 (0.36-0.72) <0.001	0.56 (0.38-0.82) 0.003
Married	0.37 (0.24-0.56) <0.001	0.43 (0.27-0.68) <0.001
Widows or divorced	0.65 (0.32-1.32) 0.234	0.56 (0.25-1.26) 0.160
Consumption of alcohol**	1.52 (1.12-2.05) 0.007	1.77 (1.24-2.52) 0.002
Consumption of drugs**	3.05 (1.76-5.27) <0.001	2.08 (1.15-3.75) 0.015
Antecedent of mental illness	3.71 (2.63-5.25) <0.001	2.94 (2.00-4.31) <0.001
Depression***		
None	Comparison category	Comparison category
Mild	2.92 (2.14-3.98) <0.001	2.48 (1.75-3.53) <0.001
Moderate	3.03 (1.73-5.28) <0.001	3.24 (1.85-5.70) <0.001

Severe	3.77 (1.56-9.09) 0.003	3.11 (1.15-8.39) 0.025
Highly severe	2.57 (0.65-10.18) 0.180	1.40 (0.22-8.70) 0.721

195 *Quantitative variable. **Household consumption in the last six months. ***Measured
196 with Hamilton's test. Prevalence ratios are shown to the left of the parentheses, 95%
197 confidence intervals within the parentheses and p-values to the right (all obtained with
198 generalized linear models: Poisson family, log link function and models adjusted for
199 robust variances).

200

201 **DISCUSSION**

202 The objective of this study was to determine the factors associated with suffering
203 domestic violence in women during the pandemic in Honduras, since, Honduras does
204 not have updated national studies, the 2017 Chavarria-Mejia study was conducted in
205 two communities in Honduras, where they found that 79% of the 153 women over 16
206 years of age with a partner, had some type of domestic violence and 40% presented
207 economic violence [10].

208 In another study on domestic violence in women in five communities in
209 Honduras, the overall prevalence was 45% (737), the main type of violence was verbal
210 52% (383), but psychological violence was also important, 4% (31), sexual violence
211 19% (140) and physical violence 21% (151), predominantly in women between 20-39
212 years of age [11].

213 The 2019 ENDESA reported among women aged 15-49 years that 16% have
214 experienced some type of violence (psychological, physical or sexual) by a
215 husband/partner, 15% have suffered psychological violence, 6%, physical violence and
216 2% sexual violence [8]. This is aggravated by the report that practically once a day a
217 woman dies in this country [9].

218 As can be seen in the three previous studies, the report of violence has been
219 obtained in different populations in Honduras, but above all among women and prior to
220 the COVID-19 pandemic, therefore, this was obtained in a population that attends
221 health facilities, which are generally those who have more access to health care and
222 have other underlying diseases. In addition, the results we present are important to be
223 able to have a current statistic and that is given after one of the most important events
224 for public health in recent decades [13].

225 The latter being very important because during the confinement that generated
226 COVID-19, domestic violence increased due to the continuous contact between
227 perpetrators and victims; an increase in cases has been reported in North America,
228 Europe, Asia, the Pacific and Africa, especially during the first week of the blockade in
229 each country [14]. A systematic review showed an increase in domestic violence
230 following lockdowns [15].

231

232 As a first important result, less than 5% of the respondents perceived some type
233 of violence. These results are similar to those described in Germany, where an online
234 survey of 3545 volunteers reported that 5% of all participants (5.1% of all women and
235 4.1% of all men) reported having experienced interpersonal violence (IPV), verbal (98%
236 of all women and 100% of all men who experienced IPV), physical (38.4% of all
237 women and 63.6% of all men who experienced IPV), or sexual (26.5% of all women
238 and 50% of all men who experienced IPV) [16]. This is relevant because of the large
239 number of people affected globally, as calculated in the United States of America, and it
240 affects the victim, families, co-workers and the community, causes a decrease in
241 psychological and physical health, a decrease in quality of life and a decrease in
242 productivity [17]. Therefore, these results should be considered exploratory, since the
243 objective was not to extrapolate the data to the entire Honduran population, but they are
244 important because they were taken from a large population in the most important
245 departments of the country and at a time when the pandemic was still present among the
246 respondents.

247 Another important result was that women had a higher frequency of perceived
248 domestic violence. Knowing that there are global reports indicating that an estimated
249 243 million women and adolescents aged 15-49 years have been victims of physical
250 and/or sexual violence at the hands of an intimate partner [18]. A study conducted in
251 2020 in the United States reported that the prevalence of domestic violence during the
252 COVID-19 pandemic was 18%; and that changes in income and employment increased
253 the incidence of domestic violence by 1.63 times [19]. In another study, also from 2020
254 in the USA, it was reported that the incidence of physical domestic violence against
255 women was 1.8 times higher than before [20].

256 This is in addition to other important results that have been reported in other
257 realities, such as in Peru in 2020 where a 48% increase in the number of calls made to
258 the Domestic Violence Control and Assistance Unit was reported [21]. A meta-analysis
259 showing an increase in psychological/emotional as well as verbal violence in the
260 general population during the COVID-19 pandemic [22]. The World Vision report on
261 domestic violence hotline calls during the outbreak of the COVID-19 pandemic shows
262 that there was an increase in domestic violence cases worldwide; more specifically, in
263 Argentina (25%), Bosnia and Herzegovina (22%), Brazil (18%), Chile (75%), Cyprus
264 (47%), France (30%), India (32%); Lebanon (50%), Mexico (25%), Montenegro (27%),
265 Singapore (33%), Spain (12%) and in the United States (22%); which were mainly
266 related to intimate partner violence [23].

267 In South Africa, it was found that the fear and stress associated with pandemics
268 can exacerbate intimate partner violence through control, manipulation and coercion
269 tactics, as well as confinement measures that give abusers greater freedom to act
270 without inspection or consequences [24]. Latin America and the Caribbean experienced
271 increases in unemployment, poverty, food insecurity, etc. [25]. The prevalence of
272 mental health symptoms during the COVID-19 crisis in Latin America reported that
273 anxiety levels were significantly higher than in other regions, such as China (25%; $p <$

274 0.001) [26]. There was also an increase in reported cases of domestic violence,
275 particularly toward child abuse and intimate partner violence against women [27].
276 Exacerbating the high rates of violence in the region, which were reported to be three
277 times the global average before the pandemic [28]. And at the national level, Honduras
278 lives in a normalized environment of violence, from the political violence that the
279 country suffers, which is also influenced by living in a conservative environment and
280 finding 60% of low educational level and 8% with no education; where it can be
281 assumed that violence may not be identified or women are not allowed to talk about it
282 [29]. All these cases give us a picture of how we can be in an environment that
283 generates greater abuse for women and other vulnerable populations, which should be
284 evaluated with greater emphasis in future research.

285 We also found that the higher the level of family income, the less domestic
286 violence there was. Kebede et al. analyzed Demographic and Health Survey data from
287 187716 women with partners aged 10-59 years from 20 low-and middle-income
288 countries; where the findings suggest that the relationship between wealth and violence
289 varies considerably in low- and middle-income countries [30]. This is in addition to the
290 fact that the economic and labor crisis created by the pandemic could increase global
291 unemployment by almost 25 million, according to a new assessment by the
292 International Labor Organization (ILO) [31]. On the other hand, during the time of
293 confinement, an increasing number of poor families do not have a source of daily
294 income, which generates frustration and feelings of helplessness, all of which creates
295 conditions for increased violence [32].

296 This is reinforced by the finding that a structured questionnaire on intimate
297 partner violence found that wealthier and more empowered women reported less
298 intimate partner violence, and that younger women and those living in rural areas
299 tended to be more exposed to intimate partner violence [33]. That, if added to the other
300 factors that were generated by movement restrictions, increases in child abuse, internet
301 addiction, domestic violence and divorce consultations, as well as suicide rates [34]. All
302 of this together tells us of a large social component that is closely linked to the whole
303 picture of violence, which should be addressed in more detail by sociologists,
304 psychologists, and other professionals.

305 We also found in this study that the consumption of alcohol or drugs in the home
306 was associated with a higher frequency of domestic violence, it should be noted that the
307 relationship between the consumption of alcohol and psychoactive substances with
308 domestic violence is already well established [35, 36]. Having found clear evidence of a
309 temporal association between episodes of alcohol consumption and subsequent episodes
310 of occurrence of verbal and physical aggression [37]. In addition, the acceptance of
311 alcohol consumption by men normalizes consumption and amplifies other behaviors
312 tolerated in macho cultures, such as the use of violence [38]. This may also be due to
313 the accumulation of stressful events, which, in the context of the current pandemic was
314 precipitated by isolation, which favored the intensification of alcohol consumption
315 within the family environment [39]. A systematic review reported that frequent alcohol

316 consumption by men increased the risk of perpetration of violence [40]. Another meta-
317 analysis showed that people at risk for domestic violence and substance abuse are
318 chronically exposed to elevated levels of anxiety, highly stressful environments, and
319 unfavorable economic situations [41]. Because of all this evidence, it is important to
320 screen families who may have alcohol, drug, or other problems that are closely related
321 to domestic violence.

322 Our results reported that less domestic violence was found among single and
323 married people. This is important to analyze within Honduran society, which, like other
324 Latin American countries, has an important male chauvinist component, where gender
325 roles influence individual behavior and relationships, due to the power and control that
326 men try to impose on the entire family [42]. Therefore, what was reported is important
327 as a starting point to determine the role that men, women and other members of the
328 household play in the violence that may be perceived.

329 Yuan K, reported that women showed a higher prevalence of depression, anxiety
330 and insomnia, women younger than 40 years or in the upper income level had a lower
331 prevalence of depression and anxiety [43]. Having a history of mental illness was also
332 associated with a higher frequency of domestic violence. COVID-19 may have a
333 substantial impact on mental health and well-being. Jung et al showed that 25% of
334 participants reported pre-existing mental health problems and showed more depression
335 and anxiety, less well-being, less sense of coherence, and worse coping skills in terms
336 of the pandemic and actions taken [16]. Thus, intimate partner violence against women
337 is frequently and strongly associated with mental health problems [44]. Vai reported
338 that female sex, having a prior psychiatric diagnosis, and psychopathology at 1-month
339 follow-up were moderators of depression in the post-COVID-19 syndrome [45].

340 Finally, depression was associated with women suffering from domestic
341 violence, Robinson 2022 reported that there was a small increase in mental health
342 symptoms shortly after the outbreak of the COVID-19 pandemic in symptoms of
343 depression and mood disorders, which tended to be higher and remained significantly
344 elevated in May-July [46]. A systematic review found that suggested risk factors for
345 violence against women and girls after disasters include increased life stressors, law
346 enforcement failure, exposure to high-risk environments, exacerbation of existing
347 gender inequalities, and unequal social norms [47]. Which is increased when there is
348 domestic conflict and violence, financial loss, anxiety and depression, and pre-existing
349 mental health conditions, all of which should be considered to prevent suicide attempts
350 and deaths secondary to the COVID-19 pandemic [48].

351 **Limitations**

352 The main limitation was selection bias, since the data obtained cannot be
353 extrapolated to the entire Honduran population; however, by obtaining good statistical
354 power, the associations and results can be used to generate important conclusions that
355 can be used for the elaboration of research that aims to have more associated variables,
356 with a larger sample size and representativeness, as well as with more advanced

357 designs. In addition, there could have been a possible bias because some may have
358 hidden or changed their answers out of fear or embarrassment; however, it is believed
359 that, if it occurred, it was minimal, since the anonymous nature of the survey helped
360 them to have the confidence to answer truthfully.

361

362 **Conclusions:**

363

364 From all the findings it is concluded that women had a higher frequency of
365 domestic violence, the consumption of alcohol or drugs in the home, having a history of
366 mental illnesses such as depression, likewise, there was less frequency of domestic
367 violence according to a higher economic income in the home, between single and
368 married people.

369 The inequality of opportunities, the responsibility of Honduran households that
370 falls on women as heads of household, extreme poverty, and low schooling make
371 women more vulnerable to domestic violence. Therefore, it is important to continue
372 with this line of research in order to have real data at the national level and make
373 decisions to address this problem.

374 **Author Contributions:** Study design: CM, EET, LIZ, and CASM Data collection: CM,
375 HNCR Data analysis: CM, DAC Writing: CM, EET, LIZ, CASM, DAC, and HNCR, all
376 authors have read and agreed to the published version of the manuscript.

377 **Funding:** The current article processing charges (publication fees) were funded by the
378 Facultad de Ciencias Médicas (FCM) (2-03-01-01), Universidad Nacional Autónoma de
379 Honduras (UNAH), Tegucigalpa, MDC, Honduras, Central America (granted to Dra.
380 Espinoza).

381 **Institutional Review Board Statement:** The study was conducted under the
382 Declaration of Helsinki. This research's preparation and execution fully complied with
383 the fundamental ethical principles of autonomy, justice, beneficence, and non-
384 maleficence. The Act Number (2019062), approved by the Ethics Committee in
385 Biomedical Research (CEIB) of the National Autonomous University of Honduras
386 (UNAH), meeting of December 02, 2021.

387 **Data Availability Statement:** The data presented in this study are available on request
388 from the corresponding author.

389 **Acknowledgments:** To all MSS of the cohort who agreed to participate in the project,
390 from September to March 2021-2022 and from October-April 2021-2022, to Mr.
391 Mauricio Gonzales for the elaboration of the database and a first filtering of the
392 database.

393 **Conflicts of Interest:** The authors declare no conflict of interest.

394 **References**

- 395 1. Organización de las Naciones Unidas. ¿Qué es el abuso doméstico? consultado 20
396 de diciembre 2022. <https://www.un.org/en/coronavirus/what-is-domestic-abuse>.
- 397 2. WHO. Violence Against Women Prevalence Estimates, 2018. 2021.
398 [https://apps.who.int/iris/bitstream/handle/10665/341337/9789240022256-](https://apps.who.int/iris/bitstream/handle/10665/341337/9789240022256-eng.pdf?sequence=1)
399 [eng.pdf?sequence=1](https://apps.who.int/iris/bitstream/handle/10665/341337/9789240022256-eng.pdf?sequence=1).
- 400 3. El periódico. Las llamadas de víctimas del machismo suben el 20%. 2020
401 consultado 22 diciembre 2022.

- 402 <https://www.elperiodico.com/es/sociedad/20200325/violencia-machista-aumento->
403 [coronavirus-7902723.](https://www.elperiodico.com/es/sociedad/20200325/violencia-machista-aumento-)
- 404 4. Kelly J MT. Coronavirus: Domestic abuse calls up 25% since lockdown, charity
405 says. 6 abril 2020 consultado 22 de diciembre 2022. [https://www.bbc.com/news/uk-](https://www.bbc.com/news/uk-52157620)
406 [52157620.](https://www.bbc.com/news/uk-52157620)
- 407 5. Radio France Internationale (RFI). Covid-19: Women trapped between a deadly
408 virus and a deadly partner. 6 april 2020 consultado 22 de december 2022. Covid-19:
409 Women trapped between a deadly virus and a deadly partner.
- 410 6. Allen-Ebrahimian B. China's domestic violence epidemic. 2020; consultado 22
411 december 2022. [https://www.axios.com/2020/03/07/china-domestic-violence-](https://www.axios.com/2020/03/07/china-domestic-violence-coronavirus-quarantine)
412 [coronavirus-quarantine.](https://www.axios.com/2020/03/07/china-domestic-violence-coronavirus-quarantine)
- 413 7. UN Women. The Shadow Pandemic: Violence against women during COVID-19.
414 consultado 22 de diciembre 2022. [https://www.unwomen.org/es/news/in-focus/in-](https://www.unwomen.org/es/news/in-focus/in-focus-gender-equality-in-covid-19-response/violence-against-women-during-covid-19)
415 [focus-gender-equality-in-covid-19-response/violence-against-women-during-covid-](https://www.unwomen.org/es/news/in-focus/in-focus-gender-equality-in-covid-19-response/violence-against-women-during-covid-19)
416 [19.](https://www.unwomen.org/es/news/in-focus/in-focus-gender-equality-in-covid-19-response/violence-against-women-during-covid-19)
- 417 8. Instituto Nacional de Estadísticas (INE). ENCUESTA NACIONAL DE
418 DEMOGRAFÍA Y SALUD / ENCUESTA DE INDICADORES MÚLTIPLES
419 POR CONGLOMERADOS (ENDESA/MICS 2019). 2019 consultado diciembre
420 2022. [https://www.ine.gov.hk/V3/imag-doc/2021/10/Informe-ENDESA-MICS-](https://www.ine.gov.hk/V3/imag-doc/2021/10/Informe-ENDESA-MICS-2019.pdf)
421 [2019.pdf.](https://www.ine.gov.hk/V3/imag-doc/2021/10/Informe-ENDESA-MICS-2019.pdf)
- 422 9. UNAH-ONV-IUDPAS. MUERTE VIOLENTA de mujeres y femicidios en
423 Honduras [Boletín Infográfico]. Tegucigalpa, Honduras: Universidad Nacional
424 Autónoma de Honduras; Nov 25, 2022.
- 425 10. Chavarria-Mejia J, Calix-Pineda D, Vasquez-Bonilla WO. CARACTERIZACIÓN
426 DE VIOLENCIA DOMÉSTICA EN MUJERES MAYORES DE 16 AÑOS CON
427 COMPAÑERO EN COMUNIDAD DE HONDURAS 2016. CIMEL 2017.
428 doi:10.23961/cimel.v22i1.740.
- 429 11. Zavala GL, Montoya-Reales DA. Violencia contra la mujer en la relación de pareja;
430 caracterización en cinco comunidades de Honduras. Rev. fac. cienc. méd. (Impr.).
431 2017:16–27.
- 432 12. Espinoza-Turcios E, Gonzales-Romero RM, Sosa-Mendoza C, Sierra-Santos M,
433 Castro-Ramos HN, Zambrano LI, et al. Factors associated with hopelessness,
434 depression and anxiety in the Honduran-Central America population during the
435 COVID-19 pandemic. Front. Psychiatry. 2023;14:1116881.
436 doi:10.3389/fpsyt.2023.1116881.
- 437 13. Arora T, Grey I, Östlundh L, Lam KBH, Omar OM, Arnone D. The prevalence of
438 psychological consequences of COVID-19: A systematic review and meta-analysis
439 of observational studies. J Health Psychol. 2022;27:805–24.
440 doi:10.1177/1359105320966639.
- 441 14. Kourti A, Stavridou A, Panagouli E, Psaltopoulou T, Spiliopoulou C, Tsofia M, et
442 al. Domestic Violence During the COVID-19 Pandemic: A Systematic Review.
443 Trauma Violence Abuse. 2021:15248380211038690.
444 doi:10.1177/15248380211038690.
- 445 15. Piquero AR, Jennings WG, Jemison E, Kaukinen C, Knaul FM. Domestic violence
446 during the COVID-19 pandemic - Evidence from a systematic review and meta-
447 analysis. J Crim Justice. 2021;74:101806. doi:10.1016/j.jcrimjus.2021.101806.

- 448 16. Jung S, Kneer J, Krüger THC. Mental Health, Sense of Coherence, and
449 Interpersonal Violence during the COVID-19 Pandemic Lockdown in Germany. *J*
450 *Clin Med* 2020. doi:10.3390/jcm9113708.
- 451 17. Houseman B, Semien G. StatPearls: Florida Domestic Violence. Treasure Island
452 (FL); 2022.
- 453 18. UN woman. COVID-19 and Ending Violence Against Women and Girls.
454 consultado 20 de diciembre 2022.
455 [https://www.unwomen.org/sites/default/files/Headquarters/Attachments/Sections/Li](https://www.unwomen.org/sites/default/files/Headquarters/Attachments/Sections/Library/Publications/2020/Issue-brief-COVID-19-and-ending-violence-against-women-and-girls-en.pdf)
456 [brary/Publications/2020/Issue-brief-COVID-19-and-ending-violence-against-](https://www.unwomen.org/sites/default/files/Headquarters/Attachments/Sections/Library/Publications/2020/Issue-brief-COVID-19-and-ending-violence-against-women-and-girls-en.pdf)
457 [women-and-girls-en.pdf](https://www.unwomen.org/sites/default/files/Headquarters/Attachments/Sections/Library/Publications/2020/Issue-brief-COVID-19-and-ending-violence-against-women-and-girls-en.pdf).
- 458 19. Jetelina KK, Knell G, Molsberry RJ. Changes in intimate partner violence during
459 the early stages of the COVID-19 pandemic in the USA. *Inj Prev*. 2021;27:93–7.
460 doi:10.1136/injuryprev-2020-043831.
- 461 20. Gosangi B, Park H, Thomas R, Gujrathi R, Bay CP, Raja AS, et al. Exacerbation of
462 Physical Intimate Partner Violence during COVID-19 Pandemic. *Radiology*.
463 2021;298:E38-E45. doi:10.1148/radiol.2020202866.
- 464 21. Agüero JM. COVID-19 and the rise of intimate partner violence. *World Dev*.
465 2021;137:105217. doi:10.1016/j.worlddev.2020.105217.
- 466 22. Thiel F, Büechl VCS, Rehberg F, Mojahed A, Daniels JK, Schellong J, Garthus-
467 Niegel S. Changes in Prevalence and Severity of Domestic Violence During the
468 COVID-19 Pandemic: A Systematic Review. *Front Psychiatry*. 2022;13:874183.
469 doi:10.3389/FPSYT.2022.874183.
- 470 23. World Vision. Repercusiones del COVID-19 UNA TORMENTA PERFECTA: MÁS
471 MILLONES DE NIÑOS CORREN EL RIESGO DE SUFRIR VIOLENCIA
472 DURANTE EL CONFINAMIENTO Y LA “NUEVA NORMALIDAD”. 2020.
473 https://www.wvi.org/sites/default/files/2020-06/COVID-19%20aftershocks_SP.pdf.
- 474 24. Bianca Dekel NA. ‘I will rather be killed by corona than by him...’: Experiences of
475 abused women seeking shelter during South Africa’s COVID-19 lockdown. *Plos*
476 *One*. 2021.
- 477 25. ECLAC. The recovery paradox in Latin America and the Caribbean Growth amid
478 persisting structural problems: inequality, poverty and low investment and
479 productivity. 2021. [https://www.cepal.org/en/publications/47059-recovery-paradox-](https://www.cepal.org/en/publications/47059-recovery-paradox-latin-america-and-caribbean-growth-amid-persisting-structural)
480 [latin-america-and-caribbean-growth-amid-persisting-structural](https://www.cepal.org/en/publications/47059-recovery-paradox-latin-america-and-caribbean-growth-amid-persisting-structural).
- 481 26. Zhang SX, Batra K, Xu W, Liu T, Dong RK, Yin A, et al. Mental disorder
482 symptoms during the COVID-19 pandemic in Latin America - a systematic review
483 and meta-analysis. *Epidemiol Psychiatr Sci*. 2022;31:e23.
484 doi:10.1017/S2045796021000767.
- 485 27. WHO. ADDRESSING VIOLENCE AGAINST CHILDREN, WOMEN AND
486 OLDER PEOPLE DURING THE COVID-19 PANDEMIC: KEY ACTIONS. 2020.
487 [https://apps.who.int/iris/bitstream/handle/10665/332458/WHO-2019-nCoV-](https://apps.who.int/iris/bitstream/handle/10665/332458/WHO-2019-nCoV-Violence_actions-2020.1-eng.pdf?sequence=1&isAllowed=y)
488 [Violence_actions-2020.1-eng.pdf?sequence=1&isAllowed=y](https://apps.who.int/iris/bitstream/handle/10665/332458/WHO-2019-nCoV-Violence_actions-2020.1-eng.pdf?sequence=1&isAllowed=y).
- 489 28. PAHO. Violence in the Americas. What must be done? 2020.
490 [https://www.paho.org/en/documents/infographic-violence-americas-what-must-be-](https://www.paho.org/en/documents/infographic-violence-americas-what-must-be-done)
491 [done](https://www.paho.org/en/documents/infographic-violence-americas-what-must-be-done).
- 492 29. Valladares D, Martínez Linares R, Pérez Ramirez I, Moncada G. Prevalencia de
493 trastornos mentales en relación con violencia, resiliencia y estrategias de

- 494 afrontamiento en Yoro, Choluteca y Cortés, junio, 2016. Portal de la Ciencia.
495 2018:139–57. doi:10.5377/pc.v0i14.6645.
- 496 30. Kebede S, van Harmelen A-L, Roman-Urrestarazu A. Wealth Inequality and
497 Intimate Partner Violence: An Individual and Ecological Level Analysis Across 20
498 Countries. *J Interpers Violence*. 2022;37:NP15568-NP15593.
499 doi:10.1177/08862605211016337.
- 500 31. International labour organization. Almost 25 million jobs could be lost worldwide
501 as a result of COVID-19, says ILO. consultado 21 de diciembre 2022.
502 [https://www.ilo.org/global/about-the-ilo/newsroom/news/WCMS_738742/lang--](https://www.ilo.org/global/about-the-ilo/newsroom/news/WCMS_738742/lang--en/index.htm)
503 en/index.htm.
- 504 32. Singh S, Roy D, Sinha K, Parveen S, Sharma G, Joshi G. Impact of COVID-19 and
505 lockdown on mental health of children and adolescents: A narrative review with
506 recommendations. *Psychiatry Res*. 2020;293:113429.
507 doi:10.1016/j.psychres.2020.113429.
- 508 33. Coll CVN, Ewerling F, García-Moreno C, Hellwig F, Barros AJD. Intimate partner
509 violence in 46 low-income and middle-income countries: an appraisal of the most
510 vulnerable groups of women using national health surveys. *BMJ Glob Health*.
511 2020;5:e002208. doi:10.1136/bmjgh-2019-002208.
- 512 34. Saeko Kamoshida, Naoto Nihonmatsu, Koubun Wakashima. The relationship
513 between family variables and family social problems during the COVID-19
514 pandemic. *Plos One*. 2022.
- 515 35. Fonseca AM, Galduróz JCF, Tondowski CS, Noto AR. Alcohol-related domestic
516 violence: a household survey in Brazil. *Rev Saude Publica*. 2009;43:743–9.
517 doi:10.1590/s0034-89102009005000049.
- 518 36. A violência doméstica a partir do discurso de mulheres agredidas; 2013.
- 519 37. Testa M, Derrick JL. A daily process examination of the temporal association
520 between alcohol use and verbal and physical aggression in community couples.
521 *Psychol Addict Behav*. 2014;28:127–38. doi:10.1037/a0032988.
- 522 38. Bonnevie E, Kigozi G, Kairania R, Ssemanda JB, Nakyanjo N, Ddaaki WG, et al.
523 Alcohol use in fishing communities and men's willingness to participate in an
524 alcohol, violence and HIV risk reduction intervention: qualitative findings from
525 Rakai, Uganda. *Cult Health Sex*. 2020;22:275–91.
526 doi:10.1080/13691058.2019.1587002.
- 527 39. Campbell AM. An increasing risk of family violence during the Covid-19
528 pandemic: Strengthening community collaborations to save lives. *Forensic Science*
529 *International: Reports*. 2020;2:100089. doi:10.1016/j.fsir.2020.100089.
- 530 40. Wright MMM, Kankkunen PM, Jokiniemi KS. Prevention interventions for
531 interpersonal violence occurring under the influence of alcohol: A mixed method
532 systematic review. *J Adv Nurs* 2022. doi:10.1111/jan.15335.
- 533 41. Abdo C, Miranda EP, Santos CS, Júnior JdB, Bernardo WM. Domestic violence
534 and substance abuse during COVID19: A systematic review. *Indian J Psychiatry*.
535 2020;62:S337-S342. doi:10.4103/psychiatry.IndianJPsychiatry_1049_20.
- 536 42. Mancera BM, Dorgo S, Provencio-Vasquez E. Risk Factors for Hispanic Male
537 Intimate Partner Violence Perpetration. *Am J Mens Health*. 2017;11:969–83.
538 doi:10.1177/1557988315579196.
- 539 43. Yuan K, Zheng Y-B, Wang Y-J, Sun Y-K, Gong Y-M, Huang Y-T, et al. A
540 systematic review and meta-analysis on prevalence of and risk factors associated

- 541 with depression, anxiety and insomnia in infectious diseases, including COVID-19:
542 a call to action. *Mol Psychiatry*. 2022;27:3214–22. doi:10.1038/s41380-022-01638-
543 z.
- 544 44. Hameed M, O'Doherty L, Gilchrist G, Tirado-Muñoz J, Taft A, Chondros P, et al.
545 Psychological therapies for women who experience intimate partner violence.
546 *Cochrane Database Syst Rev*. 2020;7:CD013017.
547 doi:10.1002/14651858.CD013017.pub2.
- 548 45. Vai B, Mazza MG, Delli Colli C, Foiselle M, Allen B, Benedetti F, et al. Mental
549 disorders and risk of COVID-19-related mortality, hospitalisation, and intensive
550 care unit admission: a systematic review and meta-analysis. *Lancet Psychiatry*.
551 2021;8:797–812. doi:10.1016/S2215-0366(21)00232-7.
- 552 46. Robinson E, Sutin AR, Daly M, Jones A. A systematic review and meta-analysis of
553 longitudinal cohort studies comparing mental health before versus during the
554 COVID-19 pandemic in 2020. *J Affect Disord*. 2022;296:567–76.
555 doi:10.1016/j.jad.2021.09.098.
- 556 47. Thurston AM, Stöckl H, Ranganathan M. Natural hazards, disasters and violence
557 against women and girls: a global mixed-methods systematic review. *BMJ Glob
558 Health* 2021. doi:10.1136/bmjgh-2020-004377.
- 559 48. Pathirathna ML, Nandasena HMRKG, Atapattu AMMP, Weerasekara I. Impact of
560 the COVID-19 pandemic on suicidal attempts and death rates: a systematic review.
561 *BMC Psychiatry*. 2022;22:506. doi:10.1186/s12888-022-04158-w.