Autoantigen profiling reveals a shared post-COVID signature in fully recovered and Long COVID patients

Aaron Bodansky,[1] Chung-Yu Wang,[2] Aditi Saxena, [2] Anthea Mitchell, [2] Saki Takahashi,[3] Khamal Anglin,[4] Beatrice Huang,[3] Rebecca Hoh,[3] Scott Lu,[4] Sarah A. Goldberg,[4] Justin Romero,[3] Brandon Tran,[3] Raushun Kirtikar,[3] Halle Grebe,[3] Matthew So,[3] Bryan Greenhouse,[3] Matthew S. Durstenfeld,[5] Priscilla Y. Hsue,[5] Joanna Hellmuth,[6] J. Daniel Kelly,[4] Jeffrey N. Martin,[4] Mark S. Anderson,[7,8] Steven G. Deeks,[3] Timothy J. Henrich,[9] Joseph L. DeRisi,[2] Michael J. Peluso[3]

Division of Pediatric Critical Care Medicine, University of California, San Francisco, San Francisco, CA, USA

Chan Zuckerberg Biohub, San Francisco, CA, USA

Division of HIV, Infectious Diseases, and Global Medicine, University of California, San Francisco, San Francisco, CA, USA

Department of Epidemiology and Biostatistics, University of California, San Francisco, San Francisco, CA, USA

Division of Cardiology, University of California, San Francisco, San Francisco, CA, USA

Department of Neurology University of California, San Francisco, San Francisco, CA, USA

Diabetes Center, University of California, San Francisco, San Francisco, CA, USA

Department of Medicine, University of California, San Francisco, San Francisco, CA, USA

Division of Experimental Medicine, University of California, San Francisco, San Francisco, CA, USA

26 Corresponding Author: Michael Peluso, MD

27   Division of HIV, Infectious Diseases, and Global Medicine

28   University of California, San Francisco

29   San Francisco, CA, USA

30   Email: michael.peluso@ucsf.edu

Keywords: SARS-CoV-2, COVID-19, post-acute sequelae of SARS-CoV-2 infection (PASC),

Long COVID, autoreactivity, autoimmunity, autoantigen

ABSTRACT

Some individuals do not return to baseline health following SARS-CoV-2 infection, leading to a

condition known as Long COVID. The underlying pathophysiology of Long COVID remains

unknown. Given that autoantibodies have been found to play a role in severity of COVID infection

and certain other post-COVID sequelae, their potential role in Long COVID is important to

investigate. Here we apply a well-established, unbiased, proteome-wide autoantibody detection

technology (PhIP-Seq) to a robustly phenotyped cohort of 121 individuals with Long COVID, 64

individuals with prior COVID-19 who reported full recovery, and 57 pre-COVID controls. While a

distinct autoreactive signature was detected which separates individuals with prior COVID

infection from those never exposed to COVID, we did not detect patterns of autoreactivity that

separate individuals with Long COVID relative to individuals fully recovered from SARS-CoV-2

infection. These data suggest that there are robust alterations in autoreactive antibody profiles

due to infection; however, no association of autoreactive antibodies and Long COVID was

apparent by this assay.

BACKGROUND

Some individuals do not fully return to baseline health following SARS-CoV-2 infection and

experience ongoing morbidity following the acute phase of COVID-19 (1, 2). There is now intense

interest in determining the underlying mechanisms of Long COVID, one type of post-acute

sequelae of SARS-CoV-2 infection (PASC) characterized by symptoms that newly developed or

worsened following infection that cannot be clearly attributed to another cause (3). Immune

dysregulation, including the generation of antibodies against self antigens, has been suggested as

one potential driver of Long COVID that warrants further investigation (4, 5).

Acute SARS-CoV-2 infection is associated with the generation of autoreactive antibodies,

particularly among individuals with severe disease requiring hospitalization (6–8). For example,

one study of 147 individuals hospitalized with COVID-19 found that autoantibodies associated with

connective tissue diseases and anti-cytokine antibodies were identified in 50% of samples, and

tracked with the humoral response to SARS-CoV-2 infection (7). Furthermore, multiple studies

have described the contribution of likely pre-existing anti-interferon antibodies to vaccine

breakthrough infections and severe manifestations of COVID-19, including death (9–12).

Although much work has been done exploring the potential virologic and immunologic factors

driving Long COVID, the evaluation of autoantibodies in this condition has been more limited.

Measurement of anti-nuclear antibodies on standard clinical tests has yielded mixed findings, with

some studies identifying a high prevalence of antinuclear antibodies (ANAs) among those with

Long COVID (13–15) and other studies finding low prevalence consistent with ANA positivity in

the general population (16–18). Although perturbations in interferon signaling pathways have been

suggested as one potential mechanism of Long COVID (4, 15), anti-interferon antibodies have not

been identified in most individuals outside of those who had severe acute infection (16). This is

consistent with other post-acute sequelae of COVID such as multisystem inflammatory syndrome

in children (MIS-C), which also has no association with anti-interferon antibodies (19).

The identification of previously reported autoantibodies can be performed using targeted assays.

However, identifying the full range of novel autoreactive antibodies, whether they are pathological

or not, requires technologies capable of high-throughput, unbiased, proteome-wide screens. In

this study, we screened a cohort of individuals with prior SARS-CoV-2 infection, many of whom

met clinical criteria for Long COVID, to determine whether a consistent pattern of autoreactivity

could be identified. This same technology has been previously utilized to discover novel

autoantibodies in a wide range of disease contexts (20–23).

RESULTS

Distinct set of autoreactive antibodies in individuals with prior SARS-CoV-2 infection

We employed a previously published proteome-wide approach using a T7 phage-display assay

with immunoprecipitation and next-generation sequencing (PhiP-Seq) (20–24). We tested sera

from 185 otherwise healthy individuals with prior SARS-CoV-2 infection in parallel to sera from 57

otherwise healthy individuals collected prior to the known existence of COVID-19 (pre-COVID).

Using an unbiased analysis, we identified a distinct pattern of autoreactivity which effectively

classifies individuals with prior SARS-CoV-2 infection from individuals not yet exposed to the virus

with a logistic regression AUC of 0.90 (Figure 1A). The protein targets from which these enriched

immunoprecipitated peptides were derived are widely varied and lack any apparent shared

biological functions or cell type (Figure 1B). Among the identified targets, a single peptide derived

from ARHGAP31 (Figure 2A) displayed the greatest amount of enrichment with 22% of individuals

with prior SARS-CoV-2 infection yielding enrichment greater than 6 standard deviations of the

mean of the pre-COVID controls. No difference in enrichment was observed for those diagnosed

with Long COVID with respect to Post-COVID (Figure 2B).

To investigate whether the ARHGAP31 peptide enrichment could be the result of cross-reactivity

with antibodies directed against SARS-CoV-2 proteins, we performed a multiple sequence

alignment of this 49-amino acid fragment against the full SARS-CoV-2 proteome. A region within

the SARS-CoV2 Orf1a polyprotein was identified with considerable physico-chemical similarity to

a portion of the autoreactive fragment in ARHGAP31(Jalview Version 2;Figure 2C), supporting the

notion that the observed human peptidome peptide enrichments in post-COVID samples are being

driven by anti-SARS-CoV-2 antibodies.

ARHGAP31 is most highly expressed in neuronal cells, Langerhans cells, and endothelial cells

(Human Protein Atlas, proteinatlas.org). We were unable to identify any clinical differences

between individuals with and without ARHGAP31 autoreactivity, such as differences in frequency

of neurologic symptoms.

Post-COVID autoreactivities are not enriched in individuals with Long COVID

To assess whether autoreactive antibodies present in individuals following SARS-CoV-2 infection

are associated with Long COVID, we compared the distribution of the enriched post-COVID

peptides amongst the 121 individuals with Long COVID and the 64 individuals with prior SARS-

CoV-2 infection but without Long COVID (Convalescent COVID). The 20 most enriched proteins

with at least 5-fold greater than background (defined as fold-change over mock-IP with protein

A/G beads) were compared (Figure 3). Seventeen of the 20 enriched proteins were present in

both Long COVID and Convalescent COVID, and none of these enrichments was observed in any

of the pre-COVID controls. Overall, there was no significant differences in enrichment between

Long COVID and Convalescent COVID. Peptides derived from three proteins, TMED10, FUCA1, and POL2RK, were only observed in Long COVID, however these were infrequently observed.

Subgroup analyses were performed to determine whether any of these observed autoreactivities were enriched in particular symptom-defined phenotypes of Long COVID. These include cardiopulmonary (cough, shortness of breath, chest pain, palpitations, and fainting), central neurologic symptoms (problems with vision, headache, difficulty with concentration or memory, dizziness, and difficulty with balance), any neurologic symptom (problems with vision, headache, difficulty with concentration or memory, dizziness, difficulty with balance, trouble with smell or taste, phantosmia, or paresthesia), gastrointestinal (diarrhea, constipation, nausea, vomiting, loss of appetite, and abdominal pain), musculoskeletal (back pain, muscle pain, pain in the arms, legs, or joints), and upper respiratory (runny nose and sore throat).

The peptide enrichments from individuals with particular Long COVID phenotypes were compared to the enrichments from individuals with Convalescent COVID (no Long COVID symptoms) using a one-sided Kolmogorov-Smirnov test. None of the top 20 autoantibodies were enriched in severe Long COVID, and only 3 autoantibodies were statistically increased in any phenotype: TTF2 and KDM3B in those with cardiopulmonary symptoms and FUCA1 in those with upper respiratory symptoms (Figure 4A). However, using a strict cutoff of 6 standard deviations above the pre-COVID controls to determine positivity, none of these autoantibodies was phenotype specific. By looking at the distribution of these antibodies in Long COVID patients with given sub-phenotype relative to the remaining Long COVID patients and all convalescent COVID patients, it was apparent that the statistical significance in a particular phenotype was driven by either a single individual with extremely high autoantibody signal in the case of KDM3B and TTF2, or higher group signal but not meeting the positive threshold cutoff in the case of FUCA1 (Figure 4B).

Absence of Long COVID specific autoreactivities

In addition to analyzing the distribution of post-COVID autoreactive peptide enrichments, we also performed additional analyses to detect those enrichments which might be present only in Long COVID or particular subcategories of Long COVID. To feature-weight enriched peptides, we applied logistic regressions to the degree of enrichment for each individual with Long COVID and each individual Long COVID symptom phenotype versus enrichment from convalescent COVID patients who did not have the particular symptoms. We included all previously discussed symptom phenotypes, as well as individuals with difficulty concentrating, worsened quality of life, new depression, and generalized anxiety disorder. We were unable to identify a set of enriched proteins specific to any of these groups of patients which could effectively distinguish the cohort from controls, as the best receiver operating characteristic (ROC) area under the curve (AUC) was 0.67, and the mean AUC was 0.44 (Supplement 1).

DISCUSSION

Autoimmunity has been proposed as one potential mechanism driving Long COVID. We applied an unbiased, proteome-wide, validated approach to assess associations between antibody autoreactivity and clinical phenotype. A clear and robust difference in autoreactivity was detected between those infected with SARS-CoV-2 and pre-COVID controls. This difference was constituted by peptides from diverse and varied proteins, most of which are intracellular, suggesting that the origin of the differential enrichment is due to cross-reactivity with SARS-CoV-2 directed antibodies in those that were exposed. A sequence comparison between a peptide from the most enriched protein, ARHGAP31, and Orf1a of SARS-CoV-2 supports this notion, but orthogonal validation through fine-scale epitope mapping and antibody cloning would be required to demonstrate this conclusively. While the clinical significance of incidental autoreactivity due to the humoral immune response to SARS-CoV-2 remains largely unknown, prior studies have

identified cross reactive autoantibodies in severe sequelae of SARS-CoV-2 infection, including

those that develop severe neurological symptoms (Song et al., 2021). Understanding the clinical

consequences of SARS-CoV-2 driven autoreactivity deserves further attention, perhaps through

long-term longitudinal studies.

We found no association to support the hypothesis that autoreactivity, as detected in this assay,

contributes to Long COVID. Despite numerous successes, PhIP-seq possesses a number of

limitations. Because the T7 phage displayed peptides are only 49 amino acids, the assay

inherently detects mostly linear epitopes. Therefore, complex conformational, post translationally

modified, or multimeric protein configurations are not predicted to be detectable by this assay, and

thus these results do not completely rule out autoimmune interactions in Long COVID that are

beyond the scope of PhIP-seq.

In addition, our classification of Long COVID and its symptom phenotypes was based exclusively

on participant self-report of symptoms. It is possible that future analyses in more homogenous

cohorts, particularly those with objectively measured physiologic perturbations now associated

with certain Long COVID phenotypes (e.g., postural orthostasis/tachycardia, neurocognitive

function deficits, abnormalities on cardiopulmonary exercise testing) may yet reveal a role of

autoantibodies in at least a subset of individuals experiencing Long COVID.

Long COVID remains a complex clinical entity. Its causes are likely multifactorial, and there is

growing consensus that different phenotypes are driven by different pathophysiologic mechanisms

(2, 3). Additional work characterizing SARS-CoV-2 specific and autoreactive immune responses in

large, well-characterized cohorts over time, during both the acute and post-acute phases of the

illness, will be necessary to delineate the biology of Long COVID and other post-acute sequelae of

SARS-CoV-2 infection, and to lead to the development of potential interventions to treat the

millions of individuals currently affected by this condition.

METHODS

Study participants and measurements

Participants were volunteers in the UCSF Long-term Impact of Infection with Novel Coronavirus

(LIINC) study (NCT04362150). The details of study design and measurement have been reported

previously (25). For the current analysis, we included 185 individuals with a history of nucleic acid-

confirmed SARS-CoV-2 infection who had a plasma sample collected between 60 and 240 days

following initial symptom onset or, if asymptomatic, first positive SARS-CoV-2 test (Supplemental

Table 1). Long COVID was defined using study instruments, which have been described in detail

elsewhere (25). Briefly, each participant was queried regarding the presence, severity, and

duration of 32 physical and mental health symptoms and quality of life. Details of their medical

history and COVID-19 infection and vaccination and treatment history were also recorded.

Symptoms that pre-dated SARS-CoV-2 infection and that were not changed following infection, as

well as those obviously attributed to another cause (e.g. ankle fracture), were not considered to

represent Long COVID. All samples included were collected prior to the subject ever receiving a

SARS-CoV-2 vaccine.

Because of challenges in objectively defining Long COVID and to thoroughly explore the data in

an unbiased manner, we utilized a number of predefined case definitions in our analysis. We

constructed case definitions based on symptom presentation and quality-of-life (QOL) responses.

Symptoms case definitions include: the presence of any new or worsening symptoms since

SARS-CoV-2 infection (Long COVID); presence of 5 or more symptoms (severe Long COVID);

specific symptom groups according to organ system involvement or phenotypic cluster; and

individual symptoms when at least 25 individuals experienced the symptom. For individual and

grouped symptom outcomes, we developed 3 potential comparisons between the symptomatic

group and (1) all individuals who reported absence of the symptoms of interest, regardless of

Long COVID status (2) only individuals who were consistently asymptomatic and (3) individuals

with Long COVID but not the symptoms of interest. Severe Long COVID was compared only to

those without any reported symptoms.

We defined 6 groups of symptoms (symptom phenotypes) based on either organ system cluster.

These include cardiopulmonary (cough, shortness of breath, chest pain, palpitations, and fainting),

CNS-specific (problems with vision, headache, difficulty with concentration or memory, dizziness,

and difficulty with balance), any neurologic symptom (problems with vision, headache, difficulty

with concentration or memory, dizziness, difficulty with balance, trouble with smell or taste,

phantosmia, or paresthesia), gastrointestinal (diarrhea, constipation, nausea, vomiting, loss of

appetite, and abdominal pain), musculoskeletal (back pain, muscle pain, pain in the arms, legs, or

joints), and upper respiratory (runny nose and sore throat).

Quality-of-life was assessed using the EuroQol-5D (EQ-5D), Patient Health Questionnaire

depression scale (PHQ-8), and Generalized Anxiety Disorder scale (GAD-7). Individuals with

Long COVID and the lowest overall quality-of-life (QOL) score measured via the visual analogue

scale of the EQ-5D were compared to individuals with the highest overall QOL scores among

those with and without Long COVID. Individuals with responses categorized as "moderate

depression" on the PHQ-8 (score higher than 10) and "moderate anxiety" on the GAD-7 (score

higher than 9) were compared to all participants with scores indicating less severe classifications

than "moderate depression" and "moderate anxiety" in the following groups: (1) all participants

regardless of Long COVID status (2) all participant without Long COVID and (3) all participants

with Long COVID.

In addition, we recently demonstrated associations between Long COVID and other chronic latent

viral infections, including serologic evidence suggesting recent Epstein-Barr virus (EBV)

reactivation (26). For this reason, we also used binary variables to create groups indicating the

presence of this condition.

Biospecimen collection

At each visit, whole blood was collected in EDTA tubes. Plasma was isolated and stored in -80F

until the time of analysis.

Phage Immunoprecipitation and Sequencing (PhIP-Seq)

PhIP-Seq was performed following our previously published vacuum-based PhIP-Seq protocol

(23) (https://www.protocols.io/view/scaled-high-throughput-vacuum-phip-protocol-

ewov1459kvr2/v1).

PhIP-Seq analysis

All analysis (except when specifically stated otherwise) was performed at the gene-level, in which

all reads for all peptides mapping to the same gene were summed, and 0.5 reads were added to

each gene to allow inclusion of genes with zero reads in mathematical analyses. Within each

individual sample, reads were normalized by converting to the percentage of total reads. To

normalize each sample against background non-specific binding, a fold-change(FC) over mock-IP

was calculated by dividing the sample read percentage for each gene by the mean read-

percentage of the same gene for the AG bead only controls. This FC signal was then used for

side-by-side comparison between samples and cohorts. Samples which had a FC of 5 or greater

were considered enriched for an antibody, and samples with a FC of 6 standard deviations above

the mean of pre-COVID controls were considered positive for an autoantibody. FC values were

also used to calculate z-scores for each disease category sample by using each respective control

(as specified in figures and results), and for each control sample by using all remaining controls.

These z-scores were used for the logistic-regression feature weighting. In the case peptide-level

analysis, raw reads were normalized by calculating the number of reads per 100,000 reads.

Statistical methods

All statistical analysis was performed in Python using the Scipy Stats package. A two-way

Kolmogorov-Smirnoff(KS) test was used for comparisons of FC PhIP-Seq data between groups of

samples, except in the case of specifically looking for those genes with increased signal only in

the disease-cohort in which a one-way KS test was employed. The logistic regression machine-

learning classifiers were performed using our recently described methods (23). Utilizing the Scikit-

learn package, logistic regression classifiers were applied to z-scored PhIP-Seq values from

individuals with a designated disease category versus the designated control. A liblinear solver

was used with L1 regularization, and the model was evaluated using a five-fold cross-validation (4

of the 5 for training, 1 of the 5 for testing).

Informed consent

Participants provided written informed consent. The study was approved by the UCSF Institutional

Review Board.

FOOTNOTES

Acknowledgements

We are grateful to the study participants and their medical providers. We acknowledge current and former LIINC clinical study team members Tamara Abualhsan, Andrea Alvarez, Khamal Anglin, Urania Argueta, Mireya Arreguin, Kofi Asare, Melissa Buitrago, Monika Deswal, Nicole DelCastillo, Emily Fehrman, Halle Grebe, Heather Hartig, Yanel Hernandez, Rebecca Hoh, Beatrice Huang, Marian Kerbleski, Raushun Kirtikar, James Lombardo, Monica Lopez, Michael Luna, Sadie Munter, Lynn Ngo, Enrique Martinez Ortiz, Antonio Rodriguez, Justin Romero, Dylan Ryder, Ruth Diaz Sanchez, Matthew So, Celina Chang Song, Viva Tai, Alex Tang, Cassandra Thanh, Fatima Ticas, Leonel Torres, Brandon Tran, Daisy Valdivieso, Deepshika Varma, and Meghann Williams; and LIINC laboratory team members Amanda Buck, Tyler-Marie Deveau, Joanna Donatelli, Jill Hakim, Nikita Iyer, Owen Janson, Brian LaFranchi, Christopher Nixon, Isaac Thomas, and Keirstinne Turcios. We thank Jessica Chen, Aidan Donovan, Carrie Forman, and Rania Ibrahim for assistance with data entry and review. We thank the UCSF AIDS Specimen Bank for processing specimens and maintaining the LIINC biospecimen repository. We are grateful to Elnaz Eilkhani and Monika Deswal for regulatory support.

Author Contributions

AB, ST, BG, JDT, JNM, SGD, TJH, JLD, and MJP designed the study. JDT, JNM, SGD, TJH, and MJP designed and oversaw the LIINC cohort, with major contributions from BG, MSD, PYH, and JH. KA, BH, and RH managed cohort operations. Clinical data were collected by JR, BT, RK, HG, and MS. SL and SG managed the data and developed the clinical dataset. ST, SL, SG, and MJP selected participants for inclusion in this study. Biospecimens were processed in the laboratories of BG and TJH. CW, AS, and AM performed the measurements in the laboratories of MSA and JLD. AB performed the data analysis. AB, TJH, JLD, and MJP wrote the first draft of the paper, with input from all other authors. All authors edited the manuscript and approved the final version.

Funding

This work was supported by NIH/NIAID 3R01AI141003-03S1 and NIH/NIAID R01AI158013 (to M

Gandhi and M Spinelli), Pediatric Scientist Development Program and the Eunice Kennedy

Shriver National Institute of Child Health and Human Development (K12-HD000850 to A.

Bodansky), and Chan Zuckerberg Biohub support for J. DeRisi.

Conflicts of Interest

TJH reports grants from Merck & Co, Gilead Sciences, and Bristol-Myers Squibb, and has

provided consulting for Roche, outside the submitted work. SGD reports grants and/or personal

fees from Gilead Sciences, Merck & Co., Viiv, AbbVie, Eli Lilly, ByroLogyx, and Enochian

Biosciences, outside the submitted work. JD reports paid compensation for consulting for the

Public Health Company and Allen & Co. MJP reports consulting fees from Gilead Sciences and

AstraZeneca, outside the submitted work. All other authors report no potential conflicts.

FIGURE LEGENDS

Figure 1: PhIP-Seq autoreactivities distinguish post-COVID sera from pre-COVID controls.

Logistic regression comparing PhIP-Seq autoreactivities in all individuals with prior COVID

infection compared to pre-COVID controls. Barplot showing autoreactivities with the top 20 logistic

regression coefficients.

Figure 2: Post-COVID anti-ARHGAP31 autoreactivities target specific region with similarity

to SARS-CoV-2

(A) Stripplots showing distribution of ARHGAP31 autoreactivities in Long COVID, individuals with

prior COVID infection but without Long COVID, and pre-COVID controls. Dotted line at 6 standard

deviations above mean of pre-COVID controls. (B) Distribution of anti-ARHGAP31 autoreactivity

signal within ARHGAP31 full length protein. One specific fragment is targeted. (C) Amino-acid

sequence of the autoreactive region of ARHGAP31 and amino-acid sequence of a region of

SARS-CoV-2 Orf1a with similarity. Shown below is the multiple sequence

alignment(ClustalOmega; asterisk=identical amino acid; colon=strongly similar properties with

Gonnet PAM 250 matrix score >0.5; period=weakly similar with Gonnet PAM 250 matrix score

between 0 and 0,5) and strong physical-chemical conservation (JalView; amino acid physical-

chemical conservation scored on a scale of 1-11, Asterisk=score of 11 and identical amino acid,

Plus=10, all properties conserved).

Figure 3: Post-COVID autoreactivities are similarly distributed among Long COVID and

controls.

Hierarchically clustered (Pearson) heatmaps showing the PhIP-Seq enrichment for the top 20

autoreactivities ranked by logistic regression coefficient in each Long COVID patient, each COVID

convalescent patient, and each pre-COVID control.

Figure 4: Few significantly increased autoreactivities in Long COVID symptom phenotypes.

(A) Heatmap with p-values (Kolmogorov-Smirnov testing) of differences in autoantigen enrichment

for all individuals with prior COVID infection with and without additional clinical factors. Top-row

compares those with and without Long COVID. Lower rows show subcategories of Long COVID.

(B) Stripplots showing the three autoantibodies with statistically significant enrichment in a post-

COVID clinical phenotype. Dotted-lines show 6 standard-deviations above the mean of pre-

COVID signal.

Supplemental Table 1: Characteristics of study participants.

Characteristics of all study participants, stratified by Long COVID status.

Supplemental Figure 1: PhIP-Seq is unable to distinguish Long COVID symptom

phenotypes from controls.

(A) Logistic regression receiver operating characteristic (ROC) curves for Long COVID patients

with different specified symptom phenotypes relative to patients previously infected with COVID

without the phenotype, or (B) with and without EBV laboratory findings.

REFERENCES

1. Nalbandian A et al. Post-acute COVID-19 syndrome [Internet]. *Nat. Med.* [published online

ahead of print: March 22, 2021]; doi:10.1038/s41591-021-01283-z

2. Davis HE, McCorkell L, Vogel JM, Topol EJ. Long COVID: major findings, mechanisms and

recommendations. *Nat. Rev. Microbiol.* 2023;1–14.

3. Peluso MJ, Deeks SG. Early clues regarding the pathogenesis of long-COVID. *Trends

Immunol.* 2022;43(4):268–270.

4. Proal AD, VanElzakker MB. Long COVID or Post-acute Sequelae of COVID-19 (PASC): An

Overview of Biological Factors That May Contribute to Persistent Symptoms [Internet]. *Frontiers in

Microbiology* 2021;12. doi:10.3389/fmicb.2021.698169

5. Peluso MJ, Donatelli J, Henrich TJ. Long-term immunologic effects of SARS-CoV-2 infection:

leveraging translational research methodology to address emerging questions. *Transl. Res.*

2022;241:1–12.

6. Zuo Y et al. Prothrombotic autoantibodies in serum from patients hospitalized with COVID-19

[Internet]. *Sci. Transl. Med.* 2020;12(570). doi:10.1126/scitranslmed.abd3876

7. Chang SE et al. New-onset IgG autoantibodies in hospitalized patients with COVID-19. *Nat.

Commun.* 2021;12(1):5417.

8. Woodruff MC et al. Dysregulated naïve B cells and de novo autoreactivity in severe COVID-19

[Internet]. *Nature* [published online ahead of print: August 31, 2022]; doi:10.1038/s41586-022-

05273-0

9. Bastard P et al. Autoantibodies against type I IFNs in patients with life-threatening COVID-19

[Internet]. *Science* 2020;370(6515). doi:10.1126/science.abd4585

10. Bastard P et al. Autoantibodies neutralizing type I IFNs are present in ~4% of uninfected

individuals over 70 years old and account for ~20% of COVID-19 deaths [Internet]. *Sci Immunol*

2021;6(62). doi:10.1126/sciimmunol.abl4340

11. Vazquez SE et al. Neutralizing Autoantibodies to Type I Interferons in COVID-19

Convalescent Donor Plasma. *J. Clin. Immunol.* 2021;41(6):1169–1171.

12. van der Wijst MGP et al. Type I interferon autoantibodies are associated with systemic

immune alterations in patients with COVID-19. *Sci. Transl. Med.* 2021;13(612):eabh2624.

13. Son K et al. Circulating anti-nuclear autoantibodies in COVID-19 survivors predict long-COVID

symptoms [Internet]. *Eur. Respir. J.* [published online ahead of print: September 22, 2022];

doi:10.1183/13993003.00970-2022

14. Seeßle J et al. Persistent symptoms in adult patients one year after COVID-19: a prospective

cohort study [Internet]. *Clin. Infect. Dis.* [published online ahead of print: July 5, 2021];

doi:10.1093/cid/ciab611

15. Su Y et al. Multiple Early Factors Anticipate Post-Acute COVID-19 Sequelae [Internet]. *Cell*

2022;0(0). doi:10.1016/j.cell.2022.01.014

16. Peluso MJ, Thomas IJ, Munter SE, Deeks SG, Henrich TJ. Lack of Antinuclear Antibodies in

Convalescent Coronavirus Disease 2019 Patients With Persistent Symptoms. *Clin. Infect. Dis.*

2022;74(11):2083–2084.

17. Schultheiß C et al. The IL-1β, IL-6, and TNF cytokine triad is associated with post-acute

sequelae of COVID-19. *Cell Rep Med* 2022;3(6):100663.

18. Klein J et al. Distinguishing features of Long COVID identified through immune profiling

[Internet]. *medRxiv* [published online ahead of print: August 10, 2022];

doi:10.1101/2022.08.09.22278592

19. Bodansky A et al. NFKB2 haploinsufficiency identified via screening for IFNα2 autoantibodies

in children and adolescents hospitalized with SARS-CoV-2-related complications [Internet]. *J.*

Allergy Clin. Immunol. [published online ahead of print: December 9, 2022];

doi:10.1016/j.jaci.2022.11.020

20. Mandel-Brehm C et al. Kelch-like Protein 11 Antibodies in Seminoma-Associated

Paraneoplastic Encephalitis. *N. Engl. J. Med.* 2019;381(1):47–54.

21. O'Donovan B et al. High-resolution epitope mapping of anti-Hu and anti-Yo autoimmunity by

programmable phage display. *Brain Commun* 2020;2(2):fcaa059.

22. Vazquez SE et al. Identification of novel, clinically correlated autoantigens in the monogenic

autoimmune syndrome APS1 by proteome-wide PhIP-Seq [Internet]. *Elife* 2020;9.

doi:10.7554/eLife.55053

23. Vazquez SE et al. Autoantibody discovery across monogenic, acquired, and COVID-19-

associated autoimmunity with scalable PhIP-seq [Internet]. *Elife* 2022;11. doi:10.7554/eLife.78550

24. Larman HB et al. Autoantigen discovery with a synthetic human peptidome. *Nat. Biotechnol.*

2011;29(6):535–541.

25. Peluso MJ et al. Persistence, magnitude, and patterns of postacute symptoms and quality of

life following onset of SARS-CoV-2 infection: cohort description and approaches for measurement.

In: *Open forum infectious diseases.* Oxford University Press US; 2022:ofab640

26. Peluso MJ et al. Impact of pre-existing chronic viral infection and reactivation on the

development of long COVID [Internet]. *J. Clin. Invest.* [published online ahead of print: December

1, 2022]; doi:10.1172/JCI163669

A Anti-ARHGAP31 Antibody Signal

Fold-Change > Mock-IP

Pre-COVID   Post-COVID   Long-COVID

B ARHGAP31 Protein: 59 Phage Fragments

Reads/100k

>20

3<

C ARHGAP31 Targeted Sequence

885- PSDCDEDDT**VTDIAQHGLEMV**EPWEEPQWVTSPLHSPTLKDAHKAQVQG -933

SARS-CoV-2 Orf1a Similar Sequence

2119- **LKTLATHGLAAV**NSVPWDTIANYAKPFLNKVVSTTTNIVTRC -2161

VTDIAQHGLEMV
LKTLATHGLAAV
 :. :* *** *

9 7 7 + * 8 * * * 4 8 *
Physical-Chemical Conservation

A

B