Cost savings in male circumcision post-operative care using two-way text-based follow-up in rural

and urban South Africa

Yanfang Su[1]¶, Rachel Mukora[2]¶, Felex Ndebele[2], Jacqueline Pienaar[2,3], Calsile Khumalo[2], Xinpeng Xu[4],

Hannock Tweya[1,5], Maria Sardini[3], Sarah Day[3,6], Kenneth Sherr[1], Geoffrey Setswe[2,7], Caryl

Feldacker[1,5*]

[1] Department of Global Health, University of Washington, Seattle, WA, USA.

[2] The Aurum Institute, Johannesburg, South Africa

[3] Centre for HIV-AIDS Prevention Studies (CHAPS), Johannesburg, South Africa

[4] School of Public Health, Nanjing Medical University, Nanjing, China

[5] International Training and Education Center for Health (I-TECH), Seattle, WA, USA;

[6] School of Public Health, Faculty of Health Sciences, University of Cape Town, South Africa

[7] Department of Health Studies, University of South Africa (UNISA), Pretoria, South Africa

¶These authors contributed equally to this work.

*Corresponding author

Email: cfeld@uw.edu (CF)

Abstract

Introduction

Voluntary medical male circumcision (VMMC) clients are required to attend multiple post-operative follow-up visits in South Africa. However, with demonstrated VMMC safety, stretched clinic staff in SA may conduct more than 400,000 unnecessary reviews for males without complications, annually. Embedded into a randomized controlled trial (RCT) to test safety of two-way, text-based (2wT) follow-up as compared to routine in-person visits among adult clients, the objective of this study was to compare 2wT and routine post-VMMC care costs in rural and urban South African settings.

Methods

Activity-based costing (ABC) estimated the costs of post-VMMC care, including counselling, follow-ups, and tracing in $US dollars. Transportation for VMMC and follow-up was provided for rural clients in outreach settings but not for urban clients in static sites. Data were collected from National Department of Health VMMC forms, RCT databases, and time-and-motion surveys. Sensitivity analysis presents different follow-up scenarios. We hypothesized that 2wT would save per-client costs overall, with higher savings in rural settings.

Results

VMMC program costs were estimated from 1,084 RCT clients: 537 in routine care and 547 in 2wT. On average, 2wT saved $3.56 per client as compared to routine care. By location, 2wT saved $7.73 per rural client and increased urban costs by $0.59 per client. 2wT would save $2.16 and $7.02 in follow-up program costs if men attended one or two post-VMMC visits, respectively.

Conclusion

Quality 2wT follow-up care reduces overall post-VMMC care costs by supporting most men to heal at home while triaging clients with potential complications to timely, in-person care. 2wT saves more

in rural areas where 2wT offsets transportation costs. Minimal additional 2wT costs in urban areas

reflect high care quality and client engagement, a worthy investment for improved VMMC service

delivery.   2wT scale-up in South Africa could significantly reduce overall VMMC costs while

maintaining service quality.

Keywords: Activity-based costing; digital health innovations; mHealth; two-way texting; 2wT;

voluntary medical male circumcision; post-operative care; South Africa.

Introduction

Voluntary medical male circumcision (VMMC) is one of the most successful biomedical prevention strategies to reduce HIV transmission risk [1, 2]. VMMC is also safe with reported rates of combined moderate and severe adverse events (AEs) in large VMMC programs operating at scale in Southern Africa below 2% [3-7], meaning that 98% of VMMC clients likely heal without any incident. The Republic of South Africa SA prioritizes VMMC as part of its comprehensive HIV prevention, treatment and care plan [8], completing over 400,000 VMMCs per year for the last 10 years [9]. National South Africa guidelines require all VMMC clients to attend two post-operative follow-up visits within 14 days after the procedure to ensure timely identification and treatment for AEs. However, these visits are costly to both the clients and the providers. To comply with guideline requirements, attendance at post-operative reviews may be overreported, potentially inflating review rates while reducing the quality of care [10, 11]. A cost-effective approach is needed to reduce unnecessary visits while maintaining care quality.

Mobile health (mHealth) is used to describe the practice of healthcare and public health supported by mobile communication devices such as cellphones. mHealth interventions are recognized as more cost-effective compared to conventional interventions and may help address persistent challenges of healthcare worker (HCW) shortages and client difficulty in accessing healthcare facilities [12, 13], including in the VMMC context [14]. From 2009 to 2019, thousands of mHealth interventions were documented and launched globally, the vast majority in resource-constrained settings, suggesting robust approval of, and desire for, digital technologies in healthcare [15-17]. When implemented optimally, mHealth interventions may help reduce healthcare costs by improving the provision of health education, identifying health concerns early when illness may be less severe, reducing the duration of therapy, and minimizing transport costs [18].

In 2018, the International Training and Education Center for Health (I-TECH) in the Department of Global Health at the University of Washington (UW) and technology partner, Medic, conducted a

randomized control trial (RCT) among VMMC clients in routine VMMC clinics in Zimbabwe using a

two-way texting (2wT) telehealth approach between clients and nurses. 2wT is an mHealth platform

for conversational messaging between VMMC clinicians and clients, providing short messaging

service (SMS) for post-operative telehealth care in lieu of scheduled, in-person visits. In Zimbabwe,

and likely in other VMMC programs operating at scale, in-person, post-operative visits may be

burdensome for patients and providers, leading to poor attendance [10, 11]. 2wT improved the

quality of post-operative care while reducing HCW visit workload [19]. The 2wT approach for VMMC

follow-up in Zimbabwe reduced VMMC program costs while improving client care engagement, with

a net savings of $2.10 using 2wT over routine follow-up visits [14]. 2wT was also found to be highly

usable and acceptable among clients and providers [20], reaching >31,000 VMMC clients in

Zimbabwe by 2023 [21].

South Africa, with reported mobile-cellular subscriptions of 161.8 per 100 population in 2020 [22] is

a good candidate for exploring cost-effective mobile phone innovations, like 2wT for VMMC. To test

the 2wT approach in South Africa, I-TECH conducted a pragmatic RCT to test 2wT in urban and rural

settings in South Africa in partnership with the Aurum Institute, the Centre for HIV-AIDS Prevention

Studies (CHAPS) and Medic. Previously, we disseminated results from clinical and usability outcomes

of the 2wT RCT in South Africa where 2wT improved the quality of post-operative care in both rural

and urban settings, increasing the ascertainment of AEs, engaging males in follow-up care, and

showing high acceptability for clients and providers [23, 24]. In growing recognition of the

importance of costing research to inform HIV prevention interventions globally and in South Africa,

we embedded a costing study within the South Africa 2wT RCT.

Costing of HIV prevention interventions is gaining recognition for its importance in informing policy

and scale-up. In recent reviews of costing research on HIV prevention interventions in sub-Saharan

Africa, 38 out of 159 studies (24%) were conducted in South Africa [25, 26]. Although there is clear

evidence for VMMC as a cost-effective HIV reduction intervention [27, 28], to our knowledge, there

is no costing research on innovations to improve efficiency of VMMC care delivery in South Africa.

This costing study is nested in the clinical 2wT RCT [23]. The objective of this embedded study was to

estimate the costs associated with 2wT mHealth follow-up as compared to routine in-person VMMC

follow-up in rural and urban South Africa.   While some clients may pay for transport for VMMC

services in urban areas where services are typically more convenient, in rural areas, it is common

that VMMC programs provide transportation for both the VMMC procedure and follow-up visits to

increase VMMC uptake [29, 30]. The VMMC program in SA operates in predominantly rural areas

[23]. As with the previous Zimbabwe study [14], the payer perspective is appropriate for this analysis

as it is largely the VMMC program (the payer) that bears these follow-up costs and stands to gain the

most from the 2wT approach to reduce unnecessary visits. As VMMC costs are largely provided by

the global donor community, including the US President's Emergency Plan for AIDS Relief (PEPFAR),

and to mirror the approach taken to assess costs benefits of 2wT for VMMC in Zimbabwe, we

assessed costs from the payer perspective. This approach is also in line with recent evidence

suggesting continuing donor support for VMMC costs for at least 5 additional years given the cost-

effectiveness of the overall VMMC intervention for HIV prevention [31].

Materials and Methods

114   We applied the activity-based costing (ABC) approach to estimate the costs associated with 2wT

115   intervention. We focused on estimating the marginal cost of follow-up for an additional VMMC

116   client using the 2wT or routine follow-up approach.

Comparators

Routine care

The VMMC implementing partner, Centre For HIV-AIDS Prevention Studies (CHAPS), followed all

National Department of Health (NDoH) protocols [32] for routine VMMC services, post-operative

wound care counselling, in-person follow-up visits on post-surgery days 2 and 7, and tracing. Routine

VMMC data collection defines the post-operative day 2 visit window from day 1 to day 4 and the day

7 visit window from day 5 to day 10. Typically, in urban CHAPS VMMC services, clients pay for their

own transportation costs to attend the VMMC procedure visit and any follow-up visits, routine or

potential AE related. In rural areas, the CHAPS team typically provides transportation both for the

procedure visit and any follow-up, routine or potential AE related. Exceptions to transportation

generalities in either setting do occur. NDoH policy requires participant tracing by phone calls and/or

home visits, for those who do not attend post-operative reviews on days 2 and 7 in routine care

(Table 1).

Table 1: Activities undertaken in the 2wT study

Activities	Routine	2wT
Day of male circumcision		
Adverse event (AEs) identification counselling	Yes	Yes
Counselling on SMS interactions and photos of potential AEs named in daily message	No	Yes
Post-operative follow-ups		
Required day 2 and day 7 in-person visits	Yes	No
In-person reviews for potential AEs on any day	Yes	Yes
Day 1-13 SMS	No	Yes
Adverse event treatment	Yes	Yes
Tracing via phone calls and home visits for lost to follow up (LTFU)	Yes	Yes

2wT intervention and specific 2wT RCT procedures

The 2wT approach for VMMC follow-up was described previously [19, 21, 23]. 2wT is a hybrid

mHealth model integrating both automated SMS and individualized mHealth messaging between

clients and a routine VMMC nurse. On day 0, 2wT clients were enrolled into a custom 2wT software

application built using the open-source Community Health Toolkit (CHT). Before leaving the VMMC

clinic, 2wT clients received enhanced post-operative counselling based on global VMMC guidelines

[33] using the study-specific 2wT flipbook with additional wound care guidance, photos of AE

warning signs, and instructions on how to respond to the daily SMS (Table 1). 2wT specific

counseling helped 2wT clients understand the automated daily message that asked about the five

common complications (i.e., bleeding, swelling, pus, redness, and wound opening), preparing men to

take ownership over their wound care. 2wT participants received an automated daily text on days 1-

13 in either English, Setswana, or isiZulu, and could respond in any language. Daily automated text

messages read, "are you experiencing any bleeding, swelling, pus, pain, redness, wound opening, or

other concern? Enter 1 = Yes; 0 (Zero) = No / I'm Ok, and press send." Receiving and sending text

messages was free for the participants. The SMS prompt encouraged the clients to reply with binary

answers (0 for No, there is not a problem and 1 for Yes, there is a problem). Nurses communicated

further via SMS with clients who reported any concerns and also made phone calls if needed. 2wT

clients were not required to attend any clinic visits but could attend the clinic at any time if they

needed. For 2wT men with a potential AE or with the need for further reassurance, the nurse triaged

the participants for clinical visits the following day or earlier if an emergency was suspected. If 2wT

clients did not respond to text messages by day 8, phone tracing was activated by the clinical VMMC

team. All 2wT participants, in both arms, were asked to return for a study-specific visit on post-

operative day 14 at which point a US$7 cell phone credit was given to all participants to compensate

for time and travel. Participants were not traced via home visit for missing the study-specific day 14

visit. The day 14 visit window included visits from day 13 until day 21. Participants were declared off-

study after day 21 or after completion of the day 14 visit.

An *Enrolled* Nurse cadre level led the 2wT interaction and patient follow-up activities at the rural site

while a *Professional* Nurse cadre level led the follow-up activities at the urban site with additional

supervisory duties across study sites.

Data collection and assumptions

While exceptions do occur, in general, it is most common that VMMC clients self-paid transportation

costs in urban sites and that the program paid for client transportation costs in rural areas. These

transportation commonalities are reflected in the payer perspective, representing CHAPS costs as

part of their overall VMMC program costs funded by the major VMMC donor, PEPFAR. This is the

most prevalent model of VMMC funding and was assumed for the current approach.

For the costing study, data were collected from three main sources: 1) RCT databases of routine

VMMC data including client visits, AEs, and follow-up tracing; 2) the 2wT database; and 3) time-and-

motion surveys designed for the costing component of the study. The first source, the RCT

databases, included de-identified routine VMMC data on the number of visits attended, the

number/type/severity of AEs, and the number of traced clients, by tracing method (phone and/or

home visit). Visit costs associated with the study day 14 visit were excluded since it was specific to

the RCT study and not considered routine VMMC follow-up care. The 2wT database contained

enrolment data including transportation costs to the clinic. The 2wT database also included counts

of inbound and outbound SMS. RCT-specific activities (e.g., additional consenting, RCT monitoring

forms) were also not included in time-and-motion data.

Based on Time and Motion Form (S1 Form), data collection tools were created using both Epicollect5

and Microsoft Excel and were piloted before data collection. Time-and-motion studies consisted of

five days of direct observations at both rural and urban sites. We collected data about distance

travelled using car odometer readings as well as time spent in travelling and conducting client

reviews in the rural district of Bojanala, South Africa, between 29[th] November 2021 and 12[th] January

2022. Nurses took notes to specify how time was spent in each activity, providing additional

qualitative data inputs for costing analysis. We collected reception time and nurse review time at the

urban district of Ekurhuleni, South Africa, between 12[th] January 2022 and 24[th] January 2022.

It is assumed that (1) all fixed costs related to post-operative follow-ups were in place, and hence we

focus on analysing marginal costs in serving an additional VMMC client in the system; (2) that full-

time employees worked a 40-hour workweek or 160 hours per month; (3) that nurse counselling

time was 5 minutes in routine counselling and 10 minutes for 2wT counselling; (4) that the nurse

spent one minute per SMS response on average, and (5) that the average phone call time with a

VMMC client was 5 minutes.

Hypotheses
191
192

193   We tested the following two hypotheses in our costing study:

194   Hypothesis 1: 2wT reduces post-VMMC care costs by triaging only those in need of in-person review

195   to care, allowing most men to heal safely at home without in-person follow-up visits.

196   Hypothesis 2: 2wT saves more costs in rural VMMC program settings than in urban program settings.

Data analysis: Activity-based costing
197

Using an ABC approach, we estimated the costs in the post-VMMC care continuum from the

perspective of the payer – the VMMC program with donor support. There were four activity

categories in costing: counselling on the day of male circumcision, SMS follow-ups, physical follow-

up visits, and tracing. All activity-based costs were estimated for 2wT care and routine care to test

the hypotheses of cost savings. We applied ranges of ±50% uncertainty interval for parameters

presented in Table 2 [14]. Input data were analysed using STATA version 14.0 and costing was

conducted in Microsoft Excel. Editable costing tool (S2 Table) presented our input data and costing

algorithms. All parameters can be adjusted to reflect new inputs and scenarios, providing a tool for

future costing studies of 2wT impact in other settings, contexts, and salary structures. All costs were

collected in the South African rand (ZAR) and converted into the United States dollar (USD) using the

exchange rate ZAR 1 = $0.066.

Counselling
209

210   Each VMMC client had a counselling session directly following circumcision. The post-operative

211   counselling cost was estimated by the average time in minutes for a counselling session and wage

212   per minute for the nurse performing the counselling.

213   Counselling cost = time * wage

SMS follow-ups

215   There were three components in SMS follow-ups per client. The first component was the 'Africa is

216   Talking' SMS aggregator monthly set-up costs. The second component was the SMS service cost, i.e.,

217   the product of SMS unit cost and average number of SMS per client. The third component was the

218   associated personnel cost of sending a manual SMS.

219   SMS cost = set-up cost + SMS service cost + manual SMS time cost

Follow-up visits and AE management

The rural costing model included the estimation of transportation fuel cost, transportation time cost,

and nurse review time cost per visit. In urban settings, we estimated reception time cost and nurse

review time cost per visit. We used the average number of follow-up visits per client by location.

Transportation cost was paid by the program in rural settings whereas patients paid for

transportation in urban settings. In the rural setting, an enrolled nurse interacted with clients via

2wT and usually drove out to meet clients at their homes or workplaces to conduct any requested

post-VMMC reviews. In urban settings, clients interacted with clients via 2wT and typically returned

to the clinic on their own for visits when needed or desired. Reception time cost only applied to

urban settings as the rural settings did not have reception.

Cumulative moderate and severe AEs as defined by global VMMC standards [33] were reported from

day 1 through day 21 by randomization group and by location. The time costs in treating AEs were

included in overall follow-up visits. We estimated the material costs for AE management separately.

Rural visit cost = # of visits * (transportation fuel cost + transportation time cost + review time cost)

Urban visit cost = # of visits * (reception time cost + review time cost)

In rural settings, transportation fuel cost = Round trip distance to client home (km) * Litres of petrol

(per km) * Price of petrol (per litre)

AE management = probability of AE * AE material cost

Tracing

In routine care, if the client missed both day 2 and day 7 visits, three phone calls were attempted to

confirm healing. In the 2wT arm, if the client had no SMS contact by day 8, the client was actively

traced via phone up to three attempts. Successful calls to clients in both arms took about 5 minutes

If the client was not reachable by phone, up to three home visits were attempted for both study

arms. The number of clients eligible for phone tracing and home tracing was reported in the RCT

[23].

Phone tracing cost = probability of phone tracing * (phone call time cost in up to 3attempts + phone

call service cost of a successful completed call)

For home tracing, transportation cost was estimated in the care model that the nurse drove to the

client's home to conduct the review. Transportation fuel cost, transportation time cost, and nurse

review time cost were also included. We considered the least expensive model, in which the nurse

also served as the driver. The urban tracing distance is 72% of the distance in rural settings

according to the survey results in our RCT [23].

Home tracing cost = probability of home tracing * (transportation time cost in all attempts +

transportation fuel cost in all attempts + review time cost in successful attempt)

For all tracing attempts, phone service costs and client review costs were only recorded for the

successful attempt. For instance, if the nurse made three home visits and the client was reached in

the last visit, the client review cost for only the one successful visit was included.

Ethics

This Multiple Principal Investigators (MPI) study was approved by the Internal Review Boards of the University of Washington (Study 00009703, PI: Feldacker) and the University of the Witwatersrand, Human Research Ethics Committee (Ethics Reference No: 200204, PI: Setswe). All RCT participants provided written informed consent for use of individual-level data collected from both 2wT study specific sources and routine VMMC program data collection. All data obtained and utilized for the costing study was de-identified data from study databases and contained no identifying information. As part of the overall RCT IRB approvals, a waiver of consent was granted for healthcare worker observations from the time-motion study.

Results

VMMC clients

A total of 1,460 VMMC clients were recruited for the RCT, and 141 (9.7%) clients were found ineligible for 2wT: 103 (45.2%) had no cell phones, 21 (5.7%) had a language barrier, 13 (3.5%) could not read or write, 2 (0.5%) were blind, and 2 (0.5%) were unfit to consent) [23]. The costing analysis included all 1,084 enrolled clients from the RCT: 537 clients in routine care and 547 clients in 2wT intervention. Table 2 indicates key costing parameters stratified by urbanicity. Average costs per client by randomization group, cost savings, and savings by location are presented in Table 3, with illustrations in Figures 1 and 2. Costing scenarios are presented in Table 4.

Table 2: Summary statistics by activity category in rural and urban South Africa

Parameters	Sum	Average	Rural	Urban	Source
	N (Uncertainty Interval)	N (UI)	N (UI)	N (UI)	
a) Arm of randomization					
Routine	537	-	268	269	RCT study logs
2wT	547	-	273	274	RCT study

Parameters	Sum	Average	Rural	Urban	Source
					logs
b) Post-operative counselling - Day 0					
Enrolled Nurse wage (per month)	-	$1,205 (602.50 to 1,807.50)	$1,205 (602.50 to 1,807.50)	$1,205 (602.50 to 1,807.50)	NDoH pay structure
Nurse counselling time in minutes (per patient) - Routine	-	5 (2.5 to 7.5)	5 (2.5 to 7.5)	5 (2.5 to 7.5)	Assumptio n
Nurse counselling time in minutes (per patient) - 2wT	-	10 (5 to 15)	10 (5 to 15)	10 (5 to 15)	Assumptio n
c) Text service for 2wT patients					
SMS aggregated set-up cost (per month)	-	$74.26 (37.13 to 111.39)	$74.26 (37.13 to 111.39)	$74.26 (37.13 to 111.39)	Africa is Talking
Cost per SMS	-	$0.02 (0.01 to 0.03)	$0.02 (0.01 to 0.03)	$0.02 (0.01 to 0.03)	Africa is Talking
Mean number of automated texts (per patient) - outbox	-	20.1 (10.05 to 30.15)	17.9 (8.95 to 26.85)	22.3 (11.15 to 33.45)	Africa is Talking
Mean number of manual texts (per patient) - outbox	-	5.65 (2.83 to 8.48)	4 (2 to 6)	7.3 (3.65 to 10.95)	Africa is Talking
Mean number of texts (per patient) - inbox	-	14.5 (7.25 to 21.75)	13 (6.5 to 19.5)	16 (8 to 24)	Africa is Talking
Duration of 2wT program (days)	-	13 (6.5 to 19.5)	13 (6.5 to 19.5)	13 (6.5 to 19.5)	RCT study logs
d) Follow-up visits					
Average number of visits attended - Routine	-	1.34 (0.67 to 2.01)	1.26 (0.63 to 1.89)	1.42 (0.71 to 2.13)	RCT study logs
Average number of visits attended - 2wT	-	0.22 (0.11 to 0.33)	0.14 (0.07 to 0.21)	0.3 (0.15 to 0.45)	RCT study logs
Unnecessary visits (Routine vs. 2wT)		1.12 (0.56 to 1.68)	1.12 (0.56 to 1.68)	1.12 (0.56 to 1.68)	RCT study logs
Administrator reception time in minutes (per patient)	-	1.34 (0.67 to 2.00)	0	2.67 (1.34 to 4.01)	Time-motion study
Nurse review time in minutes (per patient)	-	5.43 (2.71 to 8.14)	3.79 (1.90 to 5.69)	7.06 (3.53 to 10.59)	Time-motion study
Administrator wage (per month)	-	$319 (159.5 to 478.5)	0	$638 (319 to 957)	NGO pay structure
Nurse wage (per month)	-	$1676.5 (838.25 to 2514.75)	$1,205 (602.50 to 1,807.50)	$2,148 (1,074 to 3,222)	NGO pay structure
Round trip distance to patient home (km)	-	-	39.64 (19.82 to 59.46)	-	Time-motion study
Litres of petrol (per km)	-	0.06 (0.03 to 0.09)	0.06 (0.03 to 0.09)	-	Vehicle manual
Price of petrol (per litre)	-	$1.33 (0.67 to 2)	$1.33 (0.67 to 2)	-	Automobil e Associatio

Parameters	Sum	Average	Rural	Urban	Source
					n
Transportation time cost in minutes (per visit)	-	25.5 (12.75 to 38.25)	34 (17 to 51)	-	Time-motion study
Transportation cost - nurse home visits (per visit)	-	$5.61 (2.81 to 8.42)	$7.48 (3.74 to 11.22)	$3.74 (1.87 to 5.61)	RCT study logs
Patients with AEs – Routine (%)	0.01(0.005 to 0.014)	-	0	0.02 (0.01 to 0.03)	RCT study logs
Patients with AEs - 2wT (%)	0.02 (0.01 to 0.03)	-	0.01(0.005 to 0.014)	0.03 (0.02 to 0.05)	RCT study logs
Material cost per AE (e.g., antiseptic ointment)	-	$5.28 (2.64 to 7.92)	$5.28 (2.64 to 7.92)	$5.28 (2.64 to 7.92)	RCT study logs
e) Tracing - Phone calls					
Phone call service (per minute)	-	$0.08 (0.04 to 0.12)	$0.08 (0.04 to 0.12)	$0.08 (0.04 to 0.12)	Public records
Time cost per call (minutes)	-	5 (2.5 to 7.5)	5 (2.5 to 7.5)	5 (2.5 to 7.5)	Assumption
Number of phone call attempts in tracing - Routine and 2wT	-	3(1.5 to 4.5)	3(1.5 to 4.5)	3(1.5 to 4.5)	RCT study logs
Patients missed day 2 & day 7 visits - Routine (%)	0.05 (0.03 to 0.08)	-	0.02 (0.01 to 0.03)	0.09 (0.05 to 0.14)	RCT study logs
Patients without post MC contact - 2wT (%)	0.09 (0.05 to 0.14)	-	0.06 (0.03 to 0.09)	0.11 (0.06 to 0.17)	RCT study logs
f) Tracing - Home visits					
Mean distance to patient home (km)	-	29.73 (14.87 to 44.60)	39.64(19.82 to 59.46)	28.54 (14.27 to 42.81)	Time-motion study
Litres of petrol (per km)	-	0.06(0.03 to 0.09)	0.06(0.03 to 0.09)	0.06 (0.03 to 0.09)	Vehicle manual
Price of petrol (per litre)	-	$1.33(0.67 to 2)	$1.33(0.67 to 2)	$1.33(0.67 to 2)	Automobile Association
Traced patients with no post MC contact - Routine (%)	0.03 (0.02 to 0.05)	-	0.03 (0.02 to 0.05)	0.03 (0.02 to 0.05)	RCT study logs
Traced patients with no post MC contact - 2wT (%)	0.002 (0.001 to 0.003)	-	0.002 (0.001 to 0.003)	0.002 (0.001 to 0.003)	RCT study logs

Counselling

282 In both settings, an enrolled nurse with a monthly salary of $1,205 was assumed to spend 5 minutes

283 in routine post-operative counselling and 10 minutes in 2wT counselling (Table 2). There was no

284 difference in counselling time by location.

SMS follow-ups

The SMS aggregator set-up cost charged by 'Africa is Talking' was $74.25 per month for the 2wT system as detailed by Table 2. Each client was sent automated text messages for 13 days in a month of 30 days and the average cost per client was proportioned to 43% of the month. With 547 clients in 2wT, the set-up cost per client was $0.14. When 'Africa is Talking' serves more clients in future, the set-up cost is expected to reduce.

Table 2 also indicates that a standard SMS of 160 characters cost $0.02. Each 2wT client received 20.1 (10.05 to 30.15) messages on average, including 5.65 (2.83 to 8.48) manual messages from nurses. Clients sent an average of 14.5 (7.25 to 21.75) messages, total, between day 1 and Day 13 to report daily status and any AE concerns. Urban clients were more active in 2wT, sending an average of 3 more messages than rural clients. Similarly, there were 3 more messages from the nurse on average, per client, in urban settings than in rural settings.

Follow-up visits and adverse event management

Clinic visits

As previously reported in the RCT, and as expected per RCT intervention assignment, routine clients had more clinic follow-up visits than 2wT clients. Routine clients had 1.34 (0.67 to 2.01) visits on average between day 2 and day 13: 1.26 (0.63 to 1.89) in rural areas and 1.42 (0.71 to 2.13) in urban locations. 2wT clients had 0.22 (0.11 to 0.33) visits on average, with 0.14 (0.07 to 0.21) in rural areas and 0.3 (0.15 to 0.45) in urban clinics. Urban clients, in both routine and 2wT groups, had more visits than rural clients. Post-operative engagement in care was higher among 2wT arm than control: 94% of 2wT males responded to at least one 2wT message and 80.4% of control arm males attended at least 1 in-person, post-operative visit [23].

AE material cost

The study identified a total of 16 AEs across both arms. Among 719 visits in the routine arm, there were 5 AEs identified whereas 11 AEs were identified among 118 visits in 2wT arm. Material cost per AE, including bandage, paraffin gauze, and antiseptic ointment, was estimated at $5.28.

Salary and time cost

In the rural setting, Enrolled Nurses with a monthly salary of $1,205 conducted the post-VMMC reviews with 3.79 (1.90 to 5.69) minutes per client. In urban settings, administrators with a monthly salary of $638 facilitated reception with 2.67 (1.34 to 4.01) minutes per client and a *Professional Nurse* with a monthly salary of $2,148 conducted the reviews with 7.06 (3.53 to 10.59) minutes per client. The routine urban *Professional Nurse* salary was 1.78 higher than the rural Enrolled Nurse salary.

Transportation cost

In rural settings, routine VMMC teams drove to review clients or provided transport while urban clients were largely responsible for their own transportation. It was estimated that the round–trip distance to a client's home on average was 39.64 (19.82 – 9.91) kilometers in rural areas. As indicated in the vehicle manual, the average fuel consumption of the vehicle that was used for tracing was 0.06L/km (*6L/100km*). Information from the Automobile Association reported the fuel price of $1.33/L [34]. The transportation fuel cost was $3.21 (1.61 to 4.82) in rural areas.

From time-motion data, it was estimated that the program's transportation time per client visit was 34 minutes in rural settings. No client transportation was provided by the program in urban settings over the observation period.

Tracing

Among routine men, 29 clients did not attend any visit and were potentially lost-to-follow-up (LTFU). Potential LTFU were traced first by phone; 17/29 were not reached by phone and were traced by

home visit. For the participants in the 2wT arm, 47 had no contact by day 8 post-VMMC; 46 were

reached by phone and 1 was traced by home visit.

The cost of phone call was $0.08 per minute. It was assumed that an Enrolled Nurse conducted the

phone calls, and an average phone call time was 5 minutes. To trace potential LTFU clients, it was

assumed that on average 3 phone calls were attempted in both routine group and 2wT group. In

home tracing, the transportation fuel cost was the same as the clinic visit. The estimated tracing

transportation costs on average, including fuel and personnel costs, were $0.64 and $0.04 in rural

and urban areas, respectively. According to our time-motion study, it is important to note that home

visits were costly both in terms of money and time, especially in rural areas. The average travel time

to reach one client was 34 minutes (range: 17 to 51 minutes) and the average review time was 3.8

minutes (range: 1.9 to 5.7 minutes). On a typical day of rural tracing, the nurse spent 90% of the

time driving. On average, 5 clients (range: 2.5 to 7.5 clients) would be traced in one day in rural

settings.

Costs and cost savings

Table 2 shows that there were, on average, 1.12 unnecessary post-operative visits per person,

reflecting the difference between average number of in-person visits in the control arm (1.34) and

average number of elected visits in the 2wT arm (0.22). In the 2wT arm, visit attendance reflects

demand for in-person, post-VMMC care, 0.14 and 0.3 visits in rural and urban areas, respectively.

Table 3 shows that the average cost per post-operative visit was $7.75 in routine care and $4.18 in

2wT. The cost saving of $3.56 was primarily seen in rural areas ($7.73), with no savings in urban

areas (with $0.59 increased cost). Table 3 also shows the disaggregation of costs by location. The

average post-operative counselling cost was $0.63 and $1.26 in routine and 2wT groups,

respectively. SMS costs were only incurred among the 2wT group, with $1.67 average costs, of

which $2.30 was in urban areas and $1.05 in rural areas.

Table 3: Unit cost for routine care (standard of care (SoC) and 2wT ($USD)

Costs per patient	Rural			Urban			Average Overall		
	SoC	2wT	2wT vs. SoC	Routine	2wT	2wT vs. SoC	SoC	2wT	SoC vs. 2wT
Post-op counselling - day 0	$0.63	$1.26	$0.63	$0.63	$1.26	$0.63	$0.63	$1.26	$0.63
Two-way texting	$0.00	$1.05	$1.05	$0.00	$2.30	$2.30	$0.00	$1.67	$1.67
Set-up cost	$0.00	$0.06	$0.06	$0.00	$0.06	$0.06	$0.00	$0.06	$0.06
SMS service cost	$0.00	$0.49	$0.49	$0.00	$0.61	$0.61	$0.00	$0.55	$0.55
Manual SMS time cost	$0.00	$0.50	$0.50	$0.00	$1.63	$1.63	$0.00	$1.07	$1.07
Follow-up visits	$10.04	$1.12	-$8.92	$2.50	$0.53	-$1.97	$6.26	$0.82	-$5.44
Provider transportation fuel cost	$4.05	$0.45	-$3.60	$0.00	$0.00	$0.00	$2.02	$0.23	-$1.80
Provider transportation time cost	$5.39	$0.60	-$4.79	$0.00	$0.00	$0.00	$2.69	$0.30	-$2.39
Provider review time cost	$0.60	$0.07	-$0.53	$2.50	$0.53	-$1.97	$1.55	$0.30	-$1.25
AE treatment (materials)	$0.00	$0.04	$0.04	$0.10	$0.17	$0.08	$0.05	$0.11	$0.06
Tracing - Phone calls	$0.03	$0.19	$0.16	$0.30	$0.39	$0.09	$0.16	$0.29	$0.12
Tracing - Home visits	$0.73	$0.04	-$0.69	$0.56	$0.03	-$0.53	$0.64	$0.04	-$0.61
Total costs	$11.42	$3.69	-$7.73	$4.08	$4.67	$0.59	$7.75	$4.18	-$3.56

Figure 1 illustrates the cost distribution across the post-VMMC care continuum. In the 2wT group,

the costs were concentrated in early actions within the care continuum, including counselling ($1.26)

and SMS ($1.67), which accounted for 70% of the overall costs. By contrast, in the routine group,

most of the financial investment was in clinical visits, $6.26 out of $7.75 (81%), followed by

counselling ($0.63). In the post-VMMC care continuum, 2wT intervention shifts the healthcare effort

to preparing the client to actively engage in their healing process and to encourage them to

promptly seek care via consistent daily communication with the client from day 1 to day 13.

Fig 1: Costs in USD for routine care and 2wT per client (USD)

Figure 2 demonstrates costs and savings by urbanicity. Overall, there were $0.59 cost increases in

urban areas and $7.73 savings in rural area by implementing 2wT instead of utilizing routine care.

Cost savings were from reduced follow-up visits ($1.97 in urban areas and $8.92 in rural areas).

Savings from home visits in tracing were $0.53 in urban areas and $0.69 in rural areas. Increased

costs in 2wT arm by location included counselling ($0.63, the same in each location), SMS ($2.30 in

urban areas and $1.05 in rural areas), AE materials ($0.08 in urban areas and $0.04 in rural areas),

and calls ($0.09 in urban areas and $0.16 in rural areas).

Fig 2: Cost changes by implementing 2WT compared to the routine, rural vs. urban (USD)

Other scenarios

Table 4 depicts three scenarios, by location, in comparing cost savings of using 2wT compared to

routine care visits namely one visit, 1.34 visits on average, and two visits within routine care. If there

had been one required visit per client in routine care, the average cost saving was calculated at

$2.16 per client. In the scenario drawn from the pragmatic RCT of 1.34 visits within routine care, the

average cost saving was $3.56. If two follow-up visits were required per client in routine, the average

cost saving was $7.02 per client. An addition scenario with Enrolled Nurse salary applied to both

rural and urban settings had little impact, with average overall savings of $3.14 using 2wT over

routine with the observed visit patterns, showing the robustness of these savings over modest salary

changes.

Table 4: Potential cost savings from 2wT (USD)

Cost saving (2wT vs. Routine)	Rural	Urban	Average
Routine - One visit	-$5.66	$1.33	-$2.16
Routine - 1.34 visits (observed)	-$7.73	$1.33	-$3.56
Routine - Two visits	-$13.63	-$0.43	-$7.02

387   Discussion

In this costing analysis, we determined that 2wT-based VMMC follow-up saved an average of $3.56

per client across settings, with savings ranging from $2.16 to $7.02. 2wT improves the quality of

post-VMMC follow-up care at lower overall cost by providing an SMS-based mHealth option for

clients with access to cell phones, encouraging visits when needed instead of compulsory visits on

day 2 and/or day 7. Savings using the 2wT approach were higher in rural as compared to urban areas

– an important finding as the majority of VMMC program implementation occurs in rural areas [14].

Even though using 2wT costs slightly more in urban areas, the better quality of post-VMMC follow-

up care justifies these additional costs. Among 2wT costs, the largest costs were for the SMS,

themselves, an investment to actively engage men into quality follow-up care with a 2wT nurse for

reassurance or triaging to in-person review when needed. The 2wT clients on average received 20.1

text messages and sent 14.45 text messages in the post-operative period for an SMS cost of $1.67

per client. In urban areas, males interacted more via 2wT, raising costs but also increasing direct

client-to-clinician communication. Cost savings using 2wT as compared to routine post-operative

care were similar in the 2wT RCT conducted in Zimbabwe that found a net savings of $2.10 from the

payer (Ministry of Health or donor) perspective [14]. As attendance at routine post-operative visits

may be overreported in VMMC programs operating at scale , cost savings and benefits for quality

follow-up care using the 2wT approach may be underestimated.

Cost savings from 2wT vary by adherence to required attendance at post-VMMC visits and by

location. Given that moderate and severe AE rates in male circumcisions are low (1%-2%)[4], it is

costly to require all VMMC clients to attend one or two post-operative in-person clinic visits. 2wT

can save an average of $7.02 ($13.63 savings in rural and $0.43 savings in urban) compared to the

routine care with two required visits. Our study shows that routine follow-up in rural areas was

expensive as the district geography is expansive. For scheduled day 2 and day 7 post-operative visits,

the CHAPS rural VMMC team of a nurse and a dedicated driver, arranged to meet the clients at a

convenient community location for physical client reviews or a client contacted the CHAPS VMMC

team for pickup or an in-person home visit, with occasional clients returning independently to the

VMMC site. In contrast, the urban model had three teams based at three static clinics. Urban clients

were expected to commute for review visits, independently, at the dedicated static sites.

2wT may increase some costs that are likely offset by additional quality care benefits. Firstly,

enhanced counselling incurred a minimal cost of $0.63, but enabled the clients to effectively identify

and communicate AE concerns in daily SMS – a worthwhile cost. Second, daily SMS communication

improved early detection of AEs and subsequent swift referral of those in need for in-person clinical

visits. This triaging process led to identification of AEs earlier with less severity, likely averting costs

of more severe AEs [23]. Third, only the clients with AE concerns or those desiring reassurance

attended clinic visits, resulting in only 0.22 elected visits per 2wT client (Table 3), a reduction in

review workload. Lastly, reduced follow-up visits could allow providers to concentrate on cases in

need of medical attention and free up time to commence surgery early and on-time, potentially

increasing client satisfaction.

Quality assurance likely also benefits from 2wT improvements in verification of, and supervision for,

quality post-operative care. For NDoH, 2wT follow-up adhered to government regulations regarding

client privacy and complied with required NDoH documentation for VMMC client follow-up. The 2wT

system provides verification of timely, nurse-led, post-operative support in line with PEPFAR

guidelines, facilitating quality follow-up regardless of client location. 2wT documentation also

provides another layer of quality assurance to confirm program performance. Unique client

verification via the 2wT system increases confidence in program productivity, providing another data

source to prevent duplication in reporting. Although not formally considered in the costing analysis,

supervision costs of 2wT may be lower than those for routine care. Routine monitoring of care

quality via review of paper forms is time-consuming, requiring intensive efforts. However, 2wT

dashboards and system-embedded hierarchies allow for remote client oversight and access to data

for timely program monitoring. Supervisors can access 2wT messages, reports, tasks, and client

records from multiple sites or remotely from a central location. Similarly, quality assurance activities

can also take place virtually, allowing managers and program administrators to review data and

provide clinical oversight, informing improvements. In future, calculation of these costs may

demonstrate further benefit of the 2wT approach for routine VMMC follow-up.

Limitations

There are several assumptions and limitations in this study. First, this study focuses on the cost of

the activities to serve an average client presuming that 2wT is implemented within an existing

routine VMMC service. Therefore, start-up costs (e.g., developing text message library, translation

to local languages, adaptation to South Africa context) and fixed costs (e.g., full-time employment

dedicated to 2wT, cell phones, computers, vehicle purchase, insurance, maintenance cost) were

excluded. Second, we conducted costing from the perspective of the VMMC program and current

guidelines to inform feasibility of adoption by the NDoH, assuming the existing VMMC donors would

continue support for at least the 5-year time frame. However, with any payer (Ministry of Health,

donor, or program), and evolving guidelines, 2wT costs and comparators could change over time.

Third, we used the standard workweek in South Africa, i.e., 5 days per week, from Monday to Friday.

In calculating hourly wage, we used the salary of the Enrolled Nurse cadre and did not take into

account that some nurses work on weekends and public holidays or may have additional days off,

such as annual leave or sick leave – considerations that could affect costs. For AE management, only

costs related to reportable moderate and severe cases are documented and noted in this study. We

did not measure the costs to manage mild AEs such as pain. Lastly, we explored will see savings

(monetary cost) in reduced transportation fuel costs that the reduced staff time required for follow-

up (an opportunity cost not a monetary cost savings) means that overall monetary costs to the

program are likely to increase

Conclusion

Evidence from two RCTs in Zimbabwe and South Africa demonstrate that this 2wT approach

provides high-quality VMMC follow-up as compared to required in-person reviews and lowers

overall VMMC program costs. Rural savings using 2wT offset nominal increased costs in urban areas.

Additional 2wT associated improvements in care quality, supervision, and verification also likely

leading to longer-term savings. The health sector should invest in 2wT. In the context of National

VMMC targets of 350,000 to 500,000 VMMCs per annum, employing 2wT could dramatically reduce

the number of in-person post-operative reviews, resulting in concrete efficiency gains and significant

annual program savings. Investing these resources back into the VMMC program could further

expand VMMC access, improve care quality, and advance VMMC program goals of safe, efficient,

and effective VMMC scale-up.

Acknowledgements

The authors would like to thank the following: the Departments of Health of the Gauteng and

Northwest Provinces, Bojanala district and the Ekurhuleni Health District Research Committee

(EHDRC) for allowing us to conduct the study in their districts; implementing partner, Right to Care

and the CHAPS study implementation team. The authors would like to thank the Medic team and all

study participants for their involvement in the study. The authors would like to thank Lingchao Mao

and Ziwei He for independent replications of the results as well as thank Simon Ding and Emily Chu

for their comments.

Authors contributions

Conceptualization: Yanfang Su, Jacqueline Pienaar, Geoffrey Setswe, Caryl Feldacker

Formal analysis: Yanfang Su, Rachel Mukora, Hannock Tweya, Xinpeng Xu

Funding acquisition: Caryl Feldacker, Geoffrey Setswe

Investigation: Caryl Feldacker, Geoffrey Setswe, Jacqueline Pienaar

Methodology: Yanfang Su, Rachel Mukora, Felex Ndebele, Jacqueline Pienaar, Calsile Khumalo,
Hannock Tweya, Maria Sardini, Geoffrey Setswe, Caryl Feldacker

Resources: Caryl Feldacker, Geoffrey Setswe

Supervision: Yanfang Su, Geoffrey Setswe, Caryl Feldacker

Writing – original draft: Yanfang Su, Rachel Mukora, Felex Ndebele, Calsile Khumalo, Geoffrey
Setswe, Caryl Feldacker

Writing – review & editing: Yanfang Su, Rachel Mukora, Felex Ndebele, Jacqueline Pienaar, Calsile
Khumalo, Sarah Day, Hannock Tweya, Maria Sardini, Sarah Day, Kenneth Sherr, Geoffrey Setswe,
Caryl Feldacker

Funding

Research reported in this publication was supported by the National Institute of Nursing Research (NINR) of the National Institutes of Health under award number 5R01NR019229, "Expanding and Scaling Two-way Texting to Reduce Unnecessary Follow-Up and Improve Adverse Event Identification Among Voluntary Medical Male Circumcision Clients in the Republic of South Africa." There was no additional external funding received for this study. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Conflict of interest

None declared.

Abbreviations

2wT: Two-way texting

AE(s): adverse event(s)

CHAPS: Centre For HIV-AIDS Prevention Studies

VMMC: male circumcision

NDoH: National Department of Health

PEPFAR: President's Emergency Plan for AIDS Relief

SA: South Africa

WHO: World Health Organization

References

1.	Titus MJ, Moodley J. Male circumcision as a public health measure for the prevention of HIV transmission. Southern African Journal of Epidemiology and Infection. 2011;26(4):262-5.

2.	Bershteyn A, Mudimu E, Platais I, Mwalili S, Zulu JE, Mwanza WN, et al. Understanding the Evolving Role of Voluntary Medical Male Circumcision as a Public Health Strategy in Eastern and Southern Africa: Opportunities and Challenges. Curr HIV/AIDS Rep. 2022;19(6):526-36. Epub 2022/12/03. doi: 10.1007/s11904-022-00639-5. PubMed PMID: 36459306; PubMed Central PMCID: PMCPMC9759505.

3.	Hellar A, Plotkin M, Lija G, Mwanamsangu A, Mkungume S, Christensen A, et al. Adverse events in a large-scale VMMC programme in Tanzania: findings from a case series analysis. J Int AIDS Soc. 2019;22(7):e25369. Epub 2019/08/02. doi: 10.1002/jia2.25369. PubMed PMID: 31368235; PubMed Central PMCID: PMCPMC6669321.

4.	Muchiri E, Charalambous S, Ginindza S, Maraisane M, Maringa T, Vranken P, et al. Description of adverse events among adult men following voluntary medical male circumcision: Findings from a circumcision programme in two provinces of South Africa. PloS one. 2021;16(8):e0253960. Epub 2021/08/18. doi: 10.1371/journal.pone.0253960. PubMed PMID: 34403409; PubMed Central PMCID: PMCPMC8370616.

5.	Omollo V, Marongwe P, Vernon M, Thoko M, Paidamoyo G, Farai G, et al. Adverse event trends within a large-scale, routine, voluntary medical male circumcision program in Zimbabwe, 2014–2019. JAIDS Journal of Acquired Immune Deficiency Syndromes. 2021;88(2):173-80.

6.	O'Bryan G, Feldacker C, Ensminger A, Nghatanga M, Brandt L, Shepard M, et al. Adverse event profile and associated factors following surgical voluntary medical male circumcision in two regions of Namibia, 2015-2018. PloS one. 2021;16(10):e0258611. Epub 2021/10/21. doi: 10.1371/journal.pone.0258611. PubMed PMID: 34669709; PubMed Central PMCID: PMCPMC8528325.

7.	Lucas T, Cooney C, Prainito A, Godfrey C, Kiggundu V, Thomas AG, et al. Consolidated Overview of Notifiable Adverse Events in the US President's Emergency Plan for AIDS Relief's Voluntary Medical Male Circumcision Program Through 2020. Current HIV/AIDS Reports. 2022:1-8.

8.	South Africa National AIDS Council. South Africa's national strategic plan for HIV, TB and STIs 2017-2022. Pretoria: South African National AIDS Council, 2017.

9.	UNAIDS. Voluntary medical male circumcision. Steady progress in the scaleup of VMMC as an HIV prevention intervention in 15 eastern and southern African countries before the SARS-CoV2 pandemic. Geneva2021.

10.	Feldacker C, Murenje V, Makunike-Chikwinya B, Hove J, Munyaradzi T, Marongwe P, et al. Balancing competing priorities: Quantity versus quality within a routine, voluntary medical male circumcision program operating at scale in Zimbabwe. PloS one. 2020;15(10):e0240425.

11.	Marongwe P, Gonouya P, Madoda T, Murenje V, Tshimanga M, Balachandra S, et al. Trust but verify: Is there a role for active surveillance in monitoring adverse events in Zimbabwe's large-scale male circumcision program? PloS one. 2019;14(6):e0218137.

12.	Gentili A, Failla G, Melnyk A, Puleo V, Tanna GLD, Ricciardi W, et al. The cost-effectiveness of digital health interventions: A systematic review of the literature. Front Public Health. 2022;10:787135. Epub 2022/08/30. doi: 10.3389/fpubh.2022.787135. PubMed PMID: 36033812; PubMed Central PMCID: PMCPMC9403754.

13.	Iribarren SJ, Cato K, Falzon L, Stone PW. What is the economic evidence for mHealth? A systematic review of economic evaluations of mHealth solutions. PloS one. 2017;12(2):e0170581.

14.	Babigumira JB, Barnhart S, Mendelsohn JM, Murenje V, Tshimanga M, Mauhy C, et al. Cost-effectiveness analysis of two-way texting for post-operative follow-up in Zimbabwe's voluntary medical male circumcision program. PloS one. 2020;15(9):e0239915. Epub 2020/10/01. doi:

10.1371/journal.pone.0239915. PubMed PMID: 32997710; PubMed Central PMCID:
PMCPMC7526887.
15.   Griffee K, Martin R, Chory A, Vreeman R. A Systematic Review of Digital Interventions to
Improve ART Adherence among Youth Living with HIV in sub-Saharan Africa. AIDS Research and
Treatment. 2022;2022:9886306. doi: 10.1155/2022/9886306.
16.   Grundy Q. A Review of the Quality and Impact of Mobile Health Apps. Annu Rev Public
Health. 2022;43:117-34. Epub 2021/12/16. doi: 10.1146/annurev-publhealth-052020-103738.
PubMed PMID: 34910582.
17.   McCool J, Dobson R, Whittaker R, Paton C. Mobile Health (mHealth) in Low- and Middle-
Income Countries. Annu Rev Public Health. 2022;43:525-39. Epub 2021/10/15. doi:
10.1146/annurev-publhealth-052620-093850. PubMed PMID: 34648368.
18.   Iribarren SJ, Brown W, 3rd, Giguere R, Stone P, Schnall R, Staggers N, et al. Scoping review
and evaluation of SMS/text messaging platforms for mHealth projects or clinical interventions. Int J
Med Inform. 2017;101:28-40. Epub 2017/03/30. doi: 10.1016/j.ijmedinf.2017.01.017. PubMed
PMID: 28347445; PubMed Central PMCID: PMCPMC5398898.
19.   Feldacker C, Murenje V, Holeman I, Xaba S, Makunike-Chikwinya B, Korir M, et al. Reducing
Provider Workload While Preserving Patient Safety: A Randomized Control Trial Using 2-Way Texting
for Postoperative Follow-up in Zimbabwe's Voluntary Medical Male Circumcision Program. JAIDS
Journal of Acquired Immune Deficiency Syndromes. 2020;83(1):16-23.
20.   Feldacker C, Holeman I, Murenje V, Xaba S, Korir M, Wambua B, et al. Usability and
acceptability of a two-way texting intervention for post-operative follow-up for voluntary medical
male circumcision in Zimbabwe. PloS one. 2020;15(6):e0233234. Epub 2020/06/17. doi:
10.1371/journal.pone.0233234. PubMed PMID: 32544161; PubMed Central PMCID:
PMCPMC7297350.
21.   Marongwe P, Wasunna B, Gavera J, Murenje V, Gwenzi F, Hove J, et al. Transitioning a digital
health innovation from research to routine practice: Two-way texting for male circumcision follow-
up in Zimbabwe. PLOS Digit Health. 2022;1(6):e0000066. Epub 2023/02/23. doi:
10.1371/journal.pdig.0000066. PubMed PMID: 36812548; PubMed Central PMCID:
PMCPMC9931231.
22.   Statista. Number of mobile cellular subscriptions per 100 inhabitants in South Africa from
2000 to 2020 2021 [cited 2023 October 1, 2023]. Available from:
https://www.statista.com/statistics/510599/mobile-cellular-subscriptions-per-100-inhabitants-in-
south-africa/.
23.   Feldacker C, Pienaar J, Wasunna B, Ndebele F, Khumalo C, Day S, et al. Expanding the
Evidence on the Safety and Efficiency of 2-Way Text Messaging–Based Telehealth for Voluntary
Medical Male Circumcision Follow-up Compared With In-Person Reviews: Randomized Controlled
Trial in Rural and Urban South Africa. Journal of Medical Internet Research. 2023;25:e42111.
24.   Day S, Ncube V, Maja L, Wasunna B, Pienaar J, Setswe G, et al. Centering Frontline Health
Care Workers in Digital Innovation Design to Inform the Optimization of an App for Improved Male
Circumcision Follow-up in South Africa: Qualitative Usability Study. JMIR Form Res. 2023;7:e44122.
Epub 2023/03/23. doi: 10.2196/44122. PubMed PMID: 36947127; PubMed Central PMCID:
PMCPMC10131628.
25.   Cameron DB, Mustafa Diab M, Carroll LN, Bollinger LA, DeCormier Plosky W, Levin C, et al.
The state of costing research for HIV interventions in sub-Saharan Africa. African Journal of AIDS
Research. 2019;18(4):277-88.
26.   Meyer-Rath G, van Rensburg C, Chiu C, Leuner R, Jamieson L, Cohen S. The per-patient costs
of HIV services in South Africa: systematic review and application in the South African HIV
investment case. PloS one. 2019;14(2):e0210497.
27.   Kahn JG, Marseille E, Auvert B. Cost-effectiveness of male circumcision for HIV prevention in
a South African setting. PLoS Med. 2006;3(12):e517. Epub 2006/12/30. doi:

10.1371/journal.pmed.0030517. PubMed PMID: 17194197; PubMed Central PMCID:
PMCPMC1716193.
28.   Kripke K, Chen PA, Vazzano A, Thambinayagam A, Pillay Y, Loykissoonlal D, et al. Cost and
Impact of Voluntary Medical Male Circumcision in South Africa: Focusing the Program on Specific
Age Groups and Provinces. PloS one. 2016;11(7):e0157071. Epub 2016/07/15. doi:
10.1371/journal.pone.0157071. PubMed PMID: 27409079; PubMed Central PMCID:
PMCPMC4943592.
29.   Atkins K, Yeh PT, Kennedy CE, Fonner VA, Sweat MD, O'Reilly KR, et al. Service delivery
interventions to increase uptake of voluntary medical male circumcision for HIV prevention: A
systematic review. PloS one. 2020;15(1):e0227755. Epub 2020/01/14. doi:
10.1371/journal.pone.0227755. PubMed PMID: 31929587; PubMed Central PMCID:
PMCPMC6957297.
30.   Kennedy CE, Yeh PT, Atkins K, Fonner VA, Sweat MD, O'Reilly KR, et al. Economic
compensation interventions to increase uptake of voluntary medical male circumcision for HIV
prevention: A systematic review and meta-analysis. PloS one. 2020;15(1):e0227623. Epub
2020/01/16. doi: 10.1371/journal.pone.0227623. PubMed PMID: 31940422; PubMed Central
PMCID: PMCPMC6961886.
31.   Bansi-Matharu L, Mudimu E, Martin-Hughes R, Hamilton M, Johnson L, Ten Brink D, et al.
Cost-effectiveness of voluntary medical male circumcision for HIV prevention across sub-Saharan
Africa: results from five independent models. Lancet Glob Health. 2023;11(2):e244-e55. Epub
2022/12/24. doi: 10.1016/S2214-109X(22)00515-0.
32.   National Department of Health of the Republic of South Africa. South African National
Guidelines for Medical Male Circumcision. Department of Health of the Republic of South Africa,,
2016.
33.   Population Services International CoSoE, Central and Southern Africa (COSECSA), U.S.
Centers for Disease Control and Prevention, . Adverse Event Action Guide for Voluntary Medical
Male Circumcision by Surgery or Device. 2020 December 2020. Report No.
34.   Automobile Association of South Africa. Fuel pricing 2022 [cited 2023 Oct 1]. Available from:
https://aa.co.za/fuel-pricing/.