Non-linear dilution adjustment of exemplary urine arsenic

PART I: CURVING THE LINES

THOMAS CLEMENS CARMINE MD

Etzelstrasse 21 • 8808 Pfäffikon SZ • Switzerland • Tel +41 55 415 50 30 • research@dr-carmine.ch

Non-linear dilution adjustment of exemplary urine arsenic
Part I: Curving the lines

Contents

Non-linear dilution adjustment of exemplary urine arsenic
Part I: Curving the lines

Thomas Clemens Carmine M.D

Abstract

Background
Conventional dilution adjustment of spot-urinary biomarkers by correction for creatinine (uCR, CCRC), osmolality, or specific gravity remains controversial. Apart from unaccounted confounders like age, sex, muscle mass, or diet, the misperception of constant mass ratios between analyte and corrector over a wide hydration range entails deceptive correction errors foremost at both ends of the hydration spectrum. Mitigating creatinine-correction errors (CRCE) by restricting uCR-ranges to 0.3(0.4)-3 g/L generates high numbers of sample rejects and allows for misleading fluctuations within presumably tolerable uCR-ranges undermining the validity of clinical, forensic, and epidemiological investigations.

Methods
The CRCE for exemplary total weight Arsenic (TWuAs) was analyzed in n= 5599 unselected spot urine samples. After confining data to 14 – 82 years, uncorrected arsenic (uAs_{UC}) < 500 µg/l, and uCR < 4.5g/L, the remaining 5400 samples were partitioned, and a calculation method to standardize uAs_{UC} to 1 g/L uCR was developed based on uAs_{UC}-stratified power functional regression analysis (PFRA).

Findings
The obtained standardizing method of variable power-functional creatinine correction (V-PFCRC) represents a progression of reportedly proven simple power-functional modifications of CCRC (S-PFCRC). In contrast to CCRC and S-PFCRC, standardization to 1g/L uCR by V-PFCRC yielded constant uAs_N-values over the entire uCR-range in all sextiles. Residual bias was largely neutralized in all sextiles of uAs_N (R^2 9E-06 – 0.04), compared to uAs_{UC} (R^2 0.93 – 0.99) and CCRC (R^2 0.42 – 0.89). The strongest %CRCEs were found in samples at low concentrations of uCR (SBC -0.96) and/or uAs_{UC} (SBC -0.11).

Interpretation
Standardization to 1g/L uCR by V-PFRA allows for more reliable dilution adjustment of spot urinary results than CCRC and S-PFCRC. As an easily adoptable, novel approach, the calculation method developed in this study can minimize residual dilution bias in urinary biomarkers adjusted by uCR, SG, and osmolality. While minimizing sample rejects, V-PFCRC will improve reliability and comparability, particularly of lower concentrated urinary biomarkers in both conventional readings and multiple linear regression models.

Funding
The study was not funded.

Non-linear dilution adjustment by the example of urinary arsenic
Part I: Curving the lines

Thomas C. Carmine M.D.

1. Introduction

1.1. Urinary dilution correction in biomonitoring

Urine is the most used specimen in occupational and environmental biomonitoring of toxic metals.[1] For better practicability and compliance, spot urine is commonly favored to 24h urine samples which, unlike spot urine, account for diurnal variation and, to a degree, for hydration. However, spot urinary concentration of a biomarker expressed as mass per urine volume (e.g., µg/L) is an unreliable parameter. It primarily depends on hydration, and variable fluid intakes produce significant differences in urinary concentrations, with diluted urines usually yielding lower biomarker readings than concentrated ones (Fig.1a).[2] In the case of urinary arsenic (uAs), more than a third of the variation in spot urinary concentrations could be attributed to dilution.[3] To standardize intra- and interindividual concentrations over time, fluctuations in spot urine dilution are commonly adjusted for either by urinary creatinine (uCR) or less frequently by specific gravity (SG) and osmolality as proxies for urinary flow rate (UFR).[3]

All these conventional methods, however, only partially adjust for variation in diuresis.[1,3,4,5,6,7,8] Creatinine (CR), a nitrogenous waste product from muscular energy generation by breaking down creatine-phosphate, has been reported to be non-negligibly influenced by age, sex, ethnicity, lean muscle mass, meat-/creatin intake, and collection time, thus complicating inter- and intraindividual comparisons.[6,9,10,11] Furthermore, relations between analyte concentrations and UFR have been determined log-linear instead of linear, as wrongly assumed in dilution adjustments by CCRC and specific gravity/osmolality.[3,5] Despite these obvious fallacies, CCRC, specific gravity, and osmolality remain the most common default approaches to urinary dilution adjustment. All reasonable suggestions to mathematically account for excretion differences have yet to find their way into common clinical practice.[3,4]

1.2. Pitfalls in conventional creatinine correction (CCRC)

In CCRC, the analyte concentration is divided by uCR, thus converting the scale of results to the weight of analyte per weight of creatinine (analyte µg/uCR mg or g). This tends to standardize results because adults excrete CR at a relatively constant daily rate expressed as mg/kg body mass.[12] The justification of CCRC is essentially based on three implicit assumptions:

1) Individual and interindividual uCR-excretion rates were constant over time and vary only by UFR, while the influences of muscularity, age, ethnicity, exercise level, and diet (vegetarian vs. omnivorous) were negligible compared to variability in hydration.[3,12]

2) There was no confounding of CR and analyte by common causal factors nor direct causal relationships between the two.[2,13]

3) The relative ratio between the mass of the analyte and the mass of uCR was preserved with variations in hydration and UFR.

All three assumptions are inadequate and gave rise to criticism and even requirements for complete renunciation of CCRC or its substitution by multiple linear

regression models (MLRM), including uCR as a covariate.[14,15] Most objections are directed to the unaccounted factors mentioned in assumption 1, entailing uncontrolled fluctuations of uCR and analyte/uCR-ratios.[6,8,9,10,13,15] In contrast, the significant flaws in assumptions 2 and 3 appear to be less of a topic in the literature.[3]

Figure 1a/b: Relationship between uCR and TWuAs in linear and power functional regression analysis (PFRA). Subset of all samples < 40 µg/L TWuAs (n=4854) for uncorrected (1a) and conventionally CR corrected (1b) TWuAs. Compare simple linear (blue line) and simple power functional regression (red line) of unstratified data. CCRC assumes linearity, which can be understood as a particular power function with the exponent b (uAs$_{UC}$) taking on the value 1. If this assumption were adequate, the simple division of uncorrected TWuAs by uCR would eliminate the dependency of corrected TWuAs from uCR. Consequently, regression analysis in Fig.1b would produce the parallel blue line to x-Axis (here y = 8.95x/x = 8.95), yielding identical TWuAs results in all hydration states (all uCR). Conventional CR-correction (CCRC), however, produces an inverse dependency of TWuAs from uCR (red line, Fig.2b). Instead of the hypothetical parallel to X-axis (blue line, Fig.1b), another power function can be seen after regression analysis now with a negative exponent b (here -0.447, Fig.1b). The inverse relationship results from the corrective denominator (uCR[(1)]) used in CR-correction being too small for samples <1 g/L uCR and too high for urines > 1g/L uCR. The exponent b in uAs$_C$ reflects the difference between the adequate exponent b (here 0.553, Fig.1a) and the erroneous exponent 1 used in conventional CR correction: $b_{uAsC} \approx b_{AsUC} - 1$. The minor b in uAs$_C$, the more negligible the dependence of uAs$_C$ on uCR, and the more parallel the regression line to the X-axis, the more adequate the corrective function. In this case, the division of uncorrected TWuAs by uCR$^{0.553}$ (instead of uCR[1]) would have yielded a regression function in Fig.1b of y ≈ 8.68 x$^{0.553 - 0.553}$ = 8.68 x^0 = 8.68; in other words, standardization of uAs$_C$ to 1g/L uCR. The shaded areas in Fig.1b represent the uCR ranges considered ineligible for biomarker quantification, and samples within these areas will be rejected. Even in the restricted ranges of 0.3(0.4) - 3 g/L uCR, a considerable bias remains, with the regression curve deviating from the desired parallel to the abscissa.

Figure 1c: Power functional dependency of uAs$_{UC}$ (y$_0$) from uCR (x) as the algebraic base for standardization of uAs$_{UC}$ to 1g/L uCR. Normalized uAs$_N$ (= y$_1$) is yielded by simple division of uAs$_{UC}$ by (uCR)^b. This operation generates identical uAs$_N$ (= y$_1$) across the whole range of uCR (dotted red parallel to the abscissa), whereby the uAs$_N$ equals coefficient *a* at uCR = 1 g/L.

Dependent biological variables standardized by a denominator (here, uCR) generally tend to present unusual statistical properties, potentially impairing the sensitivity and precision of a marker.[11,16] To avoid ratio forms of data, it was proposed to take the denominator variable as a covariate (amongst age, sex, or ethnicity) in MLRM.[1,2,16,17] The covariate solution, however, is complicated by the eventual role of uCR as a so-called 'collider', potentially inducing noncausal associations and further confounding.[2,13] Moreover, it was pointed out that adjusting for uCR as a covariate may not adequately control for the measurement error resulting from inter-individual variations in urinary dilution.[2] While more sophisticated MLRM reportedly minimized bias in a range of simulated theoretical scenarios, their broad implementation remains complicated by demographic heterogeneity, potentially necessitating regional adjustments of MLRM, classification issues for patients of minority- or mixed ethnical backgrounds, elaborate pre-analytical questionings and exams, and an incomplete understanding of the number, nature, and complex interrelations of covariates to be included in MLRM.[2,13]

1.3. Systemic calculation errors due to misperceived linearity
While the controversy over the most accurate dilution adjustment in urinary biomonitoring continues, there seems to be only consensus on the non-validity of uCR-corrected results (uAs$_C$) in excessively dilute or concentrated urines.[1,2,3,17,18] In fact, the net excretion rate ratio between uCR (or SG/osmolality determinants) and a biomarker does neither seem to be independent of biomarker concentrations nor of UFR, as erroneously assumed in conventional dilution adjustments by CCRC and SG/osmolality. Analyte and uCR (likewise SG) usually do change their proportionality in response to UFR, and the over-simplified division of analyte by uCR generates creatinine correction errors (CRCE) either with false elevations in diluted and declines in concentrated urines, like in total weight uAs (TWuAs, Fig.1a/b), or vice versa.[3] Despite its mathematically tangible and accountable nature, this systemic methodological error inherent in all common dilution adjustments (uCR > SG > osmolality) is mitigated mainly symptomatically by restricting dilution ranges.[3] Limiting valid ranges of uCR to 0.3 (0.4) – 3 g/L uCR, however, does render a good portion of samples (in the present study, a total of 22.3%) formally uninterpretable (Fig.1b) and will eminently affect lower uCR samples of females (29% rejection rate in this study), smaller infants (30% rejection rate in < 7-year-olds), and highly diluted samples after excessive intake or recent IV administration of fluids.[10] Since it does not address the root cause of misperceived linearity, uCR-restriction neither provides hydration-independent results within the presumed acceptable range (Figs.1a/b,2c). As the remaining hydration bias will exceed the magnitude of most other uncontrolled covariates, it involves a considerable potential for misjudgment of actual exposures, underlining the need for more accurate dilution adjustment.

Several investigators have already proposed mathematical modifications of CCRC to account for excretion differences in corrective equations.[3,5,18,19,20] A more recent study by Middleton et al. (2019) compared the efficacy of various dilution corrections for spot-urinary arsenic. The authors reported significant mitigation of CRCE by mathematically accounting for the analyte-specific discordance between the respective slopes of uCR and the analyte in relation to UFR.[3] The major improvements of the simple potency-functional approach over CCRC were demonstrated in terms of both the residual dependence of As$_N$ on uCR (R) and the degree of correlation of As$_N$ with actual drinking water exposure. The authors modified CCRC by a simple power functional uCR correction (S-PFCRC) to standardize results to 1g/l uCR.[3,5] Standardized arsenic (uAs$_N$) was expressed as a function of uncorrected uAs (uAs$_{UC}$) and uCR corresponding to Eq.3 of the present study, being fundamental to the method developed herein (Fig.1):

$$uAs_N = uAs_{UC} \times uCR^{-z}$$ (z is equivalent to *b* in the present study, see Fig.1).

Theoretically, z is empirically derived by the ratio of the regression slopes of uAs (b_{uAs}) against UFR by that of uCR (b_{uCR}) against UFR ($z = b_{As}/b_{uCR}$).[3] To remove dilution bias from the sample set, the authors adopted a numeric approach to derive the prior unknown exponent z.[20] After computing uAs_N for a range of z values, z = 0.8 was considered most adequate based on the weakest linear correlation of uAs_N and uCR, with the R-value closest to zero understood as an indicator of the lowest residual dilution variation in uAs_N.[3] The authors resorted to this auxiliary method of z determination due to the absence of comprehensive empirically derived data while acknowledging that forcing a correlation of absolute zero this way may not be completely robust. Another limitation of their study was the modest sample size of 202 spot urines, which precluded systematic analyses of demographic influences like sex or age on correction performance. Nevertheless, their results clearly demonstrated substantial improvements in CCRC results by S-PFCRC, which achieved equivalent minimal residual dilution dependencies for uAs_N as for analogously modified dilution adjustments by SG and osmolality.

In the present retrospective study, the comparatively large set of n = 5599 samples permitted the development of a more sophisticated progression of S-PFCRC, providing stable dilution adjustment in the upper and lower marginal areas of uAs_{UC} (App.7b). Furthermore, the effects of age and sex on the corrective functions could be investigated in more detail (Part II).

1.4. Arsenic

1.4.1. Sources of inorganic and organic Arsenic
According to EPA and ATSDR, the metalloid Arsenic (As) is the most ubiquitous and human-health-relevant environmental toxin.[21] Inorganic As (iAs), in contrast to most organic forms of As (oAs), is highly toxic, and regulatory guidelines exist only for drinking water with the maximum contamination level defined as 10 µg/L As.[22-25] Arsenic is a Group 1 human carcinogen, its principal exposure routes are ingestion of food and water and, to a lesser extent, through air respiration in ambient (e.g., combustion of coal and burning of CCA-treated wood) or occupational settings (e.g., copper, and other metal smelting).[26] Arsenic enters the food chain from drinking water that has seeped through soil containing As derived mainly from geochemical and to a lesser extent from anthropogenic sources like herbicides or antimicrobial growth promoters. Water contaminated with As can be directly consumed, or it can enter the food chain via plants and animals and thus imperils the safety of our food supply in a multifaceted way.[27] For many human populations, seafood has been identified as the primary source of As, since concentrations of As in many fish and shellfish vastly exceed As levels of most drinking waters and other food sources as they commonly mount into the mg/kg range.[26,28-30] Marine-derived foods present As primarily in the form of less harmful, organic As-compounds (oAs), with the non-toxic, Arsenobetaine (AsB, "fish-arsenic") reportedly accounting for 75% - 100% of total oAs in seafood and 0.4% - 90% in freshwater fish.[28,29,31] AsB is rapidly excreted unmetabolized with a half-life of 18 hours.

Compared to most oAs, iAs is highly toxic, and water contaminated with iAs, used for drinking, food preparation, or irrigation of food crops, is thought to pose the greatest threat to the public from environmental As. Another critical source of iAs appears to be rice, which is ten times more absorbent of iAs than any other grain.[24,26]

Table 1: Compilation of the most abundant As species found in humans.

Name	Abbr.	Chemical Structure	Occurence/Origin[26,28,73]	Toxicity[74,75]	% of TWuAs[36,54,64]	% of total blood As[36]	pKa[56]
Arsenite (inorganic)	As(III) iAs(III)		Drinking water, Rice, Flour, various foods, Pesticides	+++	1-3 %	20.2 %	pKa1=9.23 pKa2=12.13 pKa3=13.4
Arsenate (inorganic)	As(V) iAs(V)		Drinking water, Rice, Flour, various foods, Pesticides	++	0.1-0.3%		pKa1=2.19 pKa2=6.98 pKa3=11.53
Dimethylarsinic Acid	DMA DMA(V)		Seafood (small amounts) Primary metabolite of iAs, Arsenosugars, and -lipids	DMA(V) + DMA (III) ++++	30-41%	11.7 %	pKa=6.2
Monomethylarsonic Acid	MMA MMA(V)		Seafood (small amounts) Metabolite of iAs, Arsenosugars, and -lipids	MMA(V) + MMA(III) +++++	2-5%	2.4 %	pKa1=4.1 pKa2=8.7
Arsenobetaine (75-100% of 'Fish-Arsenic')	AsB		Mollusks, Crustaceans, Cephalopods, Finfish, Shellfish, Demersal Fish	(+)?	55-65%	51.1 %	pKa=2.18
Arsenocholine	AsC		Seafood, Fish, Fish oils	(+)?	<1%	4.5 %	
Arsenosugars	AsS		Major As compound in seaweed (brown kelp). Present in mussels, oysters, clamps	(+)?	Metabolized by humans mainly into DMA[69]		
Arsenolipids 1. As-containing Hydrocarbons 2. As-containing Fatty Acids 3. As-containing Phospholipids 4. As-containing Phosphatidylcholines	1. AsHCs 2. AsFAs 3. AsPLs 4. AsPCs		1. Fish oils 2. Fish oils 3. Seaweed 4. Roe	1. (+++)? 2. (+)? 3. (+)? 4. (+)?	Metabolized by humans mainly into DMA[70]		

1.4.2. Absorption, deposition, and biotransformation

Both ingested oAs and iAs (absorption rate 95%) is very well absorbed in the intestine, whereas deposition (ca. 40%) and absorption (75-80% of the deposited As) in the lungs is firmly compound-dependent.[26] After acute intoxication, the highest As concentrations are found in the liver, followed by the kidneys and lungs.[26] Chronic exposure, however, leads to long-term retention in many tissues such as hair, skin, squamous epithelium, upper digestive tract, epididymis, thyroid, lens, and skeleton.[26,32]

The critical step in the biotransformation of absorbed inorganic arsenic is an alternate reduction followed by a sequential oxidative methylation process, which occurs primarily in the liver and to a lesser extent in kidneys and other tissues.[26,33,34] The essential co-factors of oxidative iAs methylation are S-adenosyl methionine (SAME) as the methyl donor and Glutathione (GSH) and NADH as electron donors or redox-cyclers.[2,34] DMA is generally considered the primary metabolic endpoint in the human biotransformation of iAs, accounting for the major share of iAs-derived uAs. The metabolic path from iAs to DMA leads via several methylated As intermediates, some of which (MMA$_{III}$, DMA$_{III}$) exhibit even higher toxicity than iAs (Tbl.1).[34,35] Methylation efficacy is decreased with increasing As-dose levels and in smokers.[26,36] Conversely, methylation capacity increases with age and is higher in females, possibly due to the stimulating effect of estrogen on the production of choline - a substrate for the biosynthesis of SAME in the liver and kidneys.[36] The primary route of excretion of the various inorganic and organic As-compounds is through the kidneys, with only a tiny fraction excreted into bile and feces.[26]

1.4.3. Toxicological and clinical aspects

The two biologically relevant oxidation states of arsenic are the pentavalent arsenate (As$_V$) and trivalent arsenite (As$_{III}$) forms. In solution at physiological pH, the predominant oxyanion of As$_V$ is a substrate analog of phosphate and hence a competitive inhibitor for transporters and enzymes that use or transport phosphate or have phosphorylated

intermediates.[37,38,39] Trivalent As_{III} and its methylated forms are much more toxic and abundant in most human tissues than As_V and account for most of the biohazard from As.[26,27] Trivalent inorganic and organic arsenicals strongly bind to thiolates of closely spaced cysteine residues. Consequently, they are predominantly stored in the protein fraction of mammalian tissue, where they can impede the function of many enzymes.[26,40] They also react with cellular thiols, such as reduced glutathione (GSH), thus elevating the intracellular redox potential and promoting oxidative damage.[27]

The broad clinical toxicological profile of long-term exposure to iAs from drinking water and food includes affections of the liver, gastrointestinal and nervous systems. Typical skin lesions are pigmentation changes, palmar and plantar hyperkeratosis, Blackfoot disease, Raynaud's phenomenon, acrocyanosis, or white Mee's lines in fingernails. Chronic iAs exposure has been associated with cardiovascular disease, diabetes, and various cancers (skin, lungs, and bladder, among others).[23,41-45] In fetal or early childhood exposure, As can negatively impact cognitive development and increase mortality in young adults.[43]

1.4.4. Interrelated renal excretions of uCR and TWuAs

Both creatinine and arsenic are freely glomerular filtered due to their small molecular sizes, and concerning their tubular secretion, common transport molecules, such as MATE1 and MDR1/glycoprotein 1, have been described (Tbls.2,3). Greater differences in renal handling probably exist regarding tubular reabsorption, which in the case of CR occurs at mostly negligible levels and which has been described for arsenic as a function of both urinary dilution and of the activity of a common reabsorption mechanism, i.e., via the activity of the NaPT-IIa/c transporters (Tbl.3).[37-39]

The renal clearance of CR is accomplished primarily by free glomerular filtration and – to a lesser extent – by transporter-mediated tubular secretion, which accounts for 10–40% of total creatinine clearance, depending on kidney function.[46,47] Tubular secretion of uCR becomes more relevant in chronic kidney disease (CKD) when creatinine is progressively hyper-secreted by remnant renal tubules as GFR deteriorates.[48] Apart from CKD, tubular creatinine secretion is upregulated in liver cirrhosis (via OCT2 transporters) and by testosterone.[49-51] In contrast to tubular secretion, reabsorption of creatinine is considered rare, even though it has been observed in some cases, like in newborns and the very elderly.[46]

Table 2: Renal tubular transporters of creatinine.

Membrane site of transport	Sex-dominance	Transporter	References
Basolateral Uptake (potentially bidirectional transporters)	F > M	OAT2	47,49,76,77,78
	M > F	OAT1 OAT3 OCT2 [a]	47,76,77 47,49,77,78,79 47,49,77,80,81,82,83
	equal	OCT3 (OCT1)	49,77 47,77
Apical Excretion (H+/organic cation transporters)	F > M	MDR1/P-Glycoprotein* [b]	77,83
	M > F	MATE1* [a, c]	78,81,82
	equal	MATE2-K (Kidney specific) [c]	78

MDR1/P-Glycoprotein and MATE1 transport both creatinine and very probably also arsenic.
[a] *All drugs known to affect serum creatinine inhibited OCT2 and MATE1.[49,78]*
[b] *conflicting evidence for MDR1/P-GP F>M gender dominance: M>F in rats,[77] F>M in mice,[84] F>M in rats.[85]*
[c] *hMATE1 and hMATE2-K function together by mediating the tubular secretion of intracellular ionic compounds across the brush-border membranes.*

Table 3: Renal tubular transporters of the most common urinary arsenic forms.

Membrane site of transport	Sex-dominance	Transporter	Nephron segment [a]	Arsenic-compound	References
Basolateral Uptake (Potentially bidirectional transporters)	F > M	MRP1 [b] Aquaporin 3 [c]	PT, LH, CD PT, DT, CD	iAs_{III}, iAs_V, MMA_{III}, As(GS)$_3$, MMA(GS)$_2$, DMA_V, iAs_V (?)	77,86,87,88,89,90 87,91,93,94
	M > F	GLUT1	PT	iAs_{III}, MMA_{III}	27,91,92
	not defined	GLUT5 OATP2B1 [d]	PT DT	iAs_{III}, MMA_{III} ?	91 91,98
Apical Excretion [e] (H+/organic cation transporters)	F > M	MDR1/P-Glycoprotein* [b, f] MRP2 [b]	PT PT	iAs_{III}, iAs_V, As(GS)$_3$, MMA(GS)$_2$ As(GS)$_3$, MMA(GS)$_2$	77,84,85,86,87,88 77,86,87,90,91,95,96
	M > F	MATE1* [g] Aquaporin 7	PT PCT	? iAs_{III}	81,82,91 27,87,94,97,99,100
	inconsistent	MRP4 [b, h, i]	PT	iAs_{III}, iAs_V, MMA_{III}, MMA_V, DMA_V	26,77,90,91,99
Apical Uptake	M > F	NaPT-IIa [b, j] Aquaporin 7	PT PCT	iAs_V iAs_{III}	37,87,99,102 87,94,97,100

[a] PT (Proximal tubule), PCT (Proximal convoluted tubule), DT (Distal tubule), LH (Loop of Henle), CD (Collecting Duct)

[b] wide substrate spectra overlap was reported for MRP2, MRP4, and NaPT-II,[103] and for MRP1, MRP2, and MDR1/P-glycoprotein.[90] Differences, however, exist in the transport mechanisms between the primarily active transporter P-glycoprotein on the one side and the GSH-dependent cotransporters MRP1/2/4 on the other.[86]

[c] Human embryonic kidney cell experiments suggest AQP3 facilitates intracellular Arsenic accumulation.[95]

[d] speculative, no direct evidence for As-Transport by OATP2B1 (only for Cisplatin).[90] The substrate specificity and transport rates of OATP2B are strongly pH-sensitive.

[e] apical secretion of iAs(III) may be dependent on the availability of intracellular GSH to form As(GS)$_3$ - and MAs(GS)$_2$-complexes that can be excreted more easily via apical transporters into the primary urine.[35,91]

[f] conflicting evidence for MDR1/P-GP F>M gender dominance: F>M in mice,[84] F>M in rats,[85] M>F in rats.[77]

[g] no direct evidence for Arsenic, but for MATE1-mediated tubular excretion of other metals (Cadmium, Cisplatin).[91,104,105]

[h] conflicting evidence for MRP4 gender dominance: M>F in rats,[77] F>M in mice,[89,106] upregulation by testosterone in human prostate cancer cells.[26]

[i] The protective effect of MRP4 against various Arsenic species (As$_V$, MMA$_V$, DMA$_{III}$) is completely GSH-dependent and may not be exclusively due to MRP4 transmembrane transport as MRP4-mediated excretion of cAMP into primary urine followed by Adenosine-mediated blockage of NaPT-II transporters entails reduced apical uptake of iAs$_V$, which might confound the protective properties of MRP4 against As-toxicity.[105]

[j] iAs$_V$ resembles chemically inorganic phosphate, and there is substrate competition between both for apical uptake via NaPT-IIa.[37,99]

Arsenic, like CR, undergoes free glomerular filtration and tubular secretion, with AsB, MMA, and DMA being much faster excreted in urine than inorganic arsenic.[26,52-54] The renal-tubular As-transport molecules known to date, including their differential substrates and sex preferences, are summarized in Table 3. At least some evidence exists for tubular reabsorption of arsenate (iAs$_V$) followed by its intracellular conversion to arsenite (iAs$_{III}$) and subsequent diffusion of iAs$_{III}$ across the apical and basolateral membranes of the tubular cell in the canine kidney.[55] Consistently, iAs$_{III}$ is found significantly (9.3-fold) higher than less toxic iAs$_V$ in human urinary samples.[44]

The asymmetric tubular reabsorption and net excretion of trivalent and pentavalent arsenic species might be partly explained by variations in acidity and electrical charge (Tbl.1).[56] Both iAs$_{III}$ and iAs$_V$ differ in the oxidative state and molar mass (MM iAs$_{III}$ 122.9 vs. MM iAs$_V$ 138.9). Still, far more pronounced differences exist concerning their acidity. Arsenate (iAs$_V$), featuring an acid dissociation constant of pKa$_1$ 2.19, appears to be substantially more acidic than iAs$_{III}$ (pKa$_1$ 9.23). At physiological urinary pH, iAs$_V$, unlike iAs$_{III}$, will primarily be present in its deprotonated, negatively charged, anionic form. Anionic iAs$_V$ can cross proximal tubular apical membranes using sodium/phosphate cotransporters (NaPT-IIa/c) while substituting and competitively inhibiting the reabsorption of negatively charged inorganic phosphate.[37,38,39] As the low pK$_a$ of iAs$_V$ seems to favor its tubular reabsorption, the prolongated reabsorption of more alkaline species like iAs$_{III}$, MMA (pK$_{a1}$ 4.1), and DMA (pK$_a$ 6.2) might conversely facilitate their faster urinary clearance. Besides their lower acidity, the attached methyl groups of MMA (MM 161), DMA (MM 170), and AsB (MM 178) may additionally slow down the reabsorption of these organic uAs-species by improving their solubility.

An essential factor for tubular reabsorption of at least iAs$_V$ appears to be higher urinary flow, which reportedly inhibited arsenate reabsorption in canine kidneys.[57] The activation of NaPT-II transporters with consequential higher reabsorption of PO$_4^-$ and iAs$_V$

is well known from studies on extreme dehydration, where massive increases in serum PO_4 have been observed.[58,59] Increased plasma calcium levels in the dehydrated state will suppress PTH secretion with successive rises of fractional calcium excretion and elevated tubular reabsorption of PO_4^- and iAs_V.[60]

Consistently, hydration (uCR) also appears to affect the molecular composition of TWuAs. Several studies have consistently shown positive correlations between uCR and urinary %DMA and negative correlations between uCR and %iAs in humans.[54,61-65] A longitudinal study in Bangladesh investigated the influence of uCR and other factors like age, sex, BMI, smoker status, betel-nut intake, season, and daytime fasting during Ramadan on specified uAs (TWuAs, iAs, DMA, MMA).[62] Increased uCR exerted the most substantial effect of all studied factors on uAs parameters: high uCR was positively associated with TWuAs, iAs, MMA, %MMA, DMA, and %DMA, whereas high uCR was inversely associated with %iAs. All these findings are in accordance with disproportionally stronger increases in tubular reabsorption of iAs than of the oAs-compounds in states of lower hydration/UFR.

1.4.5. Potential biochemical confounding of uCR and uAs

The solid positive associations between uCR and %DMA/MMA, paralleled by negative associations between uCR and iAs and associations of low folate levels in blood with an increased proportion of MMA, brought about the hypothesis of potential confounding of uCR and DMA by the availability of a joint biochemical methyl donor S-Adenosyl-methionine (SAME).[14,15] Being crucial for methylation capacity, SAME is an essential co-factor for both 1) the oxidative methylation of iAs to form MMA and DMA and 2) the biosynthesis of creatine, a precursor of CR.[14,15,63] Under common metabolic conditions, about 40 % of SAME is used to produce creatine, whereas arsenic methyltransferase (AS3MT) requires quantitatively much less SAME.[63,65] Considering this highly disproportional consumption of SAME, the required dose of As to implicate a significant reduction of SAME-dependent synthesis of creatine (and subsequently CR and uCR) is expected to be exceptionally high. In animal experiments, SAME was also not found to stimulate the urinary excretion of the methylated arsenic metabolites when administered prior to a challenging dose of inorganic arsenic.[66] Nonetheless, the positive associations between uCR and %MMA/DMA were hypothesized to be confounded by methylation capacity, which, besides the presence of SAME, depends on the variable efficacy of AS3MT to compete with other SAME-dependent enzymes for this essential cofactor.[65] The potential confounding of uAs by SAME led authors to abandon CCRC and use uncorrected uAs_{UC} instead or include uCR as a covariate in MLRM.[14,15] Full omission of dilution correction, however, will most likely imply a stronger urinary dilution bias of uAs_{UC} than a skewed CCRC (Figs.1,2).

Apart from biochemical confounding, several alternative explanations for the association of uCR and %DMA/MMA were proposed, with the theory of net excretions of iAs and DMA/MMA to be disproportionally dependent on UFR the most conclusive.[37] Relatively lower renal net excretion of iAs concurrent with rather constant, hydration-proportional excretion of uCR (and probably methylated uAs) in states of dehydration could likewise explain the positive association between uCR and the percentage of methylated uAs.

1.4.6. Arsenic in urine

Urine usually is the preferred specimen in epidemiological investigations assessing exposure to As.[6] In most studied populations, the methylated metabolites of iAs (DMA, MMA) predominate in urine besides arsenobetaine (AsB, so-called 'fish-arsenic').

However, there exist considerable interindividual variations in TWuAs composition (Tbl.1).[52,54,67,68] A study in a U.S. region with moderate iAs exposure through well water and diet found the distribution of As species quantities in the order AsB > DMA > MMA > iAs.[54] AsB, on average, represented 55 % of TWuAs, albeit fish consumption up to 48 h before urine collection was an exclusion criterion for analyses. Besides, AsB showed the highest variability in urinary concentration (ICR 1.7 - 29.0 vs. ICR 2.7 - 7.3 for DMA), indicating that in outliers with high unspecified TWuAs, the main As-constituent, will most likely be AsB from seafood.

Most studies either measure TWuAs, which comprises all ICP-MS-detectable forms of As in urine, or so-called 'total uAs', representing the sum of speciated iAs and their methylated metabolites (DMA, MMA) while neglecting AsB and other oAs.[24,30,44,63] This sum ('Total uAs') is pragmatically considered a parameter for chronic exposure to drinking water, albeit neglecting the contribution of marine arsenosugars or -lipids to MMA and DMA (Tbl.1) potentially incurring seafood-related misestimates of drinking water exposures.[37,44,57,61-63,69] It was therefore proposed to use the sum of urinary iAs + MMA or even iAs in blood as more adequate parameters for chronic iAs exposure instead of 'Total uAs'.[29,70]

In clinical settings, TWuAs is usually measured as a first-line test, while speciation analysis is reserved for high TWuAs samples. In the present retrospective study, TWuAs was chosen as a paradigmatic marker because of its abundant presence in human urine, providing for a sufficiently high number of samples featuring a broad enough concentration range to establish adequate numerical non-linear adjustment functions.[21,26,41,71] The imprecision due to varying As-sources and the resulting differences in TWuAs composition and, thus, CRCE is understood to be extensively compensated for by the large sample size.

1.6. Objectives of the study
The aims of the present study were:
1) assessing the dependence of TWuAs on uCR
2) developing a reliable and easily implementable mathematical correction method for standardization to uCR of 1 g/L, also applicable to other urinary biomarkers.
3) establishing the validity of the standardization function found by comparing residual dependencies in aggregate and sex-/age disaggregated data applying PFRA.

2. Study population

Anonymized data (sex, age, uCR, and TWuAs determined by GC-MS) of n = 5767 spot urine samples obtained between January 2014 and August 2022 were kindly provided by IMD Institute for Medical Diagnostics Berlin-Potsdam, Germany. The 155 samples (2.7%) presenting uAs$_{UC}$ levels below the detection limit of 1 μg/L and nine outliers with uAs$_{UC}$ > 500 μg/L were precluded from the study, which left a total of 5599 samples (2971 females and 2628 males) for analysis. For determining the standardization formula, a further six samples with extremely high creatinine values > 4.5 g/L and samples from donors outside the age band 14 – 82 years were not considered. Detailed data concerning muscle mass, dietary and exercise habits, clinical status, kidney function, exposure to toxins, ethnicity, and daytime of sample collection could not be provided by IMD. The age structure by sex is illustrated in App.1. Mean age (± SD) of females (51.5 ± 14.7) and males (50.7 ± 16.7) did not differ significantly (Mann-Whitney-U-Test). Conversely, significant male-dominant sex differences (p< 0.001) existed about uCR (F 0.82 ± 0.01, M 1.2 ± 0.014 g/L, means ± SE)

and uAs$_{UC}$ (F 17.8 ± 0.62, M 26.6 ± 0.9 µg/L), as depicted in Appendices 2 and 3. In 82% of samples, uAs$_{UC}$ was < 30 µg/L, in 95.7% < 100 µg/L and nine outliers were found with uAs$_{UC}$ > 500 µg/L (548 – 5,763 µg/L). In contrast to uAs$_{UC}$, uAs$_C$ was significantly higher (p = 0.014) in females (F 25.4 ± 0.83, M 23.9 ± 0.72 µg/g CR, App.4).

3. Algebraic standardization of uAs$_{UC}$ to 1g/L uCR

3.1. Power functional relation between uCR and TWuAs
Figures 1 and 2 illustrate the rather power functional than linear relations between uCR and uAs$_{UC}$. The power function between uAs$_{UC}$ and uCR can be formally described as:

(1) $uAs_{UC} = a \times uCR^{\,b}$ (Fig.1c)

The coefficient *a* represents the expected uAs$_{UC}$ value for uCR = 1g/L:

(2) uAs_{UC} *(µg/L)* $= a \times 1^{\,b}$ *(g/L)* $= a \times 1 = a$ *(for uCR =1 g/L)*

Any adequate corrective function for uAs$_{UC}$ is expected to generate identical corrected TWuAs values over a broad uCR concentration range, reflected by a parallel line to the abscissa (Fig.1b). This can be achieved by algebraic standardization to uAs$_{UC}$ at 1g/L creatinine (Fig.1c):

(3) $uAs_N = uAs_{UC} \times a \,/\, a \times uCR^{\,b} = uAs_{UC} \times uCR^{\,-b}$

Exponent *b* describes the slope of the power function between TWuAs and uCR. In the case of uAs$_{UC}$, *b* usually takes on values between 0 and 1 (here b =0.558, red line, Figure 1a). The resulting curve is ascending but flattening with increasing uCR, which means the uAs$_{UC}$/uCR ratio is incrementally diminished in higher concentrated urines, probably due to disproportional renal adjustment of TWuAs- and uCR- net excretions to urinary flow.[55,61] A particular case exists in *b*, taking on the value 1, which is the flawed implicit assumption of CCRC. Given the free glomerular filtration of both uCR and TWuAs, *b* = 1 would require the ratio of net secretions of both substances to be constant over a wide urinary dilution range, meaning both hydration and renal arsenic load would have to influence the respective tubular secretions of TWuAs and uCR to the same extent.

In contrast, when *b* approximates zero, a parallel to the x-axis results with no correlation between uCR and TWuAs. In the case of uAs$_C$, *b* = 0 indicates optimal correction for uCR with no remaining bias. In uAs$_{UC}$, an unrealistic *b* of 0 would render any adjustment for uCR superfluous.

3.2. Dependence of exponent b on uAs$_{UC}$
To adequately determine the dependence of exponent b on uAs$_{UC}$, the misleading effects of extreme outliers and potential age biases were mitigated by restricting data to samples of the age band 14 - 82 years with uAs$_{UC}$ ≤ 500 µg/L and uCR < 4.5 g/L. The 5400 samples fulfilling all three criteria were stratified into 15 uCR ranges and 6 uAs$_{UC}$-sextiles (Fig.2a). Power-functional regression curves were determined for each uAs$_{UC}$ sextile separately (Fig.2b). The prior partitioning and uAs$_{UC}$-sextile-stratified PFRA of data (Fig.2a) proved to be a necessary step, as S-PFCRC preferably corrects uAs$_{UC}$ values closer to the mean (App.7b). Despite of a decent overall computational correction of CRCE, there remain

substantial residual dependencies between uCR and uAs_N in the marginal quartiles 1 and 4, as revealed by PFRA (App.7b).

Alternatively, to the power function used, the relationship between uCR and uAs_{UC} could be described by a logarithmic function of the form $uAs_{UC} = a + b \times \ln(uCR)$. However, the logarithmic regression curves do not intersect the abscissa at the zero-point (as PFRA and CCRC curves do) but in the positive range leading to nonsensical negative uAs_N correction values at very high urine dilutions (here uCR < 0.11 g/L, data not shown). The coefficients *a* and exponents *b* of the 6 curves of Figure 2b are related to each other in a positive logarithmic way (Fig.3), which forms the basis for the variable power function correction derived in Eqs.1-8. The coefficient *a* of a regression curve equals its uAs_N value, which is assigned to all uAs_{UC} values of that curve along the total range of uCR (Figs.1c,2b).

Figure 2a: Apportionment of samples into 90 equinumerous rectangles (60 values/rectangle).
All 5400 samples of the age band 14 – 82 years with $uAs_{UC} \leq 500$ µg/L and uCR < 4.5 g/L were partitioned into 15 ascending uCR ranges (quindeciles) of 360 values each, and all uCR-quindecile were further divided into 6 uAs_{UC} of 60 values. This way, the area of the graph was subdivided into 90 equinumerous rectangles of variable size.

Figure 2b: Power functional regression analysis of uAs_{UC} sextile based on means of rectangles (Figure 2a).
Dots represent the arithmetical means of uAs_{UC} and uCR for each rectangle. Separate power-functional regression analyses of all sextiles yielded 6 equations taking the general form $y = a \times x^b$ with R^2 indicating the dependence of uAs_{UC} on uCR.

Figure 2c: **Power functional regression analysis of uAs_C.** Apportionment of the 5400 samples into 90 equinumerous rectangles was performed for uAs_C analog to the analysis of uAs_{UC} in Figs.2a/b. Regression functions, including R^2, were calculated for uAs_C-sextiles.

Figure 2d: **Power functional regression analysis of normalized uAs (uAs_N).** The correction for TWuAs-standardization to 1g/L uCR was determined $uAs_N = uAs_{UC} \times uCR^{-[Ln(uAsUC) + 11.82)]/[Ln(uCR) + 18.73]}$ (Sec.3.3, Eq.8). The 5400 uAs_N values were subdivided and analyzed by analogy with uAs_{UC} (Fig.2b) and uAs_C (Fig.2c). An enlarged visualization of sextiles 1-5 is given in App.7a.

The log relation between exponent *b* and coefficient *a* takes the general form of:

(4) $b_{(a)} = c \times Ln(a) + d$ *(here b = 0.053 × ln(a) + 0.631; c = 0.0534, d = 0.631)*

Solving (4) for *a* yields:

(5) $a_{(b)} = exp\ (b/c - d/c)$ *(here a = exp (18.73 × b - 11.82))*

Appendix 8a illustrates the direct proportionality of coefficient *a* to uAs$_{UC}$ for 8 different uCR concentration ranges. Congruent with both the linearity between uAs$_N$ and *a* and the positive log relationship between *a* (uAs$_N$) and *b* (Fig.3), also the dependence of exponent *b* on uAs$_{UC}$ was found to be of positive logarithmic nature (App.8b). With the apparent increase of exponent *b* as a function of uAs$_{UC}$, the relationship between uCR and uAs$_{UC}$ will approximate direct proportionality (*b* = 1) in high uAs$_{UC}$ samples.

In summary, higher TWuAs loads, represented either by *a* (Fig.3, App.8a) or by uAs$_{UC}$ (App.8b), appear to increase the fractional excretion of uAs and thus shift exponent b closer to the value of 1, implying more constant uAs$_{UC}$/uCR-ratios and lower magnitudes of CCRE (Figs.4a-c).

Figure 3: Log-Relationship between *a* (= uAs$_N$) and exponent *b*. Log regression curves were calculated for three different sample apportionments using 12, 20, or 15 uCR ranges subdivided into 30-, 9- or 6-tiles of uAs$_{UC}$ yielding 360, 180, or 90 rectangles containing 15, 30, or 60 samples each. For validation, the log relation between *a* and *b* was investigated by two alternative calculations based on 12 or 20 uCR ranges and 30 or 9 uAs$_{UC}$ levels comprising 15 or 30 samples per rectangle. These alternative regression curves (Figure 3, dotted green and black lines; squares, diamonds) widely overlapped with the original one (blue circles, straight line), indicating systematic fluctuations due to arbitrary sample apportionment to be insignificant. Alternatively, the coefficient *a* could be exponentially expressed by b in the form *a* = 0.0001 × exp (14.3 × *b)*. The stability of this curve (R^2 = 0.63), though, was far lower than the log expression of *b* by *a* (R^2 =0.93). In contrast to the uAs-dependent exponent b in V-PFCRC, both the z of S-PFCRC (green dashed line) and b=1 in CCRC (black dashed line) are invariable. The improvement of CCRC by the S-PFCRC used in is due to z = 0.8 approximating the mean of variable b in V-PFCRC.[3] The greater the deviations of uAs from mean values, the more z in S-PFCRC will differ from *b* in V-PFCRC. The resulting asymmetrical correction is illustrated in App.7b.

3.3. Algebraic standardization of uAs_{UC} to 1g/L uCR

The challenge was finding a corrective equation that eliminated any bias of uAs_C from uCR by compensating for the influences of both uCR and the magnitude of renal TWuAs-load, incurring variations of TWuAs/uCR ratios (reflected by $b \neq 1$). The first resolution step was to express b by a function of uCR and uAs_{UC} while eliminating a:

Solving (1) for a:

(1') $a = uAs_{UC}/uCR^b$

Elimination of a by the junction of (1') and (5):

(6) $uAs_{UC}/uCR^b = exp(b/c - d/c) (= a)$

Solving (6) for b:

(7) $b = [Ln(uAs_{UC}) + d/c]/[Ln(uCR) + 1/c]$

The final corrective equation is generated by plugging (7) in (3):

(8) $uAs_N = uAs_{UC} \times uCR^{-[LN(uAs_{UC}) + d/c]/[Ln(uCR) + 1/c]}$

3.4. Validating standardization

The TWuAs-specific form of Equation 8, $uAs_N = uAs_{UC} \times uCR^{-[Ln(uAs_{UC}) + 11.82]/[Ln(uCR) + 18.73]}$, was applied to the 5400 uAs_{UC} samples forming the basis of Eq.8 (Fig.2b/d, App.7a). For validation of CRCE elimination, uAs_N-stratified PFRA was performed for the uAs_N sextiles analog to uAs_{UC} (Fig.2b) and uAs_C (Fig.2c). All sextile curves of uAs_N show approximate parallelism to the x-axis with b in uAs_N (0.003 – 0.028) far closer to the desired value of zero than in uAs_C (-0.151 – -0.318). Residual dependencies on uCR were found negligible for uAs_N (R^2 9E-06 – 0.04) compared to uAs_C (R^2 0.42 – 0.89). The extensive mitigating effect of standardization on CRCE in aggregate data was confirmed in separate analyses of both sexes and seven different age groups (Part II).

Albeit significant, the absolute differences in means of the three result modes are modest (Tbl.4). In contrast, much broader variations exist in the modal values, adding evidence for higher corrective disparities in lower TWuAs samples (Tbl.4, Figs.4b/c).

The quantitative effects of standardization are illustrated by deviations of uAs_C from uAs_N (App.5). After restriction to 0.3 – 3 g/L uCR, there remains an average % deviation of uAs_C from uAs_N by 11.9 % ± 10.1 (0 – 272%), compared to 16.8% ± 16.4 in the complete, non-uCR restricted dataset. After uCR restriction to 0.3 – 3 g/L, 17% of uAs_C values (unrestricted 28.6%) still deviate from uAs_N by more than 20% and 4.7% (unrestricted 15.6%) by more than 30%. In accordance with the derivation of Eq.8, the percentage deviations of uAs_C from uAs_N strongly depend on both uAs_{UC} and uCR (Figs.4a-c). Since uCR and uAs_{UC} are both confounded primarily by hydration status and possibly by methylation capacity or other potential metabolic factors, they cannot be considered independent variables influencing %CRCE. To better estimate the individual contributions of uCR and uAs_{UC}, multiple linear regression analysis was pragmatically performed with %CRCE as the dependent variable and uCR-adjusted uAs_{UC} percentile rank (Fig.4c) besides log uCR as independent variables (Apps.6,10a). Standardized β coefficients were determined -0.96 for uCR and -0.11 for uAs_{UC}, suggesting the overall influence of uCR on

%CRCE to be substantially stronger than that of uAs$_{UC}$, even though uAs$_{UC}$, particularly at lower percentile ranks, seems to noticeably influence CRCE (Figs.4b/c).

Figure 4a: Influence of uCR on %CRCE. The percentual CRCE (%CRCE) of all (n=5599) samples < 500 µg/L uAs$_{UC}$ was calculated using the formula: %CRCE = 100 × (uAs$_C$ - uAs$_N$)/uAs$_N$. The dataset was subdivided into 13 uCR ranges, and the effects of uCR on %CRCE were analyzed by log regression of means of each uCR range. Grey-shaded areas represent conventionally restricted uCR ranges.

Figure 4b: Influence of absolute uAs$_{UC}$ on %CRCE. All n = 5599 samples were subdivided into 15 uAs$_{UC}$ ranges either without (dots, straight lines, n=5599) or with uCR restriction to 0.3 – 3 g/L (circles, dotted lines, n=4781). The mean values of each range are represented by dots (all uCR) or circles (0.3 – 3 g/L uCR). Stronger negative associations and potential contributions of uAs$_{UC}$ to %CRCE are mainly expected in low uAs$_{UC}$ < 5 µg/L. This effect, though, is strongly confounded by urinary dilution (uCR), as accounted for in Fig.4c. Above levels of 12 µg/L, the mean deviations changed the sign from positive to negative. The inlay depicts an enlarged section of the lower uAs$_{UC}$-range.

Figure 4c: Influence of relative uAs$_{UC}$ on %CRCE. To accommodate the positive confounding effect of uCR on uAs$_{UC}$, all n=5599 values were partitioned into 20 equinumerous uCR ranges of 280 samples each (except for 279 values in the highest range). All subsets were percentile-ranked separately, and each uAs$_{UC}$ value was assigned to the percentile of its subset and plotted against %CRCE. Percentile values more adequately represent renal As-loads as they are largely adjusted for hydration. The effect of uCR on the relative influence of uAs$_{UC}$ is illustrated by 7 differently colored uCR areas and their corresponding polynomial regression lines. In all 7 uCR ranges, the magnitude of %CRCE decreases with increasing percentile rank. All 7 lines converge toward the %CRCE value of 0 with increasing percentile rank irrespective of the algebraic sign of %CRCE, which is positive in the case of uCR<1 and negative in the case of uCR>1.

	Correction mode	Mean ± SE	Median	Modal	Minimum	Maximum
Linear	uAs$_{UC}$ (µg/L)	22.0 ± 0.54	8.9	1.5	1	484
	uAs$_C$ (µg/g CR)	24.7 ± 0.56	11.3	10	0.8	741
	uAs$_N$ (µg/1g CR)	23.3 ± 0.54	10.4	4.3	0.9	724
Log (LN)	uAs$_{UC}$ (µg/L)	2.3 ± 0.016	2.19	1.25	-0.04	6.18
	uAs$_C$ (µg/g CR)	2.6 ± 0.014	2.42	2.48	-0.21	6.61
	uAs$_N$ (µg/1g CR)	2.5 ± 0.014	2.34	2.2	-0.06	6.59

Table 4: Quantitative differences in uAs$_{UC}$, uAs$_C$, and uAs$_N$. Basic descriptive statistics for all n = 5400 samples of the age band 14 – 82 years, < 500 µg/L uAs$_{UC}$, and uCR < 4.5 g/L, constituting Eq.8. Arsenic values were log-transformed to approximate normal distribution. The means of each correction mode (uAs$_{UC}$ vs. uAs$_C$, uAs$_{UC}$ vs. uAs$_N$, and uAs$_C$ vs. uAs$_N$) all differed significantly within the respective log and linear datasets (p > 0.001, paired 2-tailed student t-test).

4. Discussion

4.1. Power-functional creatinine correction (PFCRC)

Power-functional approaches to dilution adjustment generally partition the graphic relation between the uncorrected analyte (here, uAs$_{UC}$) and uCR into a fan of curved lines intersecting the zero point, in contrast to the erroneously assumed straight lines in CCRC. All values on a curve are attributed to the same standardized value. The curvature of the lines as a function of uCR and uAs$_{UC}$ is determined by the disproportional dependence of the respective renal net excretion rates on UFR. In simple PFCRC (S-PFCRC), all lines of the function uAs$_N$ = uAs$_{UC}$ x uCR^{-z} share the identical exponent z, and the variation in arsenic exposure is expressed exclusively by the differences in the factors *a*. In contrast, the variable exponent b of the V-PFCRC presented herein increases in positive logarithmic dependence with uAs$_{UC}$.

Successful attempts to compensate CRCE by S-PFCRC have been well documented.[3,5,18] In S-PFCRC, the dependence of the correction on UFR is represented by the constant exponent z, which can be either empirically determined by resource-intense measurements of biomarker-specific analyte-UFR slopes or, alternatively, by adopting the somewhat asymmetric numerical approach potentially involving the observed unwanted residual dependencies at both edges of corrected data sets (App.7b).[3,20] The herein-developed progression of S-PFCRC to an analyte-stratified variable PFCRC (V-PFCRC) additionally compensates for the independent influence of analyte level on exponent z, denominated '*b*' in V-PFCRC (Fig.3). This proofed necessary, as uAs$_{UC}$, being confounded with uCR by UFR, seems to make a minor independent contribution to %CRCE, especially at lower uCR-adjusted percentile-ranks of uAs$_{UC}$ (Fig.4b/c, App.6). In contrast to S-PFCRC, V-PFCRC does not only attain overall minimization of residual correlations between uAs and uCR in uAs$_N$-data (R^2 and exponent b close to zero comparable to S-PFCRC) but additionally proved to reliably remove residual dependencies in the marginal concentration ranges as seen in unstratified data sets corrected by S-PFCRC (Fig.1d, App.7b). In V-PFCRC the degree of disproportion in UFR dependence is accounted for by two coefficients (*c* and *d*) modifying the standardizing function (Eq.8) in an analyte-specific manner (Figs.3,4). The coefficients *c* and *d* can be easily numerically identified by a larger dataset that includes uCR and analyte concentrations while avoiding more elaborate UFR measurements.

The accuracy of standardizing functions principally depends on the representativity and number of samples available for their establishment. In this study, 5400 samples proved to be enough to attain a robust log relationship between coefficient *a* and exponent *b*. The derived coefficients *c* and *d* proved stable, enabling adequate analyte-specific dilution adjustment (Fig.3). Approximate estimations on fewer samples (≥ 1000) will most likely already allow for considerable improvement in results, especially for less complex markers.

Standardization to 1g/L uCR by V-PFCRC effectively eliminated CRCE over broad urine dilution- and uAs_{UC}-ranges not only in aggregated (Fig.2d, App.7a/b) but also in age- and sex-disaggregated analyses as verified by simple PFRA (see Part II). The quite uniform mitigation of CRCE in all subgroups can be seen as evidence of its principal induction by a common fundamental physiological misconception with only minor quantitative modifications by other biasing factors like sex, age, or diet, which may nevertheless induce considerable bias by other mechanisms than dilution (Part II).

Apart from the asymmetrical net excretions, the generally higher proportions of poorly reabsorbed, methylated As-species like AsB and DMA from predominant marine sources in high TWuAs-samples could further contribute to the effect (Sec.1.4).[28,54,61] Stronger similarity in tubular behavior of methylated As-species to uCR should shift TWuAs/uCR relations closer to linearity, thus minimizing CRCE. In the present study, the imprecision due to varying As-sources and the resulting differences in TWuAs composition and, thus, CRCE is understood to be extensively compensated for by the large sample size. Future studies should determine corrective V-PFCRC formulas for all the individual TWuAs components separately to further control any TWuAs-concentration-dependent composition bias.

4.2. Standardization of other biomarkers and dilution adjustments

Apart from the more complex exemplary parameter TWuAs used herein, the general form of Eq.8, $uAs_N = uAs_{UC} \times uCR^{-[LN(uAs_{UC})+d/c]/[Ln(uCR)+1/c]}$, can be applied to all other spot urinary analytes by determining coefficients *c* and *d* for any urinary biomarker as the analyte-specific basis for adequate dilution adjustment.

The inverse correlation of %CRCE and TWuAs (Fig.4c) suggests CCRC incurring the strongest misestimates of real exposure in urinary analytes present in only minute amounts. Standardizing these low-concentrated analytes to 1 g/L uCR is a prerequisite for their precision.

The reliabilities and validities of dilution adjustments by SG or osmolality, both of which are considered less susceptible to interference than uCR, are compromised in a similar way to CCRC.[3] The negligence of the asynchronous hydration-dependent tubular handling of a biomarker on the one hand and of the urinary electrolytes and osmolytes such as urea, uCR, and uric acid that define the SG of urine on the other will induce related correction errors in both methods, albeit quantitatively weaker than in CCRC.[3,11,18] Previous promising results for simple power-functional modifications of dilution adjustments by SG and osmolality and the linear relationship between CCRC and SG correction support this assumption.[3,5,11,72] It appears consistent that variable power-functional corrections analog to V-PFCRC can likewise provide the algebraic basis for adequately compensating dilution adjustment errors by SG and osmolality.

4.3. Benefits and limitations of standardization by V-PFCRC

S-PFCRC and V-PFCRC account for the physiological hydration-dependent differences in analyte/uCR net excretion ratios. The stable results over a broad urinary dilution (or UFR) range will increase the reliability and comparability of spot urinary biomarkers and permit

the omission of symptomatic uCR-range restrictions to 0.3 – 3 g/L as a source of considerable sample rejections rates. Avoiding cumbersome pre-analytical questioning and exam, as required for more elaborate MLRM, V-PFCRC, and analog power-functional modifications of SG and osmolality, are comparably resource and cost-saving methods. They can be readily implemented in clinical and epidemiological practice by integrating the corrective functions in lab software. Its retrospective application to existing datasets will enable an adequate understanding of previously reported associations between urinary markers and proxies for UFR like uCR, osmolality, and SG.[54,61,62] That said, it should be noted that standardization primarily mitigates the systemic disproportional dependences of uCR- and analyte-excretions on UFR. It will thus strongly improve intra- and interindividual comparability and form a more rational basis for establishing normal ranges or toxic thresholds, which may still exhibit age-, sex-, diet-, ethnical-, cross-cultural or other biases due to hydration-independent toxicokinetic factors. These factors may additionally need to be compensated for, be it by differentiating reference ranges or in MLRM using PFCRC values as covariates (see Part II).[2,17]

A potential for compromise of V-PFCRC lies in the differential tubular reabsorption characteristics of various analyte components in more complex biomarkers. In TWuAs, this might become relevant in major deviations from common TWuAs composition, like in rare acute iAs-poisoning with tubular reabsorption rates substantially higher like usually seen in high TWuAs featuring higher proportions of readily excreted oAs. High doses of iAs will thus entail lower exponents b and higher CCREs, being under-compensated by normalization and leading to overestimated As-loads in dilute and underestimations in concentrated urines (Fig.1). However, this case appears to be of only minor clinical relevance, since standardized results, due to its lower exponent b, will still be more precise than CCRC and excessively high TWuAs samples will be subjected to speciation anyway. Apart from acute poisoning, major regional differences in alimentation, particularly in seafood consumption or IV-fluid- administration shortly before urine sampling, may necessitate adjustments of standardization functions.

5. Conclusions / Key Messages

- Conventional dilution adjustments of spot-urinary biomarkers are flawed with significant hydration-dependent results fluctuations. Previous studies indicated the potential benefits of power functional over conventional linear dilution adjustments.

- The results of the CRCE analysis of exemplary TWuAs conducted in this strongly support V-PFCRC to be a promising, numeric power-functional approach to abolish residual hydration bias in spot urinary biomarker results over wide ranges of uCR and analyte.

- V-PFCRC and analog corrections for specific gravity or osmolality allow for more precise dilution adjustment of spot urinary biomarkers than CCRC and S-PFCRC. The method can yield more accurate standardized readings and thus enhance the explanatory power of epidemiological, clinical, and forensic studies.

6. Abbreviations

As	Arsenic
AsB	Arsenobetaine
ATSDR	Agency for Toxic Substances and Disease Registry
CCRC	Conventional Creatinine Correction
CKD	Chronic Kidney Disease
CR	Creatinine
CRCE	Creatinine Correction Error
%CRCE	% deviation of uAs_C from uAs_N (%CRCE = 100 × (uAs_C - uAs_N)/uAs_N)
DMA	Dimethylarsinic Acid
EPA	Environmental Protection Agency (USA)
iAs	Inorganic Arsenic
ICP-MS	Inductively Coupled Plasma Mass Spectrometry
ICR	Interquartile Range
MLRM	Multiple Linear Regression Models
MMA	Monomethylarsonic Acid
oAs	Organic Arsenic
PO_4^-	Phosphate
PFCRC	Power Functional Creatinine Correction
PFRA	Power Functional Regression Analysis
SAME	S-Adenosyl-Methionine
S-PFCRC	Simple Power Functional Creatinine Correction
sP	Serum Phosphate
TWuAs	Total Weight Urinary Arsenic
uAs	Urinary Arsenic
uAs_{UC}	Uncorrected Urinary Arsenic
uAs_C	Conventionally Creatinine-corrected Urinary Arsenic
uAs_N	Standardized Urinary Arsenic, normalized to 1 g/L uCR
uCR	Urinary Creatinine
UFR	Urinary Flow Rate
V-PFCRC	Variable Power Functional Creatinine Correction

7. Acknowledgements
Dr. Katrin Huesker, department manager of Special Immunology, IMD Berlin, is thanked for providing anonymized spot urine data.
Mischa Katz GE is thanked for his timesaving help in data preparation and programming.

8. Author contribution
The idea, basic concept, procurement, preparation and evaluation of data, literature research, developing corrective functions, scrapping erroneous corrective functions, graphs, tables, labels, writing, and submitting the paper.

9. Conflict of Interest
No conflict of interest is known to me with persons, companies, institutions, myself, or any other entity known or unknown to me.

10. Ethics

The study was exempt from ethical approval by the Ethic Committee of Northwest- and Central Switzerland (EKNZ), 4056 Basel, on basis of retrospective analysis of anonymized data without clinical information other than urinary creatinine/TWuAs, age in years, sex, and date of exam. The purpose was methodological quality improvement only. No pharmacological or other medical intervention, no risk for physical or psychological harm, identifiability, or breach of confidentiality for any subject involved.

11. Funding

The study was not funded.

12. References

1) Santonen T, Schoeters G, Nordberg M. Chapter 9 - Biological monitoring of metals and biomarkers. Editor(s): Nordberg GF, Costa M. Handbook on the Toxicology of Metals (Fifth Edition), Vol 1, Academic Press, 2022: 217-35.
ISBN 9780128232927, https://doi.org/10.1016/B978-0-12-823292-7.00007-3.

2) O'Brien KM, Upson K, Cook NR, Weinberg CR. Environmental Chemicals in Urine and Blood: Improving Methods for Creatinine and Lipid Adjustment. Environ Health Perspect. 2016 Feb; 124(2): 220-27. doi: 10.1289/ehp.1509693. Epub 2015 Jul 24. PMID: 26219104; PMCID: PMC4749084.

3) Middleton DRS, Watts MJ, Polya DA. A comparative assessment of dilution correction methods for spot urinary analyte concentrations in a UK population exposed to arsenic in drinking water. Environ Int. 2019 Sep; 130: 104721. doi: 10.1016/j.envint.2019.03.069. Epub 2019 Jun 14. PMID: 31207477; PMCID: PMC6686075.

4) Akerstrom M, Barregard L, Lundh T, Sallsten G. The relationship between cadmium in kidney and cadmium in urine and blood in an environmentally exposed population. Toxicol Appl Pharmacol. 2013 May 1; 268(3): 286-93. doi: 10.1016/j.taap.2013.02.009. Epub 2013 Feb 27. PMID: 23454399.

5) Araki S, Sata F, Murata K. Adjustment for urinary flow rate: an improved approach to biological monitoring. Int. Arch. Occup. Environ. Health. 1990; 62(6): 471–77.

6) Sauvé JF, Lévesque M, Huard M, et al. Creatinine and specific gravity normalization in biological monitoring of occupational exposures. J Occup Environ Hyg. 2015; 12(2): 123-29. doi: 10.1080/15459624.2014.955179. PMID: 25192246.

7) Boeniger MF, Lowry LK, Rosenberg J. Interpretation of urine results used to assess chemical exposure with emphasis on creatinine adjustments: a review. Am Ind Hyg Assoc J. 1993 Oct; 54(10): 615-27. doi: 10.1080/15298669391355134. PMID: 8237794.

8) Suwazono Y, Akesson A, Alfvén T, Järup L, Vahter M. Creatinine versus specific gravity-adjusted urinary cadmium concentrations. Biomarkers. 2005 Mar-Jun; 10(2-3): 117-26. doi: 10.1080/13547500500159001. PMID: 16076727.

9) Alessio L, Berlin A, Dell'Orto A, Toffoletto F, Ghezzi I. Reliability of urinary creatinine as a parameter used to adjust values of urinary biological indicators. Int Arch Occup Environ Health. 1985; 55(2): 99-106. doi: 10.1007/BF00378371. PMID: 3988361.

10) Cocker J, Mason HJ, Warren ND, Cotton RJ. Creatinine adjustment of biological monitoring results. Occup Med (Lond). 2011 Aug; 61(5): 349-53.
doi: 10.1093/occmed/kqr084. PMID: 21831823.

11) Muscat JE, Liu A, Richie JP Jr. A comparison of creatinine vs. specific gravity to correct for urinary dilution of cotinine. Biomarkers. 2011 May; 16(3): 206-11.
doi: 10.3109/1354750X.2010.538084. Epub 2011 Feb 3. PMID: 21288164; PMCID: PMC3631104.

12) Allen R, Mage D, Gondy G, Christensen C, Barr D, Needham L. The use of a creatinine correction for reporting children's urinary pesticide concentrations. Epidemiology. July 2004; 15(4): 69-70.

13) Cole SR, Platt RW, Schisterman EF, et al. Illustrating bias due to conditioning on a collider. Int J Epidemiol. 2010 Apr; 39(2): 417-20. doi: 10.1093/ije/dyp334. Epub 2009 Nov 19. PMID: 19926667; PMCID: PMC2846442.

14) Gamble MV, Hall MN. Relationship of creatinine and nutrition with arsenic metabolism. Environ Health Perspect. 2012 Apr; 120(4): A145-46. doi: 10.1289/ehp.1104807. PMID: 22469551; PMCID: PMC3339472.

15) Basu A, Mitra S, Chung J, Guha Mazumder DN, Ghosh N, Kalman D, von Ehrenstein OS, Steinmaus C, Liaw J, Smith AH. Creatinine, diet, micronutrients, and arsenic methylation in West Bengal, India. Environ Health Perspect. 2011 Sep; 119(9): 1308-13. doi: 10.1289/ehp.1003393. Epub 2011 Jun 7. PMID: 21652291; PMCID: PMC3230402.

16) Anderson D, Lydic R. Ratio data and the quantification of drug effects. Biobehavioral Reviews. 1977; 1(1): 55-7. doi: 10.1016/0147-7552(77)90041-9. ISSN 0147-7552.

17) Barr DB, Wilder LC, Caudill SP, Gonzalez AJ, Needham LL, Pirkle JL. Urinary creatinine concentrations in the U.S. population: implications for urinary biologic monitoring measurements. Environ Health Perspect. 2005 Feb; 113(2): 192-200. doi: 10.1289/ehp.7337. PMID: 15687057; PMCID: PMC1277864.

18) Middleton DR, Watts MJ, Lark RM, Milne CJ, Polya DA. Assessing urinary flow rate, creatinine, osmolality and other hydration adjustment methods for urinary biomonitoring using NHANES arsenic, iodine, lead and cadmium data. Environ Health. 2016 Jun10; 15(1): 68. doi: 10.1186/s12940-016-0152-x. PMID: 27286873; PMCID: PMC4902931.

19) Middleton D. Arsenic Research and Global Sustainability: Proceedings of the Sixth International Congress on Arsenic in the Environment (As2016), June 19-23, 2016, Stockholm, Sweden. CRC Press; 2016. A modified creatinine adjustment method to improve urinary biomonitoring of exposure to arsenic in drinking water.

20) Vij HS, Howell S. Improving the specific gravity adjustment method for assessing urinary concentrations of toxic substances. Am Ind Hyg Assoc J. 1998 Jun; 59(6): 375-80. doi: 10.1080/15428119891010622. PMID: 9670467.

21) Agency for Toxic Substances and Disease Registry (ATSDR), 2019 Substance Priority List, https://www.atsdr.cdc.gov/spl/index.html#2019spl.

22) WHO. Background document for development of WHO Guidelines for Drinking-water Quality (2003). Arsenic in Drinking-water. https://apps.who.int/iris/handle/10665/75375.

23) ATSDR. Arsenic Toxicity (2011). What Are the Standards and Regulations for Arsenic Exposure? https://www.atsdr.cdc.gov/csem/arsenic/docs/arsenic.pdf

24) US Food and Drug Administration (FDA). Arsenic in rice and rice products risk assessment report (2016). https://www.regulations.gov/docket/FDA-2016-D-1099.

25) Wilson TA. FDA Proposes Limit On Inorganic Arsenic In Infant Rice Cereal. InsideHealthPolicy. com's FDA Week. 2016;22(14):14.

26) Fowler BA, Selene CH, Chou J, Jones RL, Costa M, Chen CJ. Chapter 3 - Arsenic. Editor(s): Gunnar F. Nordberg, Max Costa. Handbook on the Toxicology of Metals (Fifth Edition), Vol 2, Academic Press, 2022; 41-89, ISBN 9780128229460, https://doi.org/10.1016/B978-0-12-822946-0.00037-4.

27) Mukhopadhyay R, Bhattacharjee H, Rosen BP. Aquaglyceroporins: generalized metalloid channels. Biochim Biophys Acta. 2014 May;1840(5):1583-91. doi: 10.1016/j.bbagen.2013.11.021. Epub 2013 Nov 27. PMID: 24291688; PMCID: PMC3960311.

28) Taylor V, Goodale B, Raab B, et al. Human exposure to organic arsenic species from seafood. Science of The Total Environment. 2017; 580: 266-82, ISSN 0048-9697, doi: org/10.1016/j.scitotenv.2016.12.113.

29) Aleksandra Popowich, Qi Zhang, X. Chris Le, Arsenobetaine: the ongoing mystery, National Science Review.2016 Dec;3(4):451-58. https://doi.org/10.1093/nsr/nww061.

30) Roswall N, Hvidtfeldt UA, Harrington J, Levine KE, Sørensen M, Tjønneland A, Meliker JR, Raaschou-Nielsen O. Predictors of Urinary Arsenic Levels among Postmenopausal Danish Women. Int J Environ Res Public Health. 2018 Jun;15(7):1340. doi: 10.3390/ijerph15071340. PMID: 29949863; PMCID: PMC6068487.

31) Slejkovec Z, Bajc Z, Doganoc DZ. Arsenic speciation patterns in freshwater fish. Talanta. 2004 Apr;62(5):931-6. doi: 10.1016/j.talanta.2003.10.012. PMID: 18969382.

32) Vahter M, Marafante E, Lindgren A, et al. Tissue distribution and subcellular binding of arsenic in Marmoset monkeys after injection of ^{74}As-Arsenite. Arch Toxicol. 1982;51:65–77. https://doi.org/10.1007/BF00279322.

33) Buchet JP, Lauwerys R. Role of thiols in the in-vitro methylation of inorganic arsenic by rat liver cytosol. Biochem Pharmacol. 1988 Aug;37(16):3149-53. doi: 10.1016/0006-2952(88)90313-9. PMID: 3401245.

34) M. Lu, H. Wang, J. Geisel, X. Chris Le. Enzyme Digestion for Speciation of Arsenic, Editor(s): Janusz Pawliszyn, Comprehensive Sampling and Sample Preparation. Academic Press. 2012:421-433. ISBN 9780123813749. https://doi.org/10.1016/B978-0-12-381373-2.00150-2.

35) Mass MJ, Tennant A, Roop BC, et al. Methylated trivalent arsenic species are genotoxic. Chem Res Toxicol. 2001 Apr;14(4):355-61. doi: 10.1021/tx000251l. PMID: 11304123.

36) Choi JW, Song YC, Cheong NY, et al. Concentrations of blood and urinary arsenic species and their characteristics in general Korean population. Environ Res. 2022 Nov;214(2):113846. doi: 10.1016/j.envres.2022.113846. Epub 2022 Jul 9. PMID: 35820651.

37) Villa-Bellosta R, Sorribas V. Different effects of arsenate and phosphonoformate on P(i) transport adaptation in opossum kidney cells. Am J Physiol Cell Physiol. 2009 Sep; 297(3): C516-25. doi: 10.1152/ajpcell.00186.2009. Epub 2009 Jun 24. PMID: 19553564.

38) Villa-Bellosta R, Sorribas V. Role of rat sodium/phosphate cotransporters in the cell membrane transport of arsenate. Toxicol Appl Pharmacol. 2008 Oct;232(1):125-34. doi: 10.1016/j.taap.2008.05.026. Epub 2008 Jun 10. PMID: 18586044.

39) Villa-Bellosta R, Sorribas V. Arsenate transport by sodium/phosphate cotransporter type IIb. Toxicol Appl Pharmacol. 2010 Aug 15;247(1):36-40. doi: 10.1016/j.taap.2010.05.012. Epub 2010 May 25. PMID: 20510259.

40) Marafante E, Rade J, Sabbioni E, Bertolero F, Foà V. Intracellular interaction and metabolic fate of arsenite in the rabbit. Clin Toxicol. 1981 Nov;18(11):1335-41. doi: 10.3109/00099308109035074. PMID: 7341060.

41) Hall AH. Chronic arsenic poisoning. Toxicol Lett. 2002 Mar 10; 128(1-3): 69-72. doi: 10.1016/s0378-4274(01)00534-3. PMID: 11869818.

42) Agency for Toxic Substances and Disease Registry (ATSDR).Toxicological Profile for Arsenic. Agency for Toxic Substances and Disease Registry. U.S. Department of Health and Human Services, Atlanta, GA (2007). http://www.atsdr.cdc.gov/ToxProfiles/tp2.pdf

43) WHO. World Health Organization Fact Sheet (2018). Arsenic. https://www.who.int/news-room/fact-sheets/detail/arsenic

44) Hopenhayn-Rich C, Biggs ML, Smith AH, Kalman DA, Moore LE. Methylation study of a population environmentally exposed to arsenic in drinking water. Environ Health Perspect. 1996 Jun; 104(6): 620-28. doi: 10.1289/ehp.96104620. PMID: 8793350; PMCID: PMC1469390.

45) Straif K, Benbrahim-Tallaa L, Baan R, et al. WHO International Agency for Research on Cancer Monograph Working Group. A review of human carcinogens-Part C: metals, arsenic, dusts, and fibres. Lancet Oncol. 2009 May;10(5):453-4. doi: 10.1016/s1470-2045(09)70134-2. PMID: 19418618.

46) Tsuda A, Ishimura E, Machiba Y, et al. Increased Glomerular Hydrostatic Pressure is Associated with Tubular Creatinine Reabsorption in Healthy Subjects. Kidney Blood Press Res 2020; 45: 996-1008. DOI: 10.1159/000510838

47) Omote S, Matsuoka N, Arakawa H, Nakanishi T, Tamai I. Effect of tyrosine kinase inhibitors on renal handling of creatinine by MATE1. Sci Rep. 2018 Jun;8(1):9237. doi:10.1038/s41598-018-27672-y. PMID: 29915248; PMCID: PMC6006426.

48) Shemesh O, Golbetz H, Kriss JP, Myers BD. Limitations of creatinine as a filtration marker in glomerulopathic patients. Kidney Int. 1985 Nov; 28(5): 830-38. doi: 10.1038/ki.1985.205. PMID: 2418254.

49) Ciarimboli G, Lancaster CS, Schlatter E, et al. Proximal tubular secretion of creatinine by organic cation transporter OCT2 in cancer patients. Clin Cancer Res. 2012 Feb 15; 18(4): 1101-08. doi: 10.1158/1078-0432.CCR-11-2503. Epub 2012 Jan 5. PMID: 22223530; PMCID: PMC3288323.

50) Perucca J, Bouby N, Valeix P, Bankir L. Sex difference in urine concentration across differing ages, sodium intake, and level of kidney disease. Am J Physiol Regul Integr Comp Physiol. 2007 Feb;292(2): R700-5. doi: 10.1152/ajpregu.00500.2006. Epub 2006 Sep 21. PMID: 16990487.

51) Harvey AM, Malvin RL. Comparison of creatinine and inulin clearances in male and female rats. Am J Physiol. 1965 Oct;209(4):849-52. doi: 10.1152/ajplegacy.1965.209.4.849. PMID: 4284758.

52) Buchet JP, Lauwerys R, Roels H. Comparison of the urinary excretion of arsenic metabolites after a single oral dose of sodium arsenite, monomethylarsonate, or dimethylarsinate in man. Int Arch Occup Environ Health. 1981; 48(1): 71-9. doi: 10.1007/BF00405933. PMID: 6894292.

53) Zhang, X., Rule, A.D., McCulloch, C.E. et al. Tubular secretion of creatinine and kidney function: an observational study. BMC Nephrol. 2020;21:108. https://doi.org/10.1186/s12882-020-01736-6.

54) Gilbert-Diamond D, Li Z, Perry AE, et al. A population-based case-control study of urinary arsenic species and squamous cell carcinoma in New Hampshire, USA. Environmental Health Perspectives. 2013 Oct; 121(10): 1154-60. doi: 10.1289/ehp.1206178. PMID: 23872349; PMCID: PMC3801199.

55) Ginsburg JM. The renal mechanism for excretion and transformation of arsenic in the dog. Am J Physiol. 1965 May; 208: 832-40. doi: 10.1152/ajplegacy.1965.208.5.832. PMID: 14286849.

56) Michael S. Reid, Karen S. Hoy, Jordan R.M. Schofield, Jagdeesh S. Uppal, Yanwen Lin, Xiufen Lu, Hanyong Peng, X. Chris Le. Arsenic speciation analysis: A review with an emphasis on chromatographic separations. TrAC Trends in Analytical Chemistry. Volume 123. 2020; 115770; ISSN 0165-9936, doi: 10.1016/j.trac.2019.115770.

57) Ginsburg JM, Lotspeich WD. Interrelations of arsenate and phosphate transport in the dog kidney. Am J Physiol. 1963 Oct; 205: 707-14. doi: 10.1152/ajplegacy.1963.205.4.707. PMID: 14060809.

58) Mohanlal D, Pettifor JM, Moodley GP. Serum calcium and phosphate disturbances during rehydration in acute dehydrating gastroenteritis. J Pediatr Gastroenterol Nutr. 1987 Mar-Apr;6(2):252-6. doi: 10.1097/00005176-198703000-00016. PMID: 3694349.

59) Murtaza A, Khan SR, Butt KS, Lindblad BS, Aperia A. Hypocalcemia and hyperphosphatemia in severely dehydrated children with and without convulsions. Acta Paediatr Scand. 1988 Mar;77(2):251-6. doi: 10.1111/j.1651-2227.1988.tb10638.x. PMID: 3354336.

60) Acharya R, Winters DM, Rowe C, Buckley N, Kafle S, Chhetri B. An unusual case of severe hypercalcemia: as dehydrated as a bone. J Community Hosp Intern Med Perspect. 2021 Jan 26;11(1):135-138. doi: 10.1080/20009666.2020.1851859. PMID: 33552436; PMCID: PMC7850409.

61) Peters BA, Hall MN, Liu X, et al. Creatinine, arsenic metabolism, and renal function in an arsenic-exposed population in Bangladesh. PLoS One. 2014 Dec 1; 9(12): e113760. doi: 10.1371/journal.pone.0113760. PMID: 25438247; PMCID: PMC4249915.

62) Kile ML, Hoffman E, Hsueh YM, Afroz S, et al. Variability in biomarkers of arsenic exposure and metabolism in adults over time. Environ Health Perspect. 2009 Mar; 117(3): 455-60. doi: 10.1289/ehp.11251. Epub 2008 Nov 19. PMID: 19337522; PMCID: PMC2661917.

63) Peters BA, Hall MN, Liu X, et al. Renal function is associated with indicators of arsenic methylation capacity in Bangladeshi adults. Environ Res. 2015 Nov; 143(Pt A): 123-30. doi: 10.1016/j.envres.2015.10.001. Epub 2015 Oct 19. PMID: 26476787; PMCID: PMC4740972.

64) Hata A, Yamanaka K, Habib MA, Endo Y, Fujitani N, Endo G. Arsenic speciation analysis of urine samples from individuals living in an arsenic-contaminated area in Bangladesh. Environ Health Prev Med. 2012 May; 17(3):235-45. doi: 10.1007/s12199-011-0247-5. Epub 2011 Nov 3. PMID: 22048870; PMCID: PMC3348244.

65) Stead LM, Brosnan JT, Brosnan ME, Vance DE, Jacobs RL. Is it time to reevaluate methyl balance in humans? Am J Clin Nutr. 2006 Jan;83(1):5-10. doi: 10.1093/ajcn/83.1.5. PMID: 16400042.

66) Buchet JP, Lauwerys R. Study of factors influencing the in vivo methylation of inorganic arsenic in rats. Toxicol Appl Pharmacol. 1987 Oct;91(1):65-74. doi: 10.1016/0041-008x(87)90194-3. PMID: 3672518.

67) Vahter M. Genetic polymorphism in the biotransformation of inorganic arsenic and its role in toxicity. Toxicol Lett. 2000 Mar 15; 112-113: 209-17. doi: 10.1016/s0378-4274(99)00271-4. PMID: 10720733.

68) Lai VW, Sun Y, Ting E, Cullen WR, Reimer KJ. Arsenic speciation in human urine: are we all the same? Toxicol Appl Pharmacol. 2004 Aug;198(3):297-306. doi: 10.1016/j.taap.2003.10.033. PMID: 15276409.

69) Calderon RL, Hudgens E, Le XC, Schreinemachers D, Thomas DJ. Excretion of arsenic in urine as a function of exposure to arsenic in drinking water. Environ Health Perspect. 1999 Aug;107(8):663-7. doi: 10.1289/ehp.99107663. PMID: 10417365; PMCID: PMC1566491.

70) Hata A, Kurosawa H, Endo Y, Yamanaka K, Fujitani N, Endo G. A biological indicator of inorganic arsenic exposure using the sum of urinary inorganic arsenic and monomethylarsonic acid concentrations. J Occup Health. 2016 May;58(2):196-200. doi: 10.1539/joh.15-0241-OA. Epub 2016 Mar 24. PMID: 27010090; PMCID: PMC5356966.

71) Apostoli P, Bartoli D, Alessio L, Buchet JP. Biological monitoring of occupational exposure to inorganic arsenic. Occup Environ Med. 1999 Dec;56(12):825-32. DOI: 10.1136/oem.56.12.825. PMID: 10658539; PMCID: PMC1757692.

72) Ozdemir S, Sears CG, Harrington JM, et al. Relationship between Urine Creatinine and Urine Osmolality in Spot Samples among Men and Women in the Danish Diet Cancer and Health Cohort. Toxics. 2021 Nov 1;9(11):282. doi: 10.3390/toxics9110282. PMID: 34822673; PMCID: PMC8625939.

73) Molin M, Ulven SM, Meltzer HM, Alexander J. Arsenic in the human food chain, biotransformation, and toxicology - Review focusing on seafood arsenic. J Trace Elem Med Biol. 2015; 31: 249-59. doi: 10.1016/j.jtemb.2015.01.010. Epub 2015 Jan 28. PMID: 25666158.

74) Petrick JS, Ayala-Fierro F, Cullen WR, Carter DE, Vasken AH. Monomethylarsonous acid (MMA(III)) is more toxic than arsenite in Chang human hepatocytes. Toxicol Appl Pharmacol. 2000 Mar 1; 163(2): 203-07. doi: 10.1006/taap.1999.8872. PMID: 10698679.

75) Vahter M, Concha G. Role of metabolism in arsenic toxicity. Pharmacol Toxicol. 2001 Jul; 89(1): 1-5. doi: 10.1034/j.1600-0773.2001.d01-128.x. PMID: 11484904.

76) Eisner C, Faulhaber-Walter R, Wang Y, et al. Major contribution of tubular secretion to creatinine clearance in mice. Kidney Int. 2010 Mar;77(6):519-26. doi: 10.1038/ki.2009.501. Epub 2009 Dec 23. PMID: 20032962; PMCID: PMC3160625.

77) Sabolić I, Asif AR, Budach WE, Wanke C, Bahn A, Burckhardt G. Gender differences in kidney function. Pflugers Arch. 2007 Dec;455(3):397-429. doi: 10.1007/s00424-007-0308-1. Epub 2007 Jul 19. PMID: 17638010.

78) Lepist EI, Zhang X, Hao J, et al. Contribution of the organic anion transporter OAT2 to the renal active tubular secretion of creatinine and mechanism for serum creatinine elevations caused by cobicistat. Kidney Int. 2014 Aug;86(2):350-7. doi: 10.1038/ki.2014.66. Epub 2014 Mar 19. PMID: 24646860; PMCID: PMC4120670.

79) Kobayashi Y, Hirokawa N, Ohshiro N, et al. Differential gene expression of organic anion transporters in male and female rats. Biochem Biophys Res Commun. 2002 Jan 11;290(1):482-7. doi: 10.1006/bbrc.2001.6180. PMID: 11779196.

80) Wang K, Kestenbaum B. Proximal Tubular Secretory Clearance: A Neglected Partner of Kidney Function. Clin J Am Soc Nephrol. 2018 Aug 7;13(8):1291-1296. doi: 10.2215/CJN.12001017. Epub 2018 Feb 28. PMID: 29490976; PMCID: PMC6086711.

81) Ligen A, Dandan Z, Wolin H, et al. Gonadal hormone regulates the expression MATE1 and OCT2 in type 2 diabetes mice. Endocrine Abstracts. 2015; 37: 376 DOI: 10.1530/endoabs.37.EP376.

82) He R, Ai L, Zhang D, et al. Different effect of testosterone and oestrogen on urinary excretion of metformin via regulating OCTs and MATEs expression in the kidney of mice. J Cell Mol Med. 2016 Dec;20(12):2309-17. doi: 10.1111/jcmm.12922. Epub 2016 Jul 29. PMID: 27469532; PMCID: PMC5134372.

83) Brown CD, Sayer R, Windass AS, et al. Characterisation of human tubular cell monolayers as a model of proximal tubular xenobiotic handling. Toxicol Appl Pharmacol. 2008 Dec 15;233(3):428-38. doi: 10.1016/j.taap.2008.09.018. Epub 2008 Oct 1. PMID: 18930752.

84) Kanado Y, Tsurudome Y, Omata Y, et al. Estradiol regulation of P-glycoprotein expression in mouse kidney and human tubular epithelial cells, implication for renal clearance of drugs. Biochem Biophys Res Commun. 2019 Nov;519(3):613-619. doi: 10.1016/j.bbrc.2019.09.021. Epub 2019 Sep 17. PMID: 31540689.

85) Suzuki T, Zhao YL, Nadai M, et al. Gender-related differences in expression and function of hepatic P-glycoprotein and multidrug resistance-associated protein (Mrp2) in rats. Life Sci. 2006 Jun;79(5):455-61. doi: 10.1016/j.lfs.2006.01.024. Epub 2006 Feb 17. PMID: 16483613.

86) Bridges CC, Zalups RK. Molecular and ionic mimicry and the transport of toxic metals. Toxicol Appl Pharmacol. 2005 May;204(3):274-308. doi: 10.1016/j.taap.2004.09.007. PMID: 15845419; PMCID: PMC2409291.

87) Bhattacharjee H, Rosen BP, Mukhopadhyay R. Aquaglyceroporins and metalloid transport: implications in human diseases. Handb Exp Pharmacol. 2009;(190):309-25. doi: 10.1007/978-3-540-79885-9_16. PMID: 19096785; PMCID: PMC2729095.

88) Banerjee M, Kaur G, Whitlock BD, Carew MW, Le XC, Leslie EM. Multidrug Resistance Protein 1 (MRP1/ABCC1)-Mediated Cellular Protection and Transport of Methylated Arsenic Metabolites Differs between Human Cell Lines. Drug Metab Dispos. 2018 Aug;46(8):1096-1105. doi: 10.1124/dmd.117.079640. Epub 2018 May 11. PMID: 29752257.

89) Maher JM, Slitt AL, Cherrington NJ, Cheng X, Klaassen CD. Tissue distribution and hepatic and renal ontogeny of the multidrug resistance-associated protein (Mrp) family in mice. Drug Metab Dispos. 2005 Jul;33(7):947-55. doi: 10.1124/dmd.105.003780. Epub 2005 Mar 31. PMID: 15802388.

90) Leslie EM, Deeley RG, Cole SP. Multidrug resistance proteins: role of P-glycoprotein, MRP1, MRP2, and BCRP (ABCG2) in tissue defense. Toxicol Appl Pharmacol. 2005 May 1;204(3):216-37. doi: 10.1016/j.taap.2004.10.012. PMID: 15845415.

91) Orr SE, Bridges CC. Chronic Kidney Disease and Exposure to Nephrotoxic Metals. International Journal of Molecular Sciences. 2017; 18(5):1039. https://doi.org/10.3390/ijms18051039

92) Gonzalez-Menendez P, Hevia D, Mayo JC, Sainz RM. The dark side of glucose transporters in prostate cancer: Are they a new feature to characterize carcinomas? Int J Cancer. 2018 Jun;142(12):2414-2424. doi: 10.1002/ijc.31165. Epub 2017 Dec 4. PMID: 29159872.

93) Tang S, Leung JC, Lam CW, Lai FM, Chan TM, Lai KN. In vitro studies of aquaporins 1 and 3 expression in cultured human proximal tubular cells: upregulation by transferrin but not albumin. Am J Kidney Dis. 2001 Aug;38(2):317-30. doi: 10.1053/ajkd.2001.26097. PMID: 11479158.

94) Loh SY, Giribabu N, Gholami K, Salleh N. Effects of testosterone on mean arterial pressure and aquaporin (AQP)-1, 2, 3, 4, 6 and 7 expressions in the kidney of orchidectomized, adult male Sprague-Dawley rats. Arch Biochem Biophys. 2017 Jan; 614:41-49. doi: 10.1016/j.abb.2016.12.008. Epub 2016 Dec 23. PMID: 28024836.

95) Lee TC, Ho IC, Lu WJ, Huang JD. Enhanced expression of multidrug resistance-associated protein 2 and reduced expression of aquaglyceroporin 3 in an arsenic-resistant human cell line. J Biol Chem. 2006 Jul 7;281(27):18401-07. doi: 10.1074/jbc.M601266200. Epub 2006 May 3. Erratum in: J Biol Chem. 2007 Apr 13;282(15):11612. PMID: 16672223.

96) Simon FR, Iwahashi M, Hu LJ, et al. Hormonal regulation of hepatic multidrug resistance-associated protein 2 (Abcc2) primarily involves the pattern of growth hormone secretion. Am J Physiol Gastrointest Liver Physiol. 2006 Apr;290(4):G595-608. doi: 10.1152/ajpgi.00240.2005. PMID: 16537972

97) Liu Z, Shen J, Carbrey JM, Mukhopadhyay R, Agre P, Rosen BP. Arsenite transport by mammalian aquaglyceroporins AQP7 and AQP9. Proc Natl Acad Sci USA. 2002 Apr; 99(9): 6053-58. doi: 10.1073/pnas.092131899. Epub 2002 Apr 23. PMID: 11972053; PMCID: PMC122900.

98) Kinzi J, Grube M, Meyer Zu Schwabedissen HE. OATP2B1 - The underrated member of the organic anion transporting polypeptide family of drug transporters? Biochem Pharmacol. 2021 Jun;188:114534. doi: 10.1016/j.bcp.2021.114534. Epub 2021 Mar 29. PMID: 33794186.

99) Garbinski LD, Rosen BP, Chen J. Pathways of arsenic uptake and efflux. Environ Int. 2019 May;126:585-97. doi: 10.1016/j.envint.2019.02.058. Epub 2019 Mar 8. PMID: 30852446; PMCID: PMC6472914.

100) Ishibashi K, Imai M, Sasaki S. Cellular localization of aquaporin 7 in the rat kidney. Exp Nephrol. 2000 Jul-Oct;8(4-5):252-57. doi: 10.1159/000020676. PMID: 10940724.

101) Ho LL, Kench JG, Handelsman DJ, et al. Androgen regulation of multidrug resistance-associated protein 4 (MRP4/ABCC4) in prostate cancer. Prostate. 2008 Sep;68(13): 1421-29. doi: 10.1002/pros.20809. PMID: 18615486.

102) Veiras LC, Girardi ACC, Curry J, et al. Sexual Dimorphic Pattern of Renal Transporters and Electrolyte Homeostasis. J Am Soc Nephrol. 2017 Dec;28(12):3504-17. doi: 10.1681/ASN.2017030295. Epub 2017 Aug 3. PMID: 28774999; PMCID: PMC5698077.

103) van Aubel RAMH, Smeets PHE, Peters JGP, Bindels RJM, Russel FGM. The MRP4/ABCC4 gene encodes a novel apical organic anion transporter in human kidney proximal tubules: putative efflux pump for urinary cAMP and cGMP. J Am Soc Nephrol. 2002 Mar;13(3): 595-603. doi: 10.1681/ASN.V133595. PMID: 11856762.

104) Nakamura T, Yonezawa A, Hashimoto S, Katsura T, Inui K. Disruption of multidrug and toxin extrusion MATE1 potentiates cisplatin-induced nephrotoxicity. Biochem Pharmacol. 2010 Dec;80(11):1762-67. doi: 10.1016/j.bcp.2010.08.019. Epub 2010 Sep 9. PMID: 20813096.

105) Yonezawa A, Inui K. Importance of the multidrug and toxin extrusion MATE/SLC47A family to pharmacokinetics, pharmacodynamics/toxicodynamics and pharmacogenomics. Br J Pharmacol. 2011 Dec;164(7):1817-25. doi: 10.1111/j.1476-5381.2011.01394.x. PMID: 21457222; PMCID: PMC3246706.

106) Maher JM, Cheng X, Tanaka Y, Scheffer GL, Klaassen CD. Hormonal regulation of renal multidrug resistance-associated proteins 3 and 4 (Mrp3 and Mrp4) in mice. Biochem Pharmacol. 2006 May;71(10):1470-78. doi: 10.1016/j.bcp.2006.02.005. Epub 2006 Mar 9. PMID: 16529719.

13. Supplementary Material/Appendices

Appendix 1: Age structure of study population by sex. Female and male study groups were similar with respect to age structure. n=5599 samples ≤ 500 mcg/L uAs_{UC} (from a total of 5608 samples) were analyzed, and 9 outliers with uAs_{UC} >500 (548 – 5763 mcg/L) were precluded. A Mann-Whitney U Test did reveal no significant age difference between the female and male groups.

Appendix 2: Urinary creatinine by sex. Female patients have significantly lower uCR than men ($p < 0.001$, Mann-Whitney U Test).

Appendix 3: Uncorrected urinary Arsenic (uAs_{UC}) by sex. Female patients have significantly lower uAs than men ($p < 0.001$, Mann-Whitney U Test).

Appendix 4: Conventionally CR-corrected urinary Arsenic (uAs) by sex. Female patients have significantly higher corrected uAs than men (p = 0.014, Mann-Whitney U Test).

a) Absolute deviation of uAs_C from uAs_N (µg uAs/g uCR)		
uCR range	0.3-3 g/L	All uCR
$|\Delta|$(µg/g $_{uCR}$); $\bar{x} \pm$ SD	1.9 ± 2.5	2.7 ± 3.8
Minimum $|\Delta|$	0	0
Maximum $|\Delta|$	33.3	56.9

b) Percentual deviation of uAs_C from uAs_N (%)		
uCR range	0.3-3 g/L	All uCR
$|\Delta\%|$ ($\bar{x} \pm$ SD)	11.9 ± 10.1	16.8 ± 16.4
Minimum $|\Delta\%|$	0	0
Maximum $|\Delta\%|$	271.8	271.8

c) uAs_C values deviating from uAs_N by more than 20% or 30%				
uCR range	0.3-3 g/L uCR		All uCR	
n total	4781		5599	
Limit	20%	30%	20%	30%
n > Limit	814	223	1603	873
% n > Limit	17.0	4.7	28.6	15.6

Appendix 5: Distribution of absolute and percentual deviations of uAs_N from uAs_C. All samples with < 500 µg uAs_{UC}. a) Absolute differences were calculated for all by $|\Delta| = |uAs_N - uAs_C|$. Overlapping histograms are plotted for the full range of uCR (dark green, n = 5599) and after uCR-restriction adopting the usual exclusion limits, i.e., all samples within the range from 0.3 - 3 g/L uCR (light green, n = 4781). b) Percentual deviations were calculated by $|\Delta\%| = |\Delta| \times 100/uAs_{UC}$. They are either plotted as complete or uCR-restricted dataset analog to histogram a. c) Absolute numbers and percentages of samples with uAs_c deviating more than either 20% or 30% from uAs_N.

Multiple linear regression model	Unstandardized Coefficients	Standardized Coefficients				95% confidence interval for B	
	B	Beta	SE	t	p	Lower bound	Upper bound
Constant	6.52		0.2	32.72	<.001	6.13	6.91
LN uCR (g/L)	-24.7	-0.9	0.11	-222.21	<.001	-24.92	-24.48
LN uAsUC (µg/L)	-2.07	-0.11	0.07	-28.28	<.001	-2.21	-1.92

ANOVA / Model summary	df	F	p	R	R²	SE of the estimate
	2	37,271.71	<.001	0.96	0.93	5.65

Appendix 6: Multilinear regression analysis of contributions of uCR and uAs$_{UC}$ to %CRCE.
ANOVA established significance. Standardized regression beta coefficients (β) were calculated for uAs$_{UC}$ and uCR. Formally, multiple-linear regression is not quite appropriate here because %CRCE, as the dependent variable, is not unrelated to the independent variables uCR and uAs. Also, all three variables are not normally distributed. Despite these limitations, the beta coefficient, which is significantly higher for uCR in all 3 cases, signals a stronger influence of uCR than uAs$_{UC}$ on %CRCE.

Appendix 7a: Enlarged illustration of uCR/uAs-relations in sextiles 1-5 for uAs$_{UC}$, uAs$_C$, and uAs$_N$.
Section of Fig. 2b-d with ordinate truncated to 40.

Appendix 7b: Compare quartiles of uAs$_{UC}$, uAs$_C$, and uAs$_N$ for S-PFCRC (c) and V-PFCRC (d).
Preliminary analysis of all n=298 samples from in-house patients with uAs$_{UC}$ < 100 µg/L.
S-PFCRC (c) still exhibits stronger residual dependencies (high R^2 and exponent b) in the marginal quartiles
1 and 4, which is resolved by V-PFCRC (d).

Appendix 8a: The uCR-differential linear relationship between coefficient *a* and uAs$_{UC}$.
Data were apportioned in 8 uCR-ranges and 5 uAsUC-quintiles and analyzed by power regression analog to
Figs.2a/b, yielding 5 curves of the form uAs$_{UC}$ = a × uCR [b]. Linear relation between uAs$_{UC}$ (mean of respective
quintile) and *a* of the corresponding curve were established by linear regression for all 8 uCR ranges (uCR-
means in the legend of App.8b). The gradual decrease of line slopes with urinary concentrations (uCR)
signifies uAs$_N$ (= *a*) to take on smaller values than uAs$_{UC}$ in samples with uCR > 1 g/L and higher than uAs$_{UC}$
with uCR < 1 g/L.

Appendix 8b: The uCR-differential logarithmical relationship between exponent *b* and uAs$_{UC}$.
The positive logarithmic dependence of exponent *b* on uAs$_{UC}$ is illustrated by the 5 power equations of
App.8a for all 8 uCR. Higher renal As-loads may raise the fractional excretion rate of uAs, thus converging it
with the fractional excretion of uCR, which under ordinary conditions is not reabsorbed at all. The
consequences are more negligible CRCEs at higher uAs$_{UC}$ due to the approximation of *b* to the value of 1. In
addition to the uAs$_{UC}$-dependent increment of exponent *b*, a negative dependence of the curve slopes on
uCR was observed. In all eight uCR ranges, the curves become gradually steeper with urinary dilution,
indicating that closer to linear relations between uCR and uAS$_{UC}$ are attained already at lower uAs$_{UC}$.

Appendix 9a: Effect of logarithmization on data distribution of uCR. Histograms and Quantile-Quantile plots of all n=5599 samples with uAs$_{UC}$ ≤ 500 μg/L are given in linear (left) and logarithmized (right) form for uCR (a), uAs$_{UC}$ (b), uAs (c).

C   uAs$_C$

Linear ——— | ——— Logarithmized

Appendix 9b-d: Effect of logarithmization on data distribution of uAs. Histograms and Quantile-Quantile plots of all n=5599 samples with uAs$_{UC}$ ≤ 500 µg/L are given in linear (left) and logarithmized (right) form for uCR(a), uAs$_{UC}$(b), uAs(c).