1   **Title: The genetic architecture of human cerebellar morphology supports a key role for the**
2   **cerebellum in human evolution and psychopathology**
3
4   **Short title: Genetic architecture of cerebellar morphology**
5

6   **Authors:** Torgeir Moberget*[1,2], Dennis van der Meer†[1,3], Shahram Bahrami†[1], Daniel Roelfs[1],
7   Oleksandr Frei[1], Tobias Kaufmann[1,4], Sara Fernandez-Cabello[1], Milin Kim[1], Thomas Wolfers[1,4],
8   Joern Diedrichsen[5], Olav B. Smeland[1], Alexey Shadrin[1], Anders Dale[6], Ole A. Andreassen[1] & Lars
9   T. Westlye[1,7].
10
11  *Corresponding author
12  †These two authors contributed equally
13
**Affiliations:**
1: Norwegian Centre for Mental Disorders Research (NORMENT), Division of Mental Health and
Addiction, Oslo University Hospital & Institute of Clinical Medicine, University of Oslo, Oslo,
Norway
2: Department of Behavioral Science, School of Health Sciences, Oslo Metropolitan University -
OsloMet, Oslo, Norway.
3: School of Mental Health and Neuroscience, Faculty of Health, Medicine and Life Sciences,
Maastricht University, The Netherlands
4: Department of Psychiatry and Psychotherapy, Tübingen Center for Mental Health, University of
Tübingen, Germany
5: Brain and Mind Institute, University of Western Ontario, London, Canada
6: Center for Multimodal Imaging and Genetics, University of California at San Diego, La Jolla, CA,
USA
7: Department of Psychology, Faculty of Social Sciences, University of Oslo, Norway

**Abstract current: 154 (limit: 160)**

The functional domain of the cerebellum has expanded beyond motor control to also include cognitive and affective functions. In line with this notion, cerebellar volume has increased over recent primate evolution, and cerebellar alterations have been linked to heritable mental disorders. To map the genetic architecture of human cerebellar morphology, we here studied a large imaging genetics sample from the UK Biobank (n discovery = 27,302; n replication: 11,264) with state-of-the art neuroimaging and biostatistics tools. Multivariate GWAS (MOSTest) on empirically derived regional cerebellar MRI features yielded 351 significant genetic loci (228 novel, 94% replicated). Candidate SNPs showed significant positive enrichment for relatively recent genetic mutations over the last 300k years (i.e., coinciding with the emergence of Homo sapiens), while gene level analyses revealed enrichment for genes associated with human-specific evolution over the last ~6-8 million years. Finally, we observed genetic overlap with major mental disorders, reinforcing the notion of cerebellar involvement in psychopathology.

**Teaser:** Genome-wide analysis of cerebellar morphology reveals links to recent human evolution and psychopathology

## Introduction

The cerebellum contains ~80% of all neurons in the human brain[1] and has rapidly expanded in volume over recent primate evolution[2]. Indeed, the surface area of the cerebellar cortex extends to almost 80% of the surface area of the cerebral cortex[3]. Comparative genetic analyses suggest that protein coding genes with known roles in cerebellar development have been subject to a similar, or even greater, rate of hominid evolution as compared to cerebro-cortical developmental genes[4]. Thus, the evolution of the cerebellum may have played a key role in the emergence of human cognition, including language[5].

A growing number of neuroimaging and clinical studies in humans also link cerebellar structure and function to a wide range of cognitive and affective functions[6-8], as well as to a number of heritable developmental[9] and psychiatric[10] disorders where these abilities either fail to develop properly or are compromised later in life. However, compared to supra-tentorial brain structures such as the cerebral cortex[11] and the hippocampus[12], few studies have mapped the genetic architecture of the cerebellum.

Of note, the few existing cerebellar genome-wide association studies (GWAS) have mostly been restricted to total cerebellar volume[13, 14], thus largely ignoring regional variation in cerebellar morphology. Importantly, such variation in the relative volumes of cerebellar subregions (i.e., variation in cerebellar shape independent of total cerebellar volume) has been associated with variation in behavioral repertoires in several species[15, 16], including domain-general cognition in primates[16].

We here performed a GWAS of MRI-derived regional cerebellar morphological features in a large population-based sample from the UK biobank (n discovery = 27,302; n replication:11,264), functionally characterized the genetic signal, tested for enrichment of SNPs and genes linked to human evolution, and assessed genetic overlap with major mental disorders.

## Results

**Data-driven decomposition of cerebellar grey matter maps reveals highly reproducible morphological features.**

Since traditional atlases of the cerebellar cortex based on gross anatomical landmarks (i.e., lobules) only partially overlap with more recent functional parcellations of the cerebellum[17-25], we first used a data-driven approach (non-negative matrix factorization, NNMF[26-28]) to parcellate voxel-based morphometry (VBM) based maps of cerebellar grey matter volume from 28,212 participants into robustly reproducible sub-regions. NNMF yielded highly reproducible decompositions (across split-half datasets) for model orders (i.e., number of components specified) ranging from 2 to 100 (see Online Methods for details about the study sample, MRI processing and quality control procedures). After observing that the improved fit to the original data seen with higher model orders tended to

112 level off between 15 and 30 components (indicating that the intrinsic dimensionality of the data might
113 have been reached), we decided on a model order of 23 based on its good split-half reproducibility
114 (see Fig. 1B, Online Methods for details regarding model order selection, as well as Supplementary
115 Figure 1 for summary maps of all tested model orders).

*Figure 1: Data driven decomposition of cerebellar grey matter maps yields highly reproducible and moderately heritable morphological features. **A**: Binarized winner-takes-all map for the 23-component solution based on data-driven decomposition of cerebellar grey matter maps from 28,212 participants. Note that empirically derived boundaries between cerebellar regions only partially follow traditional lobular borders (marked with dotted black lines); **B:** Five distinct components overlapping cerebellar Crus I derived from the split-half reliability analyses. While one of these components emerged as bilateral in Split 2, the remaining four components were almost identical, despite being derived from two independent samples; **C:** Narrow sense (SNP-based) heritability of the 23 components (see Supplementary Data 5 for numerical values). **D:** Hierarchical clustering of the 23 components (derived from the full sample decomposition) based on their pairwise genetic correlations revealed a primary division between the anterior and posterior cerebellum, with additional separations between medial and lateral regions. The full genetic correlation matrix can be found in Supplementary Data 6.*

Of note, our data-driven decomposition differed markedly from the standard cerebellar atlas based
on gross anatomical features, shown as dotted lines in Fig 1A (see Supplementary Figure 2 for all
23 components and Supplementary Data 1 for quantification of overlap between NNMF-derived
components and standard cerebellar anatomical regions, i.e. lobules). For instance, five distinct
components overlapped Crus I of cerebellar lobule VII (shown in Fig 1B), an anatomical region which
already started to split into separate components at a model order of three. We further observed
only partial overlap with task-based functional parcellations of the cerebellar cortex (Supplementary
Data 2). While some components clearly overlapped cerebellar regions previously associated with
hand movements, eye-movements/saccades or autobiographical recall, other data-driven
components overlapped multiple functionally defined cerebellar regions (Supplementary Data 2).

**Cerebellar morphological features are heritable and reveals a distinct anterior-posterior pattern based on their bivariate genetic correlations.**

After removal of one of each genetically related pair of individuals (n = 910), 27,302 participants remained for the genetic analyses. In addition to the 23 regional cerebellar morphometric features of primary interest, we also included total cerebellar volume, estimated total intracranial volume and 9 cerebral brain phenotypes to serve as covariates and/or comparison phenotypes. Prior to the genetic analyses, all morphological features were adjusted for effects of scanner site, sex, age, estimated total intracranial volume, 40 genetic population components, genetic analysis batch and a quantitative structural MRI quality index (the Euler number(*29*)) using general additive models, before finally being rank-order normalized (see Online Methods for details).

To validate our analysis approach, we computed genetic correlations (using LD-score regression, LDSC(*30*)) between univariate GWAS results on the comparison features and previously neuroimaging GWAS studies. Results showed a mean $r_g$ of .90 (range: .80-.99, see Supplementary Data 3). For the 23 cerebellar morphological features, we computed genetic correlations between discovery (n = 27,302) and replication (n = 11,264) samples using LDSC(*30*). These genetic correlations were high (mean rG: .92; range: .83-1), indicating reliable genetic signals (see Supplementary Data 4).

Genetic complex trait analysis (GCTA(*31*)) revealed SNP-based heritability estimates ($h^2$) ranging from .33 to .44 (Figure 1C and Supplementary Data 5). Analyses of total cerebellar volume ($h^2$ = .35), estimated total intracranial volume ($h^2$ = .41) and the 9 cerebral comparison phenotypes ($h^2$ range: .26 to .45) gave a similar range of heritability estimates (Supplementary Data 5).

Hierarchical clustering of cerebellar features based on their bivariate genetic correlation matrix (GCTA-based rG ranging from 0.35 to 0.98, Supplementary Data 6) revealed a primary anterior-posterior division running along the horizontal fissure separating Crus I and Crus II, with secondary divisions grouping features into more lateral or medial, as well as more anterior and posterior features within the major anterior and posterior regions (Fig 1D). Of note, the primary division along the horizontal fissure was also evident from the (genetically naïve) two-component NNMF decomposition, while medial-to-lateral divisions already began to emerge with a model order of three (see Supplementary Figure 1).

In order to examine whether this phenotypic and genetic correlation structure was also reflected in regional cerebellar gene expression patterns, we used the abagen toolbox(*32*) to extract Allen Human Brain Atlas(*33*) gene expression profiles for 22 of the 23 morphological features and computed their bivariate Pearson correlations (across 15,631 genes; Supplementary Data 7) and hierarchical clustering solution (Supplementary Figure 4). The anterior-posterior boundary across the horizontal fissure was also evident in the gene expression data, which in addition highlighted distinct gene expression patterns for the posterior midline (grouped together with the horizontal fissure), as well as for the most lateral regions of the cerebellar cortex.

**Multivariate GWAS reveals 351 genetic loci associated with cerebellar morphology.** Figure 2A shows the main results for the multivariate GWAS across the 23 cerebellar morphological features applying MOSTest(*34*). We observed 35,098 genome-wide significant (GWS) SNPs, which were mapped by FUMA(*35*) to a total of 51,803 candidate SNPs by adding reference panel SNPs in high LD (r >.6) with GWS SNPs. The 51,803 candidate SNPs (see Supplementary Data 8) were represented by 1,936 independent significant SNPs and 560 lead SNPs in 351 genomic loci (see Supplementary Data 10). The QQ-plot of the MOSTest results did not suggest inflation, while the QQ-plot of results based on permuted data (under the null hypothesis) confirmed the validity of the MOSTest analytical approach (Supplementary Figure 5).

*Figure 2: Multivariate GWA analysis of the 23 cerebellar morphological features revealed 351 independent genome-wide significant (GWS) loci. A: The upper half of the Miami plot shows the main results from the multivariate analysis. The lower half displays results from a series of 23 univariate analyses (corrected for multiple comparisons using the standard min-P approach), as well as results from a univariate analysis of total cerebellar grey matter (marked in orange). B-D: Euler diagrams showing the relative numbers of - and overlaps between - candidate SNPs mapped by the three analysis approaches employed in the current study (B), the current and four recent studies*

*reporting genetic associations with cerebellar morphology (C), as well as results from multivariate*
*GWASs on hippocampal and cerebrocortical morphology (D).*

Annotation of all candidate SNPs using ANNOVAR(*36*) as implemented in FUMA[35] revealed that the
majority of candidate SNPs were intronic (57.8%) or intergenic (38.3%). While only 0.7% were
exonic, about 81% of the candidate SNPs were assigned minimal chromatin states between 1 and
7 (i.e., open chromatin states), implying effects on active transcription[37] (see Supplementary Data 8-
9).

We evaluated the robustness of these multivariate results using a multivariate replication
procedure established in Loughnan et al(*37*), which computes a composite score from the mass-
univariate *z*-statistics (i.e., applying multivariate weights from the discovery sample to the replication
sample input data) and then tests for associations between this composite score and genotypes in
the replication sample (for mathematical formulation see Loughnan et al(*37*)). Results showed that
97% of loci lead SNPs present in both samples replicated at a nominal significance threshold of p <
.05 (Supplementary Figure 6 and Supplementary Data 8), and that 74% remained significant after
Bonferroni correction for the 339 replication tests conducted (Supplementary Data 10). Moreover,
99% of loci lead SNPs showed the same effect direction across discovery and replication samples
(Supplementary Figure 6 and Supplementary Data 10). Thus, 329 (94%) of the 351 reported loci
were replicated.

In addition, we assessed the robustness of the multivariate patterns by computing bivariate
correlations between feature z-score vectors assigned to the discovery sample lead SNPs in an
independent multivariate GWAS (MOSTest) performed on the replication sample. These correlations
(restricted to the 339 loci lead SNPs present in both samples) were relatively high (mean r: .70, see
Supplementary Data 10 and Supplementary Figure 7). Figure 3 and Supplementary Figure 7 give
some examples of discovery and replication sample multivariate patterns projected back onto the
cerebellar cortex.

To compare our main multivariate MOSTest results to univariate approaches, the lower part
of the Miami plot in Figure 2A, and Figure 2B, displays results from a set of univariate GWASs on
the cerebellar morphological features (which yielded 8370 candidate SNPs and 57 genomic loci,
corrected for multiple comparisons using the min-P approach(*38, 39*)), as well as the 4044 candidate
SNPs and 10 significant loci resulting from the univariate GWAS on total cerebellar grey matter
volume (marked in orange). 52 of the 57 loci identified in the univariate analyses of regional
cerebellar features overlapped 55 of the 351 loci identified using the multivariate method, while 9 out
of 10 loci identified in the univariate analysis of total cerebellar volume were also identified in the
multivariate analysis. Thus, our multivariate analysis of regional cerebellar features increased the
locus yield ~35-fold relative to analyzing total cerebellar volume and ~6-fold relative to performing a
set of univariate analyses on the same regional features.

We next compared the current findings with previously reported genetic loci for cerebellar
morphology by extracting summary statistics from two recent GWAS studies using the UKBB sample
(n = 19k(*40*) and 33k(*41*)) that included regional cerebellar volumes among the full set of analyzed
brain imaging-derived phenotypes (101(*40*) and 3,144(*41*), respectively), as well as two recent
GWASs on total cerebellar volume (n = 33k(*13*) and 27k(*14*), respectively). Candidate SNPs and
independent GWS loci were identified in FUMA using the same settings as for our primary analyses
and employing a liberal *p*-value threshold of 5e-8 (i.e., not correcting for the total number of brain
imaging features analyzed). Results are displayed in Figure 2c and Supplementary Data 8 and 10.
In brief, we found that 19,527 of the 51,803 candidate SNP (i.e., 36%) and 123 of the 351 identified
genomic loci (i.e., 35%) overlapped with candidate SNPs and loci extracted from these three
previous studies. Thus, 228 of the 351 (i.e., 65%) genetic loci reported here are novel to the literature
on of cerebellar morphology genetics (see Supplementary Data 10.

Overlap of cerebellar candidate SNPs and genetic loci with results from recent multivariate
analyses of hippocampal and cerebrocortical morphology are displayed in Figure 2D and
Supplementary Data 8 and 10 (final columns). Of note, we found that 32 and 29 percent of the
candidate SNPs discovered here for the cerebellum overlapped with candidate SNPs for vertex-wise
cerebrocortical surface area and thickness(*42*), respectively, while 11.4 percent overlapped with
candidate SNPs found for hippopcampal subregions(*43*). 95 of the 351 genetic loci overlapped loci
linked to the other multivariate brain phenotypes (Supplementary Data 10). Thus, 64% of the
candidate SNPs and 73% of genetic loci appeared to be specifically associated with cerebellar
morphology.

**Significant genetic variants show heterogeneous effects across the cerebellar cortex,**
**influencing both regional and total volumes.**

A major advantage of our multivariate analysis approach is its sensitivity to both highly localized and
more generally distributed effects of SNPs on cerebellar morphology. This is illustrated in Figure 3,
which displays the 351 loci lead SNPs as a function of both the most extreme individual Z-score
across all cerebellar features (e.g, analogous to the strongest "local" effect) and of the mean Z-score
across these features (i.e. analogous to the main effect on overall cerebellar volume).

Method    ● MOSTest and minP    ● MOSTest only    ● MOSTest, minP and total cerebellum

**Figure 3: Lead SNPs show spatially heterogeneous – and replicable - effects across the cerebellar cortex.** *The 351 loci lead SNPs identified by MOSTest are plotted as a function of main overall effect across all cerebellar features (x-axis: mean Z-score) and most extreme effect for any single cerebellar feature (y-axis: most extreme Z-score across features), and colour coded by SNP discovery method. The cerebellar flat-maps show discovery (left) and replication (right) sample regional distributions of Z-scores (color-scale range from -10 to 10) for a few selected lead SNPs (see Supplementary Data 10 for individual feature Z-scores for all 351 discovery sample loci lead SNPs). SNP rs7877685 was only present in the discovery sample.*

As can be seen, some lead SNPs (e.g. rs13107325; rs76934732) show pronounced positive or negative mean z-scores, indicating a relatively consistent direction of effect across cerebellar features. See also inset figures displaying feature Z-scores projected back onto the cerebellar cortex. Many of these SNPs also emerged in the univariate analysis of total cerebellar volume and were recently reported in GWASs on total cerebellar volume(*13, 14*).

Many other lead SNPs, however, show strong "local" signals with opposite effect directions across features, yielding very weak global signals (e.g. rs117332043; rs78777685). Thus, while several of the most significant SNPs in this category have previously been reported in GWASs including local cerebellar morphological features(*40, 41*), they did not emerge from analyses of total cerebellar volume, neither in the current nor in previous studies.

Finally, our multivariate MOSTest approach is also sensitive to SNPs displaying weaker
effects distributed across several cerebellar features, often with opposite effect directions (e.g.
rs12464825; rs2388334). For this category of SNPs, neither of the univariate methods have sufficient
power at the current sample size. In contrast, these two example SNPs robustly emerged from the
multivariate analysis (discovery sample p-values < 1e-56; replication sample p-values <1e-15).

**Genetic variants associated with cerebellar morphology are enriched for evolutionary recent**
**mutations in the human genome**
We next mapped the evolutionary age of cerebellar candidate SNPs by merging them with a recently
published dataset on dated mutations in the human genome(*44*). Following the analysis procedure
established by Libedinsky et al.(*45*), we plotted the histogram of dated candidate SNPs over the last
2 million years in bins of 20.000 years (Figure 4A and Supplementary Data 11), and tested for
positive or negative enrichment by comparing them to empirical null distributions derived from 10,000
randomly drawn and equally sized sets of all SNPs in the full human genome dating dataset (after
matching them to brain-related SNPs in terms of minor allele frequency; see Online Methods for
details).

*Figure 4: Genetic variants associated with cerebellar morphology are enriched for*
*evolutionary recent mutations in the human genome. A: Histogram of estimated SNP age*
*(ranging from 0 to 2 million years, in bins on 20,000 years) for 41,549 candidate SNPS associated*
*with cerebellar morphology. The solid black line and grey ribbon denote the mean and Bonferroni-*
*corrected 95% confidence intervals derived from a null model constructed from 10,000 of equally*
*sized sets of SNPs randomly drawn from the Human Genome Dating Atlas of Variant Age (after*
*matching these to brain-related SNPs in terms of minor allele frequencies). Red bars denote time-*
*bins of significant positive enrichment. B: Enrichment Z-scores (calculated from means and standard*
*deviations of the corresponding null models) for candidate SNPs associated with cerebellar,*
*hippocampal and cerebrocortical surface area and cortical thickness morphology. Time bins with*
*significant positive enrichment are highlighted and marked with black borders, while circle size reflect*
*z-score (see Supplementary Data 11-12 for full numerical results).*

Results revealed two main peaks of positive enrichment, the most prominent ranging from 260-300k

years ago (enrichment z-scores ranging from 23.7-28.6; see Figure 4B and Supplementary Data 11)

, i.e., coinciding with the emergence of anatomically modern humans (*46, 47*). The second strongest

enrichment peak (z-score range: 6.1-12.5) ranged from 40-120k years ago, i.e., overlapping the

human migration out of Africa, as well as the first evidence of several uniquely human behaviors

(often referred to as behavioral modernity), such as tool-making, recording of information onto

objects and purposeful burials(*48*).

For comparison, Figure 4B also shows results from analyses on candidate SNPs identified

in previous multivariate GWAS studies of hippocampal(*43*) and cerebrocortical(*42*) morphology.

These brain features also showed significant enrichment peaks, partially overlapping the cerebellar

results. Specifically, SNPs associated with hippocampal morphology also displayed the most

prominent enrichment peak between 260 and 300k years ago, while SNPs linked to cerebrocortical

morphology (surface area and thickness) tended to show the most significant enrichment for more

recent time bins (i.e., 40-120k years ago, see full results in Supplementary Data 12).

**Genes associated with cerebellar morphology show selective expression in cerebellar and**

**prenatal brain tissue, as well as enrichment for genes linked to human accelerated regions.**

To functionally characterize the multivariate GWAS signal, we mapped the full set of GWAS p-values

to 19,329 protein coding genes using MAGMA(*49*) and used the resulting gene-level p-values to test

for 1) GWS genes, 2) gene expression in brain tissue; and 3) enrichment for genes linked to human

accelerated regions (HARs),  i.e. sections of DNA that have remained relatively conserved

throughout mammalian evolution, before being subject to  a burst of changes in humans since

divergence of humans from chimpanzees(*50, 51*). These analyses yielded a total of 534 GWS genes

(i.e., 2.78% of all protein coding genes, see Figure 5A and Supplementary Data 13). Using the full

set of 19,329 gene-level p-values in MAGMA gene property analyses revealed significant and

specific gene expression in cerebellar and prenatal brain tissue (Figure 5B-C and Supplementary

Data 14-15), with the selective cerebellar expression seen in two independent datasets (Allen

Human Brain Atlas(*33*) and The Genotype-Tissue Expression (GTEx) Project).

The MAGMA gene set analysis of HAR-linked genes revealed significant enrichment (*p* =

7.09e-08) for genes associated with cerebellar morphology (Figure 5D, Supplementary Data 16). Of

note, running this same HAR gene set analysis on summary statistics from recent multivariate GWAS

studies on cerebrocortical(*42*) or hippocampal(*43*) morphological features yielded similar (for cortical

features) or significantly weaker (hippocampal features) enrichment effects (Figure 5d, see Online

Methods for processing pipeline).

*Figure 5: Gene mapping reveals selective enrichment for cerebellar and prenatal brain tissue, as well as for genes linked to human accelerated regions. A: MAGMA mapped the full range of SNPs from the multivariate GWAS to 19.329 protein-coding genes, 534 of which were genome-wide significant (Bonferroni corrected threshold marked with red line), with the 30 most significant genes labeled in the gene-based Manhattan plot. B-C: MAGMA gene-property analyses revealed selective brain expression in the cerebellum and during prenatal developmental stages. D: MAGMA gene-set analysis revealed significant enrichment for sets of genes previously linked to human accelerated regions (HARs). Figure 4d also shows results from comparative analyses using summary statistics from other multivariate GWASs of MRI-based brain, as well as significant results from statistical tests comparing beta-weights for HAR-linked genes across multivariate brain features. Horizontal red lines mark the Bonferroni-corrected significance threshold for each subplot. See Supplementary Data 13-16 for full results.*

**Genes linked to human cerebellar morphology show enrichment for gene sets linked to selective cerebellar gene expression, altered cerebellar morphology in mouse models, human clinical/anthropometric traits, as well as specific biological processes and pathways.** In line with the continuous brain tissue gene expression results described above, we also observed significant and relatively selective enrichment for smaller curated sets of genes previously found to be highly and selectively expressed in mouse (Figure 6A, Supplementary Data 17) and human (Figure 6C, Supplementary Data 19) cerebellar brain tissue, as well as for sets of genes previously

shown to affect cerebellar morphology in mouse gene perturbation experiments (Figure 6B,
Supplementary Data 18) and various clinical conditions and anthropometric traits in humans (Figure
6D, Supplementary Data 20). The most significant gene ontology and curated gene sets from the
MSigDB(*52, 53*) database were related to brain development (e.g., neurogenesis, axon guidance
and neuron differentiation, Figure 6E, Supplementary Data 21), and highlighted the reelin signaling
pathway (Figure 6F, Supplementary Data 22).

**Figure 6: Gene mapping reveals selective enrichment across brain tissues and curated gene
sets.** *MAGMA gene-set analyses revealed significant enrichment for sets of genes previously linked
to preferential expression in human **(A)** and mouse **(C)** cerebellar tissue, as well as effects on
cerebellar morphology (and other brain phenotypes) in mouse gene perturbation experiments **(B)**
and human clinical disorders and phenotypes **(D)**. Significant gene ontology terms were related to
neural development **(E)**, while curated gene sets highlighted the Reelin signaling pathway **(F)**.
Across all subplots the x-axis shows the -log10 p-value, the y-axis marks the number of gene sets,
and the top 10 most significant gene-sets are labelled. Red lines mark the Bonferroni-corrected
significance threshold for each subplot. See Supplementary Data 17-22 for full results.*

388    As can be seen in Figure 5A (and Supplementary Data 13), *RELN* (encoding the protein Reelin) was
389    also the most significant gene mapped by MAGMA. See also Supplementary Figure 8 for a regional
390    locus plot showing the 12 lead SNPs mapped to *RELN* and their associated z-score maps.
391
**Gene mapping reveals sets of plausible causal genes**
In addition to the gene-based mapping strategy using all SNPs described above (MAGMA), we also
mapped candidate SNPs to genes using two complementary gene mapping strategies: 1) positional
mapping of deleterious SNPs (defined as having a CADD-score(*54*) > 12.37); and 2) mapping of
SNPs previously shown to affect gene expression in cerebellar tissues (i.e., eQTL mapping). Across
all three strategies, we mapped a total of 674 unique genes; 531 using MAGMA, 310 using positional
and 197 using eQTL mapping (see Figure 7A and Supplementary Data 2~~3~~). 298 genes were
identified by at least two strategies, while 65 genes were mapped by all three strategies. Out of these
674 genes, 61 have previously been associated with cerebellar pathology in humans and/or altered
cerebellar morphology in mouse gene perturbation experiments, while 121 have been linked to
human accelerated regions (Supplementary Data 23). As can be seen in Figure 7B and
Supplementary Data 23, the 674 genes mapped to cerebellar morphology showed some overlap
with genes mapped to hippocampal(*43*) and cerebrocortical(*42*) morphology using the same
mapping strategies, but 264 genes (39%) appeared relatively specific to the cerebellum.
Results from gene set analyses on the 674 mapped genes (Figure 7, Supplementary Data
24-28) largely mirrored results from the MAGMA analyses described above, but in addition revealed
that this set of 674 mapped genes was also enriched for gene sets associated with several complex
clinical phenotypes and anthropometric traits in humans, including cognitive ability, neuroticism and
schizophrenia (Figure 7F, Supplementary Data 28).
Of note, while the full set of mapped genes showed significant enrichment for sets of genes
known to alter cerebellar morphology in mouse mutation or knock-down experiments
(Supplementary Data 26), we also note that 613 of the 674 mapped genes have not to our knowledge
previously been linked to cerebellar development, anatomy or pathology in mice or humans
(Supplementary Data 23), and thus constitute potential targets for future gene perturbation
experiments in animal models.
Restricting the above analyses to the 298 genes mapped across at least two strategies did
not markedly affect the results (Supplementary Data 24-28, final columns).
419
420
421
422

**Figure 7: Mapping and functional characterization of plausible causal genes.** *The 674 plausible causal genes mapped using three complementary strategies (A) show partial overlap with genes mapped to hippocampal(43) and cerebrocortical(42) morphology (B); significant enrichment for sets of genes linked to human accelerated regions (HAR) (C); selective expression in the cerebellum in mice (D) and humans (E), as well as effects on cerebellar morphology in mouse gene perturbation studies (F) and human disorders and anthropometric phenotypes (G). See Supplementary Data 23-28 for full results.*

**Cerebellar morphology shows significant genetic overlap with psychiatric disorders**

We finally tested for overlap between the multivariate genetic profile for cerebellar morphology and genetic profiles for five major developmental/psychiatric disorders (attention deficit hyperactivity disorder: ADHD; autism spectrum disorder: ASD; bipolar disorder: BIP; major depressive disorder:

MDD; and schizophrenia: SCZ) using conditional/conjunctional FDR analysis(*55*). As can be seen
in the conditional QQ-plots in Figure 8, these analyses revealed clear patterns of enriched
association with the clinical phenotypes when selecting subsets of SNPs with increasingly stronger
association with cerebellar morphology (Fig. 8A and figure insets in (C-F).

*Figure 8: Genetic variants influencing cerebellar morphology overlap with variants*
*associated with five major mental disorders. Conditional QQ plots (A and figure insets in C-F)*
*show an incremental incidence of association with five mental disorders (leftward deflection) as a*
*function of the significance of association with cerebellar morphology. Manhattan plots show SNPs*
*with significant association with both traits, thresholded at a conjunctional FDR threshold of p > .01*
*(red dotted line). SCZ: Schizophrenia; BIP: Bipolar Disorder; MDD: Major Depressive Disorder;*
*ADHD: Attention Deficit Hyperactivity Disorder; ASD: Autism Spectrum Disorders.*

See also Supplementary Figure 9 for QQ-plots depicting the reverse association, i.e., enriched association with cerebellar morphology when conditioning on the association with developmental/psychiatric disorders. Specific genetic variants jointly influencing the two phenotypes were identified using conjunctional FDR analyses at a conservative statistical threshold of p<.01. Results revealed shared genetic loci with all disorders; namely 48 with SCZ, 22 with BIP, 2 with MDD, 5 with ADHD and 5 with ASD (Figure 6; Supplementary Data 29-33). We mapped lead and candidate SNPs for each of these loci to genes using positional and eQTL mapping and checked for gene overlap across disorders (Supplementary Data 34). Of note, the *LRP8* gene (a HAR-linked gene(*50, 51*) encoding a Reelin receptor) emerged from the conjunctional FDR analyses of both BIP and SCZ, thus again highlighting the Reelin signaling pathway.

To complement these multivariate analyses, we also conducted a set of univariate analyses, using LD-score regression(*30*) to calculate the genetic correlations between each individual cerebellar feature and the five developmental/psychiatric disorders.

*Figure 9: Univariate genetic correlations between cerebellar morphological features and five major mental disorders are negative. The top row display unthresholded genetic correlations, while these are filtered at increasingly strict statistical thresholds in the following rows, i.e. nominal p-value < .05 (second row); Bonferroni correction for the 23 cerebellar features tested (third row); and Bonferroni correction for both 23 features and 5 clinical conditions (bottom row). Black dotted lines denote lobular boundaries.*

As can be seen in the first row of Figure 9 (and Supplementary Data 35), genetic correlations with
cerebellar morphological features were predominantly negative across diagnoses, indicating that
genetic variants associated with a clinical diagnosis tended to also be associated with reduced
cerebellar volumes (with 42 out of 115 tested associations showing nominally significant negative
correlations; Fig 9, second row). However, all univariate genetic correlations were relatively weak,
and only a few negative genetic correlations with BIP and SCZ survived Bonferroni correction for the
23 features tested. When also correcting for the five clinical conditions, only the negative correlation
between BIP and cerebellar feature 23 (primarily overlapping vermal lobules VIIIa and VIIIb)
remained significant.

Supplementary Data 36 show genetic correlations between the five disorders and the ten
comparison brain phenotypes, as well as total cerebellar volume. In general agreement with the
observed pattern for regional cerebellar features, total cerebellar grey matter volume showed
nominally significant negative genetic correlations with BIP (rg: -.10; p = 0.0052) and SCZ (rg: -.09;
p = 0.0133). Pallidal volume also showed nominally significant negative genetic correlations with
these two disorders (BIP: rg: -.10; p = 0.0087; SCZ: rg: -.08; p = 0.0172, while ADHD displayed
negative genetic correlations with estimated total intracranial volume (rg: -.15; p = 0.0003) and total
cortical surface area (rg:-.14 ; p = 0.0065), as well as a positive genetic correlation with hippocampal
volume (rg: .13; p = 0.0164). Finally, MDD showed nominally significant negative genetic correlations
with volumes of the hippocampus (rg -.08: p = 0.025) and thalamus (rg: -.09; p = 0.0258). Only the
negative genetic correlation between ADHD and estimated total intracranial volume survived
Bonferroni correction across the 55 tests performed.

492
493
494  **Discussion**
495
The current study identified novel features of the genetic architecture of cerebellar morphology,
supported the notion of recent changes over human evolution, implicated specific neurobiological
pathways, and demonstrated genetic overlap with major mental disorders.

With respect to the main features of cerebellar morphology, it is worth noting that results from
our data-driven decomposition of cerebellar grey matter maps (which was not informed by genetic
data), the genetic correlation analyses of the chosen 23-feature solution and gene expression data
from the Allen Human Brain Atlas all converge on a similar general pattern. The first boundary to
emerge in the data-driven decompositions ran along the horizontal fissure separating Crus I and II
of lobule VII, reflecting its centrality in characterizing phenotypic variability in cerebellar morphology
at the population level. Of note, this boundary also emerged from clustering of the 23 cerebellar
morphological features based on their bivariate genetic correlations or gene expression profiles
(based on the Allen Human Brain Atlas). This latter finding essentially mirrors results from a recent
report using a different analysis strategy on the same gene expression data(56). Interestingly, the
horizontal fissure has been suggested to mark the border between two separate cerebellar

representations of the cerebral cortex, with a possible third representation in lobules IX-X(*57*). The current cerebellar results thus complement previous work on the hierarchical genetic organization of the cerebral cortex, which has identified the Rolandic fissure (separating the frontal and parietal lobes) as a main boundary with respect to genetic effects of effects of surface area(*58*), as well as a superior-inferior gradient for genetic influences on cortical thickness(*59*)

Our multivariate GWAS using MOSTest identified 351 independent GWS loci associated with cerebellar morphology, increasing the yield ~35-fold relative to analyzing total cerebellar volume and ~6-fold relative to performing a set of univariate analyses on the same regional features in our current sample. 329 (94%) loci from the multivariate analyses were replicated in an independent sample, indicating robust results. After applying a liberal threshold to summary statistics from previous well-powered studies, we find that 228 (65%) of our reported loci are novel. Importantly, among the genes mapped to these novel loci we find several that are known to play important roles in cerebellar development in mice (e.g., *RORA*(*60*), *FGF8*(*61*) and *BAHRL1*(*62*), see Supplementary Data 22). While candidate SNPs associated with cerebellar morphology partially overlap with SNPs previously mapped to other multivariate brain phenotypes, we note that a substantial number of SNPs appear to be relatively selectively linked to cerebellar morphology, a finding that is in in line with the distinct gene expression profile found for the cerebellum(*33*).

SNP- and gene-level results from the current study also bolster – and refine – the notion of relatively recent changes in cerebellar morphology over human evolution(*2, 4, 15, 16*). Specifically, we found that candidate SNPs associated with cerebellar morphology were particularly enriched for SNPs with an estimated age of 260-300 thousand years, i.e., coinciding with the emergence of Homo Sapiens(*46, 47*). Significant, but weaker, enrichment was also seen for time bins stretching from 40-120 thousand years ago, i.e., overlapping the human migration out of Africa, as well as the first evidence of several uniquely human behaviors (often referred to as behavioral modernity), such as tool-making, recording of information onto objects and purposeful burials(*48*). Similar patterns of enrichment was also found when analyzing candidate SNPs derived from previous studies on multivariate hippocampal(*43*) and cerebro-cortical(*42*) morphology Gene level analyses further showed that genes associated with inter-individual variation in cerebellar morphology are enriched for genes linked to human accelerated regions (HARs)(*50*) of the genome. HARs denote previously conserved regions of the genome that were subject to a burst of changes in humans after the divergence of humans from chimpanzees about 6-8 million years ago(*50*). Of note, a recent GWAS on total cerebellar volume found no enrichment for HAR-linked genes(*14*), suggesting that SNPs associated with regional cerebellar variation may be driving this effect. While the comparisons with cerebral features should be interpreted with care since different image preprocessing pipelines were employed, it is nonetheless striking that HAR gene enrichment was nominally stronger for cerebellar morphology than for vertex-wise cerebrocortical thickness and area, and significantly stronger than for hippocampal regional volumes (Figure 5D).

547     Together, these SNP- and gene-level results suggest that genetic variants influencing
548     cerebellar morphology in modern humans have been subject to selection over relatively recent
549     human evolution, and that cerebellar changes – in concert with other brain regions - may thus have
550     played a central part in the emergence of uniquely human cognitive abilities.

551                     Results from the MAGMA gene property and gene set analyses bolster our
confidence in the genetic signal by showing selective gene expression in human cerebellar brain
tissue across two independent datasets (Allen Human Brain Atlas and GTEx.8) and significant
enrichment for sets of genes which have previously been shown to affect cerebellar morphology in
mouse gene perturbation experiments. The MAGMA results further show significant enrichment for
sets of genes known to play key roles in neurodevelopment (e.g., neurogenesis & axon guidance)
and preferential expression in prenatal brain tissue, thus supporting a primarily developmental origin
of genetically determined effects on adult cerebellar morphology.Of note, our MAGMA enrichment
analysis of curated gene sets strongly implicated the Reelin signaling pathway. Indeed, both the
gene coding for the Reelin protein (*RELN*) and genes coding for its two receptors (*LRP8* & *VLDLR*)
were identified across at least two gene mapping strategies, with *RELN* emerging as the single most
significant gene by MAGMA. The Reelin pathway is known to play important roles in
neurodevelopment (e.g. neuronal migration), and mutations in the *RELN* and VLDLR(*63*) (and to a
lesser extent LRP8; also known as ApoER2(*64*)) have been associated with cerebellar
malformations and/or dysfunction. LRP8 is also among the genes linked to human accelerated
regions (HARs) of the genome.

The sets of genes mapped by the three complementary mapping strategies provide a
database for future studies investigating the genetic architecture of cerebellar morphology. For
instance, we mapped 616 genes associated with inter-individual variability in human cerebellar
morphology that have not yet to our knowledge been examined in mouse gene perturbation
experiments and/or associated with cerebellar pathology in humans. Among these, we highlight
*MAP2K5* and *GRB14*, two HAR-linked genes mapped across all strategies and associated with lead
SNP p-values <1e-50, but whose functions in the brain are largely unknown.

The reported results for previously discovered variants, loci and genes add important
information regarding regional effects on cerebellar morphology. For instance, while genetic variants
linked to the *RELN* gene have previously been associated with volumes of cerebellar vermal lobules
VI-X and hemispheric lobule IX(*40, 41*), we here mapped 12 lead SNPs to *RELN* showing
heterogeneous effects across the entire cerebellar cortex (but with peak effects overlapping
previously described midline and posterior cerebellar regions, see Supplementary Figure 8).

The observed genetic overlap between cerebellar morphology and the five mental disorders
reinforces the recent notion of the cerebellum as a key brain structure in complex clinical traits and
disorders(*6-10*). Across the five diagnoses, the strongest evidence for genetic overlap with cerebellar
morphology was found for SCZ and BIP, likely at least in part because these disorder GWASs were
relatively well-powered. While an in-depth discussion of genetic loci jointly influencing psychiatric

disorders and cerebellar morphology is beyond the scope of this report, we note that the Reelin
pathway again emerges in the genetic overlap analyses for SCZ and BIP. Specifically, the current
finding of *LRP8* (a reelin receptor, and HAR-linked gene) as one of the genes jointly associated with
cerebellar morphology and the aforementioned severe mental disorders points towards a potential
molecular pathway involved in the cerebellar abnormalities previously reported in SCZ(*10, 65, 66*)
and BIP(*65, 66*). Indeed, in line with its key importance for brain development and function, the
Reelin pathway is also increasingly seen as relevant for a wide range of neurodevelopmental,
psychiatric and neurodegenerative disorders(*67*). Of particular relevance to the current findings,
converging evidence supports LRP8 as a key susceptibility gene for psychosis(*68*).

The main limitations of the current study concern the ancestral homogeneity of the sample,
the sample size and the exclusion of very rare genetic variants. Limiting the sample to participants
of European ancestry was deemed necessary considering the current state of the multivariate
GWAS methods used but may limit the generalizability of our findings. Second, while the current
sample size is large in comparison with previous imaging genetics studies, it is still relatively small
in comparison to GWASs of other complex human phenotypes (e.g., intelligence, with a current n of
> 3 million(*69*)). Finally, very rare genetic variants (MAF < 0.005) were excluded from the current
multivariate GWAS, but are likely to include a number of variants with relatively large effect sizes on
complex human traits (*70*). Thus, future studies using larger and more diverse samples – as well as
whole exome and/or genome sequencing - are likely to discover more of the genetic variants
associated with cerebellar structure.

In conclusion, the current results enhance our understanding of the genetic architecture of
human cerebellar morphology, provide supporting evidence for cerebellar morphological changes
during the last ~6-8 million years of human evolution, and reinforce the notion of cerebellar
involvement in several mental disorders by demonstrating significant genetic overlap.

**Materials and Methods:**

**Participants**

For our main analyses, T1-weighted MR images, demographic and genetic data from 39,178 UK Biobank participants were accessed using access number 27412. After removing 1043 participants who either had missing genetic data or had withdrawn consent (as of 19.11.2019), data from 38,135 participants remained for the main analysis (age range: 44.6-82.1; mean age: 64.1, 51.9% female). Following quality control procedures (QC, see below), 28,212 UK Biobank participants of European descent remained for the main analyses (age range: 45.1-82.1; mean age: 64.1, 55.1% female), 910 of which were identified as close relatives and removed prior to genetic analyses (see below), leaving a final sample for the primary analyses of 27,302 (age range: 45.1-82.1; mean age: 64.1, 54.9% female). For the replication sample, we accessed a newer release of UK Biobank participants (n = 48,045) and removed the 27,302 participants included in the primary analyses. After running through identical QC procedures as for the main sample (although applied only to the cerebellar features of primary interest and covariates included when analyzing these), the replication sample consisted of 11,264 unrelated UK Biobank participants of European descent (age range: 46.1-83.7; mean age: 66.8, 46.9% female). The UK Biobank was approved by the National Health Service National Research Ethics Service (ref. 11/NW/0382),

**Initial MR image processing**

MRI data was first processed using the recon-all pipeline in Freesurfer 5.3(*71*), yielding a large number of brain features. Of these, we retained measures of estimated total intracranial volume (eTIV), total cerebro-cortical surface area, average cerebro-cortical thickness as well as the volumes of seven subcortical structures (hippocampus, amygdala, thalamus, pallidum, putamen, caudate nucleus and nucleus accumbens) as cerebral (or global, in the case of eTICV) comparison regions for our main cerebellar analyses. These anatomical features were averaged across hemispheres, yielding a total of ten comparison phenotypes.

Next, the bias-field corrected T1-images from the FreeSurfer analyses were analyzed using the cerebellum-optimized SUIT-toolbox(*72*). In brief, SUIT isolates the cerebellum and brain stem from T1-images, segments cropped images into grey and white matter, and normalizes these tissue probability maps to a cerebellum-specific anatomical template. After multiplying the grey matter maps with the Jacobian of the transformation matrix (i.e. preserving information about absolute volumes), we extracted grey matter intensity values overlapping the 28 cerebellar lobular labels in the probabilistic SUIT-atlas. Since cerebellar volumetric indices showed high correlations across the two hemispheres (range: .87-.96; mean: .94), we created mean volumetric measures by averaging across hemispheres. Finally, we also combined the two smallest regions in the SUIT-atlas (Vermis

Crus I and II located in the midline, average volumes: 2.9 and 293.4 mm$^2$, respectively) to create a

new Vermis Crus region (average volume: 296.2 mm$^2$). This procedure reduced the 28 cerebellar

lobular volumes to 16 morphological indices. Total cerebellar grey matter volume was defined as the

sum of all 28 lobular labels in the SUIT-atlas.

**Quality control procedures**

After excluding 4,818 UKBB participants of non-European origin, anatomical indices from the

remaining 33,317 participants went through an iterative quality control (QC) procedure. First, we

excluded 639 subjects with a mean Euler number below -217, indicating poor MRI quality(*29*), as

well as 12 subjects with missing and 90 subjects with zero values for any of the key cerebellar or

cerebral brain measures of interest. Next, we used general additive models (GAM, implemented in

the R-package "mgcv") in order to model the effects of age (estimated as smooth functions for males

and females separately, using cubic splines with 10 knots), sex, and scanner site on estimated total

intracranial volume (eTIV) and mean thickness of the cerebral cortex. Specifically, we used the

following formula:

*GAM_model<-gam(Anatomical feature ~ s(Age, bs = "cs", by = Sex, k = 10) + as.factor(Sex)*

*+ as.factor(Scanner), data = UKBB_cerebellum_GWAS, select = TRUE, method = "REML")*

Adjusted eTIV and mean cortical thickness indices were then created by reconstructing the data

using the intercept and residuals from this model (i.e., removing effects of age, sex and scanner),

before identifying and rejecting potential outliers, defined as +/- 3 median absolute deviations

(MAD)(*73*) from the median of these adjusted values. Data from 350 subjects were rejected based

on these criteria.

For the remaining cerebral anatomical measures, as well as total cerebellar volume, this

procedure was then repeated, with scaled eTIV as an additional predictor. Since previous studies

have demonstrated that the relationship between regional brain volumes and intracranial volume is

not strictly allometric(*74, 75*), we estimated the effect of eTIV using cubic splines with 10 knots.

Specifically, we used the following formula:

*GAM_model<-gam(Anatomical feature ~ s(Age, bs = "cs", by = Sex, k = 10) + as.factor(Sex)*

*+ as.factor(Scanner) + s(eTIV_scaled, bs = "cs", k = 10), data = UKBB_cerebellum_GWAS,*

*select = TRUE, method = "REML")*

In order to be maximally sensitive to outliers in relative cerebellar volumes, we replaced eTIV with

total cerebellar volume in the GAM models of cerebellar regions of interest (i.e., SUIT atlas regions),

using the following formula:

*GAM_model<-gam(Anatomical feature ~ s(Age, bs = "cs", by = Sex, k = 10) + as.factor(Sex)*

*+ as.factor(Scanner) + total cerebellar volume, data = UKBB_cerebellum_GWAS, select =*

*TRUE, method = "REML")*

Adjusted cerebral and cerebellar indices were then created by reconstructing the data using the

intercept and residuals from these models (i.e., removing estimated effects of age, sex and eTIV or

total cerebellar volume), before rejecting 1792 participants with potential outlier cerebral indices and

2222 participants with potential outlier cerebellar indices (MAD > +/- 3).

This iterative QC procedure resulted in the rejection of 5,105 (i.e., 15.3%) of the original 33,317

datasets, leaving 28,212 datasets for further analysis.

**Non-negative matrix factorization**

Cerebellar grey matter maps from 28,212 subjects passing the iterative QC procedure were

smoothed with a 4mm full-with-half-maximum gaussian kernel in SPM12(*76*), concatenated across

all subjects and multiplied with a binary mask to isolate voxels located in the cerebellar cortex. This

cerebellar cortical mask was constructed by multiplying a binary mask containing all 28 cerebellar

lobules of the SUIT-atlas with the thresholded (at a value of 0.1) mean (unsmoothed) grey matter

segmentation across all 28,212 participants.

The smoothed, concatenated and masked grey matter maps were then subjected to

orthogonal projective non-negative matrix-factorization (OPNMF)(*26*), in order to derive data-driven

parcellations of regional cerebellar grey matter volume.

Non-negative matrix factorization (NNMF) is a blind source separation technique that allows

structural brain networks to be described in a hypothesis-free, data-driven way by identifying patterns

of covariation in the data. In contrast to alternative techniques, such as principal component analysis

and independent component analysis, which yield components with both positive and negative

weights that are often difficult to interpret, NNMF produces a sparse, positive-only, parts-based

representation of the data. Importantly, NNMF has previously been proven effective in estimating

covariance patterns in neuroimaging data while providing an easier interpretation of the results than

other matrix decomposition techniques such as principal component analysis (PCA) or independent

component analysis (ICA)(*26-28, 77, 78*).

Briefly, NNMF decomposes an input matrix (voxels×subjects) into two matrices; a component

matrix W (voxels×k) and a weight matrix H (k×subjects) where *k* is the number of components that

needs to be specified by the user. Here, we used an implementation of orthogolnal projective non-

negative matrix factorization previously used in a number of publications(*26-28, 77, 78*) and

downloaded from [https://github.com/asotiras/brainparts](https://github.com/asotiras/brainparts). Given the large number of participants, we

used the opnmf_mem.m function, which has been optimized for high-dimensional data. The function

was run with default parameter settings, exept for maximal number of iterations, which was increased
to 200k, in order to ensure full convergence across all tested model orders.
738
**Model order selection**
Since the resulting parcellations are highly dependent on the requested number of components (or
model order) specified, we tested model orders ranging from 2 to 30 in steps of 1, as well as from
30 to 100 in steps of 10. Binarized winner-winner-takes-all (i.e. assigning each voxel to the NNMF
component with the highest loading) maps of the resulting decompositions projected onto a flattened
representation of the cerebellar cortex are shown in Supplementary Figure 1.
Resulting NNMF decompositions were then evaluated based on two criteria; 1) how much of the
variance in the original input data a given NNMF solution (i.e., component maps and subject weights)
could explain and on 2) how reproducible the component maps were. As a metric of change in
explained variance we used the change of the Frobenius norm of the reconstruction error. With
increasing model orders the variance explained will always increase and the reconstruction error
decrease, but if the decrease in the reconstruction error (or gradient) levels off, this indicates that
the intrinsic dimensionality of the data might have been approximated (and that the subsequent
increase in explained variance can largely be attributed to fitting random noise in the input data). In
order to assess reproducibility, we split the full dataset into two equal sets (matched with respect to
scanner site, n = 14,105 and 14,107, respectively), and ran NNMF on each split-half sample. For
each set of independent NNMF parcellations, we computed two reproducibility indices. First, for each
model order we matched components across split-half runs using the Hungarian algorithm(*79*),
before computing the spatial correlations between matched component maps, and extracting the
median correlation across all matched components as our first reproducibility index. For our second
reproducibility index, we first computed one overall – and categorical – component map using a
"winner-takes-all"-approach, i.e., assigning each voxel to the NNMF component with the highest
loading. Next, we calculated the adjusted Rand index (ranging from 0 to 1, with higher values
indicating greater similarity(*80*), across the two categorical parcellation maps for each model order
as our second metric of reproducibility.
As can be seen in Supplementary Figure 2A, for lower model orders, increasing the model
order resulted in a sharp decrease in the reconstruction error, indicating that models with more
components resulted in a significantly better fit to the data. However, after reaching model orders
between 15 and 30, the incremental improvement in fit from adding another component appeared to
level off. As expected, the reproducibility results showed a largely inverse pattern, with both
reproducibility indices decreasing with increasing model orders (Supplementary Figure 2B). Of note,
for model orders up to 8, median spatial correlations across all matched components was above .99,
indicating almost identical parcellations derived from the two independent samples. However, even
for a model order of 100, the median pairwise correlation was still pretty high (.85), with 60% of
components showing pairwise spatial correlations above 0.8, suggesting a reasonable level of

reproducibility even for the most fine-grained parcellation. Our reproducibility index for the categorical parcellations showed very similar results; the adjusted Rand index remained above .9 for model orders up to 8, and then decreased to 0.56 at a model order of 100.

Given that the change in reconstruction error appeared to stabilize between 15 and 30 components, indicating that the intrinsic dimensionality of the data had been approximated, we searched for the most reproducible parcellations within this range. Since the reliability estimates for model orders of 16 and 23 were very similar, with the 23-component solution explaining more of the variance in the original data, we used the 23-component parcellation for all further analyses.

**Adjustment and rank-order normalization of anatomical indices**

Prior to being subjected to genome-wide association analyses, we adjusted all anatomical indices for effects of age, sex, estimated total intracranial volume, scanner site, 40 genetic population components, genetic batch and mean Euler number (i.e, an index of MRI image quality(*29*), averaged across hemispheres). Finally, all adjusted anatomical indices were inverse rank normalized(*81*).

**Pre-processing of genetic data**

For all genetic analyses we made use of the UKB v3 imputed data, which has undergone extensive quality control procedures as described by the UKB genetics team(*82*). After converting the BGEN format to PLINK binary format, we additionally carried out standard quality check procedures. We first selected White Europeans, as determined by a combination of self-identification as 'White British' and similar genetic ancestry based on genetic principal components, that had undergone the neuroimaging protocol. We then filtered out individuals with more than 10% missingness, removed SNPs with low imputation quality scores (INFO <.5), SNPs with more than 10% missingness, and SNPs failing the Hardy-Weinberg equilibrium test at p=1*10-9. We further set a minor allele frequency threshold of 0.005 leaving 9,061,238 SNPs. After estimating the genetic relationship matrix (GRM) using genetic complex trait analysis (GCTA(*31*)), we finally removed 910 participants defined as close relatives using a threshold of 0.05 (approximately corresponding to 3rd cousins).

**Heritability estimation and genetic correlation analyses**

SNP-based heritability estimates for all morphological features – as well as the pairwise genetic correlations between cerebellar features - were estimated using genetic complex trait analysis (GCTA(*31*)).

**Univariate genome-wide association analyses**

Univariate analyses of total cerebellar grey matter volume and the ten cerebral comparison phenotypes were conducted using Plink v1.9.

**Genetic correlation analyses across discovery and replication samples and with previous published GWASs**

Genetic correlation analyses across different samples were conducted using LD-score regression(*30*).

**Multivariate genome-wide association analyses**

For our main analysis, we used a recently developed multivariate analysis method (MOSTest(*34*)), to conduct a multivariate genome-wide association (GWA) analysis on cerebellar morphological features. MOSTest identifies genetic effects across multiple phenotypes, yielding a multivariate GWAS summary statistic across all 23 features, and provides robust (permutation based) test statistics. For mathematical details of the implementation, see van der Meer et al. (2020)(*34*), for details on the software implementation see github.com/precimed/mostest. MOSTest has been extensively validated in the original methods paper, including simulations and comparisons with other methods that have confirmed its solid discovery performance as well as an order of magnitude shorter runtime compared to other tools(*34*). For comparison to standard univariate approaches, we also performed univariate GWASs (extracted from the univariate stream of MOSTest(*34*) and identical to results from analyses using Plink).

**Multivariate replication analysis**

To ensure that not only single lead SNP associations replicate but that also the multivariate pattern of these associations are consistent in the discovery and replication sample, we implemented a multivariate replication procedure established in Loughnan et al.(*37*). In brief, for each lead SNP identified in the multivariate analysis in the discovery sample, this procedure derives a composite score from the mass-univariate *z*-statistics and tests for associations of the composite score with the genotype in the replication sample (for mathematical formulation see Loughnan et al.(*37*)). 19 of the 560 lead SNPs could not be tested as they were not available in the replication sample. For the remaining SNPs we report the percent of loci replicating at $P < 0.05$, the percent remaining significant after Bonferroni correction for 541 conducted tests, and the percent of lead SNPs showing the same effect direction. In addition, we report the percentage of genetic loci replicated (defined as containing at least one replicating lead SNP).

**Multivariate cerebral comparison phenotypes**

To compare key multivariate cerebellar results with other multivariate brain phenotypes, we downloaded summary statistics from two recent studies on cerebrocortical(*42*) and hippocampal(*43*) regional morphology. These comparison summary statistics were next analyzed using FUMA(*35*) as described for the main cerebellar results below.

**Locus identification and SNP annotation**

To identify genetic loci we uploaded summary statistics to the FUMA platform v1.4.1(*35*) Using the
1000GPhase3 EUR as reference panel, we identified independent SNPs as SNPs below the
significance threshold of $P < 5e{-}8$ that were also in linkage equilibrium with each other at $r^2 < 0.6$.
For each independent SNP, all candidate variants are identified as variants with LD r2≥ 0.6 with the
independent SNP. A fraction of the independent significant SNPs in approximate linkage equilibrium
with each other at $r^2 < 0.1$ were considered as lead SNPs. For a given lead SNP, the borders of the
genomic locus are defined as min/max positional coordinates over all corresponding candidate
SNPs. Finally, loci are merged if they are separated by less than 250kb.

FUMA further annotates associated SNPs based on functional categories, Combined
Annotation Dependent Depletion (CADD) scores which predicts the deleteriousness of SNPs on
protein structure/function(*54*), RegulomeDB scores which predicts regulatory functions(*83*); and
chromatin states that shows the transcription/regulation effects of chromatin states at the SNP
locus(*84*). For all these analyses, we used default FUMA parameters.

**Genome-wide gene-based association and gene-set analyses**
We conducted genome-wide gene-based association and gene-set analyses using MAGMA
v.1.10(*49*) (http://ctg.cncr.nl/software/magma). MAGMA performs multiple linear regression to map
the input SNPs to 19,190 protein coding genes and estimates the significance value of that gene.
Genes were considered significant if the $P$ value was <0.05 after Bonferroni correction for 19,190
genes. The same procedure was used for MAGMA analysis of summary statistics for the three
multivariate cerebral comparison phenotypes (cerebrocortical thickness and surface area(*42*) and
regional hippocampal volumes(*43*)).

MAGMA gene-level statistics were next used as input to gene-property and gene-set
analyses in MAGMA. Gene-property analyses test for associations between tissue specific gene
expression profiles and disease-gene associations. The gene-property analysis is based on the
regression model: $Z \sim \beta_0 + E_t\beta_E + A\beta_A + B\beta_B + \epsilon$, where Z  is a gene-based Z-score converted from
the gene-based P-value, B is a matrix of several technical confounders included by default (e.g.,
gene size, gene density, sample size), $E_t$ is the gene expression value of a testing tissue type and A
is the average expression across all tissue types in a data set (ensuring a test of expression
specificity). We performed a one-sided test ($\beta_E > 0$) which is essentially testing the positive
relationship between tissue specificity and genetic association of genes.

We tested associations with two regional brain gene expression datasets (Allen Human
Brain Atlas(*33*) and GTEx) and one developmental brain gene expression dataset (BrainSpan). For
extraction and processing of gene expression data, see below.

**Extraction and processing of gene expression data**
*Allen Human Brain Atlas:* Regional microarray expression data were obtained from 6 post-mortem
brains (1 female, ages 24.0--57.0, 42.50 +/- 13.38) provided by the Allen Human Brain Atlas(*33*).

Data were processed with the abagen toolbox (version 0.1.3) (*32*) using two volumetric atlases; 1)

the binarized 23-region NNMF-derived parcellation of the cerebellar cortex; and 2) a modified version

of the Desikan atlas were ROIs were merged to construct 9 bilateral regions: cerebellum, cerebral

cortex, pallidum, caudate, putamen, thalamus, amygdala, nucleus accumbens and hippocampus.

First, microarray probes were reannotated using data provided by Arnatkeviciute, Fulcher

and Fornito(*85*); probes not matched to a valid Entrez ID were discarded. Next, probes were filtered

based on their expression intensity relative to background noise(*86*), such that probes with intensity

less than the background in >=50.00% of samples across donors were discarded, yielding 31,569

probes. When multiple probes indexed the expression of the same gene, we selected and used the

probe with the most consistent pattern of regional variation across donors (i.e., differential

stability(*87*)), calculated with:

$$\Delta_S(p) = \frac{1}{\binom{N}{2}} \sum_{i=1}^{N-1} \sum_{j=i+1}^{N} \rho[B_i(p), B_j(p)]$$

where *p* is Spearman's rank correlation of the expression of a single probe, p, across regions

in two donors $B_i$ and $B_j$, and N is the total number of donors.

Here, regions correspond to the structural designations provided in the ontology from the

AHBA. The MNI coordinates of tissue samples were updated to those generated via non-linear

registration using the Advanced Normalization Tools (ANTs; https://github.com/chrisfilo/alleninf).

Samples were assigned to brain regions in the provided atlas if their MNI coordinates were within 2

mm of a given parcel. To reduce the potential for misassignment, sample-to-region matching was

constrained by hemisphere and gross structural divisions (i.e., cortex, subcortex/ brainstem, and

cerebellum, such that e.g., a sample in the left cortex could only be assigned to an atlas parcel in

the left cortex(*85*)). All tissue samples not assigned to a brain region in the provided atlas were

discarded. \n\nInter-subject variation was addressed by normalizing tissue sample expression

values across genes using a robust sigmoid function(*88*):

$$x_{norm} = \frac{1}{1+\exp(-\frac{(x-\langle x\rangle)}{\text{IQR}_x})}$$

913        where (x) is the median and $\text{IQR}_z$ is the normalized interquartile range of the expression of a

914    single tissue sample across genes. Normalized expression values were then rescaled to the unit

915    interval:

$$x_{scaled} = \frac{x_{norm}-\min(x_{norm})}{\max(x_{norm})-\min(x_{norm})}$$

Gene expression values were then normalized across tissue samples using an identical

procedure. Samples assigned to the same brain region were averaged separately for each donor

and then across donors, yielding two regional expression matrices with 23 and 9 rows, corresponding

to brain regions, and 15,631 and 15,633 columns, corresponding to the retained genes. Prior to

inclusion in MAGMA gene property analyses, we converted gene names for the modified Desikan

atlas to ENSMBL IDs, and calculated the mean expression value across tissue types (in order to

include this as a covariate in MAGMA analyses testing for gene expression specificity), resulting in

a 10 (regions) by (15,490) gene expression matrix.

*GTeX:* Text files containing median transcript per millimeter (TPM) values for 53 tissue types

(GTEx_Analysis_2017-06-05_v8_RNASeQCv1.1.9_gene_median_tpm.gct.gz) were down-loaded

from the GTEx portal (https://gtexportal.org/home/datasets/). After selecting only expression data

from the seven (out of nine) comparison brain regions present in the GTEx dataset (i.e., amygdala,

caudate, cerebellum, cortex, hippocampus, nucleus accumbens and putamen), we filtered the data

by only including genes with median TPM values above 1 for at least one of these tissue type

(retaining 19,578 of 56,200 annotated genes). Following the procedure used by FUMA(*35*), we next

winsorized median TPM values at 50 (i.e., replaced TPM>50 with 50), before log transforming TPM

with pseudocount 1 (log2(RPKM+1)). Finally, we calculated the mean expression value across tissue

types (in order to include this as a covariate in MAGMA analyses testing for gene expression

specificity).

*BrainSpan data:* The analysis of BrainSpan data testing for developmentally specific brain

expression was performed entirely within FUMA v1.4.1., using default parameters.

**Analysis of Human Dating Genome Data.** The Atlas of Variant Age for chromosomes 1-22 was

downloaded from the Human Genome Dating (HGD) website: https://human.genome.dating/. This

atlas contains more than 45 million SNPs which has been assigned dates of origin based on a

recombination clock and mutation clock applied to two large-scale sequencing datasets (1000

Genomes Project(*89*) and The Simons Genome Diversity Project(*90*)), with no assumptions made

about demographic or selective processes(*44*). The current study used the median joint age

estimates from both clocks when analyzing SNPs present in both datasets in combination (i.e.,

13,694,493 SNPs).

After merging dated SNPs with the 40,405,505 SNPs also present in the Haplotype

Reference Consortium reference data (to add minor allele frequencies, MAF) and removing 715,083

(5.2%) SNPs with missing MAF values as well as the very few (14,549, 0.1%) SNPs dated older

than 2 million years, 12,960,066 SNPs remained. 146,977 of these dated SNPs were matched to

candidate SNPs linked to cerebellar morphology or any of the comparison brain phenotypes,

hereafter referred to as brain-SNPs. Partially because very rare variants (Minor Allele

Frequency/MAF < 0.005) had been removed prior to the multivariate GWAS analysis, MAF was not

equally distributed between these brain-SNPs and the full range of dated SNPs in the HGD dataset.

Importantly, MAF has been shown to be systematically related to the estimated age of SNPs, with a

higher proportion of low-MAF SNPs in more recent than in older time-bins(44, 45). Consequently,

following the analysis approach established by Libedinsky et al. (45), we first determined the MAF-

distribution of brain-SNPs (across eight bins: <0.0001; 0.0001-0.001; 0.001-0.01; 0.01-0.1; 0.1-0.2;

0.2-0.3; 0.3-0.4; 0.4-0.5) and selected random set of 2,645,586 SNPs from the HGD dataset that

were matched to the brain-SNPs in terms of MAF-bin distribution.

963           For statistical inference we constructed null models (separate for each brain phenotype) by

randomly drawing sets of SNPs (of equal size to the number of phenotype-linked candidate SNPs)

from the MAF-matched HGD-dataset and computing the histograms of estimated dates from 0 to 2

million years ago (divided into 100 bins of 20.000 years) over 10,000 iterations. From these null

models we extracted bin means and standard deviations (used to assign z-scores to the bin values

of brain phenotype histograms), as well as significance thresholds (defined as the upper and lower

99.95$^{th}$ percentile of the null model (i.e. corresponding to a Bonferroni corrected threshold of 0.05

across 100 bins).

**Positional and eQTL mapping of SNPs to plausible causal genes**

In addition to using MAGMA, we also mapped candidate SNPs to plausible causal genes using two

complementary gene mapping strategies implemented ion FUMA(35): 1) Positional mapping of

deleterious SNPs (defined as having a CADD-score > 12.37) and 2) eQTL-mapping of SNPs

previously shown to alter gene expression in cerebellar tissue (from the BRAINEAC and GTEx v8

databases). These analyses were run with default FUMA parameters. For the three multivariate

cerebral comparison phenotypes (i.e., cerebrocortical thickness, cerebrocortical surface area and

hippocampal regional volumes), we employed identical gene mapping procedures to our cerebellar

morphology results, except for the tissues chosen for eQTL mapping (cerebrocortical and

hippocampal, respectively).

**Gene set analyses using lists of mapped genes**

All gene set analyses using mapped genes were conducted using the hypeR R-package (91). This

package implements the hypergeometric test (also known as Fisher's exact test), which assigns a

p-value to gene-set overlaps given gene set sizes and the number of background genes. This R-

package also contains functions for downloading and/or formatting gene sets. The following gene

sets were accessed using hypeR: "Allen_Brain_Atlas_up" (regional overexpression in the Allen

Muse Brain Atlas), "MGI_Mammalian_Phenotype_Level_4_2021", "DisGeNet" (all from the enrichR

platform. In addition, we downloaded sets of genes with regional overexpression in the Allen Human

Brain Atlas from the Harmonizome platform(92) and accessed a list of genes mapped to human

accelerated regions(93). For all gene-set analyses we employed Bonferroni-correction by dividing

the p-value threshold of 0.05 by the number of gene sets included in each analysis.

**Genetic overlap between cerebellar morphology and brain disorders**

We accessed GWAS summary statistics for attention deficit hyperactivity disorder (ADHD)(*94*), autism spectrum disorder (ASD)(*95*), bipolar disorder (BIP)(*96*) and major depressive disorder (MDD)(*97*) from the Psychiatric Genomics Consortium. In order to avoid sample overlap, for MDD we used summary statistics based on a sample with the UK Biobank participants removed. 23&me participants included in the original MDDGWAS were also excluded, since these data are not freely available). Finally, we included data from a recent study of schizophrenia (SCZ)(*98*). Shared variants associated with cerebellar morphology and each of the above-mentioned brain disorders were identified using conjunctional FDR statistics (FDR < 0.05)(*99, 100*). In contrast to genetic correlation analysis, conjunctional FDR does not require effect directions and can therefore be applied to summary statistics from multivariate GWAS, which do not contain effect directions. Two genomic regions, the extended major histocompatibility complex genes region (hg19 location Chr 6: 25119106–33854733) and chromosome 8p23.1 (hg19 location Chr 8: 7242715–12483982) for all cases and *APOE* region for ASD, were excluded from the FDR-fitting procedures because complex correlations in regions with intricate LD can bias FDR estimation. We submitted the results from conjunctional FDR to FUMA v1.3.7(*35*) to annotate the genomic loci with conjFDR value < 0.10 having an $r^2 \geq 0.6$ with one of the independent significant lead SNPs. Genetic correlations between univariate results for the 23 cerebellar features, the 10 comparison brain phenotypes, and each of the five mental disorders were computed using LD-score regression as described above.

**References for Online Methods:**

1. S. Herculano-Houzel, *The human advantage : a new understanding of how our brain became remarkable* (The MIT Press, Cambridge, Massachusetts, 2016), pp. xiii, 256 pages.
2. R. A. Barton, C. Venditti, in *Curr. Biol.* (2014).
3. M. I. Sereno, J. Diedrichsen, M. Tachrount, G. Testa-Silva, H. d'Arceuil, C. De Zeeuw, The human cerebellum has almost 80% of the surface area of the neocortex. *Proc Natl Acad Sci U S A* **117**, 19538-19543 (2020).
4. P. W. Harrison, S. H. Montgomery, Genetics of Cerebellar and Neocortical Expansion in Anthropoid Primates: A Comparative Approach. *Brain Behav Evol* **89**, 274-285 (2017).
5. H. C. Leiner, A. L. Leiner, R. S. Dow, in *Trends in Neurosciences*. (1993), vol. 16, pp. 444-447.
6. P. L. Strick, R. P. Dum, J. A. Fiez, in *Annu. Rev. Neurosci.* (2009), vol. 32, pp. 413-434.
7. R. L. Buckner, in *Neuron*. (2013), vol. 80, pp. 807-815.
8. M. King, C. R. Hernandez-Castillo, R. A. Poldrack, R. B. Ivry, J. Diedrichsen, Functional boundaries in the human cerebellum revealed by a multi-domain task battery. *Nat Neurosci* **22**, 1371-1378 (2019).
9. S. S. Wang, A. D. Kloth, A. Badura, The cerebellum, sensitive periods, and autism. *Neuron* **83**, 518-532 (2014).
10. T. Moberget, N. T. Doan, D. Alnaes, T. Kaufmann, A. Cordova-Palomera, T. V. Lagerberg, J. Diedrichsen, E. Schwarz, M. Zink, S. Eisenacher, P. Kirsch, E. G. Jonsson, H. Fatouros-Bergman, L. Flyckt, KaSp, G. Pergola, T. Quarto, A. Bertolino, D. Barch, A. Meyer-Lindenberg, I. Agartz, O. A. Andreassen, L. T. Westlye, Cerebellar volume and cerebellocerebral structural covariance in schizophrenia: a multisite mega-analysis of 983 patients and 1349 healthy controls. *Mol Psychiatry* **23**, 1512-1520 (2018).
11. H. H. Adams, D. P. Hibar, V. Chouraki, J. L. Stein, P. A. Nyquist, M. E. Renteria, S. Trompet, A. Arias-Vasquez, S. Seshadri, S. Desrivieres, A. H. Beecham, N. Jahanshad, K. Wittfeld, S. J. Van der Lee, L. Abramovic, S. Alhusaini, N. Amin, M. Andersson, K. Arfanakis, B. S. Aribisala, N. J. Armstrong, L. Athanasiu, T. Axelsson, A. Beiser, M. Bernard, J. C. Bis, L. M. Blanken, S. H. Blanton, M. M. Bohlken, M. P. Boks, J. Bralten, A. M. Brickman, O. Carmichael, M. M. Chakravarty, G. Chauhan, Q. Chen, C. R. Ching, G. Cuellar-Partida, A. D. Braber, N. T. Doan, S. Ehrlich, I. Filippi, T. Ge, S. Giddaluru, A. L. Goldman, R. F. Gottesman, C. U. Greven, O. Grimm, M. E. Griswold, T. Guadalupe, J. Hass, U. K. Haukvik, S. Hilal, E. Hofer, D. Hoehn, A. J. Holmes, M. Hoogman, D. Janowitz, T. Jia, D. Kasperaviciute, S. Kim, M. Klein, B. Kraemer, P. H. Lee, J. Liao, D. C. Liewald, L. M. Lopez, M. Luciano, C. Macare, A. Marquand, M. Matarin, K. A. Mather, M. Mattheisen, B. Mazoyer, D. R. McKay, R. McWhirter, Y. Milaneschi, N. Mirza-Schreiber, R. L. Muetzel, S. M. Maniega, K. Nho, A. C. Nugent, L. M. Loohuis, J. Oosterlaan, M. Papmeyer, I. Pappa, L. Pirpamer, S. Pudas, B. Putz, K. B. Rajan, A. Ramasamy, J. S. Richards, S. L. Risacher, R. Roiz-Santianez, N. Rommelse, E. J. Rose, N. A. Royle, T. Rundek, P. G. Samann, C. L. Satizabal, L. Schmaal, A. J. Schork, L. Shen, J. Shin, E. Shumskaya, A. V. Smith, E. Sprooten, L. T. Strike, A. Teumer, R. Thomson, D. Tordesillas-Gutierrez, R. Toro, D. Trabzuni, D. Vaidya, J. Van der Grond, D. Van der Meer, M. M. Van Donkelaar, K. R. Van Eijk, T. G. Van Erp, D. Van Rooij, E. Walton, L. T. Westlye, C. D. Whelan, B. G. Windham, A. M. Winkler, G. Woldehawariat, C. Wolf, T. Wolfers, B. Xu, L. R. Yanek, J. Yang, A. Zijdenbos, M. P. Zwiers, I. Agartz, N. T. Aggarwal, L. Almasy, D. Ames, P. Amouyel, O. A. Andreassen, S. Arepalli, A. A. Assareh, S. Barral, M. E. Bastin, D. M. Becker, J. T. Becker, D. A. Bennett, J. Blangero, H. van Bokhoven, D. I. Boomsma, H. Brodaty, R. M. Brouwer, H. G. Brunner, R. L. Buckner, J. K. Buitelaar, K. B. Bulayeva, W. Cahn, V. D. Calhoun, D. M. Cannon, G. L. Cavalleri, C. Chen, C. Y. Cheng, S. Cichon, M. R. Cookson, A. Corvin, B. Crespo-Facorro, J. E. Curran, M. Czisch, A. M. Dale, G. E. Davies, E. J. De Geus, P. L. De Jager, G. I. de Zubicaray, N. Delanty, C. Depondt, A. L. DeStefano, A. Dillman, S. Djurovic, G. Donohoe, W. C. Drevets, R. Duggirala, T. D. Dyer, S. Erk, T. Espeseth, D. A. Evans, I. O. Fedko, G. Fernandez, L. Ferrucci, S. E. Fisher, D. A. Fleischman, I. Ford, T. M. Foroud, P. T. Fox, C. Francks, M. Fukunaga, J. R. Gibbs, D. C. Glahn, R. L. Gollub, H. H. Goring, H. J. Grabe, R. C. Green, O. Gruber, V. Gudnason, S. Guelfi, N. K.

Hansell, J. Hardy, C. A. Hartman, R. Hashimoto, K. Hegenscheid, A. Heinz, S. Le Hellard, D. G. Hernandez, D. J. Heslenfeld, B. C. Ho, P. J. Hoekstra, W. Hoffmann, A. Hofman, F. Holsboer, G. Homuth, N. Hosten, J. J. Hottenga, H. E. Pol, M. Ikeda, M. K. Ikram, C. R. Jack, Jr., M. Jenkinson, R. Johnson, E. G. Jonsson, J. W. Jukema, R. S. Kahn, R. Kanai, I. Kloszewska, D. S. Knopman, P. Kochunov, J. B. Kwok, S. M. Lawrie, H. Lemaitre, X. Liu, D. L. Longo, W. T. Longstreth, Jr., O. L. Lopez, S. Lovestone, O. Martinez, J. L. Martinot, V. S. Mattay, C. McDonald, A. M. McIntosh, K. L. McMahon, F. J. McMahon, P. Mecocci, I. Melle, A. Meyer-Lindenberg, S. Mohnke, G. W. Montgomery, D. W. Morris, T. H. Mosley, T. W. Muhleisen, B. Muller-Myhsok, M. A. Nalls, M. Nauck, T. E. Nichols, W. J. Niessen, M. M. Nothen, L. Nyberg, K. Ohi, R. L. Olvera, R. A. Ophoff, M. Pandolfo, T. Paus, Z. Pausova, B. W. Penninx, G. B. Pike, S. G. Potkin, B. M. Psaty, S. Reppermund, M. Rietschel, J. L. Roffman, N. Romanczuk-Seiferth, J. I. Rotter, M. Ryten, R. L. Sacco, P. S. Sachdev, A. J. Saykin, R. Schmidt, P. R. Schofield, S. Sigurdsson, A. Simmons, A. Singleton, S. M. Sisodiya, C. Smith, J. W. Smoller, H. Soininen, V. Srikanth, V. M. Steen, D. J. Stott, J. E. Sussmann, A. Thalamuthu, H. Tiemeier, A. W. Toga, B. J. Traynor, J. Troncoso, J. A. Turner, C. Tzourio, A. G. Uitterlinden, M. C. Hernandez, M. Van der Brug, A. Van der Lugt, N. J. Van der Wee, C. M. Van Duijn, N. E. Van Haren, T. E. D. Van, M. J. Van Tol, B. N. Vardarajan, D. J. Veltman, M. W. Vernooij, H. Volzke, H. Walter, J. M. Wardlaw, T. H. Wassink, M. E. Weale, D. R. Weinberger, M. W. Weiner, W. Wen, E. Westman, T. White, T. Y. Wong, C. B. Wright, H. R. Zielke, A. B. Zonderman, I. J. Deary, C. DeCarli, H. Schmidt, N. G. Martin, A. J. De Craen, M. J. Wright, L. J. Launer, G. Schumann, M. Fornage, B. Franke, S. Debette, S. E. Medland, M. A. Ikram, P. M. Thompson, Novel genetic loci underlying human intracranial volume identified through genome-wide association. *Nat Neurosci* **19**, 1569-1582 (2016).

12. D. van der Meer, J. Rokicki, T. Kaufmann, A. Cordova-Palomera, T. Moberget, D. Alnaes, F. Bettella, O. Frei, N. T. Doan, I. E. Sonderby, O. B. Smeland, I. Agartz, A. Bertolino, J. Bralten, C. L. Brandt, J. K. Buitelaar, S. Djurovic, M. van Donkelaar, E. S. Dorum, T. Espeseth, S. V. Faraone, G. Fernandez, S. E. Fisher, B. Franke, B. Haatveit, C. A. Hartman, P. J. Hoekstra, A. K. Haberg, E. G. Jonsson, K. K. Kolskar, S. Le Hellard, M. J. Lund, A. J. Lundervold, A. Lundervold, I. Melle, J. Monereo Sanchez, L. C. Norbom, J. E. Nordvik, L. Nyberg, J. Oosterlaan, M. Papalino, A. Papassotiropoulos, G. Pergola, D. J. F. de Quervain, G. Richard, A. M. Sanders, P. Selvaggi, E. Shumskaya, V. M. Steen, S. Tonnesen, K. M. Ulrichsen, M. P. Zwiers, O. A. Andreassen, L. T. Westlye, I. Alzheimer's Disease Neuroimaging, N. Pediatric Imaging, S. Genetics, Brain scans from 21,297 individuals reveal the genetic architecture of hippocampal subfield volumes. *Mol Psychiatry*, (2018).

13. T. Chambers, V. Escott-Price, S. Legge, E. Baker, K. D. Singh, J. T. R. Walters, X. Caseras, R. J. L. Anney, Genetic common variants associated with cerebellar volume and their overlap with mental disorders: a study on 33,265 individuals from the UK-Biobank. *Mol Psychiatry* **27**, 2282-2290 (2022).

14. E. Tissink, S. C. de Lange, J. E. Savage, D. P. Wightman, C. A. de Leeuw, K. M. Kelly, M. Nagel, M. P. van den Heuvel, D. Posthuma, Genome-wide association study of cerebellar volume provides insights into heritable mechanisms underlying brain development and mental health. *Commun Biol* **5**, 710 (2022).

15. S. Macri, Y. Savriama, I. Khan, N. Di-Poi, Comparative analysis of squamate brains unveils multi-level variation in cerebellar architecture associated with locomotor specialization. *Nat Commun* **10**, 5560 (2019).

16. J. B. Smaers, A. H. Turner, A. Gomez-Robles, C. C. Sherwood, A cerebellar substrate for cognition evolved multiple times independently in mammals. *Elife* **7**, (2018).

17. K.-H. E, S.-H. A. Chen, M.-H. R. Ho, J. E. Desmond, in *Hum Brain Mapp*. (2014), vol. 35, pp. 593-615.

18. C. J. Stoodley, J. D. Schmahmann, in *NeuroImage*. (2009), vol. 44, pp. 489-501.

19. R. L. Buckner, F. M. Krienen, A. Castellanos, J. C. Diaz, B. T. T. Yeo, in *Journal of Neurophysiology*. (2011), vol. 106, pp. 2322-2345.

20. L. Sang, W. Qin, Y. Liu, W. Han, Y. Zhang, T. Jiang, C. Yu, in *NeuroImage*. (Elsevier Inc., 2012), vol. 61, pp. 1213-1225.
21. F. M. Krienen, R. L. Buckner, in *Cerebral Cortex*. (2009), vol. 19, pp. 2485-2497.
22. J. A. Bernard, R. D. Seidler, K. M. Hassevoort, B. L. Benson, R. C. Welsh, J. L. Wiggins, S. M. Jaeggi, M. Buschkuehl, C. S. Monk, J. Jonides, S. J. Peltier, in *Front Neuroanat*. (2012), vol. 6, pp. 31.
23. J. A. Kipping, W. Grodd, V. Kumar, M. Taubert, A. Villringer, D. S. Margulies, in *NeuroImage*. (2013), vol. 83, pp. 837-848.
24. C. Wang, J. Kipping, C. Bao, H. Ji, A. Qiu, Cerebellar Functional Parcellation Using Sparse Dictionary Learning Clustering. *Front Neurosci* **10**, 188 (2016).
25. J. A. Kipping, T. A. Tuan, M. V. Fortier, A. Qiu, Asynchronous Development of Cerebellar, Cerebello-Cortical, and Cortico-Cortical Functional Networks in Infancy, Childhood, and Adulthood. *Cereb Cortex*, (2016).
26. A. Sotiras, S. M. Resnick, C. Davatzikos, Finding imaging patterns of structural covariance via Non-Negative Matrix Factorization. *Neuroimage* **108**, 1-16 (2015).
27. A. Sotiras, J. B. Toledo, R. E. Gur, R. C. Gur, T. D. Satterthwaite, C. Davatzikos, Patterns of coordinated cortical remodeling during adolescence and their associations with functional specialization and evolutionary expansion. *Proc Natl Acad Sci U S A* **114**, 3527-3532 (2017).
28. D. P. Varikuti, S. Genon, A. Sotiras, H. Schwender, F. Hoffstaedter, K. R. Patil, C. Jockwitz, S. Caspers, S. Moebus, K. Amunts, C. Davatzikos, S. B. Eickhoff, Evaluation of non-negative matrix factorization of grey matter in age prediction. *Neuroimage* **173**, 394-410 (2018).
29. A. F. G. Rosen, D. R. Roalf, K. Ruparel, J. Blake, K. Seelaus, L. P. Villa, R. Ciric, P. A. Cook, C. Davatzikos, M. A. Elliott, A. Garcia de La Garza, E. D. Gennatas, M. Quarmley, J. E. Schmitt, R. T. Shinohara, M. D. Tisdall, R. C. Craddock, R. E. Gur, R. C. Gur, T. D. Satterthwaite, Quantitative assessment of structural image quality. *Neuroimage* **169**, 407-418 (2018).
30. B. K. Bulik-Sullivan, P. R. Loh, H. K. Finucane, S. Ripke, J. Yang, C. Schizophrenia Working Group of the Psychiatric Genomics, N. Patterson, M. J. Daly, A. L. Price, B. M. Neale, LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat Genet* **47**, 291-295 (2015).
31. J. Yang, S. H. Lee, M. E. Goddard, P. M. Visscher, GCTA: a tool for genome-wide complex trait analysis. *Am J Hum Genet* **88**, 76-82 (2011).
32. R. D. Markello, A. Arnatkeviciute, J. B. Poline, B. D. Fulcher, A. Fornito, B. Misic, Standardizing workflows in imaging transcriptomics with the abagen toolbox. *Elife* **10**, (2021).
33. M. J. Hawrylycz, E. S. Lein, A. L. Guillozet-Bongaarts, E. H. Shen, L. Ng, J. A. Miller, L. N. van de Lagemaat, K. A. Smith, A. Ebbert, Z. L. Riley, C. Abajian, C. F. Beckmann, A. Bernard, D. Bertagnolli, A. F. Boe, P. M. Cartagena, M. M. Chakravarty, M. Chapin, J. Chong, R. A. Dalley, B. David Daly, C. Dang, S. Datta, N. Dee, T. A. Dolbeare, V. Faber, D. Feng, D. R. Fowler, J. Goldy, B. W. Gregor, Z. Haradon, D. R. Haynor, J. G. Hohmann, S. Horvath, R. E. Howard, A. Jeromin, J. M. Jochim, M. Kinnunen, C. Lau, E. T. Lazarz, C. Lee, T. A. Lemon, L. Li, Y. Li, J. A. Morris, C. C. Overly, P. D. Parker, S. E. Parry, M. Reding, J. J. Royall, J. Schulkin, P. A. Sequeira, C. R. Slaughterbeck, S. C. Smith, A. J. Sodt, S. M. Sunkin, B. E. Swanson, M. P. Vawter, D. Williams, P. Wohnoutka, H. R. Zielke, D. H. Geschwind, P. R. Hof, S. M. Smith, C. Koch, S. G. N. Grant, A. R. Jones, An anatomically comprehensive atlas of the adult human brain transcriptome. *Nature* **489**, 391-399 (2012).
34. D. van der Meer, O. Frei, T. Kaufmann, A. A. Shadrin, A. Devor, O. B. Smeland, W. K. Thompson, C. C. Fan, D. Holland, L. T. Westlye, O. A. Andreassen, A. M. Dale, Understanding the genetic determinants of the brain with MOSTest. *Nat Commun* **11**, 3512 (2020).
35. K. Watanabe, E. Taskesen, A. van Bochoven, D. Posthuma, Functional mapping and annotation of genetic associations with FUMA. *Nat Commun* **8**, 1826 (2017).
36. K. Wang, M. Li, H. Hakonarson, ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic Acids Res* **38**, e164 (2010).
37. R. J. Loughnan, A. A. Shadrin, O. Frei, D. van der Meer, W. Zhao, C. E. Palmer, W. K. Thompson, C. Makowski, T. L. Jernigan, O. A. Andreassen, C. C. Fan, A. M. Dale,

Generalization of cortical MOSTest genome-wide associations within and across samples.
*Neuroimage* **263**, 119632 (2022).

38.    S. van der Sluis, D. Posthuma, C. V. Dolan, TATES: efficient multivariate genotype-phenotype
analysis for genome-wide association studies. *PLoS Genet* **9**, e1003235 (2013).

39.    P. F. O'Reilly, C. J. Hoggart, Y. Pomyen, F. C. Calboli, P. Elliott, M. R. Jarvelin, L. J. Coin,
MultiPhen: joint model of multiple phenotypes can increase discovery in GWAS. *PLoS One* **7**,
e34861 (2012).

40.    B. Zhao, T. Luo, T. Li, Y. Li, J. Zhang, Y. Shan, X. Wang, L. Yang, F. Zhou, Z. Zhu, I. Alzheimer's
Disease Neuroimaging, N. Pediatric Imaging, Genetics, H. Zhu, Genome-wide association
analysis of 19,629 individuals identifies variants influencing regional brain volumes and refines
their genetic co-architecture with cognitive and mental health traits. *Nat Genet* **51**, 1637-1644
(2019).

41.    S. M. Smith, G. Douaud, W. Chen, T. Hanayik, F. Alfaro-Almagro, K. Sharp, L. T. Elliott, An
expanded set of genome-wide association studies of brain imaging phenotypes in UK Biobank.
*Nat Neurosci* **24**, 737-745 (2021).

42.    D. van der Meer, T. Kaufmann, A. A. Shadrin, C. Makowski, O. Frei, D. Roelfs, J. Monereo-
Sanchez, D. E. J. Linden, J. Rokicki, D. Alnaes, C. de Leeuw, W. K. Thompson, R. Loughnan,
C. C. Fan, L. T. Westlye, O. A. Andreassen, A. M. Dale, The genetic architecture of human
cortical folding. *Sci Adv* **7**, eabj9446 (2021).

43.    S. Bahrami, K. Nordengen, A. A. Shadrin, O. Frei, D. van der Meer, A. M. Dale, L. T. Westlye,
O. A. Andreassen, T. Kaufmann, Distributed genetic architecture across the hippocampal
formation implies common neuropathology across brain disorders. *Nat Commun* **13**, 3436
(2022).

44.    P. K. Albers, G. McVean, Dating genomic variants and shared ancestry in population-scale
sequencing data. *PLoS Biol* **18**, e3000586 (2020).

45.    I. Libedinsky, Y. Wei, C. d. Leeuw, J. Rilling, D. Posthuma, M. P. v. d. Heuvel, Genetic timeline
of human brain and cognitive traits. *bioRxiv*, 2023.2002.2005.525539 (2023).

46.    J. J. Hublin, A. Ben-Ncer, S. E. Bailey, S. E. Freidline, S. Neubauer, M. M. Skinner, I. Bergmann,
A. Le Cabec, S. Benazzi, K. Harvati, P. Gunz, New fossils from Jebel Irhoud, Morocco and the
pan-African origin of Homo sapiens. *Nature* **546**, 289-292 (2017).

47.    D. Richter, R. Grun, R. Joannes-Boyau, T. E. Steele, F. Amani, M. Rue, P. Fernandes, J. P.
Raynal, D. Geraads, A. Ben-Ncer, J. J. Hublin, S. P. McPherron, The age of the hominin fossils
from Jebel Irhoud, Morocco, and the origins of the Middle Stone Age. *Nature* **546**, 293-296
(2017).

48.    S. Davies, Behavioral Modernity in Retrospect. *Topoi* **40**, 221-232 (2021).

49.    C. A. de Leeuw, J. M. Mooij, T. Heskes, D. Posthuma, MAGMA: generalized gene-set analysis
of GWAS data. *PLoS Comput Biol* **11**, e1004219 (2015).

50.    Y. Wei, S. C. de Lange, L. H. Scholtens, K. Watanabe, D. J. Ardesch, P. R. Jansen, J. E. Savage,
L. Li, T. M. Preuss, J. K. Rilling, D. Posthuma, M. P. van den Heuvel, Genetic mapping and
evolutionary analysis of human-expanded cognitive networks. *Nat Commun* **10**, 4839 (2019).

51.    H. Won, J. Huang, C. K. Opland, C. L. Hartl, D. H. Geschwind, Human evolved regulatory
elements modulate genes involved in cortical expansion and neurodevelopmental disease
susceptibility. *Nat Commun* **10**, 2396 (2019).

52.    A. Liberzon, C. Birger, H. Thorvaldsdottir, M. Ghandi, J. P. Mesirov, P. Tamayo, The Molecular
Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst* **1**, 417-425 (2015).

53.    A. Subramanian, P. Tamayo, V. K. Mootha, S. Mukherjee, B. L. Ebert, M. A. Gillette, A.
Paulovich, S. L. Pomeroy, T. R. Golub, E. S. Lander, J. P. Mesirov, Gene set enrichment
analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc
Natl Acad Sci U S A* **102**, 15545-15550 (2005).

54.    M. Kircher, D. M. Witten, P. Jain, B. J. O'Roak, G. M. Cooper, J. Shendure, A general framework
for estimating the relative pathogenicity of human genetic variants. *Nat Genet* **46**, 310-315
(2014).

55. O. A. Andreassen, W. K. Thompson, A. J. Schork, S. Ripke, M. Mattingsdal, J. R. Kelsoe, K. S. Kendler, M. C. O'Donovan, D. Rujescu, T. Werge, P. Sklar, C. Psychiatric Genomics, D. Bipolar, G. Schizophrenia Working, J. C. Roddey, C. H. Chen, L. McEvoy, R. S. Desikan, S. Djurovic, A. M. Dale, Improved detection of common variants associated with schizophrenia and bipolar disorder using pleiotropy-informed conditional false discovery rate. *PLoS Genet* **9**, e1003455 (2013).

56. M. Z. King, Zonglei; Ivry, Richard B.; Weiner, Kevin S., Transcriptomic Gradients Of The Human Cerebellum.

57. X. Guell, J. D. E. Gabrieli, J. D. Schmahmann, Triple representation of language, working memory, social and emotion processing in the cerebellum: convergent evidence from task and seed-based resting-state fMRI analyses in a single large cohort. *Neuroimage* **172**, 437-449 (2018).

58. C. H. Chen, E. D. Gutierrez, W. Thompson, M. S. Panizzon, T. L. Jernigan, L. T. Eyler, C. Fennema-Notestine, A. J. Jak, M. C. Neale, C. E. Franz, M. J. Lyons, M. D. Grant, B. Fischl, L. J. Seidman, M. T. Tsuang, W. S. Kremen, A. M. Dale, Hierarchical genetic organization of human cortical surface area. *Science* **335**, 1634-1636 (2012).

59. C. H. Chen, M. Fiecas, E. D. Gutierrez, M. S. Panizzon, L. T. Eyler, E. Vuoksimaa, W. K. Thompson, C. Fennema-Notestine, D. J. Hagler, Jr., T. L. Jernigan, M. C. Neale, C. E. Franz, M. J. Lyons, B. Fischl, M. T. Tsuang, A. M. Dale, W. S. Kremen, Genetic topography of brain morphology. *Proc Natl Acad Sci U S A* **110**, 17089-17094 (2013).

60. C. Guissart, X. Latypova, P. Rollier, T. N. Khan, H. Stamberger, K. McWalter, M. T. Cho, S. Kjaergaard, S. Weckhuysen, G. Lesca, T. Besnard, K. Ounap, L. Schema, A. G. Chiocchetti, M. McDonald, J. de Bellescize, M. Vincent, H. Van Esch, S. Sattler, I. Forghani, I. Thiffault, C. M. Freitag, D. S. Barbouth, M. Cadieux-Dion, R. Willaert, M. J. Guillen Sacoto, N. P. Safina, C. Dubourg, L. Grote, W. Carre, C. Saunders, S. Pajusalu, E. Farrow, A. Boland, D. H. Karlowicz, J. F. Deleuze, M. H. Wojcik, R. Pressman, B. Isidor, A. Vogels, W. Van Paesschen, L. Al-Gazali, A. M. Al Shamsi, M. Claustres, A. Pujol, S. J. Sanders, F. Rivier, N. Leboucq, B. Cogne, S. Sasorith, D. Sanlaville, K. Retterer, S. Odent, N. Katsanis, S. Bezieau, M. Koenig, E. E. Davis, L. Pasquier, S. Kury, Dual Molecular Effects of Dominant RORA Mutations Cause Two Variants of Syndromic Intellectual Disability with Either Autism or Cerebellar Ataxia. *Am J Hum Genet* **102**, 744-759 (2018).

61. C. L. Chi, S. Martinez, W. Wurst, G. R. Martin, The isthmic organizer signal FGF8 is required for cell survival in the prospective midbrain and cerebellum. *Development* **130**, 2633-2644 (2003).

62. S. Li, F. Qiu, A. Xu, S. M. Price, M. Xiang, Barhl1 regulates migration and survival of cerebellar granule cells by controlling expression of the neurotrophin-3 gene. *J Neurosci* **24**, 3104-3114 (2004).

63. S. Valence, C. Garel, M. Barth, A. Toutain, C. Paris, D. Amsallem, M. A. Barthez, M. Mayer, D. Rodriguez, L. Burglen, RELN and VLDLR mutations underlie two distinguishable clinico-radiological phenotypes. *Clin Genet* **90**, 545-549 (2016).

64. M. Larouche, U. Beffert, J. Herz, R. Hawkes, The Reelin receptors Apoer2 and Vldlr coordinate the patterning of Purkinje cell topography in the developing mouse cerebellum. *PLoS One* **3**, e1653 (2008).

65. T. Wolfers, N. T. Doan, T. Kaufmann, D. Alnaes, T. Moberget, I. Agartz, J. K. Buitelaar, T. Ueland, I. Melle, B. Franke, O. A. Andreassen, C. F. Beckmann, L. T. Westlye, A. F. Marquand, Mapping the Heterogeneous Phenotype of Schizophrenia and Bipolar Disorder Using Normative Models. *JAMA Psychiatry* **75**, 1146-1155 (2018).

66. T. Wolfers, J. Rokicki, D. Alnaes, P. Berthet, I. Agartz, S. M. Kia, T. Kaufmann, M. Zabihi, T. Moberget, I. Melle, C. F. Beckmann, O. A. Andreassen, A. F. Marquand, L. T. Westlye, Replicating extensive brain structural heterogeneity in individuals with schizophrenia and bipolar disorder. *Hum Brain Mapp* **42**, 2546-2555 (2021).

67. N. C. Armstrong, R. C. Anderson, K. W. McDermott, Reelin: Diverse roles in central nervous system development, health and disease. *Int J Biochem Cell Biol* **112**, 72-75 (2019).

68. M. Li, L. Huang, M. Grigoroiu-Serbanescu, S. E. Bergen, M. Landen, C. M. Hultman, A. J.
Forstner, J. Strohmaier, J. Hecker, T. G. Schulze, B. Muller-Myhsok, A. Reif, P. B. Mitchell, N.
G. Martin, S. Cichon, M. M. Nothen, A. Alkelai, B. Lerer, S. Jamain, M. Leboyer, F. Bellivier, B.
Etain, J. P. Kahn, C. Henry, M. Rietschel, D. S. C. Moo, G. Swedish Bipolar Study, Convergent
Lines of Evidence Support LRP8 as a Susceptibility Gene for Psychosis. *Mol Neurobiol* **53**, 6608-
6619 (2016).
69. A. Okbay, Y. Wu, N. Wang, H. Jayashankar, M. Bennett, S. M. Nehzati, J. Sidorenko, H. Kweon,
G. Goldman, T. Gjorgjieva, Y. Jiang, B. Hicks, C. Tian, D. A. Hinds, R. Ahlskog, P. K. E.
Magnusson, S. Oskarsson, C. Hayward, A. Campbell, D. J. Porteous, J. Freese, P. Herd, T.
andMe Research, C. Social Science Genetic Association, C. Watson, J. Jala, D. Conley, P. D.
Koellinger, M. Johannesson, D. Laibson, M. N. Meyer, J. J. Lee, A. Kong, L. Yengo, D. Cesarini,
P. Turley, P. M. Visscher, J. P. Beauchamp, D. J. Benjamin, A. I. Young, Polygenic prediction of
educational attainment within and between families from genome-wide association analyses in
3 million individuals. *Nat Genet* **54**, 437-449 (2022).
70. Q. Wang, R. S. Dhindsa, K. Carss, A. R. Harper, A. Nag, I. Tachmazidou, D. Vitsios, S. V. V.
Deevi, A. Mackay, D. Muthas, M. Huhn, S. Monkley, H. Olsson, I. AstraZeneca Genomics, S.
Wasilewski, K. R. Smith, R. March, A. Platt, C. Haefliger, S. Petrovski, Rare variant contribution
to human disease in 281,104 UK Biobank exomes. *Nature* **597**, 527-532 (2021).
71. B. Fischl, D. H. Salat, E. Busa, M. Albert, M. Dieterich, C. Haselgrove, A. van der Kouwe, R.
Killiany, D. Kennedy, S. Klaveness, A. Montillo, N. Makris, B. Rosen, A. M. Dale, Whole brain
segmentation: automated labeling of neuroanatomical structures in the human brain. *Neuron* **33**,
341-355 (2002).
72. J. Diedrichsen, in *NeuroImage*. (2006), vol. 33, pp. 127-138.
73. C. Leys, C. Ley, O. Klein, P. Bernard, L. Licata, Detecting outliers: Do not use standard deviation
around the mean, use absolute deviation around the median. *Journal of Experimental Social
Psychology* **49**, 764-766 (2013).
74. L. W. de Jong, J. S. Vidal, L. E. Forsberg, A. P. Zijdenbos, T. Haight, I. Alzheimer's Disease
Neuroimaging, S. Sigurdsson, V. Gudnason, M. A. van Buchem, L. J. Launer, Allometric scaling
of brain regions to intra-cranial volume: An epidemiological MRI study. *Hum Brain Mapp* **38**, 151-
164 (2017).
75. C. Mankiw, M. T. M. Park, P. K. Reardon, A. M. Fish, L. S. Clasen, D. Greenstein, J. N. Giedd,
J. D. Blumenthal, J. P. Lerch, M. M. Chakravarty, A. Raznahan, Allometric Analysis Detects Brain
Size-Independent Effects of Sex and Sex Chromosome Complement on Human Cerebellar
Organization. *J Neurosci* **37**, 5221-5231 (2017).
76. K. J. Friston, *Statistical parametric mapping : the analysis of funtional brain images*
(Elsevier/Academic Press, Amsterdam ; Boston, ed. 1st, 2007), pp. vii, 647 p.
77. R. Nassar, A. N. Kaczkurkin, C. H. Xia, A. Sotiras, M. Pehlivanova, T. M. Moore, A. Garcia de
La Garza, D. R. Roalf, A. F. G. Rosen, S. A. Lorch, K. Ruparel, R. T. Shinohara, C. Davatzikos,
R. C. Gur, R. E. Gur, T. D. Satterthwaite, Gestational Age is Dimensionally Associated with
Structural Brain Network Abnormalities Across Development. *Cereb Cortex* **29**, 2102-2114
(2019).
78. M. Pehlivanova, D. H. Wolf, A. Sotiras, A. N. Kaczkurkin, T. M. Moore, R. Ciric, P. A. Cook, A.
Garcia de La Garza, A. F. G. Rosen, K. Ruparel, A. Sharma, R. T. Shinohara, D. R. Roalf, R. C.
Gur, C. Davatzikos, R. E. Gur, J. W. Kable, T. D. Satterthwaite, Diminished Cortical Thickness
Is Associated with Impulsive Choice in Adolescence. *J Neurosci* **38**, 2471-2481 (2018).
79. H. W. Kuhn, The Hungarian method for the assignment problem. *Naval Research Logistics
Quarterly* **2**, 14 (1955).
80. J. M. Santos, M. Embrechts. (Springer Berlin Heidelberg, Berlin, Heidelberg, 2009), pp. 175-184.
81. T. M. Beasley, S. Erickson, D. B. Allison, Rank-based inverse normal transformations are
increasingly used, but are they merited? *Behav Genet* **39**, 580-595 (2009).
82. C. Bycroft, C. Freeman, D. Petkova, G. Band, L. T. Elliott, K. Sharp, A. Motyer, D. Vukcevic, O.
Delaneau, J. O'Connell, A. Cortes, S. Welsh, A. Young, M. Effingham, G. McVean, S. Leslie, N.

| | | |
|---|---|---|
| | | Allen, P. Donnelly, J. Marchini, The UK Biobank resource with deep phenotyping and genomic data. *Nature* **562**, 203-209 (2018). |
| 83. | | A. P. Boyle, E. L. Hong, M. Hariharan, Y. Cheng, M. A. Schaub, M. Kasowski, K. J. Karczewski, J. Park, B. C. Hitz, S. Weng, J. M. Cherry, M. Snyder, Annotation of functional variation in personal genomes using RegulomeDB. *Genome Res* **22**, 1790-1797 (2012). |
| 84. | | Z. Zhu, F. Zhang, H. Hu, A. Bakshi, M. R. Robinson, J. E. Powell, G. W. Montgomery, M. E. Goddard, N. R. Wray, P. M. Visscher, J. Yang, Integration of summary data from GWAS and eQTL studies predicts complex trait gene targets. *Nat Genet* **48**, 481-487 (2016). |
| 85. | | A. Arnatkeviciute, B. D. Fulcher, A. Fornito, A practical guide to linking brain-wide gene expression and neuroimaging data. *Neuroimage* **189**, 353-367 (2019). |
| 86. | | J. Quackenbush, Microarray data normalization and transformation. *Nat Genet* **32 Suppl**, 496-501 (2002). |
| 87. | | M. Hawrylycz, J. A. Miller, V. Menon, D. Feng, T. Dolbeare, A. L. Guillozet-Bongaarts, A. G. Jegga, B. J. Aronow, C. K. Lee, A. Bernard, M. F. Glasser, D. L. Dierker, J. Menche, A. Szafer, F. Collman, P. Grange, K. A. Berman, S. Mihalas, Z. Yao, L. Stewart, A. L. Barabasi, J. Schulkin, J. Phillips, L. Ng, C. Dang, D. R. Haynor, A. Jones, D. C. Van Essen, C. Koch, E. Lein, Canonical genetic signatures of the adult human brain. *Nat Neurosci* **18**, 1832-1844 (2015). |
| 88. | | B. D. Fulcher, M. A. Little, N. S. Jones, Highly comparative time-series analysis: the empirical structure of time series and their methods. *J R Soc Interface* **10**, 20130048 (2013). |
| 89. | | C. Genomes Project, A. Auton, L. D. Brooks, R. M. Durbin, E. P. Garrison, H. M. Kang, J. O. Korbel, J. L. Marchini, S. McCarthy, G. A. McVean, G. R. Abecasis, A global reference for human genetic variation. *Nature* **526**, 68-74 (2015). |
| 90. | | S. Mallick, H. Li, M. Lipson, I. Mathieson, M. Gymrek, F. Racimo, M. Zhao, N. Chennagiri, S. Nordenfelt, A. Tandon, P. Skoglund, I. Lazaridis, S. Sankararaman, Q. Fu, N. Rohland, G. Renaud, Y. Erlich, T. Willems, C. Gallo, J. P. Spence, Y. S. Song, G. Poletti, F. Balloux, G. van Driem, P. de Knijff, I. G. Romero, A. R. Jha, D. M. Behar, C. M. Bravi, C. Capelli, T. Hervig, A. Moreno-Estrada, O. L. Posukh, E. Balanovska, O. Balanovsky, S. Karachanak-Yankova, H. Sahakyan, D. Toncheva, L. Yepiskoposyan, C. Tyler-Smith, Y. Xue, M. S. Abdullah, A. Ruiz-Linares, C. M. Beall, A. Di Rienzo, C. Jeong, E. B. Starikovskaya, E. Metspalu, J. Parik, R. Villems, B. M. Henn, U. Hodoglugil, R. Mahley, A. Sajantila, G. Stamatoyannopoulos, J. T. Wee, R. Khusainova, E. Khusnutdinova, S. Litvinov, G. Ayodo, D. Comas, M. F. Hammer, T. Kivisild, W. Klitz, C. A. Winkler, D. Labuda, M. Bamshad, L. B. Jorde, S. A. Tishkoff, W. S. Watkins, M. Metspalu, S. Dryomov, R. Sukernik, L. Singh, K. Thangaraj, S. Paabo, J. Kelso, N. Patterson, D. Reich, The Simons Genome Diversity Project: 300 genomes from 142 diverse populations. *Nature* **538**, 201-206 (2016). |
| 91. | | A. Federico, S. Monti, hypeR: an R package for geneset enrichment workflows. *Bioinformatics* **36**, 1307-1308 (2020). |
| 92. | | A. D. Rouillard, G. W. Gundersen, N. F. Fernandez, Z. Wang, C. D. Monteiro, M. G. McDermott, A. Ma'ayan, The harmonizome: a collection of processed datasets gathered to serve and mine knowledge about genes and proteins. *Database (Oxford)* **2016**, (2016). |
| 93. | | R. N. Doan, B. I. Bae, B. Cubelos, C. Chang, A. A. Hossain, S. Al-Saad, N. M. Mukaddes, O. Oner, M. Al-Saffar, S. Balkhy, G. G. Gascon, A. Homozygosity Mapping Consortium for, M. Nieto, C. A. Walsh, Mutations in Human Accelerated Regions Disrupt Cognition and Social Behavior. *Cell* **167**, 341-354 e312 (2016). |
| 94. | | D. Demontis, R. K. Walters, J. Martin, M. Mattheisen, T. D. Als, E. Agerbo, G. Baldursson, R. Belliveau, J. Bybjerg-Grauholm, M. Baekvad-Hansen, F. Cerrato, K. Chambert, C. Churchhouse, A. Dumont, N. Eriksson, M. Gandal, J. I. Goldstein, K. L. Grasby, J. Grove, O. O. Gudmundsson, C. S. Hansen, M. E. Hauberg, M. V. Hollegaard, D. P. Howrigan, H. Huang, J. B. Maller, A. R. Martin, N. G. Martin, J. Moran, J. Pallesen, D. S. Palmer, C. B. Pedersen, M. G. Pedersen, T. Poterba, J. B. Poulsen, S. Ripke, E. B. Robinson, F. K. Satterstrom, H. Stefansson, C. Stevens, P. Turley, G. B. Walters, H. Won, M. J. Wright, A. W. G. o. t. P. G. Consortium, L. Early, C. Genetic Epidemiology, T. andMe Research, O. A. Andreassen, P. Asherson, C. L. Burton, D. I. Boomsma, B. Cormand, S. Dalsgaard, B. Franke, J. Gelernter, D. Geschwind, H. Hakonarson, |

J. Haavik, H. R. Kranzler, J. Kuntsi, K. Langley, K. P. Lesch, C. Middeldorp, A. Reif, L. A. Rohde, P. Roussos, R. Schachar, P. Sklar, E. J. S. Sonuga-Barke, P. F. Sullivan, A. Thapar, J. Y. Tung, I. D. Waldman, S. E. Medland, K. Stefansson, M. Nordentoft, D. M. Hougaard, T. Werge, O. Mors, P. B. Mortensen, M. J. Daly, S. V. Faraone, A. D. Borglum, B. M. Neale, Discovery of the first genome-wide significant risk loci for attention deficit/hyperactivity disorder. *Nat Genet* **51**, 63-75 (2019).

95.	J. Grove, S. Ripke, T. D. Als, M. Mattheisen, R. K. Walters, H. Won, J. Pallesen, E. Agerbo, O. A. Andreassen, R. Anney, S. Awashti, R. Belliveau, F. Bettella, J. D. Buxbaum, J. Bybjerg-Grauholm, M. Baekvad-Hansen, F. Cerrato, K. Chambert, J. H. Christensen, C. Churchhouse, K. Dellenvall, D. Demontis, S. De Rubeis, B. Devlin, S. Djurovic, A. L. Dumont, J. I. Goldstein, C. S. Hansen, M. E. Hauberg, M. V. Hollegaard, S. Hope, D. P. Howrigan, H. Huang, C. M. Hultman, L. Klei, J. Maller, J. Martin, A. R. Martin, J. L. Moran, M. Nyegaard, T. Naerland, D. S. Palmer, A. Palotie, C. B. Pedersen, M. G. Pedersen, T. dPoterba, J. B. Poulsen, B. S. Pourcain, P. Qvist, K. Rehnstrom, A. Reichenberg, J. Reichert, E. B. Robinson, K. Roeder, P. Roussos, E. Saemundsen, S. Sandin, F. K. Satterstrom, G. Davey Smith, H. Stefansson, S. Steinberg, C. R. Stevens, P. F. Sullivan, P. Turley, G. B. Walters, X. Xu, C. Autism Spectrum Disorder Working Group of the Psychiatric Genomics, Bupgen, C. Major Depressive Disorder Working Group of the Psychiatric Genomics, T. andMe Research, K. Stefansson, D. H. Geschwind, M. Nordentoft, D. M. Hougaard, T. Werge, O. Mors, P. B. Mortensen, B. M. Neale, M. J. Daly, A. D. Borglum, Identification of common genetic risk variants for autism spectrum disorder. *Nat Genet* **51**, 431-444 (2019).

96.	N. Mullins, A. J. Forstner, K. S. O'Connell, B. Coombes, J. R. I. Coleman, Z. Qiao, T. D. Als, T. B. Bigdeli, S. Borte, J. Bryois, A. W. Charney, O. K. Drange, M. J. Gandal, S. P. Hagenaars, M. Ikeda, N. Kamitaki, M. Kim, K. Krebs, G. Panagiotaropoulou, B. M. Schilder, L. G. Sloofman, S. Steinberg, V. Trubetskoy, B. S. Winsvold, H. H. Won, L. Abramova, K. Adorjan, E. Agerbo, M. Al Eissa, D. Albani, N. Alliey-Rodriguez, A. Anjorin, V. Antilla, A. Antoniou, S. Awasthi, J. H. Baek, M. Baekvad-Hansen, N. Bass, M. Bauer, E. C. Beins, S. E. Bergen, A. Birner, C. Bocker Pedersen, E. Boen, M. P. Boks, R. Bosch, M. Brum, B. M. Brumpton, N. Brunkhorst-Kanaan, M. Budde, J. Bybjerg-Grauholm, W. Byerley, M. Cairns, M. Casas, P. Cervantes, T. K. Clarke, C. Cruceanu, A. Cuellar-Barboza, J. Cunningham, D. Curtis, P. M. Czerski, A. M. Dale, N. Dalkner, F. S. David, F. Degenhardt, S. Djurovic, A. L. Dobbyn, A. Douzenis, T. Elvsashagen, V. Escott-Price, I. N. Ferrier, A. Fiorentino, T. M. Foroud, L. Forty, J. Frank, O. Frei, N. B. Freimer, L. Frisen, K. Gade, J. Garnham, J. Gelernter, M. Giortz Pedersen, I. R. Gizer, S. D. Gordon, K. Gordon-Smith, T. A. Greenwood, J. Grove, J. Guzman-Parra, K. Ha, M. Haraldsson, M. Hautzinger, U. Heilbronner, D. Hellgren, S. Herms, P. Hoffmann, P. A. Holmans, L. Huckins, S. Jamain, J. S. Johnson, J. L. Kalman, Y. Kamatani, J. L. Kennedy, S. Kittel-Schneider, J. A. Knowles, M. Kogevinas, M. Koromina, T. M. Kranz, H. R. Kranzler, M. Kubo, R. Kupka, S. A. Kushner, C. Lavebratt, J. Lawrence, M. Leber, H. J. Lee, P. H. Lee, S. E. Levy, C. Lewis, C. Liao, S. Lucae, M. Lundberg, D. J. MacIntyre, S. H. Magnusson, W. Maier, A. Maihofer, D. Malaspina, E. Maratou, L. Martinsson, M. Mattheisen, S. A. McCarroll, N. W. McGregor, P. McGuffin, J. D. McKay, H. Medeiros, S. E. Medland, V. Millischer, G. W. Montgomery, J. L. Moran, D. W. Morris, T. W. Muhleisen, N. O'Brien, C. O'Donovan, L. M. Olde Loohuis, L. Oruc, S. Papiol, A. F. Pardinas, A. Perry, A. Pfennig, E. Porichi, J. B. Potash, D. Quested, T. Raj, M. H. Rapaport, J. R. DePaulo, E. J. Regeer, J. P. Rice, F. Rivas, M. Rivera, J. Roth, P. Roussos, D. M. Ruderfer, C. Sanchez-Mora, E. C. Schulte, F. Senner, S. Sharp, P. D. Shilling, E. Sigurdsson, L. Sirignano, C. Slaney, O. B. Smeland, D. J. Smith, J. L. Sobell, C. Soholm Hansen, M. Soler Artigas, A. T. Spijker, D. J. Stein, J. S. Strauss, B. Swiatkowska, C. Terao, T. E. Thorgeirsson, C. Toma, P. Tooney, E. E. Tsermpini, M. P. Vawter, H. Vedder, J. T. R. Walters, S. H. Witt, S. Xi, W. Xu, J. M. K. Yang, A. H. Young, H. Young, P. P. Zandi, H. Zhou, L. Zillich, H. A.-I. Psychiatry, R. Adolfsson, I. Agartz, M. Alda, L. Alfredsson, G. Babadjanova, L. Backlund, B. T. Baune, F. Bellivier, S. Bengesser, W. H. Berrettini, D. H. R. Blackwood, M. Boehnke, A. D. Borglum, G. Breen, V. J. Carr, S. Catts, A. Corvin, N. Craddock, U. Dannlowski, D. Dikeos, T. Esko, B. Etain, P. Ferentinos, M. Frye, J. M. Fullerton, M. Gawlik, E. S. Gershon, F. S. Goes, M.

J. Green, M. Grigoroiu-Serbanescu, J. Hauser, F. Henskens, J. Hillert, K. S. Hong, D. M. Hougaard, C. M. Hultman, K. Hveem, N. Iwata, A. V. Jablensky, I. Jones, L. A. Jones, R. S. Kahn, J. R. Kelsoe, G. Kirov, M. Landen, M. Leboyer, C. M. Lewis, Q. S. Li, J. Lissowska, C. Lochner, C. Loughland, N. G. Martin, C. A. Mathews, F. Mayoral, S. L. McElroy, A. M. McIntosh, F. J. McMahon, I. Melle, P. Michie, L. Milani, P. B. Mitchell, G. Morken, O. Mors, P. B. Mortensen, B. Mowry, B. Muller-Myhsok, R. M. Myers, B. M. Neale, C. M. Nievergelt, M. Nordentoft, M. M. Nothen, M. C. O'Donovan, K. J. Oedegaard, T. Olsson, M. J. Owen, S. A. Paciga, C. Pantelis, C. Pato, M. T. Pato, G. P. Patrinos, R. H. Perlis, D. Posthuma, J. A. Ramos-Quiroga, A. Reif, E. Z. Reininghaus, M. Ribases, M. Rietschel, S. Ripke, G. A. Rouleau, T. Saito, U. Schall, M. Schalling, P. R. Schofield, T. G. Schulze, L. J. Scott, R. J. Scott, A. Serretti, C. Shannon Weickert, J. W. Smoller, H. Stefansson, K. Stefansson, E. Stordal, F. Streit, P. F. Sullivan, G. Turecki, A. E. Vaaler, E. Vieta, J. B. Vincent, I. D. Waldman, T. W. Weickert, T. Werge, N. R. Wray, J. A. Zwart, J. M. Biernacka, J. I. Nurnberger, S. Cichon, H. J. Edenberg, E. A. Stahl, A. McQuillin, A. Di Florio, R. A. Ophoff, O. A. Andreassen, Genome-wide association study of more than 40,000 bipolar disorder cases provides new insights into the underlying biology. *Nat Genet* **53**, 817-829 (2021).

97. N. R. Wray, S. Ripke, M. Mattheisen, M. Trzaskowski, E. M. Byrne, A. Abdellaoui, M. J. Adams, E. Agerbo, T. M. Air, T. M. F. Andlauer, S. A. Bacanu, M. Baekvad-Hansen, A. F. T. Beekman, T. B. Bigdeli, E. B. Binder, D. R. H. Blackwood, J. Bryois, H. N. Buttenschon, J. Bybjerg-Grauholm, N. Cai, E. Castelao, J. H. Christensen, T. K. Clarke, J. I. R. Coleman, L. Colodro-Conde, B. Couvy-Duchesne, N. Craddock, G. E. Crawford, C. A. Crowley, H. S. Dashti, G. Davies, I. J. Deary, F. Degenhardt, E. M. Derks, N. Direk, C. V. Dolan, E. C. Dunn, T. C. Eley, N. Eriksson, V. Escott-Price, F. H. F. Kiadeh, H. K. Finucane, A. J. Forstner, J. Frank, H. A. Gaspar, M. Gill, P. Giusti-Rodriguez, F. S. Goes, S. D. Gordon, J. Grove, L. S. Hall, E. Hannon, C. S. Hansen, T. F. Hansen, S. Herms, I. B. Hickie, P. Hoffmann, G. Homuth, C. Horn, J. J. Hottenga, D. M. Hougaard, M. Hu, C. L. Hyde, M. Ising, R. Jansen, F. Jin, E. Jorgenson, J. A. Knowles, I. S. Kohane, J. Kraft, W. W. Kretzschmar, J. Krogh, Z. Kutalik, J. M. Lane, Y. Li, Y. Li, P. A. Lind, X. Liu, L. Lu, D. J. MacIntyre, D. F. MacKinnon, R. M. Maier, W. Maier, J. Marchini, H. Mbarek, P. McGrath, P. McGuffin, S. E. Medland, D. Mehta, C. M. Middeldorp, E. Mihailov, Y. Milaneschi, L. Milani, J. Mill, F. M. Mondimore, G. W. Montgomery, S. Mostafavi, N. Mullins, M. Nauck, B. Ng, M. G. Nivard, D. R. Nyholt, P. F. O'Reilly, H. Oskarsson, M. J. Owen, J. N. Painter, C. B. Pedersen, M. G. Pedersen, R. E. Peterson, E. Pettersson, W. J. Peyrot, G. Pistis, D. Posthuma, S. M. Purcell, J. A. Quiroz, P. Qvist, J. P. Rice, B. P. Riley, M. Rivera, S. Saeed Mirza, R. Saxena, R. Schoevers, E. C. Schulte, L. Shen, J. Shi, S. I. Shyn, E. Sigurdsson, G. B. C. Sinnamon, J. H. Smit, D. J. Smith, H. Stefansson, S. Steinberg, C. A. Stockmeier, F. Streit, J. Strohmaier, K. E. Tansey, H. Teismann, A. Teumer, W. Thompson, P. A. Thomson, T. E. Thorgeirsson, C. Tian, M. Traylor, J. Treutlein, V. Trubetskoy, A. G. Uitterlinden, D. Umbricht, S. Van der Auwera, A. M. van Hemert, A. Viktorin, P. M. Visscher, Y. Wang, B. T. Webb, S. M. Weinsheimer, J. Wellmann, G. Willemsen, S. H. Witt, Y. Wu, H. S. Xi, J. Yang, F. Zhang, eQtlgen, andMe, V. Arolt, B. T. Baune, K. Berger, D. I. Boomsma, S. Cichon, U. Dannlowski, E. C. J. de Geus, J. R. DePaulo, E. Domenici, K. Domschke, T. Esko, H. J. Grabe, S. P. Hamilton, C. Hayward, A. C. Heath, D. A. Hinds, K. S. Kendler, S. Kloiber, G. Lewis, Q. S. Li, S. Lucae, P. F. A. Madden, P. K. Magnusson, N. G. Martin, A. M. McIntosh, A. Metspalu, O. Mors, P. B. Mortensen, B. Muller-Myhsok, M. Nordentoft, M. M. Nothen, M. C. O'Donovan, S. A. Paciga, N. L. Pedersen, B. Penninx, R. H. Perlis, D. J. Porteous, J. B. Potash, M. Preisig, M. Rietschel, C. Schaefer, T. G. Schulze, J. W. Smoller, K. Stefansson, H. Tiemeier, R. Uher, H. Volzke, M. M. Weissman, T. Werge, A. R. Winslow, C. M. Lewis, D. F. Levinson, G. Breen, A. D. Borglum, P. F. Sullivan, C. Major Depressive Disorder Working Group of the Psychiatric Genomics, Genome-wide association analyses identify 44 risk variants and refine the genetic architecture of major depression. *Nat Genet* **50**, 668-681 (2018).

98. A. F. Pardinas, P. Holmans, A. J. Pocklington, V. Escott-Price, S. Ripke, N. Carrera, S. E. Legge, S. Bishop, D. Cameron, M. L. Hamshere, J. Han, L. Hubbard, A. Lynham, K. Mantripragada, E. Rees, J. H. MacCabe, S. A. McCarroll, B. T. Baune, G. Breen, E. M. Byrne, U. Dannlowski, T.

C. Eley, C. Hayward, N. G. Martin, A. M. McIntosh, R. Plomin, D. J. Porteous, N. R. Wray, A. Caballero, D. H. Geschwind, L. M. Huckins, D. M. Ruderfer, E. Santiago, P. Sklar, E. A. Stahl, H. Won, E. Agerbo, T. D. Als, O. A. Andreassen, M. Baekvad-Hansen, P. B. Mortensen, C. B. Pedersen, A. D. Borglum, J. Bybjerg-Grauholm, S. Djurovic, N. Durmishi, M. G. Pedersen, V. Golimbet, J. Grove, D. M. Hougaard, M. Mattheisen, E. Molden, O. Mors, M. Nordentoft, M. Pejovic-Milovancevic, E. Sigurdsson, T. Silagadze, C. S. Hansen, K. Stefansson, H. Stefansson, S. Steinberg, S. Tosato, T. Werge, G. Consortium, C. Consortium, D. A. Collier, D. Rujescu, G. Kirov, M. J. Owen, M. C. O'Donovan, J. T. R. Walters, Common schizophrenia alleles are enriched in mutation-intolerant genes and in regions under strong background selection. *Nat Genet* **50**, 381-389 (2018).

99.  O. A. Andreassen, S. Djurovic, W. K. Thompson, A. J. Schork, K. S. Kendler, M. C. O'Donovan, D. Rujescu, T. Werge, M. van de Bunt, A. P. Morris, M. I. McCarthy, G. International Consortium for Blood Pressure, R. Diabetes Genetics, C. Meta-analysis, G. Psychiatric Genomics Consortium Schizophrenia Working, J. C. Roddey, L. K. McEvoy, R. S. Desikan, A. M. Dale, Improved detection of common variants associated with schizophrenia by leveraging pleiotropy with cardiovascular-disease risk factors. *Am J Hum Genet* **92**, 197-209 (2013).

100. O. B. Smeland, O. Frei, A. Shadrin, K. O'Connell, C. C. Fan, S. Bahrami, D. Holland, S. Djurovic, W. K. Thompson, A. M. Dale, O. A. Andreassen, Discovery of shared genomic loci using the conditional false discovery rate approach. *Hum Genet* **139**, 85-94 (2020).

**Acknowledgements**

This work was performed on the Tjeneste for Sensitive Data (TSD) facilities, owned by the University of Oslo, operated and developed by the TSD service group at the University of Oslo, IT-Department (USIT) and on resources provided by UNINETT Sigma2—the National Infrastructure for High Performance Computing and Data Storage in Norway. The research has been conducted using the UK Biobank Resource (access code 27412) and using summary statistics for various brain disorders. We would like to thank the research participants and employees of UK Biobank and the consortia contributing summary statistics for making this work possible.

**Funding:** The authors were funded by the South-Eastern Norway Regional Health Authority (TM: 2021-040; OAA: 2013-123, 2017-112, 2019-108. LTW: 2014-097, 2015-073, 2016-083), the Research Council of Norway (TK: 276082, 323961. OAA: 213837, 223273, 248778, 273291, 262656, 229129, 283798, 311993. LTW: 204966, 249795, 273345), Norwegian Health Association (SB: 22731), Stiftelsen Kristian Gerhard Jebsen, and the European Research Council (LTW: ERCStG 802998). The funding bodies had no role in the analysis or interpretation of the data; the preparation, review or approval of the manuscript; nor in the decision to submit the manuscript for publication.

**Author contributions:** TM originally conceived of the project. TM, DvdM and SB performed the analyses. TM, LTW and OAA wrote the initial draft of the manuscript. All authors discussed the results and contributed to the final manuscript.

**Competing interests:** O.A.A. has received speaker's honorarium from Lundbeck and is a consultant to HealthLytix. A.M.D. is the founder of and holds equity in CorTechs Labs Inc. and serves on its Scientific Advisory Board. A.M.D. is also a member of the Scientific Advisory Board of Human Longevity Inc. and receives funding through research agreements with General Electric Healthcare and Medtronic Inc. The terms of these arrangements have been reviewed and approved by UCSD in accordance with its conflict-of-interest policies. A.M.D. is an inventor on a patent related to this work, filed by CorTechs Labs Inc. (9 US-7324842B2, filed 22 January 2002, published 29-01-2008). The other authors declare that they have no competing interests.

## Data availability

In this study we used brain imaging and genetics data from the UK Biobank [https://www.ukbiobank.ac.uk/], and GWAS summary statistics obtained from the Psychiatric Genomics Consortium [https://www.med.unc.edu/pgc/shared-methods/] and GWAS catalog (https://www.ebi.ac.uk/gwas/). The summary statistics for cerebellar morphology derived in this study are available in our github repository [https://github.com/norment/open-science] (*will make this public upon acceptance*). FUMA results are available online (*will make this public upon acceptance*). The Human Genome Dating Dataset (HGD) is available at https://human.genome.dating.

## Code availability

All code and software needed to generate the results is available as part of public resources, specifically FreeSurfer (https://surfer.nmr.mgh.harvard.edu), SUIT (https://github.com/jdiedrichsen/suit), OPNMF (https://github.com/asotiras/brainparts), MOSTest (https://github.com/precimed/mostest), FUMA (https://fuma.ctglab.nl/), MAGMA (https://ctg.cncr.nl/software/magma), abagen (https://github.com/rmarkello/abagen), conjunctional FDR (https://github.com/precimed/pleiofdr/) and LD score regression (https://github.com/bulik/ldsc).

**Supplementary Materials:**

1. Supplementary Figures
2. Supplementary Information – Data Tables 1-36