Public toilets have reduced enteric pathogen hazards in San Francisco

Troy Barker[1]*, Drew Capone[2]*, Heather K. Amato[3], Ryan Clark[1], Abigail Henderson[3], David A. Holcomb[1], Elizabeth Kim[1], Jillian Pape[3], Emily Parker[3], Thomas VanderYacht[3], Jay Graham[3], and Joe Brown[1]**

[1]Department of Environmental Sciences and Engineering, Gillings School of Global Public Health, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

[2]Department of Environmental and Occupational Health, School of Public Health, Indiana University Bloomington, Bloomington, IN, USA

[3]Department of Environmental Health, School of Public Health, University of California – Berkeley, Berkeley, CA, USA

*These authors contributed equally to the manuscript.

**Corresponding author: Joe Brown, Department of Environmental Sciences and Engineering, Gillings School of Global Public Health, University of North Carolina at Chapel Hill. CB7431 Rosenau Hall, Chapel Hill, NC 27599-7431, USA. Tel: +1 919 360 8752. Email: joebrown@unc.edu

KEYWORDS: homelessness, urban sanitation, mtDNA, microbial hazards, environmental exposures, environmental microbiology, dPCR, qPCR

SYNOPSIS: This paper describes enteric pathogen hazards from discarded feces on the streets

of San Francisco and estimates their reduction following a public toilet intervention.

TOC/Abstract art:

Created with BioRender and a photograph by author Jay Graham

ABSTRACT

Uncontained fecal wastes in cities may present exposure risks to the public. We collected

discarded feces from public spaces in San Francisco for analysis by RT-qPCR for a range of

enteric pathogens. Out of 59 samples, we found 12 (20%) were of human origin and 47 (80%)

were non-human; 30 of 59 stools were positive for ≥1 of the 35 pathogens assessed, including

pathogenic *E. coli*, *Shigella*, norovirus, *Cryptosporidium*, and *Trichuris*. Using quantitative

enteric pathogen estimates and data on observed fecal waste from a public reporting system, we

modeled pathogens removed from the environment attributable to a recently implemented

program of public toilet construction. We estimated that each new public toilet reduced the

annual number of enteric pathogens released into the immediate environment (within 500 m

walking distance), including 6.3×10^{12} enteropathogenic *E. coli* (95% CI: 4.0×10^{12} – $7.9 \times$

10^{12}), 3.2 x 10^{11} enteroaggregative *E. coli* (95% CI: 1.3 x 10^{11} – 6.3 x 10^{11}), and 3.2 x 10^{8}

Shigella (6.3 x 10^{7} – 2.5 x 10^{9}). Improving access to public sanitation can reduce enteric

pathogen hazards in cities. Interventions must also consider the hygienic disposal of animal

waste to reduce microbial hazards with zoonotic infection potential.

INTRODUCTION

People experiencing homelessness are more likely than the general population to suffer from

communicable diseases[1], partly because they lack consistent access to basic infrastructure

including safe water and sanitation[2, 3]. Nearly one million persons in US cities have insufficient

access to basic sanitation and over 600,000 cannot consistently access basic water infrastructure[2],

primarily those lacking stable housing.

Nearly 130,000 people in California experience homelessness each day, and they are

disproportionately unsheltered compared to those in other states[1, 4]. Like many US cities, San

Francisco does not have enough public toilets to meet demand[5]. Consistent access to sanitation

and soap and water for proper handwashing is necessary to prevent the spread of enteric

pathogens that may cause diarrheal disease[6]. Without safe, hygienic, and publicly accessible

toilets when and where people need them, open defecation is common[7-9]. As a result,

uncontained fecal waste can accumulate near where people live, work, and play, creating

opportunities for exposure to enteric pathogens through well-known direct and indirect

pathways[10-13].

Fecal contamination on the streets of San Francisco is so common that the Department of Public

Works (DPW) created a well-used system to report feces in public spaces, where instances of

fecal waste can be reported by dialing 311 from a telephone (or, now, via a website or

Twitter)[11,14]. Data since 2008 are publicly available at https://datasf.org/opendata/. As a step

toward addressing the well-publicized problem, DPW implemented the Pit Stop program

beginning in 2014, aiming to reduce open defecation in public spaces by installing staffed public

toilets in areas of high need.[11,12] The intervention also includes animal waste bags and waste bins

for disposal. A full description of the program and its history is available at the Pit Stop website:

https://sfpublicworks.wixsite.com/pitstop. A recent impact assessment of the program estimated

that the installation of public latrines reduced 311 reports of fecal waste on the street by a mean

of 12.5 stools per week within 500 meters (walking distance) of newly installed Pit Stop

locations in the six months following installation, compared with a pre-intervention baseline[15].

The greatest reduction in reports of fecal waste occurred in the Tenderloin neighborhood (i.e., 18

stools per week per new facility), which had 10 public toilets, the most of any neighborhood in

the program. The Tenderloin and South of Market (SoMa) had the highest counts of 311 reports

in the study period; together, these neighborhoods make up District 6, which had the highest

count of people experiencing homelessness (n=3,656) in SF as of 2019[16]. With only 3 Pit Stop

public facilities in SoMa, there was no significant reduction of 311 reports associated with the Pit

Stop intervention in this neighborhood. A recent independent analysis by the San Francisco

Chronicle of 311 data concluded that, while feces-related service requests for every other

neighborhood in San Francisco have gone up an average of 400% in the period from 2012-2021,

reports for the Tenderloin reduced by 29%, with pronounced further decreases in the area

immediately around three Pit Stop interventions[14].

These interventions may affect exposures to enteric pathogen hazards to the public from human and animal fecal waste, including among people experiencing homelessness who may bear the greatest direct risks. Our primary aim was to estimate the degree to which Pit Stop interventions have reduced enteric pathogens in the immediate environment surrounding new toilet facilities. To this end, we: (1) conducted a systematic survey of discarded feces in a pre-defined area; (2) determined whether each fecal sample was of human or non-human origin; (3) quantified a range of enteric pathogens in recovered fecal samples, using molecular methods; and (4) used the intervention-attributable reduction in observed feces from 311 reports[15] to estimate reductions of the enteric pathogens we detected.

MATERIALS AND METHODS

A previous longitudinal study estimated that the installation of public toilets in two San Francisco neighborhoods resulted in a mean reduction of 18 stools per week within 500 meters (walking distance) of each facility[15]. We systematically collected discarded stools in this area and quantified a range of enteric pathogens in each sample. We used these data to model the reduction of pathogen hazards attributable to each facility.

Sample Collection. We used 311 reports of fecal waste in San Francisco, CA in August 2020 to identify hotspots for open defecation (OD) reports and design a systematic survey of discarded stools for enteric pathogen analysis. The available 311 data include reports of any discarded feces, including both suspected human and non-human fecal waste. The Tenderloin and SoMa neighborhoods had the highest number of 311 reports and were therefore selected for sampling;

we matched observed feces with 311 reports to verify that 311 data reliably indicated instances

of fecal waste (Text S1). We prioritized 20 blocks, including both sidewalks on either side of the

street for collection (Figure S1, Figure S2). We generated a perimeter around the selected blocks,

with any block inside the perimeter potentially utilized if samples were not available on the 20

selected blocks. We collected biospecimens on four Wednesday mornings in September and

October of 2020 before street cleaning began. Stool samples were collected into one-liter

biohazard bags and stored in a cooler with ice packs. We transferred samples into 1.5 mL

cryotubes and stored them at -20°C within 4 hours of collection. Any confirmed animal stool

(e.g., if the team observed an animal defecating) was not collected.

Sample preparation and analysis. We used the QIAamp 96 Virus QIAcube HT Kit (Qiagen,

Hilden, Germany) to extract nucleic acids from 100 mg of stool using a pre-treatment step

previously validated for molecular detection of multiple enteric pathogens with both DNA and

RNA genomes (Text S2)[17, 18]. We proceeded with extraction following the manufacturer's

protocol for the QIAamp 96 Virus QIAcube HT Kit, which we automated on the QIAcube

(Qiagen, Hilden, Germany). We measured the concentration of dsDNA using the dsDNA HS

assay with a Qubit 4 Fluorometer (Invitrogen, Waltham, Massachusetts).

We quantified human mitochondrial DNA (mtDNA) in each sample by dPCR to determine

whether feces were of presumptive human origin (Text S5, Figure S3) using an assay that

previously demonstrated 100% sensitivity and 97% specificity to human stool[19]. We normalized

mtDNA gene copy estimates to ng of dsDNA, and compared results against values reported in

the literature to categorize samples as human or non-human.[19] Positive and negative PCR

controls[20] were run each day of analysis via qPCR (Text S3) and dPCR (Text S5, Table S7). We

developed and used custom TaqMan Array Cards (TAC) (Thermo Fisher Scientific, Waltham,

MA) using published primer and probe sequences (Tables S2-S4) for a range of enteric bacteria,

viruses, protozoa, helminths, and controls. TAC is a 384-well array card with 8 ports for loading

samples and each 1.5 µL well contains lyophilized primers and probes for the detection of pre-

defined targets. We analyzed extracted nucleic acids via TAC on the QuantStudio 7, generating

real-time RT-qPCR curves for each target for each sample (Text S3). Standard curve details and

95% limits of detection are presented in Table S3. We visually compared exponential curves and

multicomponent plots with positive control plots to validate positive amplification[21, 22]. We

manually set thresholds to the point of inflection and considered targets amplifying at or below

35 cycles positive[23]. We re-ran samples that did not amplify DNA/RNA extraction positive

controls as expected at a 1:10 dilution, and samples that did not then amplify controls were

excluded from analysis. We performed additional confirmatory analysis of samples positive for

soil-transmitted helminths via microscopy using the mini-FLOTAC method (Text S4, Figure

S4).[16] We transformed cycle quantification (Cq) values into gene copy concentrations per gram

feces using pathogen-specific standard curves.

Stochastic model. We estimated the annual number of pathogens diverted from the environment

attributable to the Pit Stop intervention program using R version 4.1.0[24]. We transformed gene

copies (gc) into genomic units (i.e., discrete pathogens) using published values for gene copies

per genome (Table S5, model parameters)[23]. We treated non-detects as true zeroes. Using the

fitdistrplus package in R[25], we used maximum likelihood estimation (MLE) to fit log-normal

distributions to the quantity of pathogens per gram of stool in fecal samples, obtaining separate

estimates of the mean and standard deviation of log-pathogens per gram of presumptively human

and non-human stool, respectively, for each pathogen detected. We thus generated an estimated

mean and standard deviation for the number of pathogens per gram of feces. However, we

assigned upper bound thresholds to the distributions of pathogen concentrations in feces based

on biological plausibility (i.e., bacteria: 10^9/gram; viruses: 10^{12}/gram; protozoa: 10^7/gram;

helminths: 10^6/gram) and to prevent outliers in the distribution from driving the overall annual

estimate. We modeled an estimate of 18 stools per week that were diverted from the

environment[15] due to each new toilet facility across 52 weeks, because this was the reduction

reported by Amato *et al*. 2022[15] for the Tenderloin and SOMA neighborhoods. Using a Monte

Carlo Simulation, we applied a binomial distribution to estimate whether each unique diverted

stool (of 18) contained a specific pathogen. For each stool simulated to contain a given pathogen,

we estimated the concentration using the corresponding MLE-generated distribution. We fit a

log-normal distribution to the mass of human defecation events reported in Cummings *et al*.

1992 to stochastically estimate the mass of each stool[26]. For pathogens detected in only one of

the collected samples, we used the number of pathogens per gram of stool for the single stool in

place of an MLE-generated mean and the average MLE standard deviation from plurally

detected pathogens (Table S5). We repeated the process 52 times to estimate pathogens reduced

over a year. We then summed the number of estimated pathogens across the entire year to

estimate the number of pathogens diverted annually, repeating the process 1000 times to generate

95% confidence intervals.

RESULTS

We tested 60 stool samples, but excluded one whose assays lacked positive control

amplification, leaving 59 samples for analysis. Positive controls exhibited consistent

amplification (Cq~20) and no amplification was observed in our negative controls, except for the

16S assay (Cq~35, Text S6), which is a known contaminant in mastermix containing the

TaqMan polymerase[27,28]. Pathogen prevalence disaggregated by human mtDNA results is shown

in Table 1; we determined that 12 samples were of human origin and 47 were from non-human

sources. Out of 35 pathogens analyzed using TAC, 30/59 samples (51%) contained one or more

pathogens, 21/59 (36%) contained two or more pathogens, 8/59 (14%) contained 3 or more

pathogens, and 1/59 (2%) contained 5 pathogens. The most prevalent pathogens were atypical

enteropathogenic *E. coli* (EPEC) at 37% (22/59), typical EPEC and *Acanthamoeba spp.* each at

12% (7/59), *Cryptosporidium* at 8% (5/59), and *Giardia* at 7% (4/59).

The mtDNA dPCR assay indicated 12 samples were likely human in origin and 47 were likely

animal in origin (Figure S3). The prevalence of atypical EPEC was higher in human stools (42%)

than in the non-human stools (36%). Typical EPEC, tied for second most abundant pathogen

found, was absent in human stools, as were *Salmonella*, Norovirus, *Cryptosporidium*, *Giardia*,

and *Balantidium coli*. The pathogens *Helicobacter pylori*, *Shigella* spp./EIEC (enteroinvasive

Escherichia coli), *Plesiomonas shigelloides*, and *Yersinia entercolitica* were only found in

human fecal samples.

Two samples were TAC-positive for *Trichuris*, one of which was presumptively of human

origin. Due to a borderline initial Cq value (34.4), a negative microscopy result, and inhibited or

negative results on subsequent TAC runs for *Trichuris* for this sample, we excluded the result in

our prevalence calculations. Microscopy for the other (non-human) *Trichuris*-positive sample

revealed *Trichuris vulpis*, *Toxocara canis*, and hookworm ova (Figure S4).

We estimated the annual number of pathogens prevented from release into the local environment

as a result of each Pit Stop intervention under three scenarios (Table 2): (i) that fecal waste

reduced was both human and animal, since Pit Stop interventions also include animal waste bags

and bins (ii) that all fecal waste reduced was of human origin; and (3) (i) that all fecal waste

reduced was of animal origin. There results of the three scenarios were similar due to the high

concentration of pathogen shedding in feces, except for the instances where we only detected a

pathogen in human or non-human feces (e.g., *Cryptosporidium*) (Table 2). The estimated

reduction in pathogens released to environment in the scenario that considered both fecal waste

sources varied from 1×10^7 (95% CI: 1.6×10^6, 1.3×10^8) for *Trichuris* to 6.3×10^{12} (95% CI:

4.0×10^{12}, 7.9×10^{12}) for atypical enteropathogenic *E. coli*.

Because the Pit Latrine intervention includes animal waste bags and bins, it is plausible that

these facilities may also result in reductions in enteric pathogens observed only in animal feces,

including EPEC (typical), *Salmonella* spp., norovirus, *Cryptosporidium* spp., *Giardia* spp.,

Balantidium coli, and *Trichuris* spp. Assuming that the Pit Stop facilities reduce both human and

animal feces in the same proportion that they appear in our fecal samples (20% human, 80%

non-human), we estimate quantitative reductions of the broad range of pathogens represented

across all samples (Table 2).

Pathogen	All fecal samples (n=59)	Presumptively human[e] (n=12)	Presumptively non-human[e] (n=47)
Bacteria			
EPEC[a] (atypical)	37%	42%	36%
EPEC[a] (typical)	12%	0%	15%
EAEC[b] (aaiC/aatA)	3%	8%	2%
ETEC[c] (LT/ST)	3%	8%	2%
Helicobacter pylori	3%	17%	0%
*Shigella spp./*EIEC[d] (ipaH)	2%	8%	0%
Plesiomonas shigelloides	2%	8%	0%
Salmonella spp.	2%	0%	2%
Yersinia enterocolitica	2%	8%	0%
Virus			
norovirus (GI/II)	2%	0%	2%
Protozoa			
Acanthamoeba spp.	12%	17%	11%
Cryptosporidium spp.	8%	0%	11%
Giardia spp.	7%	0%	9%
Blastocystis spp.	3%	8%	2%
Balantidium coli	2%	0%	2%
Soil-transmitted helminth			
Trichuris spp.	2%	0%	2%

[a]Enteropathogenic *Escherichia coli*
[b]Enteroaggregative *Escherichia coli*
[c]Enterotoxigenic *Escherichia coli*
[d]Enteroinvasive *E. coli*
[e]Classified based on the concentration of human mitochondrial DNA (Supporting Information, Text S5)[19]

Not detected in any sample: *Campylobacter jejuni/coli, Clostridium difficile, E. coli* O157:H7, astrovirus, rotavirus, sapovirus (I/II/IV/V), SARS-CoV-2, *Cystoisospora belli, Cyclospora cayetanensis, Enterocytozoon bieneusi, Encephalitozoon intestinalis, Entamoeba histolytica, Entamoeba* spp., *Ancyclostoma duodenale, Ascaris lumbricoides, Enterobius vermicularis, Hymenolepis nana, Necator americanus,* and *Strongyloides stercolaris*

230 **Table 1.** Prevalence of pathogens in collected stool samples.

Pathogen	All fecal samples (n=59)		Presumptively human[e] (n=12)		Presumptively non-human[e] (n=47)	
	prevalence	median (95% CI)	prevalence	median (95% CI)	prevalence	median (95% CI)
EPEC[a] (atypical)	22/59	6.3×10^{12} (4.0×10^{12}, 7.9×10^{12})	5/12	6.3×10^{12} (5.0×10^{12}, 7.9×10^{12})	17/47	5.0×10^{12} (4.0×10^{12}, 7.9×10^{12})
EPEC[a] (typical)	7/59	1.0×10^{11} (4.0×10^{10}, 2.5×10^{11})	2/12	1.6×10^{11} (7.9×10^{10}, 3.2×10^{11})	5/47	1.0×10^{11} (4.0×10^{10}, 2.5×10^{11})
Acanthamoeba spp.	7/59	7.9×10^{6} (6.3×10^{6}, 1.0×10^{7})	1/12	5.0×10^{6} (4.0×10^{6}, 7.9×10^{6})	6/47	7.9×10^{6} (6.3×10^{6}, 1.0×10^{7})
Cryptosporidium spp.	5/59	1.6×10^{10} (1.0×10^{10}, 2.5×10^{10})	0/12	0	5/47	2.0×10^{10} (1.3×10^{10}, 3.2×10^{10})
Blastocystis spp.	2/59	5.0×10^{11} (1.3×10^{11}, 1.0×10^{12})	1/12	1.3×10^{12} (6.3×10^{11}, 2.0×10^{12})	1/47	3.2×10^{11} (4.0×10^{10}, 7.9×10^{11})
Giardia spp.	4/59	4.0×10^{9} (1.6×10^{9}, 7.9×10^{9})	0/12	0	4/47	5.0×10^{9} (2.0×10^{9}, 7.9×10^{9})
EAEC[b]	2/59	3.2×10^{11} (1.3×10^{11}, 6.3×10^{11})	1/12	7.9×10^{11} (5.0×10^{11}, 1.3×10^{12})	1/47	2.0×10^{11} (6.3×10^{10}, 4.0×10^{11})
ETEC[c]	2/59	2.5×10^{7} (1.3×10^{7}, 4.0×10^{7})	1/12	4.0×10^{7} (3.2×10^{7}, 7.9×10^{7})	1/47	1.6×10^{7} (6.3×10^{6}, 2.5×10^{7})
Helicobacter pylori	2/59	7.9×10^{8} (4.0×10^{8}, 1.3×10^{9})	0/12	0	2/47	1.0×10^{9} (6.3×10^{8}, 1.6×10^{9})
Shigella spp./EIEC[d]	1/59	3.2×10^{8} (6.3×10^{7}, 2.5×10^{9})	1/12	2.0×10^{9} (1.0×10^{9}, 6.3×10^{9})	0/47	0
Plesiomonas shigelloides	1/59	3.2×10^{9} (6.3×10^{8}, 2.5×10^{10})	0/12	0	1/47	5.0×10^{9} (1.3×10^{9}, 3.2×10^{10})
Salmonella spp.	1/59	1.0×10^{9} (2.5×10^{8},	0/12	0	1/47	1.6×10^{9} (3.2×10^{8},

		6.3x10^9)				7.9x10^9)
Yersinia enterocolitica	1/59	1.6x10^{10} (3.2x10^9, 1.0x10^{11})	1/12	1.0x10^{11} (4.0x10^{11}, 2.5x10^{11})	0/47	0
norovirus (GI/II)	1/59	4.0x10^8 (7.9x10^7, 3.2x10^9)	0/12	0	1/47	6.3x10^8 (1.3x10^8, 4.0x10^9)
Balantidium coli	1/59	2.5x10^9 (5.0x10^8, 1.6x10^{10})	0/12	0	1/47	4.0x10^9 (7.9x10^8, 2.0x10^{10})
Trichuris spp.	1/59	1.0x10^7 (1.6x10^6, 1.3x10^8)	0/12	0	1/47	1.6x10^7 (2.5x10^6, 1.3x10^8)

[a]Enteropathogenic *Escherichia coli*
[b]Enteroaggregative *Escherichia coli*
[c]Enterotoxigenic *Escherichia coli*
[d]Enteroinvasive *E. coli*
[e]Classified based on the concentration of human mitochondrial DNA (Supporting Information, Text S5)[19]

Table 2. Modeled annual pathogens prevented from release into the environment per Pit Stop in study area (within 500 m walking distance).

DISCUSSION

We detected a wide range of enteric pathogens in the fecal samples we collected, with approximately half (51%) of all samples positive for one or more of the pathogens we sought. Based on our pathogen analysis of fecal wastes and previously estimated reductions in fecal wastes on the street[15], the Pit Stop intervention has likely reduced the number of pathogens released into the environment within 500 m walking distance of each new toilet facility installed. Reducing pathogen hazards in a densely populated environment can prevent disease transmission, especially for the most vulnerable population: people experiencing homelessness, particularly unsheltered people living in the study area who previously lacked accessible public sanitation and hygiene facilities.

Four out of five samples were non-human in origin, with *Giardia*, *Cryptosporidium*, and EPEC (typical) occurring only in animal stools, though each of these has zoonotic potential[12, 29].

Microscopy results also revealed a high burden of helminth infection in a presumptive canine

sample. Humans are not the definitive host of *Toxocara* and canine hookworm, but humans can

be infected by them[30, 31]. While provision of public toilets has the potential to reduce human open

defecation, control of animal feces requires different interventions. Although the Pit Stop

intervention included animal waste bag distribution and disposal bins, further measures are

probably required, including public education, enforcement, environmental controls, or other

measures. Stray or feral animals may also contribute fecal waste. Based on our detection of a

range of potentially zoonotic enteric pathogens in non-human fecal waste with the potential for

human contact, control of animal feces should be considered in this setting[12, 29].

Our findings should be considered alongside some limitations and caveats. First, though we

tested for a range of important enteric pathogens, we selected these targets *a priori* and they are a

subset of pathogens that may be relevant in this context, especially considering the widespread

presence of non-human fecal wastes. Other potential zoonoses, including *Toxoplasma gondii*,

Toxocara, and canine hookworm may have been present and future studies should consider

them[12, 29]. Second, the frequency of detection of these pathogens cannot be assumed to represent

prevalence of infection in any population: multiple fecal samples in the study area may well have

been from a single individual, and our quantitative estimations are based on a limited number of

samples. Third, detection of pathogen-associated nucleic acids does not and cannot indicate

viability or infectivity. These data cannot be used directly in assessing risk of exposure, without

further assumptions beyond the scope of our analysis. Fecal samples were apparently fresh when

sampled, but pathogens can be inactivated in the environment. Fourth, even without public

toilets, many discarded stools will go on to be collected and safely disposed of through street and

sidewalk cleaning, being effectively removed from the environment and therefore unlikely to

result in exposure. For this reason and others, we cannot conclude that the reduction in pathogen

hazards associated with feces would necessarily result in changes to human exposures, infection,

or disease, only that the potential exists.

Fifth, we treated non-detects on TAC as true zeroes, so our estimates might be conservative

given the lower limit of detection for targets using these assays (Table S3). While TAC uses the

same highly sensitive and specific probe-based RT-qPCR chemistry that has been widely

adopted for pathogen detection and quantification in both clinical and research settings across a

range of sample matrices, the physical constraints of the platform result in much smaller reaction

volumes (~1.5 µL) than for traditional tube-based approaches (20 µL – 50 µL). The reduced

reaction volume may negatively impact analytical sensitivity (i.e., increase the probability of

false negatives at low target concentrations) and likewise increase the variability of estimated

target quantities at lower concentrations. However, TAC has previously been shown to compare

favorably with traditional qPCR approaches in terms of quantification linearity as well as

pathogen-specific sensitivity and specificity in stool samples[21, 23, 32]. We observed similar

linearity of quantification with our standard curves for all but two assays, which were excluded

from the analysis (Table S3), suggesting that our estimated pathogen gene copy concentrations in

positive samples were sufficiently reliable for the purposes of parametrizing the log-normal

pathogen concentration distributions (which operate on the scale of order-of-magnitude

differences in pathogen quantity) we employed in our stochastic models.

Finally, in our quantitative model estimating pathogen reductions attributable to the Pit Stop

intervention, we assumed that our stool samples (their pathogen content over time, and

human/animal origin) are representative of the fecal wastes reduced due to public toilet

construction. This may or may not be the case. Our samples were from a narrow window in time,

when some pathogens may be more prevalent than others, and this may not be representative of

what is being shed over time in the populations contributing fecal wastes to the streets in our

study area. We assumed a mean reduction of 18 instances of OD within 500 meters of newly

installed Pit Stop locations[15] in our study area, per week, throughout the modeled period of one

year based on six months of observational data. We first estimated reductions in pathogen

hazards assuming all contained waste was of human origin, given the Pit Stop's primary

ostensible role in serving people. We also estimated hazard reductions based on enteric pathogen

quantification across all samples, which from our collection effort were determined to be 20%

human origin and 80% animal origin. The Pit Stop interventions included both toilet facilities as

well as bags and bins for animal waste control, so both types of stools could plausibly be reduced

in the immediate surroundings. We observed differences between pathogens detected according

to presumptive human versus animal sources, with some pathogens appearing in only human

stools and others occurring in non-human stool only. For example, *Helicobacter pylori* and

Shigella/EIEC were detected only in presumptively human stools.

Clean water, safe sanitation, and adequate hygiene are not universal in American cities[2, 33, 34],

with gaps most apparent among those experiencing homelessness[2, 7]. San Francisco deserves

credit for proactively working to solve this problem, which is not inexpensive[14, 15] and can be

politically contentious as cities grapple with the growing crisis of homelessness. Water and

sanitation are human rights[35] that are essential to living a dignified life. Construction of publicly

accessible, safe toilets is a commonsense approach to reducing enteric pathogen hazards in

cities[9], though the primary reason to continue to invest in public sanitation facilities is to support

the physical, mental, and social well-being of people – much of which is difficult or impossible

to measure in practice – and because it is the right and humane thing to do. Moreover, sanitation

is a biological necessity that is needed wherever people live. The waste must go somewhere: an

adult weighing 80 kg (near the mean body mass in North America) will produce an average of

approximately 38 kg of feces per year[36, 37], 100% of which should be effectively and safely

managed to protect all members of the community from infectious disease. Cities should also

consider interventions aimed at reducing animal feces, which are an underappreciated source of

enteric pathogen hazards in urban spaces. Our findings demonstrate that a wide range of

pathogens with zoonotic potential may be present in discarded animal waste, with uncertain

implications for human exposure and disease transmission.

REFERENCES

1.   Fazel, S.; Geddes, J. R.; Kushel, M., The health of homeless people in high-income countries:
descriptive epidemiology, health consequences, and clinical and policy recommendations. *Lancet* **2014,**
384, (9953), 1529-40.
2.   Capone, D.; Cumming, O.; Nichols, D.; Brown, J., Water and Sanitation in Urban America, 2017–
2019. *American Journal of Public Health* **2020,** *110*, (10), 1567-1572.
3.   Yin, S.; Barker, L.; Ly, K. N.; Kilmer, G.; Foster, M. A.; Drobeniuc, J.; Jiles, R. B., Susceptibility to
Hepatitis A Virus Infection in the United States, 2007-2016. *Clin Infect Dis* **2020,** *71*, (10), e571-e579.
4.   Liu, C. Y.; Chai, S. J.; Watt, J. P., Communicable disease among people experiencing
homelessness in California. *Epidemiol Infect* **2020,** *148*, e85.
5.   Park, Y. S.; Bliss, D. Z., Availability of Public Toilets in Parks and Recreational Sites in Selected US
Cities. *Journal of Wound Ostomy & Continence Nursing* **2019,** *46*, (3), 235-239.
6.   Bartram, J.; Cairncross, S., Hygiene, sanitation, and water: forgotten foundations of health. *PLoS*
Med **2010,** *7*, (11), e1000367.

7.   Capone, D.; Ferguson, A.; Gribble, M. O.; Brown, J., Open Defecation Sites, Unmet Sanitation
Needs, and Potential Sanitary Risks in Atlanta, Georgia, 2017–2018. *American Journal of Public Health*
2018, *108*, (9), 1238-1240.
8.   Frye, E. A., Capone, D., and Evans, D.P., Open Defecation in the United States: Perspectives from
the Streets. *Environmental Justice* **2019,** *12*, (5), 226-230.
9.   Moreira, F. D.; Rezende, S.; Passos, F., On-street toilets for sanitation access in urban public
spaces: A systematic review. *Utilities Policy* **2021,** *70*, 101186.
10.   Brown, J.; Cairncross, S.; Ensink, J. H., Water, sanitation, hygiene and enteric infections in
children. *Arch Dis Child* **2013,** *98*, (8), 629-34.
11.   Ginn, O.; Rocha-Melogno, L.; Bivins, A.; Lowry, S.; Cardelino, M.; Nichols, D.; Tripathi, S. N.; Soria,
F.; Andrade, M.; Bergin, M.; Deshusses, M. A.; Brown, J., Detection and Quantification of Enteric
Pathogens in Aerosols Near Open Wastewater Canals in Cities with Poor Sanitation. *Environ Sci Technol*
2021, *55*, (21), 14758-14771.
12.   Penakalapati, G.; Swarthout, J.; Delahoy, M. J.; McAliley, L.; Wodnik, B.; Levy, K.; Freeman, M. C.,
Exposure to Animal Feces and Human Health: A Systematic Review and Proposed Research Priorities.
Environ Sci Technol **2017,** *51*, (20), 11537-11552.
13.   Wagner, E. G.; Lanoix, J. N., Excreta disposal for rural areas and small communities. *Monogr Ser*
World Health Organ **1958,** *39*, 1-182.
14.   Rezal, A., Poop complaints have swelled in all San Francisco neighborhoods — except this one.
San Francisco Chronicle 23 August 2022, 2022.
15.   Amato, H. K.; Martin, D.; Hoover, C. M.; Graham, J. P., Somewhere to go: assessing the impact of
public restroom interventions on reports of open defecation in San Francisco, California from 2014 to
2020. *BMC Public Health* **2022,** *22*, (1), 1673.
16.   Research, A. S. *San Francisco Homeless Count & Survey Comprehensive Report*;
www.appliedsurveyresearch.org: San Jose, California, 2019; p 76 pages.
17.   Capone, D.; Berendes, D.; Cumming, O.; Knee, J.; Nalá, R.; Risk, B. B.; Stauber, C.; Zhu, K.; Brown,
J., Analysis of Fecal Sludges Reveals Common Enteric Pathogens in Urban Maputo, Mozambique.
Environmental Science & Technology Letters **2020,** *7*, (12), 889-895.
18.   Knee, J.; Sumner, T.; Adriano, Z.; Anderson, C.; Bush, F.; Capone, D.; Casmo, V.; Holcomb, D.;
Kolsky, P.; MacDougall, A.; Molotkova, E.; Braga, J. M.; Russo, C.; Schmidt, W. P.; Stewart, J.; Zambrana,
W.; Zuin, V.; Nalá, R.; Cumming, O.; Brown, J., Effects of an urban sanitation intervention on childhood
enteric infection and diarrhea in Maputo, Mozambique: A controlled before-and-after trial. *eLife* **2021,**
10, e62278.
19.   Zhu, K.; Suttner, B.; Pickering, A.; Konstantinidis, K. T.; Brown, J., A novel droplet digital PCR
human mtDNA assay for fecal source tracking. *Water Res* **2020,** *183*, 116085.
20.   Borchardt, M. A.; Boehm, A. B.; Salit, M.; Spencer, S. K.; Wigginton, K. R.; Noble, R. T., The
Environmental Microbiology Minimum Information (EMMI) Guidelines: qPCR and dPCR Quality and
Reporting for Environmental Microbiology. *Environmental Science & Technology* **2021,** *55*, (15), 10210-
10223.
21.   Lappan, R.; Henry, R.; Chown, S. L.; Luby, S. P.; Higginson, E. E.; Bata, L.; Jirapanjawat, T.; Schang,
C.; Openshaw, J. J.; O'Toole, J.; Lin, A.; Tela, A.; Turagabeci, A.; Wong, T. H. F.; French, M. A.; Brown, R.
R.; Leder, K.; Greening, C.; McCarthy, D., Monitoring of diverse enteric pathogens across environmental
and host reservoirs with TaqMan array cards and standard qPCR: a methodological comparison study.
Lancet Planet Health **2021,** *5*, (5), e297-e308.
22.   Lappan, R.; Jirapanjawat, T.; Williamson, D. A.; Lange, S.; Chown, S. L.; Greening, C.,
Simultaneous detection of multiple pathogens with the TaqMan Array Card. *MethodsX* **2022,** *9*, 101707.
23.   Liu, J.; Gratz, J.; Amour, C.; Nshama, R.; Walongo, T.; Maro, A.; Mduma, E.; Platts-Mills, J.;
Boisen, N.; Nataro, J.; Haverstick, D. M.; Kabir, F.; Lertsethtakarn, P.; Silapong, S.; Jeamwattanalert, P.;

Bodhidatta, L.; Mason, C.; Begum, S.; Haque, R.; Praharaj, I.; Kang, G.; Houpt, E. R., Optimization of
Quantitative PCR Methods for Enteropathogen Detection. *PLOS ONE* **2016,** *11*, (6), e0158199.
24. Team, R. C., R: A language and environment for statistical computing (Version 4.0. 5)[Computer
software]. *R Foundation for Statistical Computing* **2021**.
25. Delignette-Muller, M. L.; Dutang, C., fitdistrplus: An R Package for Fitting Distributions. *Journal*
of Statistical Software **2015,** *64*, (4), 1 - 34.
26. Cummings, J. H.; Bingham, S. A.; Heaton, K. W.; Eastwood, M. A., Fecal weight, colon cancer risk,
and dietary intake of nonstarch polysaccharides (dietary fiber). *Gastroenterology* **1992,** *103*, (6), 1783-9.
27. Corless, C. E.; Guiver, M.; Borrow, R.; Edwards-Jones, V.; Kaczmarski, E. B.; Fox, A. J.,
Contamination and sensitivity issues with a real-time universal 16S rRNA PCR. *J Clin Microbiol* **2000,** *38*,
(5), 1747-52.
28. Philipp, S.; Huemer, H. P.; Irschick, E. U.; Gassner, C., Obstacles of Multiplex Real-Time PCR for
Bacterial 16S rDNA: Primer Specifity and DNA Decontamination of Taq Polymerase. *Transfus Med*
Hemother **2010,** *37*, (1), 21-28.
29. Prendergast, A. J.; Gharpure, R.; Mor, S.; Viney, M.; Dube, K.; Lello, J.; Berger, C.; Siwila, J.;
Joyeux, M.; Hodobo, T.; Hurt, L.; Brown, T.; Hoto, P.; Tavengwa, N.; Mutasa, K.; Craddock, S.; Chasekwa,
B.; Robertson, R. C.; Evans, C.; Chidhanguro, D.; Mutasa, B.; Majo, F.; Smith, L. E.; Hirai, M.; Ntozini, R.;
Humphrey, J. H.; Berendes, D., Putting the "A" into WaSH: a call for integrated management of water,
animals, sanitation, and hygiene. *Lancet Planet Health* **2019,** *3*, (8), e336-e337.
30. Borg, O. A.; Woodruff, A. W., Prevalence of infective ova of Toxocara species in public places. *Br*
Med J **1973,** *4*, (5890), 470-2.
31. Woodruff, A. W., Toxocariasis. *British Medical Journal* **1970,** *3*, (5724), 663-669.
32. Liu, J.; Gratz, J.; Amour, C.; Kibiki, G.; Becker, S.; Janaki, L.; Verweij, J. J.; Taniuchi, M.; Sobuz, S.
U.; Haque, R.; Haverstick, D. M.; Houpt, E. R., A laboratory-developed TaqMan Array Card for
simultaneous detection of 19 enteropathogens. *J Clin Microbiol* **2013,** *51*, (2), 472-80.
33. Mueller, J. T.; Gasteyer, S., The widespread and unjust drinking water and clean water crisis in
the United States. *Nature Communications* **2021,** *12*, (1), 3544.
34. Meehan, K.; Jurjevich, J. R.; Chun, N. M. J. W.; Sherrill, J., Geographies of insecure water access
and the housing–water nexus in US cities. *Proceedings of the National Academy of Sciences*
2020, *117*, (46), 28700-28707.
35. Nations, U., The Human Rights to Safe Drinking Water and Sanitation: Resolution In Council, H.
R., Ed. UN General Assembly: 2018.
36. Berendes, D. M.; Yang, P. J.; Lai, A.; Hu, D.; Brown, J., Estimation of global recoverable human
and animal faecal biomass. *Nature Sustainability* **2018,** *1*, (11), 679-685.
37. Yang, P. J.; LaMarca, M.; Kaminski, C.; Chu, D. I.; Hu, D. L., Hydrodynamics of defecation. *Soft*
Matter **2017,** *13*, (29), 4960-4970.