Sleep-Disordered Breathing Destabilizes Ventricular Repolarization

Soroosh Solhjoo, PhD[1], Mark C. Haigney, MD[1,2], Trishul Siddharthan, MD[3], Abigail Koch, MD[3], Naresh M. Punjabi, MD, PhD[3]

[1] F. Edward Hébert School of Medicine, Bethesda, Maryland, USA
[2] Military Cardiovascular Outcomes Research (MiCOR), Bethesda, Maryland, USA
[3] University of Miami Miller School of Medicine, Miami, Florida, USA

Short Title: Sleep apnea, intermittent hypoxemia, and QTVI

Total word count: 5,697

Corresponding Author:

Soroosh Solhjoo, PhD
F. Edward Hébert School of Medicine
4301 Jones Bridge Road
Bethesda, MD 20814
soroosh@jhmi.edu

ABSTRACT

BACKGROUND: Sleep-disordered breathing (SDB) is associated with an increased risk of

cardiac arrhythmias, sudden cardiac death, and all-cause mortality. This study investigates the

mechanisms underlying this association by examining the effects of prevalent and incident SDB

and intermittent hypoxia on ventricular repolarization lability.

METHODS: Electrocardiographic recordings from three cohorts were used for this study:

(1) a cross-sectional sample of 61 participants with severe SDB and a matched group without

SDB; (2) a longitudinal sample of 26 participants without SDB at baseline who developed SDB

after a 5-year follow-up, and a matched group without SDB; and (3) a cohort of 19 healthy adults

exposed to intermittent hypoxia or ambient air. Mean heart rate, heart rate variability (SDNN), and

QT variability index (QTVI), a measure of ventricular repolarization lability, were calculated from

one-lead ECG recordings.

RESULTS: In the cross-sectional sample, participants with severe SDB had larger QTVI

($P = 0.027$), heart rate ($P = 0.028$), SDNN ($P = 0.018$), and hypoxemia burden as assessed by

the total sleep time with oxygen saturation less than 90% (TST_{90}; $P < 0.001$) than those without

SDB. TST_{90}, but not the frequency of arousals, was an independent predictor of QTVI ($P = 0.017$).

Both heart rate and QTVI were predictive of all-cause mortality. In the longitudinal sample with

incident SDB, QTVI increased from -1.23 ± 0.15 to -0.86 ± 0.14 ($P = 0.017$) over the 5-year follow-

up period. Finally, in the cohort of healthy adults, 4 hours of exposure to intermittent hypoxia

increased heart rate ($P = 0.001$) and QTVI ($P = 0.016$).

CONCLUSIONS: Prevalent and incident SDB are associated with increased ventricular

repolarization instability which predisposes to ventricular arrhythmias and sudden cardiac death.

Intermittent hypoxemia can destabilize ventricular repolarization and is a likely mediator of

increased mortality in SDB.

KEY WORDS: QT variability index, QTVI, intermittent hypoxia, Sleep Heart Health Study

CLINICAL PERSPECTIVE

What Is New?

- Sleep-disordered breathing, a prevalent condition in the general population, is associated with unstable ventricular repolarization.

- Incident sleep-disordered breathing destabilizes ventricular repolarization, likely due to the effects of intermittent hypoxemia.

- Acute exposure to intermittent hypoxia destabilizes ventricular repolarization in healthy adults indicating a causal effect of sleep-disordered breathing on ventricular repolarization and increased mortality risk.

What Are the Clinical Implications?

Identifying and treating sleep-disordered breathing in patients at risk for sudden cardiac death, such as those with congestive heart failure, can improve clinical outcomes.

INTRODUCTION

Sleep-disordered breathing (SDB) is a common disorder with prevalence estimates of 9%-38%.[1, 2] It is more prevalent in older adults affecting approximately 90% of men and 78% of women between the ages of 65 to 80 years.[3] SDB is characterized by repetitive collapse of the upper airway during sleep. The ensuing periods of decreased airflow (hypopnea) or no airflow (apnea) are associated with swings in intrathoracic pressure, hypoxemia, hypercapnia, and arousals from sleep. SDB is a well-established risk factor for hypertension and cardiovascular disease.[4-8] There is unequivocal evidence that SDB is associated with a predisposition for cardiac arrhythmias.[9-11] Conduction abnormalities, bradyarrhythmias, and both atrial and ventricular tachyarrhythmias are more prevalent in SDB.[11] A growing body of empirical evidence also indicates that SDB is a risk factor for sudden cardiac death.[12-15]

The role of SDB in triggering malignant arrhythmias can be assessed through its association with abnormal electrocardiographic (ECG) features known to indicate the risk of sudden cardiac death. It is well established that the beat-to-beat QT variability index (QTVI), a measure of ventricular repolarization lability, is predictive of ventricular arrhythmias and sudden cardiac death.[16-21] Elevated QTVI has also been linked to a higher propensity for all-cause and cardiovascular mortality.[22] An increase in QTVI indicates that the QT interval is varying out of proportion to heart rate variability. SDB-associated nocturnal hypoxemia, hypercapnia, and sleep fragmentation can increase the sympathetic nervous system's activity which, in turn, can modulate the QT interval through mechanisms independent of its effect on heart rate[23-25] and result in an increase in QT variability that is out of proportion to the variability in heart rate.[26-28] Therefore, QTVI

is expected to increase in patients with SDB. However, the data are limited, and significant

gaps remain in our understanding of the effects of SDB on QTVI. First, available data are

conflicting with studies showing no association,[29] a positive association,[30] or an inverse

association[31] between SDB severity and QTVI. Second, it remains to be determined

whether incident SDB leads to longitudinal changes in ventricular repolarization. Finally,

there are no data as to whether intermittent hypoxemia or sleep fragmentation, the two

pathophysiological concomitants of SDB, can independently alter overall QTVI.

Therefore, the current study sought to determine whether prevalent and incident SDB are

associated with ventricular repolarization instability and to characterize the role of

intermittent hypoxemia and arousals in increasing QTVI.

METHODS

Sample Selection

Three distinct samples were selected for this study. The first two samples were from the

Sleep Heart Health Study (SHHS), a prospective cohort study on the cardiovascular

consequences of SDB.[32] Of the 6,441 participants who completed the baseline SHHS

examination, 3,296 had a follow-up polysomnogram approximately five years later. The

home polysomnogram was conducted using a portable monitor (P-Series, Compumedics,

Abbotsville, AU). The following physiologic signals were recorded: C3 and C4

electroencephalograms, right and left electrooculograms, a single bipolar ECG, a chin

electromyogram, oxyhemoglobin saturation (SpO_2) by pulse oximetry, chest and

abdominal excursion by inductance plethysmography, airflow by an oronasal

thermocouple, and body position by a mercury gauge. Sleep was manually scored as

follows. An apnea was identified if the airflow was absent or nearly absent for at least ten

seconds. Hypopnea was defined as at least a 30% reduction in airflow or

thoracoabdominal movement below baseline values for at least 10 seconds associated

with an oxyhemoglobin desaturation of 4%. Details regarding cohort recruitment and

sleep and ECG data collection have been previously published.[33]

To assess whether prevalent SDB is associated with a higher QTVI, a cross-

sectional sample from the baseline SHHS examination was selected that included

participants with severe SDB (apnea/hypopnea index (AHI) $\geq$ 33 events/hour, i.e., 95[th]

percentile) along with a group of age (±1 year), sex, BMI (±1 kg/m^2), and race-matched

participants without SDB (AHI < 1.33 events/hour; i.e., 25[th] percentile). A second sample

of SHHS participants was selected to assess whether incident SDB and associated

nocturnal hypoxemia increase QTVI over time. The second sample consisted of

participants initially without SDB with a TST_{90} < 0.22% (i.e., 50[th] percentile) who then

developed SDB and had a TST_{90} $\geq$ 10.4% (i.e., 90[th] percentile) at the follow-up

polysomnogram five years later. An age (±1 year), sex, BMI (±1 kg/m^2), and race-matched

group of comparator participants was also identified who did not develop SDB and had a

TST_{90} < 0.22% during both the baseline and follow-up polysomnograms. Additional

criteria for the cross-sectional and longitudinal samples required that the participants were

free of prevalent myocardial infarction, coronary artery bypass surgery, heart failure, or

angina and were not on beta-blockers, calcium channel blockers, or antiarrhythmic

agents. Informed consent was obtained from all SHHS participants, and the study

protocol was approved by the institutional review committee on human research

(Approval Number: NA_00010790). Anonymized data are available from National Sleep

Research Resource.[34]

A third sample of healthy adult volunteers (N=19) was recruited from the local

community as part of an experimental study on the acute effects of intermittent hypoxia,

as previously described.[35] All subjects had normal polysomnography and demonstrated

at least 7 hours of habitual sleep on actigraphy. For each subject, the experiment was

done on two days, separated by one week. On one day, the subject was exposed to

intermittent hypoxia, and on the other, to normoxia. The exposure to intermittent hypoxia

or normoxia was randomly determined. Intermittent hypoxia was induced using a

standard nasal mask connected to a three-way valve by alternating the airflow between

hypoxic gas (5% O_2, 95% N_2) and ambient air (21% O_2, 79% N_2). Exposure to hypoxic

gas was maintained until the SpO_2 reached 85%. Approximately 20-25 hypoxic

events/hour were elicited. On the normoxia day, the subject was on 21% O_2 for the entire

experimental period by replacing the hypoxic gas tank with a pressurized ambient air tank.

SpO_2 was recorded using the Biox 3700 pulse oximeter (Ohmeda, Englewood, CO) at

2 Hz. A one-lead ECG signal was recorded continuously at 200 Hz using an Embla N-

7000 system (Natus Medical, Pleasanton, CA). The first four hours of each experiment

were used to assess the effects of intermittent hypoxia on QT variability. Informed consent

was obtained from all participants in the study, and the study protocol was approved by

the institutional review committee on human research (Approval Number: NA_00002563).

Anonymized data are available from the corresponding author upon reasonable request.

ECG Signal Processing

The ECG signal was analyzed using customized software developed in MATLAB (Mathworks, Natick, MA) to derive measures of heart rate (HR), heart rate variability, and QTVI over a predefined epoch length. The standard deviation of the intervals between successive normal beats (SDNN) was calculated as the time-domain measure of heart rate variability. Spectral analysis was used to derive frequency-domain measures of heart rate variability. QT variability was measured using a template-matching algorithm,[16, 36] and was calculated as $QTVI = \log_{10}(QTVN/HRVN)$, where $QTVN = QT_{variance}/QT_{mean}^2$ and $HRVN = HR_{variance}/HR_{mean}^2$.

Statistical Analysis

For each participant, repeated measurements of electrocardiographic parameters over consecutive epochs were summarized using medians, which were subsequently compared using the *t* test for independent or paired samples as appropriate. Normality was tested using the Lilliefors test. *P* values < 0.05 were considered significant. Multivariable linear regression based on least-squares estimation was used to investigate the association between QTVI and hypoxic burden. The least absolute shrinkage and selection operator (lasso)[37] was used to determine the main contributors to mortality among the demographic, polysomnographic, and electrocardiographic parameters based on multivariable logistic regression models. Statistical analyses and graphing were done using the R statistical package, glmnet package,[38] MATLAB, and Origin (OriginLab, Northampton, MA).

RESULTS

SDB Is Associated with Instability of Ventricular Repolarization

The cross-sectional sample comprised 61 SHHS participants (45 males, 16 females) in each group. The average AHI was 46.28 and 0.62 events/hour in participants with and without SDB, respectively. The two groups were matched by age, sex, race, and BMI (Table 1). TST_{90} was greater in those with than without SDB (11.39 ± 1.42% vs. 1.32 ± 0.61%, $P < 0.001$). The SDB group had a higher mean heart rate ($P = 0.028$) and SDNN ($P = 0.018$) during sleep than those without SDB. QTVI was also higher with SDB, indicating greater ventricular repolarization lability ($P = 0.027$). Stratified analyses by rapid eye movement sleep (REM) and non-REM sleep showed similar findings with a higher QTVI with SDB irrespective of sleep stage (Table S1 and Figure 1).

Logistic regression analyses were used to determine whether electrocardiographic (heart rate, SDNN, and QTVI), demographic (age, sex, BMI, smoking status), and polysomnographic measures (AHI, frequency of arousals, and TST_{90}) were associated with all-cause mortality over an average follow-up of 8.2 years. Heart rate and QTVI were the only two predictors of mortality selected by the lasso analysis with an area under the receiver operating characteristic curve (AUC) of 0.75 (95% confidence interval: 0.63-0.87) based on 10-fold cross-validation. Including all other covariates in addition to heart rate and QTVI resulted in no significant increase in AUC. Multivariable linear regression analysis showed that higher TST_{90} values were associated with greater QTVI independent of age, sex, BMI, and smoking status. For a10% increase in TST_{90}, QTVI increased by 0.13 ($P = 0.017$). No association was noted between the frequency of arousals and QTVI.

Development of SDB Destabilizes Ventricular Repolarization

Based on the finding that TST_{90} was independently predictive of QTVI in the cross-sectional cohort, the role of incident SDB in increasing QTVI was examined. The longitudinal sample comprised 26 participants (8 males, 18 females) in each group. Table 2 shows the demographic, polysomnographic, and electrocardiographic data. In participants who developed SDB, AHI increased from 2.54 ± 0.72 events/hour at baseline to 13.74 ± 3.5 events/hour at follow-up (P = 0.001). The AHI remained unchanged in those who did not develop SDB (2.79 ± 0.58 vs. 3.36 ± 0.67 events/hour, P = 0.386). With incident SDB, mean heart rate and SDNN did not change between the two visits. However, QTVI increased from -1.23 ± 0.15 to -0.86 ± 0.154 (P = 0.017, Figure 2). Heart rate, SDNN, and QTVI did not significantly change in the group without SDB. Stratified analysis by sleep stage revealed that incident SDB was associated with increased QTVI during NREM but not REM sleep (Table S2).

Experimental Intermittent Hypoxia Acutely Destabilizes Ventricular Repolarization

To assess the causal effects of hypoxemia in adversely affecting QTVI, ECG data from 19 healthy volunteers exposed to intermittent hypoxia or normoxia were used. The average age and BMI of the 19 healthy volunteers were 25.5 (Range: 18.0–37.0) years and 25.2 (21.7–33.0) kg/m^2, respectively. On the day of intermittent hypoxia, the participants were exposed to the hypoxic gas mixture 24.59 ± 0.78 times/hour for an average of 67.52 ± 2.79 seconds per event. Consequently, mean SpO_2 was significantly lower with intermittent hypoxia (90.90 ± 0.20%) than with normoxia (97.37 ± 0.20%, $P <$ 10^{-13}, Table S3). Heart rate transiently increased with each episode of hypoxia (Figure 3), leading to increased heart rate variability. SDNN was higher (P = 0.014) during the first

hour of intermittent hypoxia than normoxia (Figure 4). Other electrocardiographic

measures (i.e., NN interval, LF/(HF+LF), and QTVI) were similar between the two

conditions during the first hour. No significant changes were observed in any

electrocardiographic measures between the first and the fourth hour of normoxia (Table

3, Figure 4). However, between the first and the last hour of intermittent hypoxia, heart

rate and the LF/(LF+HF) ratio increased while log(HF) decreased, indicating a shift in

sympathovagal balance towards sympathetic activity (Table 6). QTVI also increased with

intermittent hypoxia over the four hours (Table 6). By the 4th hour of the experiments,

SDNN ($P = 0.042$), heart rate ($P = 0.018$), LF/(LF+HF) ($P = 0.009$), and QTVI ($P = 0.021$)

were all significantly higher with intermittent hypoxia compared to normoxia (Figure 4).

DISCUSSION

The data presented herein demonstrate that SDB is associated with ventricular

repolarization instability. In a cohort of middle-aged and older adults, compared to those

without SDB, severe SDB was associated with a higher QTVI. The severity of nocturnal

hypoxemia, but not the frequency of arousals, was associated with a higher QTVI, which

was predictive of all-cause mortality. Longitudinal data showed that incident SDB was

associated with increased QTVI over the five-year follow-up period. Finally, in a sample

of healthy adults, experimental exposure to intermittent hypoxia progressively increased

QTVI.

The last decade has seen enormous growth in the evidence linking SDB to

ventricular arrhythmias, and sudden cardiac death.[12, 14, 39-43] Epidemiological data have

shown that SDB is associated with a higher burden of complex ventricular ectopy (i.e.,

bigeminy, trigeminy, and quadrigeminy), non-sustained ventricular tachycardia, and

sudden cardiac death.[11] A meta-analysis of 22 studies (42,099 participants) reported that

the SDB-related relative risk for sudden death was 1.74 (95% CI: 1.40-2.10).[43] Subgroup

analysis of studies with polysomnography data revealed a dose-response association

between SDB severity and the risk of sudden cardiac death. Several biological

mechanisms can explicate the observed clinical and epidemiological associations

between SDB and sudden cardiac death. Upper airway collapse during sleep can induce

ventricular electromechanical decoupling and thereby create an arrhythmogenic

substrate, as evident by transient prolongation of the QT interval.[44] In an observational

cohort study of 10,701 adults undergoing a diagnostic clinical polysomnogram,

resuscitated or fatal sudden cardiac arrest was best predicted by an AHI ≥ 20 events/hour,

mean nocturnal oxygen saturation < 93%, and lowest oxygen saturation ≤ 78%.[14] Thus,

SDB may increase the risk of sudden cardiac death partly through the effects of nocturnal

hypoxemia. Furthermore, *in vivo* experimental studies show that chronic intermittent

hypoxemia can augment sympathetic activity, alter ventricular repolarization, and

increase the risk of sudden cardiac death.[45] Mild hypoxia has been reported to increase

the late component of the sodium current and alter the L-type calcium current *in vitro*.[46-

48] Further, hypoxia regulates the L-type calcium channels by increasing their sensitivity

to β-adrenergic stimulation.[49] Significant instability in action potential duration, including

early afterdepolarizations, was detected when hypoxia was combined with simulated

activation of the sympathetic nervous system using isoproterenol in adult guinea pig

myocytes or the Luo-Rudy *in silico* model.[50] Therefore, the hypoxemia and sympathetic

activation associated with SDB would be expected to increase QT variability.

Beat-to-beat variability of the QT interval, which is incorporated in QTVI, was

proposed by Berger et al.[36] as a measure of ventricular repolarization lability to forecast

the occurrence of reentrant ventricular arrhythmias in patients with dilated

cardiomyopathy. In the absence of cardiac pathology, QTVI increases with age and is

higher in women than men.[51] Studies that have characterized QTVI have consistently

found that a higher QTVI is associated with greater risk for ventricular arrhythmias,

sudden cardiac death, and both all-cause and cardiovascular mortality.[17-22] There are

major gaps in our understanding of the association between SDB and QT variability, and

the available limited data are conflicting due to the methodological differences in study

design and the lack of control for confounding factors.[29-31] The data presented herein

provide unequivocal support for the role of SDB in adversely influencing QTVI given the

availability of cross-sectional, longitudinal, and experimental evidence. By minimizing

confounding in carefully selected samples, the current study also provides support for

ventricular repolarization instability as a potential mediator of all-cause mortality in SDB.

The argument for a causal link between SDB, ventricular repolarization instability,

and mortality is furthered by the observation in the current study that experimental

exposure to intermittent hypoxemia for as little as four hours destabilized ventricular

repolarization. The lack of an association between the frequency of arousals from sleep

and QTVI in the current study could be related to the small sample sizes in the cross-

sectional and longitudinal analyses. Recurrent arousals from sleep increase sympathetic

nervous system activity,[52] which can increase QT variability and arrhythmogenicity. In

fact, data from an epidemiological study of older men showed that excessive QTVI during

arousals predicted all-cause and cardiovascular mortality.[53] Such arousal-induced

increases in QTVI appeared to be transient, as there were no associations between pre-

arousal QTVI and mortality. Our results, however, suggest a cumulative impact of

intermittent hypoxemia on cardiac repolarization. Indeed, in the longitudinal analysis, the

mean QTVI increased to -0.86 in those with incident SDB without any concurrent medical

conditions, including heart failure. The high level of repolarization instability present with

incident SDB is comparable to that seen in heart failure patients with an increased risk

for mortality.[22] Additionally, the QTVI levels seen with incident SDB were similar to that of

patients on chronic methadone therapy, who experience nocturnal hypoxemia and a

predisposition for high density of premature ventricular contractions during sleep.[54]

Methadone, an opioid that inhibits both hERG and I_{K1} currents,[55] is associated with a

significant risk for arrhythmias and sudden nocturnal death.[56] The present study suggests

that a similar proarrhythmic effect might be associated with intermittent hypoxemia and

sympathetic stimulation.

Several strengths and limitations of the current study should be discussed.

Strengths include the matching of study participants with severe SDB to those without

SDB for the cross-sectional comparisons on QTVI, thus minimizing the effects of

confounding covariates, assessing the effects of incident SDB on QTVI, demonstrating

the impact of experimentally induced intermittent hypoxemia, and the use of full-montage

polysomnography to characterize SDB and ventricular repolarization abnormalities during

sleep. Weaknesses included the limited sample size, the lack of data on a dose-response

association between SDB and QTVI, and the lack of an assessment of potential

interactions between SDB and other factors (i.e., age, sex) that increase QTVI. These

issues notwithstanding, the current study motivates investments in future efforts on

whether treatment of SDB with positive airway pressure therapy is associated with

improvements in ventricular repolarization and downstream effects on cardiovascular and

all-cause mortality, particularly in at-risk patients such as those with congestive heart

failure.

ARTICLE INFORMATION

Acknowledgments

The authors are grateful to Drs. Habil Zare (The University of Texas Health Science Center, San Antonio, TX) and Courtney M. Johnson (Johns Hopkins University School of Medicine, Baltimore, MD) for thoughtful discussions and suggestions.

Sources of Funding

This work was supported by the Defense Health Agency (HU00011920029 to MCH and SS), the Jay P Sanford Award (to SS), and the National Institutes of Health (HL117167, HL146709, HL167121, P50MD017347 to NMP).

Disclaimer

The opinions and assertions expressed herein are those of the authors. They do not reflect the official policy or position of the Uniformed Services University of the Health Sciences or the Department of Defense.

Supplemental Material

Supplemental Tables S1–S3

Disclosures

None.

REFERENCES

1.   Punjabi NM. The epidemiology of adult obstructive sleep apnea. *Proc Am Thorac Soc*. 2008;5:136-143

2.   Senaratna CV, Perret JL, Lodge CJ, Lowe AJ, Campbell BE, Matheson MC, et al. Prevalence of obstructive sleep apnea in the general population: A systematic review. *Sleep Med Rev*. 2017;34:70-81

3.   Heinzer R, Vat S, Marques-Vidal P, Marti-Soler H, Andries D, Tobback N, et al. Prevalence of sleep-disordered breathing in the general population: The hypnolaus study. *Lancet Respir Med*. 2015;3:310-318

4.   Peppard PE, Young T, Palta M, Skatrud J. Prospective study of the association between sleep-disordered breathing and hypertension. *N Engl J Med*. 2000;342:1378-1384

5.   Leung RS, Bradley TD. Sleep apnea and cardiovascular disease. *Am J Respir Crit Care Med*. 2001;164:2147-2165

6.   Drager LF, McEvoy RD, Barbe F, Lorenzi-Filho G, Redline S, Initiative I. Sleep apnea and cardiovascular disease: Lessons from recent trials and need for team science. *Circulation*. 2017;136:1840-1850

7.   Gottlieb DJ. Sleep apnea and cardiovascular disease. *Curr Diab Rep*. 2021;21:64

8.   Brown J, Yazdi F, Jodari-Karimi M, Owen JG, Reisin E. Obstructive sleep apnea and hypertension: Updates to a critical relationship. *Curr Hypertens Rep*. 2022;24:173-184

9.   Mehra R, Benjamin EJ, Shahar E, Gottlieb DJ, Nawabit R, Kirchner HL, et al. Association of nocturnal arrhythmias with sleep-disordered breathing: The sleep heart health study. *Am J Respir Crit Care Med*. 2006;173:910-916

10.   Monahan K, Storfer-Isser A, Mehra R, Shahar E, Mittleman M, Rottman J, et al. Triggering of nocturnal arrhythmias by sleep-disordered breathing events. *J Am Coll Cardiol*. 2009;54:1797-1804

11.   Mehra R, Chung MK, Olshansky B, Dobrev D, Jackson CL, Kundel V, et al. Sleep-disordered breathing and cardiac arrhythmias in adults: Mechanistic insights and clinical implications: A scientific statement from the american heart association. *Circulation*. 2022;146:e119-e136

12.   Gami AS, Howard DE, Olson EJ, Somers VK. Day-night pattern of sudden death in obstructive sleep apnea. *N Engl J Med*. 2005;352:1206-1214

13.   Yaggi HK, Concato J, Kernan WN, Lichtman JH, Brass LM, Mohsenin V. Obstructive sleep apnea as a risk factor for stroke and death. *N Engl J Med*. 2005;353:2034-2041

14.   Gami AS, Olson EJ, Shen WK, Wright RS, Ballman KV, Hodge DO, et al. Obstructive sleep apnea and the risk of sudden cardiac death: A longitudinal study of 10,701 adults. *J Am Coll Cardiol*. 2013;62:610-616

15.   Blackwell JN, Walker M, Stafford P, Estrada S, Adabag S, Kwon Y. Sleep apnea and sudden cardiac death. *Circ Rep*. 2019;1:568-574

16.   Dobson CP, Kim A, Haigney M. Qt variability index. *Prog Cardiovasc Dis*. 2013;56:186-194

17.   Haigney MC, Zareba W, Gentlesk PJ, Goldstein RE, Illovsky M, McNitt S, et al. Qt interval variability and spontaneous ventricular tachycardia or fibrillation in the multicenter automatic defibrillator implantation trial (madit) ii patients. *J Am Coll Cardiol*. 2004;44:1481-1487

18.   Piccirillo G, Magri D, Matera S, Magnanti M, Torrini A, Pasquazzi E, et al. Qt variability strongly predicts sudden cardiac death in asymptomatic subjects with mild or moderate left ventricular systolic dysfunction: A prospective study. *Eur Heart J*. 2007;28:1344-1350

19.   Haigney MC, Zareba W, Nasir JM, McNitt S, McAdams D, Gentlesk PJ, et al. Gender differences and risk of ventricular tachycardia or ventricular fibrillation. *Heart Rhythm*. 2009;6:180-186

20. Tereshchenko LG, Fetics BJ, Domitrovich PP, Lindsay BD, Berger RD. Prediction of ventricular tachyarrhythmias by intracardiac repolarization variability analysis. *Circ Arrhythm Electrophysiol*. 2009;2:276-284

21. Tereshchenko LG, Cygankiewicz I, McNitt S, Vazquez R, Bayes-Genis A, Han L, et al. Predictive value of beat-to-beat qt variability index across the continuum of left ventricular dysfunction: Competing risks of noncardiac or cardiovascular death and sudden or nonsudden cardiac death. *Circ Arrhythm Electrophysiol*. 2012;5:719-727

22. Dobson CP, La Rovere MT, Pinna GD, Goldstein R, Olsen C, Bernardinangeli M, et al. Qt variability index on 24-hour holter independently predicts mortality in patients with heart failure: Analysis of gruppo italiano per lo studio della sopravvivenza nell'insufficienza cardiaca (gissi-hf) trial. *Heart Rhythm*. 2011;8:1237-1242

23. Ahnve S, Vallin H. Influence of heart rate and inhibition of autonomic tone on the qt interval. *Circulation*. 1982;65:435-439

24. Browne KF, Zipes DP, Heger JJ, Prystowsky EN. Influence of the autonomic nervous system on the q-t interval in man. *Am J Cardiol*. 1982;50:1099-1103

25. Magnano AR, Holleran S, Ramakrishnan R, Reiffel JA, Bloomfield DM. Autonomic nervous system influences on qt interval in normal subjects. *J Am Coll Cardiol*. 2002;39:1820-1826

26. Yeragani VK, Pohl R, Jampala VC, Balon R, Kay J, Igel G. Effect of posture and isoproterenol on beat-to-beat heart rate and qt variability. *Neuropsychobiology*. 2000;41:113-123

27. Seethala S, Shusterman V, Saba S, Mularski S, Nemec J. Effect of beta-adrenergic stimulation on qt interval accommodation. *Heart Rhythm*. 2011;8:263-270

28. Nayyar S, Roberts-Thomson KC, Hasan MA, Sullivan T, Harrington J, Sanders P, et al. Autonomic modulation of repolarization instability in patients with heart failure prone to ventricular tachycardia. *Am J Physiol Heart Circ Physiol*. 2013;305:H1181-1188

29. Baumert M, Smith J, Catcheside P, McEvoy RD, Abbott D, Sanders P, et al. Variability of qt interval duration in obstructive sleep apnea: An indicator of disease severity. *Sleep*. 2008;31:959-966

30. Viigimae M, Karai D, Pilt K, Pirn P, Huhtala H, Polo O, et al. Qt interval variability index and qt interval duration during different sleep stages in patients with obstructive sleep apnea. *Sleep Med*. 2017;37:160-167

31. Schmidt M, Baumert M, Penzel T, Malberg H, Zaunseder S. Nocturnal ventricular repolarization lability predicts cardiovascular mortality in the sleep heart health study. *Am J Physiol Heart Circ Physiol*. 2019;316:H495-H505

32. Quan SF, Howard BV, Iber C, Kiley JP, Nieto FJ, O'Connor GT, et al. The sleep heart health study: Design, rationale, and methods. *Sleep*. 1997;20:1077-1085

33. Redline S, Sanders MH, Lind BK, Quan SF, Iber C, Gottlieb DJ, et al. Methods for obtaining and analyzing unattended polysomnography data for a multicenter study. Sleep heart health research group. *Sleep*. 1998;21:759-767

34. Zhang GQ, Cui L, Mueller R, Tao S, Kim M, Rueschman M, et al. The national sleep research resource: Towards a sleep data commons. *J Am Med Inform Assoc*. 2018;25:1351-1358

35. Louis M, Punjabi NM. Effects of acute intermittent hypoxia on glucose metabolism in awake healthy volunteers. *J Appl Physiol (1985)*. 2009;106:1538-1544

36. Berger RD, Kasper EK, Baughman KL, Marban E, Calkins H, Tomaselli GF. Beat-to-beat qt interval variability: Novel evidence for repolarization lability in ischemic and nonischemic dilated cardiomyopathy. *Circulation*. 1997;96:1557-1565

37. Tibshirani R. Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society: Series B (Methodological)*. 1996;58:267-288

38. Friedman J, Hastie T, Tibshirani R. Regularization paths for generalized linear models via coordinate descent. *J Stat Softw*. 2010;33:1-22

39. Ludka O, Konecny T, Somers V. Sleep apnea, cardiac arrhythmias, and sudden death. *Tex Heart Inst J*. 2011;38:340-343

40. Aizawa Y, Kohno T, Yuasa S, Fukuda K. The role of circadian rhythms in fatal arrhythmias and the potential impact of intervention for sleep-disordered breathing. *Curr Pharm Des*. 2015;21:3512-3522

41. Kerns ES, Kim ED, Meoni LA, Sozio SM, Jaar BG, Estrella MM, et al. Obstructive sleep apnea increases sudden cardiac death in incident hemodialysis patients. *Am J Nephrol*. 2018;48:147-156

42. Bencardino G, Vitulano N, Bisignani A, Gabrielli FA, Pelargonio G, Narducci ML, et al. Severity of spleep apnea syndrome and life-threatening tachyarrhythmias in patients with implantable cardioverter defibrillator. *Pacing Clin Electrophysiol*. 2021;44:1657-1662

43. Heilbrunn ES, Ssentongo P, Chinchilli VM, Oh J, Ssentongo AE. Sudden death in individuals with obstructive sleep apnoea: A systematic review and meta-analysis. *BMJ Open Respir Res*. 2021;8

44. Linz B, Sattler SM, Flethoj M, Hotbjerg Hansen ME, Hesselkilde EM, Saljic A, et al. Arrhythmogenic mechanisms of acute obstructive respiratory events in a porcine model of drug-induced long qt. *Heart Rhythm*. 2021;18:1384-1391

45. Morand J, Arnaud C, Pepin JL, Godin-Ribuot D. Chronic intermittent hypoxia promotes myocardial ischemia-related ventricular arrhythmias and sudden cardiac death. *Sci Rep*. 2018;8:2997

46. Ju YK, Saint DA, Gage PW. Hypoxia increases persistent sodium current in rat ventricular myocytes. *J Physiol*. 1996;497 (Pt 2):337-347

47. Fearon IM, Palmer AC, Balmforth AJ, Ball SG, Mikala G, Schwartz A, et al. Hypoxia inhibits the recombinant alpha 1c subunit of the human cardiac l-type ca2+ channel. *J Physiol*. 1997;500 (Pt 3):551-556

48. Smani T, Hernandez A, Urena J, Castellano AG, Franco-Obregon A, Ordonez A, et al. Reduction of ca(2+) channel activity by hypoxia in human and porcine coronary myocytes. *Cardiovasc Res*. 2002;53:97-104

49. Hool LC. Hypoxia increases the sensitivity of the l-type ca(2+) current to beta-adrenergic receptor stimulation via a c2 region-containing protein kinase c isoform. *Circ Res*. 2000;87:1164-1171

50. Gaur N, Rudy Y, Hool L. Contributions of ion channel currents to ventricular action potential changes and induction of early afterdepolarizations during acute hypoxia. *Circ Res*. 2009;105:1196-1203

51. Baumert M, Porta A, Vos MA, Malik M, Couderc JP, Laguna P, et al. Qt interval variability in body surface ecg: Measurement, physiological basis, and clinical value: Position statement and consensus guidance endorsed by the european heart rhythm association jointly with the esc working group on cardiac cellular electrophysiology. *Europace*. 2016;18:925-944

52. Taylor KS, Murai H, Millar PJ, Haruki N, Kimmerly DS, Morris BL, et al. Arousal from sleep and sympathetic excitation during wakefulness. *Hypertension*. 2016;68:1467-1474

53. Shahrbabaki SS, Linz D, Redline S, Stone K, Ensrud K, Baumert M. Sleep arousal-related ventricular repolarization lability is associated with cardiovascular mortality in older community-dwelling men. *Chest*. 2022

54. Solhjoo S, Punjabi NM, Ivanescu AE, Crainiceanu C, Gaynanova I, Wicken C, et al. Methadone destabilizes cardiac repolarization during sleep. *Clin Pharmacol Ther*. 2021;110:1066-1074

55. Klein MG, Krantz MJ, Fatima N, Watters A, Colon-Sanchez D, Geiger RM, et al. Methadone blockade of cardiac inward rectifier k(+) current augments membrane

instability and amplifies u waves on surface ecgs: A translational study. *J Am Heart Assoc*. 2022;11:e023482

56.   Kao DP, Haigney MC, Mehler PS, Krantz MJ. Arrhythmia associated with buprenorphine and methadone reported to the food and drug administration. *Addiction*. 2015;110:1468-1475

FIGURE LEGENDS

Figure 1. QTVI was higher in participants with severe SDB. The diagram shows the mean QTVI measurements for participants without SDB (N = 61, red) and with severe SDB (N = 61, blue).

Figure 2: Incident SDB destabilized ventricular repolarization. The diagram shows QTVI for participants with and without incident SDB. QTVI significantly increased in participants who developed SDB. N = 26 in each group. *P* values are for the paired samples *t* test between baseline and follow-up measurements for each group.

Figure 3. Temporal effects of intermittent hypoxia on electrocardiographic measures. The left panel shows data on one subject during ~ 20 minutes of exposure to intermittent hypoxia. Cyclical changes were noted in the heart rate and QT interval with intermittent hypoxia. The right panel shows the NN interval, SDNN, LF/(HF+LF) ratio, and QTVI of the 19 subjects for each hour of the four hours of exposure to intermittent hypoxia and normoxia. * indicates *P* < 0.05 for comparing the two conditions during the corresponding hour.

Table 1. Demographic, polysomnographic, and electrocardiographic data for the cross-sectional sample comparing severe SDB versus no SDB.

Parameters	No SDB	Severe SDB	*P* value
Demographic			
Age (years)	60.54 (1.00)	60.72 (1.01)	0.899
BMI (kg/m^2)	29.28 (0.48)	29.45 (0.50)	0.802
Polysomnographic			
AHI (/hour)	0.62 (0.05)	46.28 (1.59)	< 0.001
Mean SpO$_2$ (%)	95.07 (0.23)	93.75 (0.20)	< 0.001
TST$_{90}$ (%)	1.32 (0.61)	11.39 (1.42)	< 0.001
Electrocardiographic			
QTVI	-1.43 (0.06)	-1.19 (0.08)	0.027
HR (beats/min)	64.92 (1.11)	68.34 (1.08)	0.028
SDNN (msec)	32.01 (2.21)	41.48 (3.25)	0.018
QTc (msec)	442.58 (4.80)	457.86 (4.20)	0.018

Values are the mean (SEM). *P* values are for the Student's *t* test comparing matched participants with and without SDB. N = 61 for each group.

Table 2. Demographic, polysomnographic, and electrocardiographic data for the longitudinal sample of the participants with and without incident SDB.

Parameters	Baseline		P value	Follow-up		P value
	Without SDB	Incident SDB		Without SDB	Incident SDB	
Demographic						
Age (years)	63.88 (1.86)	64.08 (1.87)	0.942	69.15 (1.83)	69.12 (1.84)	0.988
BMI (kg/m^2)	27.24 (0.87)	27.70 (1.00)	0.732	26.87 (0.81)	28.45 (1.18)	0.272
Polysomnographic						
AHI (/hour)	2.79 (0.58)	2.54 (0.72)	0.793	3.36 (0.67)	13.74 (3.50)	0.005
Mean SpO$_2$ (%)	95.81 (0.21)	94.98 (0.24)	0.012	95.59 (0.23)	91.26 (0.15)	< 0.001
TST$_{90}$ (%)	0.05 (0.01)	0.05 (0.01)	0.905	0.03 (0.01)	21.79 (2.08)	< 0.001
Electrocardiographic						
QTVI	-1.26 (0.09)	-1.23 (0.15)	0.876	-1.24 (0.10)	-0.86 (0.14)	0.034
HR (beats/min)	64.16 (1.51)	64.68 (1.74)	0.824	64.10 (1.75)	66.86 (2.08)	0.315
SDNN (msec)	28.39 (2.64)	27.66 (2.69)	0.846	27.43 (2.34)	24.41 (3.01)	0.432
QTc (msec)	436.85 (5.43)	447.39 (8.40)	0.297	444.93 (5.27)	450.34 (6.03)	0.503

Values are the mean (SEM). *P* values are for the Student's *t* test comparing participants who developed SDB (N = 26) to those who did not (N = 26).

Table 3. Electrographic measures during normoxia versus intermittent hypoxia.

Electrocardiographic Parameters	1st hour	4th hour	*P* value
Normoxia			
QTVI	-1.89 (0.09)	-1.86 (0.06)	0.412
HR (beats/min)	63.61 (2.43)	64.73 (2.15)	0.183
SDNN (msec)	94.28 (8.99)	90.56 (7.65)	0.336
Log(HF)	3.11 (0.11)	3.06 (0.10)	0.189
LF/(HF+LF)	0.59 (0.02)	0.59 (0.03)	0.715
Intermittent hypoxia			
QTVI	-1.85 (0.04)	-1.64 (0.07)	0.016
HR	63.31 (1.51)	69.21 (2.14)	0.001
SDNN	111.15 (7.99)	104.19 (8.29)	0.317
Log(HF)	3.17 (0.09)	3.01 (0.09)	0.046
LF/(HF+LF)	0.61 (0.03)	0.67 (0.02)	0.014

Values are the mean (SEM). *P* values are for the paired *t* test between normoxia and intermittent hypoxia experiments. N = 19.

Figure 1. QTVI was higher in participants with severe SDB. The diagram shows the

mean QTVI measurements for participants without SDB (N = 61, red) and with severe

SDB (N = 61, blue).

Figure 2. Incident SDB destabilized ventricular repolarization. The diagram shows QTVI for participants with and without incident SDB. QTVI significantly increased in participants who developed SDB. N = 26 in each group. *P* values are for the paired samples *t* test between baseline and follow-up measurements for each group.

Figure 3. Temporal effects of intermittent hypoxia on electrocardiographic measures. The left panel shows data on one subject during ~ 20 minutes of exposure to intermittent hypoxia. Cyclical changes were noted in the heart rate and QT interval with intermittent hypoxia. The right panel shows the NN interval, SDNN, LF/(HF+LF) ratio, and QTVI of the 19 subjects for each hour of the four hours of exposure to intermittent hypoxia and normoxia. * indicates *P* < 0.05 for comparing the two conditions during the corresponding hour.

Supplemental Material

Sleep-Disordered Breathing Destabilizes Ventricular Repolarization

Soroosh Solhjoo, PhD[1], Mark C. Haigney, MD[1,2], Trishul Siddharthan, MD[3], Abigail Koch, MD[3], Naresh M. Punjabi, MD, PhD[3]

[1] F. Edward Hébert School of Medicine, Bethesda, Maryland, USA
[2] Military Cardiovascular Outcomes Research (MiCOR), Bethesda, Maryland, USA
[3] University of Miami Miller School of Medicine, Miami, Florida, USA

Short Title: Sleep apnea, intermittent hypoxemia, and QTVI

Corresponding Author:

Soroosh Solhjoo, PhD
F. Edward Hébert School of Medicine
4301 Jones Bridge Road
Bethesda, MD 20814
soroosh@jhmi.edu

Table S1. Stratified analysis of electrocardiographic parameters and oxygen saturation levels for the cross-sectional sample of the SHHS participants with severe SDB versus no SDB.

Parameters	No SDB	Severe SDB	*P* value
NREM			
Mean SpO_2 (%)	94.98 (0.23)	93.80 (0.20)	< 0.001
Minimum SpO_2 (%)	94.53 (0.23)	91.53 (0.27)	< 0.001
QTVI	-1.43 (0.07)	-1.19 (0.08)	0.033
HR (beats/min)	65.03 (1.17)	68.14 (1.02)	0.048
SDNN (msec)	32.88 (2.36)	42.12 (3.42)	0.028
REM			
Mean SpO_2	95.27 (0.26)	93.22 (0.29)	< 0.001
Minimum SpO_2	94.58 (0.28)	90.00 (0.50)	< 0.001
QTVI	-1.41 (0.06)	-1.15 (0.09)	0.019
HR	65.69 (1.11)	68.90 (1.22)	0.053
SDNN	32.63 (2.31)	40.62 (3.04)	0.038

Values are the mean (SEM). *P* values are for the Student's *t* test comparing matched participants with and without SDB. N = 61 for each group.

Table S2. Stratified analysis of electrocardiographic parameters and oxygen saturation levels for the longitudinal sample of the participants with and without incident SDB.

Parameters	Baseline		P value	Follow-up		P value
	No SDB	Incident SDB		No SDB	Incident SDB	
NREM						
Mean SpO$_2$ (%)	95.75 (0.22)	94.96 (0.24)	0.018	95.54 (0.23)	91.34 (0.15)	< 0.001
QTVI	-1.28 (0.10)	-1.23 (0.15)	0.778	-1.24 (0.10)	-0.86 (0.14)	0.031
HR (beats/min)	63.76 (1.50)	64.63 (1.73)	0.706	63.93 (1.75)	66.87 (2.07)	0.283
SDNN (msec)	29.71 (3.07)	27.19 (2.48)	0.526	27.31 (2.36)	24.11 (3.28)	0.434
QTc (msec)	436.27 (5.45)	447.47 (8.50)	0.273	444.66 (5.29)	450.70 (6.13)	0.459
REM						
Mean SpO$_2$	96.11 (0.24)	95.15 (0.24)	0.007	95.84 (0.26)	90.51 (0.25)	< 0.001
QTVI	-1.18 (0.10)	-1.31 (0.17)	0.493	-1.27 (0.10)	-1.01 (0.15)	0.163
HR	65.35 (1.45)	65.17 (1.96)	0.943	65.75 (1.97)	65.55 (2.20)	0.945
SDNN	25.93 (2.14)	30.54 (3.86)	0.301	28.06 (2.94)	25.99 (3.22)	0.636
QTc	440.08 (5.83)	447.43 (8.72)	0.484	446.94 (5.91)	445.69 (5.03)	0.873

Values are the mean (SEM). P values are for the Student's t test comparing participants with and without incident SDB.

Table S3. Oxygen saturation levels during normoxia and intermittent hypoxia.

SpO$_2$ (%)	Normoxia	Intermittent Hypoxia	*P* value
Mean	97.37 (0.20)	90.90 (0.20)	< 0.001
Minimum	96.11 (0.29)	81.05 (0.50)	< 0.001
SD	0.56 (0.03)	4.53 (0.14)	< 0.001

Values are the mean (SEM). *P* values are for the paired samples *t* test between normoxia and intermittent hypoxia experiments. N = 19.