**Title Page**

**Title**: Relationship between adherence to the 2019 Canada's Food Guide recommendations on healthy food choices and nutrient intakes in older adults

**Author names:**

Brassard, Didier (1)

Chevalier, Stéphanie (1, 2)

**Affiliations:**

1)  School of Human Nutrition, McGill University, Québec, Canada
2)  Research Institute of the McGill University Health Centre, McGill University, Québec, Canada

**Sources of support:** This research received no external funding.

**Conflict of Interest and Funding Disclosure:** DB was a casual employee of Health Canada (2019-2020), held a doctoral training award from the *Fonds de recherche du Québec – Santé* (2019-2021) and holds a Canadian Institutes of Health Research (CIHR) Fellowship award (MFE-181852). SC receives research funding from the CIHR, Dairy Farmers of Canada and the Canadian Foundation for Dietetics Research. None of these agencies has funded nor was involved in this work. DB and SC have no conflicts of interest.

**Corresponding author:**

Stéphanie Chevalier, PhD,

School of Human Nutrition, McGill University

21111 Lakeshore Rd, Sainte-Anne-de-Bellevue, QC H9X 3V9

Tel : 514-398-8603

Email: stephanie.chevalier@mcgill.ca

**Running title**: Nutrient adequacy of older adults adhering to CFG

**Abbreviations list:** CCHS: Canadian Community Health Survey; CFG: Canada's Food Guide; DRI, Dietary Reference Intake; HEFI-2019: Healthy Eating Food Index-2019; NCI: National Cancer Institute.

**Abstract**

**Background:** Following Canada's food guide (CFG) recommendations should ensure adequate
nutrient intakes for persons aged 2 years and more. Older adults have increased needs in certain
nutrients and the extent to which adherence to CFG recommendations can help reduce inadequate
nutrient intakes is unknown.

**Objective:** To assess the relationship between adherence to CFG recommendations on healthy food
choices and intake of key nutrients in adults 65 years and older from the Canadian Community
Health Survey (CCHS) 2015 - Nutrition.

**Methods:** Secondary analysis of data from 4,093 adults of the CCHS 2015 - Nutrition (mean age,
73.6 y, 54% females). Dietary intakes were measured using an interviewer-administered 24-hour
dietary recall including one repeat in a subsample (42%). The National Cancer Institute
multivariate method was used to estimate usual (i.e., long-term) dietary intakes. Adherence to CFG
recommendations was measured using the Healthy Eating Food Index (HEFI)-2019 score. Simple
linear and logistic regression models estimated the effect of increased HEFI-2019 score on usual
nutrient intakes and the prevalence of inadequate nutrient intakes (i.e., below the estimated
average requirements), respectively.

**Results:** Compared with the prevalence of inadequate intakes at median HEFI-2019 score (46.4/80
points), a higher HEFI-2019 (+11 points) was associated with reductions in the prevalence of
inadequate intakes of magnesium, vitamin B6, and protein (-19.8% [95%CI: -30.8, -8.9], -12.7%
[95%CI: -22.5, -3.0], and -4.7% [95%CI: -9.4, -0.1], respectively). In contrast, data for higher HEFI-
2019 scores were compatible with increased prevalence of inadequate intakes of folate, vitamin D,
and calcium (4.0% [95%CI: -8.4, 16.3], 2.6% [95%CI: 1.1, 4.0], and 2.3% [95%CI: -3.0, 7.5],
respectively).

**Conclusions:** Based on dietary intakes of Canadian older adults in 2015, increasing the degree of
adherence to CFG recommendations on healthy food choices may reduce nutrient intake
inadequacy for most key nutrients except folate, vitamin D and calcium.

**Keywords:** older adults, 24-hour dietary recalls; CFG; HEFI-2019; Canada's Food Guide; dietary
guidelines; healthy eating food index; Canadian Community Health Survey; CCHS.

## Introduction

Previous editions of the Canada's Food Guide (CFG) informed on food choices to prevent malnutrition as the primary goal. Recommendations of the CFG-2007 were expressed in terms of number of servings to consume every day for broad categories of foods (Vegetables and fruits, Grains, Dairy and Alternatives, Meat and Alternatives) (1, 2). The CFG-2007 recommended daily servings based on comprehensive diet simulations, thus ensuring that Dietary Reference Intake (DRI) were met (2). The latest CFG (2019) adopted a different approach and primarily aims at chronic disease risk reduction (3-6). In addition, recommendations in CFG-2019 (e.g., "eat more often ...") are more flexible than those of CFG-2007 and rely on proportions of food categories to constitute a healthy plate, without recommended quantitative daily servings. Finally, the revision process of CFG-2019 was more transparent and more evidence-based than that of CFG-2007 (7).

The release of the CFG-2019 elicited positive reactions (8, 9), but also raised concerns on nutrient adequacy. For example, preliminary modeling of eating patterns consistent with the CFG-2019 plate snapshot revealed that adherence may be insufficient to meet calcium and vitamin D requirements (10). The ability of consumers and policymakers to properly implement CFG-2019 recommendations on protein foods was also found unclear (11). In the Canadian Community Health Survey (CCHS) 2015 – Nutrition, animal-based protein foods contributed more than two thirds of total protein intake (12), which contrasts with CFG-2019 recommendations aiming at increasing intake of plant-based foods. Accordingly, an inadequate implementation of CFG-2019 recommendations could lead to amplify the proportion of the population with inadequate nutrient intakes (e.g., substituting animal-based foods rich in protein with plant-based alternatives low in protein), especially for certain age groups at higher risk. In that regard, maintaining adequate nutrient intakes is a challenge for older adults (13, 14). Older adults face social (e.g., loneliness, inability to buy or prepare foods) and physiological (e.g., changes in taste, loss of appetite, malabsorption) barriers to consuming a nutrient-dense diet (14). Notably, calcium, vitamin D, fibers, potassium and protein were identified as nutrients of concern among older adults in the United States 2015 Dietary Guidelines report (15). In sum, the one-size-fits-all recommendations in CFG-2019 may not be adapted for older adults who face unique challenges to healthy eating compared with other strata of the population.

Whether adherence to CFG-2019 recommendations on healthy food choices fulfils nutritional needs of older Canadians is currently unknown; no evidence from contemporary eating patterns exists. The aim of this study was therefore to assess the relationship between adherence to CFG-2019 recommendations, using the Healthy Eating Food Index-2019 (HEFI-2019) (16, 17), and intakes of key nutrients of adults aged ≥65 years from the CCHS 2015 – Nutrition. We hypothesized that higher adherence to CFG-2019 recommendations shows an inverse relationship with intakes of nutrients more commonly found or eaten in animal-based foods (i.e., protein, calcium, vitamin D, iron, zinc, vitamin B6 and B12), and a positive relationship with intakes of nutrients more commonly found in plant-based foods (i.e., fibers, folate, magnesium, potassium, vitamin A).

**Methods**

**Study design and participants**

This study is based on a sample of older adults aged 65 years or older from the CCHS 2015 –
Nutrition (18). The CCHS 2015 -Nutrition is a nationally representative survey of individuals aged 1
year and older living in private dwellings in the 10 Canadian provinces. Full-time members of the
Canadian Forces and individuals living in the Territories, on reserves, in remote areas, and in
institutions were not included. Data collection occurred between January 1st to December 31, 2015.
Respondents aged less than 65 years (n=16,394) and those reporting zero energy on their first 24-
hour dietary recall intake were excluded (n=4), yielding a final sample of 4,089 respondents. The
public use microdata file (PUMF) of CCHS 2015 – Nutrition was obtained from Statistics Canada.

**Data collection and dietary assessment**

Interviewer-administered 24-hour dietary recalls were used to assess dietary intakes. The
interviews were mostly (98%) conducted in person for the first 24-hour recall and by telephone for
the second 24-hour recall. All 24-hour recall interviews were structured according to the
Automated Multiple Pass Method (18). Portion size estimation of foods and beverages in plates,
bowls, glasses and mugs was facilitated using a food booklet designed for the survey (18). All
respondents completed one 24-hour recall and a subsample of 1,706 respondents (42%) completed
a second 24-hour recall. Nutrient intakes were calculated based on the Canadian Nutrient File 2015
(19), except for intakes of free sugars which were recently published by Health Canada (20). Total
food intakes expressed in reference amounts (RA) (21) were calculated for each respondent, 24-
hour dietary recall and HEFI-2019 food and beverage categories. The interviewers measured body
weight of respondents with a standard scale (LifeSource Scales Model US-321).

*The Healthy Eating Food Index (HEFI) 2019*

The HEFI-2019 is a continuous score which measures the degree of adherence between dietary
intakes and Canada's Food Guide 2019 recommendations on healthy food choices. Complete details
about the development and the evaluation of the HEFI-2019 are available elsewhere (16, 17).
Briefly, the HEFI-2019 has 10 components including 5 based on intake of foods (*Vegetables and*
*fruits, Whole-grain foods, Grain foods ratio, Protein foods, Plant-based protein foods*), 1 on beverages
(*Beverages*), and 4 on nutrients (*Fatty acids ratio, Saturated fats, Free sugars, and Sodium*). The total
HEFI-2019 score sums up to 80 points with higher scores indicating greater adherence to
recommendations. Details about each component and scoring standards are presented in
**Supplemental Table 1**.

*Nutrient intakes*

Key nutrients for older adults were selected according to data availability in CCHS 2015 – Nutrition
as well as previous literature (13, 22). The nutrients were protein, calcium, vitamin D, iron, zinc,
vitamin B6, vitamin B12, dietary folate equivalent (folate hereafter), magnesium, potassium, fibers,

and retinol activity equivalent (vitamin A hereafter). The proportion of respondents with
inadequate intakes was estimated using the cut-point method and the Estimated Average
Requirements (EAR) of the Dietary Reference Intake (23). The EAR for each nutrient assessed are
presented in **Supplemental Table 2**. Because there is no EAR for potassium and fibers, the
prevalence of intake inadequacy cannot be assessed, but only the proportion of respondents with
intakes above the adequate intake (AI) value. Recognizing the consensus that recommended
protein intakes for older adults should be higher than current recommendations (24-26), we also
estimated the proportion of respondents with inadequate protein intakes at cut-offs higher than the
current EAR. The hypothetical EAR cut-offs of 0.8 and 1.0 grams of protein per kg of bodyweight
were selected to reflect hypothetical higher recommended daily allowance (RDA) of 1.0 and 1.2
g/kg, respectively.

### Statistical analyses

Sampling weights provided by Statistics Canada were used in all analyses to reflect the Canadian
population of older adults in 2015 as well as bootstrap replicate weights for variance estimation.
The sampling weights accounting for missing data on body weight were also used where
appropriate. Analyses were performed in SAS Studio v3.8 (SAS Institute) and R v4.2.2 (R
Foundation for Statistical Computing).

To assess the relationship between the HEFI-2019 score and nutrient intakes, analyses involved
four steps, detailed below: 1) measurement error correction of dietary intakes using the National
Cancer Institute (NCI) Markov Chain Monte Carlo multivariate method; 2) linear regression of
continuous nutrient intakes on total HEFI-2019 scores; 3) logistic regression of the proportion of
individuals below EAR on total HEFI-2019 scores; and 4) variance estimation using bootstrap
replicate weights provided by Statistics Canada.

First, the 24-hour dietary recalls are mainly affected by within-individual random errors, which
require correction to estimate distribution of intakes (27, 28). To account for correlated random
errors of all dietary constituents of the HEFI-2019 (e.g., vegetables and fruits, free sugars) and those
of nutrient intakes (e.g., protein), the NCI multivariate method was used (29, 30). Briefly, the
multivariate method uses Monte Carlo simulation to estimate distribution of usual intakes (i.e.,
long-term average) of multiple dietary constituents correlated with each other. The multivariate
methods also accounts for systematic differences due to "nuisance" factors (day of the week,
sequence of recall) and considers foods that are episodically consumed (e.g., plant-based protein
foods) (29). Nutrients with common food sources were modelled together to have parsimonious
measurement error correction models (see Supplemental Methods and Supplemental Table 2). The
15 dietary constituents of the HEFI-2019 were included in each model. Whole-grain foods, refined
grain foods, plant-based protein foods, beverages not recommended (i.e., sugary drinks, artificially
sweetened beverages, vegetable and fruit juices, sweetened milk and plant-based beverages,
alcohol) and unsweetened milk were considered episodically consumed. All the remaining foods
and nutrients were considered as consumed daily. The measurement error correction models were
stratified by sex to reflect sex-specific random variations in dietary intakes and to derive sex-
specific associations (30, 31). The models also included the covariates age (indicator for 71 years or
older), sequence of recall (indicator for second recall) and weekend (indicator for a 24-hour recall

of intakes on Friday, Saturday or Sunday). In addition, the model for protein intake also included
body weight (kg) as a covariate to derive protein intake per kg and assess the proportion of
individuals below the EAR. A total of 500 pseudo-individuals were generated in the Monte Carlo
simulation step of the NCI multivariate method. Simulations from each stratum (i.e., males and
females) were pooled together. The HEFI-2019 scoring algorithm was then applied using the
modelled dietary constituents among the 500 pseudo-individuals to derive HEFI-2019 scores.

Second, simple linear regression models were used to assess the relationship between the
continuous HEFI-2019 score, as the independent variable, and continuous nutrient intakes, as the
dependent variable. A restricted cubic spline transformation with 5 knots (percentiles 5, 27, 50, 73
and 95) was applied *a priori* to the HEFI-2019 score to assess potential non-linearity (32). To
estimate change according to feasible increases in adherence, expected nutrient intake differences
were calculated according to an increase in HEFI-2019 score from the median score to the 90[th]
percentile of the usual intake distribution. In other words, we estimated nutrient intake difference
according to a hypothetical change where respondents would have had high HEFI-2019 scores
compared with the HEFI-2019 score respondents had on average, taken as the median of this
sample.

Third, logistic regression models were used to assess the odds of having nutrient intake below the
age- and sex-specific EAR and the continuous HEFI-2019 score. A restricted cubic spline
transformation was also applied to the HEFI-2019 score. Predicted odds of nutrient intake
inadequacy were generated for the 90[th] percentile and the median. Both predicted odds were then
re-expressed as risk of inadequacy ($risk = \frac{e^X}{1+e^X}$; where $X$ corresponds to predicted odds at a given
HEFI-2019 score percentile). The expected change in the prevalence of inadequate nutrient intakes,
i.e., risk difference, was calculated by the difference between the estimated risk of inadequacy at the
90[th] percentile vs. the median HEFI-2019 score.

Fourth, steps 1 to 3 were repeated 500 times using bootstrap replicate weights to generate
standard errors and 95% CI via normal approximation. The convergence of bootstrap standard
errors and the normality of bootstrap estimates were confirmed graphically. Data from one
bootstrap replicate was removed for potassium due to non-convergence.

**Results**

**Total HEFI-2019 score**

The mean (SD) HEFI-2019 score among all adults 65 years and older was 46.0 (8.9) (/80 points).
Among age and sex subgroups, the mean HEFI-2019 score was the highest in females, 65 to 70
years (48.2 points) and the lowest in males, 65 to 70 years (44.9 points; **Figure 1**).

**Prevalence of inadequate nutrient intake**

The prevalence of inadequate nutrient intakes ranged from 1% (iron) to 96% (vitamin D; **Figure 2**.
The prevalence of inadequate intakes was high for vitamin D, calcium, magnesium, vitamin B6, and
vitamin A (96%, 83%, 64%, 38%, and 36%, respectively; Supplemental Figure 1. Results were
similar for most nutrients when stratified by DRI age and sex group (**Supplemental Figure 1**).

**Relationship between HEFI-2019 score and nutrient intakes**

Relationships between the HEFI-2019 score and nutrient intakes are illustrated in **Figure 3**.
Respondents with higher HEFI-2019 score had higher intakes of fibers, magnesium, vitamin B6,
potassium, and protein, but lower intakes of vitamin D, vitamin B12, dietary folate equivalent, and
iron. The HEFI-2019 score had null associations or associations that plateaued at higher scores for
vitamin A, calcium, and zinc. **Table 1** presents the expected differences in nutrient intakes
associated with an increase in HEFI-2019 score to the 90[th] percentile of the score distribution
compared with median HEFI-2019 score. For example, increasing HEFI-2019 from the median to
the 90[th] percentile of the score distribution was associated with a 3.4 g/day higher fiber intake
(95%CI: 2.0, 4.8; Table 1). In contrast, the same HEFI-2019 score increase was associated with a 0.5
μg/day lower vitamin D intake (95%CI: -1.0, 0.0; Table 1).

**Relationship between the HEFI-2019 score and nutrient intake adequacy**

**Inadequate nutrient intakes**

**Figure 4** presents the prevalence and the difference in the prevalence of inadequate nutrient intake
according to the HEFI-2019 score. An increase of HEFI-2019 score to the 90[th] percentile of the score
distribution compared with median HEFI-2019 score was associated with reduction of 20, 13, and 5
percentage point in the prevalence of inadequate nutrient intakes for magnesium, vitamin B6, and
protein, respectively. Inversely, the same HEFI-2019 score increase was associated with a minor
increase in the prevalence of inadequate vitamin D intakes (prevalence difference, +2.6%; 95%CI:
1.1, 4.0%). For dietary folate equivalent, the 95%CI was compatible with an increase in the
prevalence of inadequate intakes as large as +16.3%, but also compatible with a reduction as low as
-8.4%. For calcium, data were compatible with a negligible reduction in the prevalence of

inadequacy (as low as -3%), but also an increase in the prevalence of inadequacy as large as +7.5%
(Figure 3).

***Hypothetical higher Estimated Average Requirements for protein***

Compared with the median HEFI-2019 score, scores at the 90[th] percentile decreased the proportion
of respondents with protein intakes below hypothetical EAR of 0.8 g/kg/day and 1.0 g/kg/day
(**Supplemental Figure 2**). For 0.8 g/kg/day, the prevalence of inadequacy decreased from 29.3%
to 18.5% (prevalence difference: -10.8; 95%CI: -20.0, -1.6). For 1.0 g/kg/day, the prevalence
decreased from 64.7% to 50.8% (prevalence difference: -13.9; 95%CI: -25.0, -2.8).

**Adequate intakes for fibers and potassium**

Compared with the median HEFI-2019 score, score at the 90[th] percentile increased the proportion
of respondents with intakes above the adequate intake for both fibers and potassium. For fibers, the
prevalence increased from 6.5% to 20.5% (prevalence difference: +14.1; 95%CI: 5.1, 23.0). For
potassium, the prevalence increased from 25.1% to 36.8% (prevalence difference: +11.7; 95%CI:
2.2, 21.1).

**Discussion**

The objective of this study was to describe the relationship between adherence to CFG-2019 recommendations on healthy food choices, measured using the HEFI-2019, and nutrient intakes in adults 65 years and older from Canada in 2015. We found that higher adherence was associated with higher intakes of fibres, magnesium, vitamin B6, potassium and protein. Had respondents had a higher adherence, we estimated that the prevalence of nutrient inadequacy based on foods for magnesium, vitamin B6 and protein would have been 20%, 13% and 5% lower, respectively. However, a higher adherence was not associated with higher intakes of calcium, zinc, iron, folate, vitamin B12 and vitamin D. In turn, we estimated that the high prevalence of calcium inadequacy would not have changed and that of food-based vitamin D would have been 3% higher in 2015. Overall, these results indicate that CFG-2019 recommendations are insufficient to mitigate nutrient intake inadequacy for certain key nutrients based on the eating patterns of adults aged 65 years or more from Canada in 2015. These findings partially confirm our hypothesis that nutrients typically found in animal-based foods (e.g., iron, vitamin B12, calcium, vitamin D) were inversely associated with adherence to CFG.

Few studies have examined nutrient intake adequacy according to adherence to the CFG-2019 recommendations. Barr (10) assessed the probability of intake inadequacy upon adherence to an eating pattern consistent with foods depicted in the CFG-2019 plate snapshot. The probability of inadequacy was near 100% for calcium and vitamin D for males and females aged 71 years or more. This result is consistent with findings in the present study, where a higher adherence to recommendations on healthy food choices was insufficient to mitigate the prevalence of inadequacy for calcium and vitamin D. The lack of specific recommendations regarding dairy foods and alternatives in CFG-2019, and consequently the absence of points in the HEFI-2019, may partly explain these findings. Indeed, the main food source of calcium and vitamin D in Canadians' diet was the "Milk & Alternatives" food group in 2015 (33-35). Of note, the prevalence of inadequacy for vitamin D was very high in this sample (>95%), supporting the notion that meeting vitamin D recommendations through foods alone is an important challenge (35). Similarly, data from the CCHS indicate that calcium intake inadequacy in supplement non-users increased from 58% to 68% between 2004 and 2015, respectively (34). All in all, higher adherence to CFG-2019 recommendations on food choices was insufficient to mitigate the high prevalence of inadequate calcium and vitamin D intakes. Thus, meeting calcium and vitamin D needs would require additional strategies such as specific recommendations in CFG in addition to supplementation or food fortification.

A review by Fernandez et al. highlighted potential gaps in the application of CFG-2019 recommendations, notably whether the CFG-2019 permitted the adequate consumption of nutrient-rich protein foods or not (11). Our results revealed that a higher adherence to CFG-2019 recommendations slightly reduced the low proportion of older adults with intakes below the current EAR (0.66 g/kg) and reduced by 11% and 14% the proportion of those below the

hypothetical EAR of 0.8 g/kg and 1.0 g/kg. Notwithstanding, the prevalence of inadequacy for the
hypothetical EAR would remain considerable, with 25% and 50% inadequate protein intakes,
respectively. Furthermore, these findings only reflect the total daily protein intake and the impact
on overall protein food quality or distribution during the day is unknown. Future studies should
investigate the relationship between higher adherence to CFG-2019 and overall protein food
quality and should examine how to increase consumption of high-quality plant-based protein foods.
In that regard, a qualitative study among older adults highlighted that health benefits and food
preparation skills would facilitate consumption of plant-based protein foods (36). Finally, we stress
that the CCHS 2015 – Nutrition excluded individuals living in institutions who may have additional
needs as well as barriers to consuming higher protein diets (14). The inverse associations between
the HEFI-2019 and intakes of iron, folate, and vitamin B12 also raise concerns. On one hand, the
prevalence of inadequacy for these nutrients did not increase at the population level, partly because
the EAR can be met at relatively low intakes for these nutrients. On the other hand, at the individual
level, meeting nutrient requirements while following CFG-2019 recommendations may be more
difficult in individuals at risk of inadequate intakes, e.g., older adults with loss of appetite (13, 14).
Further, vitamin B12 may be particularly important for older adults and recent evidence support
that food groups contribute differently at reducing risk of deficiency (37). In sum, CFG-2019
recommendations should be carefully implemented to ensure meeting nutrient intakes both at the
population and individual level.

The use of national survey data to estimate the impact of higher adherence on the prevalence of
nutrient inadequacy in Canada is a strength of this study. But it must be recognized that
relationships between adherence to CFG-2019 recommendations and nutrient intakes reflect the
underlying eating patterns of adults 65 years or older from Canada in 2015. A variety of underlying
food and beverage consumption patterns may be highly consistent with CFG-2019
recommendations yet, have diverging associations with intakes of certain nutrients. As well, eating
patterns of older adults, though generally stable, may evolve over time which would result in
different associations between adherence to CFG-2019 and nutrients intakes. Another key strength
of this work is the use of the NCI multivariate method to account for random errors in dietary
intakes measured with 24-hour recall (29, 30). Using the multivariate method permitted the joint
assessment of the relationship between usual nutrient intakes and adherence to CFG-2019
recommendations on healthy food choices, as measured with the HEFI-2019, and the modelling of
higher adherence to CFG-2019 recommendations compared with average adherence. Limitations
must be addressed. First, self-reported dietary intakes with 24-hour dietary recall are affected by
systematic errors (38, 39). The assumption that intakes are unbiased cannot be verified, although
evidence indicates that diet quality based on 24-hour dietary recall may not be prone to large bias
(40). Second, the food composition database, the Canadian Nutrient File 2015, may not adequately
reflect the nutrient profile of foods actually consumed (41). Third, our results cannot be used to
extrapolate nutrient intake inadequacy at the national level. Nutrient intakes from dietary
supplements were not included since the objective was to assess the relationship between
nutrients and adherence to CFG-2019 recommendations on healthy food choices (4, 5, 16).

In conclusion, higher adherence to CFG-2019 recommendations on healthy food choices was
associated with greater intakes of many nutrients considered, based on eating patterns of adults
aged 65 years or more from the CCHS 2015 – Nutrition. However, higher adherence was associated
with lower intakes of iron, folate, vitamin B12 and vitamin D, and not associated with calcium and
zinc intakes, thus insufficient to mitigate dietary inadequacy in these key nutrients. Knowledge of
shortfalls of CFG-2019 recommendations regarding nutrient intake adequacy are relevant to help
government, policymakers and registered dietitians in providing more comprehensive and
appropriate recommendations to older adults. Future studies should investigate how eating
patterns changed after the publication of CFG-2019 and how these changes may affect the
relationship between adherence to recommendations and nutrient intakes.

**Acknowledgments**

**Data availability statement**: The Canadian Community Health Survey 2015 – Nutrition Public Use
Microdata File are available upon request at
https://www150.statcan.gc.ca/n1/en/catalogue/82M0024X2018001 . The analytic code will be
made freely available at https://github.com/didierbrassard.

**Authors' contributions to the manuscript**: DB and SC designed the research; DB performed
statistical analysis; DB wrote the first draft of the manuscript; DB and SC gave final approval and
critically reviewed the manuscript.

# References

1. Health Canada. Eating Well with Canada's Food Guide. Ottawa: Health Canada, 2007:1-6.

2. Katamay SW, Esslinger KA, Vigneault M, et al. Eating well with Canada's Food Guide (2007): development of the food intake pattern. Nutr Rev 2007;65(4):155-66.

3. Health Canada. Food, Nutrients and Health: Interim Evidence Update 2018 for Health Professionals and Policy Makers. Ottawa: Health Canada, 2019.

4. Health Canada. Canada's food guide. Ottawa: Health Canada, 2019.

5. Health Canada. Canada's Dietary Guidelines - for Health Professionals and Policy Makers. 2019.

6. Brassard D, Manikpurage HD, Thériault S, et al. Greater adherence to the 2019 Canada's Food Guide recommendations on healthy food choices reduces the risk of cardiovascular disease in adults: a prospective analysis of UK Biobank data. Am J Clin Nutr 2022;116(6):1748-58. doi: 10.1093/ajcn/nqac256.

7. Dai Z, Kroeger CM, Lawrence M, et al. Comparison of methodological quality between the 2007 and 2019 Canadian dietary guidelines. Public Health Nutr 2020;23(16):2879-85. doi: 10.1017/S1368980020000956.

8. Asher KE, Doucet S, Luke A. Registered dietitians' perceptions and use of the plant-based recommendations in the 2019 Canada's Food Guide. J Hum Nutr Diet 2021;34(4):715-23. doi: 10.1111/jhn.12845.

9. Barco Leme AC, Laila A, Hou S, et al. Perceptions of the 2019 Canada's Food Guide: a qualitative study with parents from Southwestern Ontario. Appl Physiol Nutr Metab 2021. doi: 10.1139/apnm-2021-0414.

10. Barr SI. Is the 2019 Canada's Food Guide Snapshot nutritionally adequate? Appl Physiol Nutr Metab 2019;44(12):1387-90. doi: 10.1139/apnm-2019-0432.

11. Fernandez MA, Bertolo RF, Duncan AM, et al. Translating "protein foods" from the new Canada's Food Guide to consumers: knowledge gaps and recommendations. Appl Physiol Nutr Metab 2020;45(12):1311-23. doi: 10.1139/apnm-2020-0192.

12. Auclair O, Burgos SA. Protein consumption in Canadian habitual diets: usual intake, inadequacy, and the contribution of animal- and plant-based foods to nutrient intakes. Appl Physiol Nutr Metab 2021;46(5):501-10. doi: 10.1139/apnm-2020-0760.

13. Shlisky J, Bloom DE, Beaudreault AR, et al. Nutritional Considerations for Healthy Aging and Reduction in Age-Related Chronic Disease. Adv Nutr 2017;8(1):17-26. doi: 10.3945/an.116.013474.

14. Remond D, Shahar DR, Gille D, et al. Understanding the gastrointestinal tract of the elderly to develop dietary solutions that prevent malnutrition. Oncotarget 2015;6(16):13858-98. doi: 10.18632/oncotarget.4030.

15. Millen BE, Abrams S, Adams-Campbell L, et al. The 2015 dietary guidelines advisory committee scientific report: development and major conclusions. Adv Nutr 2016;7(3):438-44. doi: 10.3945/an.116.012120.

16. Brassard D, Elvidge Munene LA, St-Pierre S, et al. Development of the Healthy Eating Food Index (HEFI)-2019 measuring adherence to Canada's Food Guide 2019 recommendations on healthy food choices. Appl Physiol Nutr Metab 2022;47(5):595-610. doi: 10.1139/apnm-2021-0415.

17. Brassard D, Elvidge Munene LA, St-Pierre S, et al. Evaluation of the Healthy Eating Food Index (HEFI)-2019 measuring adherence to Canada's Food Guide 2019 recommendations on healthy food choices. Appl Physiol Nutr Metab 2022;47(5):582-94. doi: 10.1139/apnm-2021-0416.

18. Health Canada. Reference Guide to Understanding and Using the Data: 2015 Canadian Community Health Survey - Nutrition. 2017:1-91.

19. Health Canada. Canadian Nutrient File (CNF) 2015. Internet: https://food-nutrition.canada.ca/cnf-fce/index-eng.jsp.

20.	Rana H, Mallet M-C, Gonzalez A, et al. Free Sugars Consumption in Canada. Nutrients 2021;13(5):1471. doi: 10.3390/nu13051471.

21.	Health Canada. Nutrition Labelling: Table of Reference Amounts for Food. Ottawa, 2016.

22.	Dorrington N, Fallaize R, Hobbs DA, et al. A Review of Nutritional Requirements of Adults Aged >/=65 Years in the UK. J Nutr 2020;150(9):2245-56. doi: 10.1093/jn/nxaa153.

23.	Otten JJ, Hellwig JP, Meyers LD. Dietary reference intakes : the essential guide to nutrient requirements. Washington, D.C.: National Academies Press, 2006.

24.	Phillips SM, Chevalier S, Leidy HJ. Protein "requirements" beyond the RDA: implications for optimizing health. Appl Physiol Nutr Metab 2016;41(5):565-72. doi: 10.1139/apnm-2015-0550.

25.	Bauer J, Biolo G, Cederholm T, et al. Evidence-based recommendations for optimal dietary protein intake in older people: a position paper from the PROT-AGE Study Group. J Am Med Dir Assoc 2013;14(8):542-59. doi: 10.1016/j.jamda.2013.05.021.

26.	Traylor DA, Gorissen SHM, Phillips SM. Perspective: Protein Requirements and Optimal Intakes in Aging: Are We Ready to Recommend More Than the Recommended Daily Allowance? Adv Nutr 2018;9(3):171-82. doi: 10.1093/advances/nmy003.

27.	Dodd KW, Guenther PM, Freedman LS, et al. Statistical methods for estimating usual intake of nutrients and foods: a review of the theory. J Am Diet Assoc 2006;106(10):1640-50. doi: 10.1016/j.jada.2006.07.011.

28.	Thompson FE, Kirkpatrick SI, Subar AF, et al. The National Cancer Institute's Dietary Assessment Primer: A Resource for Diet Research. J Acad Nutr Diet 2015;115(12):1986-95. doi: 10.1016/j.jand.2015.08.016.

29.	Zhang S, Midthune D, Guenther PM, et al. A New Multivariate Measurement Error Model with Zero-Inflated Dietary Data, and Its Application to Dietary Assessment. Ann Appl Stat 2011;5(2B):1456-87. doi: 10.1214/10-AOAS446.

30.	Kirkpatrick SI, Guenther PM, Subar AF, et al. Using short-term dietary intake data to address research questions related to usual dietary intake among populations and subpopulations: Assumptions, statistical techniques, and considerations. J Acad Nutr Diet 2022;122(7):1246-62. doi: 10.1016/j.jand.2022.03.010.

31.	Herrick KA, Rossen LM, Parsons R, et al. Estimating Usual Dietary In take From National Health and Nut rition Examination Survey Data Using the National Cancer Institute Method. Vital Health Stat 2 2018(178):1-63.

32.	Harrell FE. General Aspects of Fitting Regression Models. Edtion ed. Regression Modeling Strategies: With Applications to Linear Models, Logistic and Ordinal Regression, and Survival Analysis. Cham: Springer International Publishing, 2015:13-44.

33.	Vatanparast H, Islam N, Shafiee M. Consumption of Milk and alternatives decreased among Canadians from 2004 to 2015: evidence from the Canadian community health surveys. BMC Nutr 2021;7(1):63. doi: 10.1186/s40795-021-00465-9.

34.	Vatanparast H, Islam N, Patil RP, et al. Calcium Intake from Food and Supplemental Sources Decreased in the Canadian Population from 2004 to 2015. J Nutr 2020;150(4):833-41. doi: 10.1093/jn/nxz318.

35.	Vatanparast H, Patil RP, Islam N, et al. Vitamin D Intake from Supplemental Sources but Not from Food Sources Has Increased in the Canadian Population Over Time. J Nutr 2020;150(3):526-35. doi: 10.1093/jn/nxz291.

36.	Drolet-Labelle V, Laurin D, Bedard A, et al. Beliefs underlying older adults' intention to consume plant-based protein foods: A qualitative study. Appetite 2022:106346. doi: 10.1016/j.appet.2022.106346.

37.	Huang HH, Cohen AA, Gaudreau P, et al. Vitamin B-12 Intake from Dairy But Not Meat is Associated with Decreased Risk of Low Vitamin B-12 Status and Deficiency in Older Adults from Quebec, Canada. J Nutr 2022;152(11):2483-92. doi: 10.1093/jn/nxac143.

38.     Freedman LS, Commins JM, Moler JE, et al. Pooled results from 5 validation studies of dietary self-report instruments using recovery biomarkers for potassium and sodium intake. Am J Epidemiol 2015;181(7):473-87. doi: 10.1093/aje/kwu325.

39.     Freedman LS, Commins JM, Moler JE, et al. Pooled results from 5 validation studies of dietary self-report instruments using recovery biomarkers for energy and protein intake. Am J Epidemiol 2014;180(2):172-88. doi: 10.1093/aje/kwu116.

40.     Kirkpatrick SI, Dodd KW, Potischman N, et al. Healthy Eating Index-2015 Scores Among Adults Based on Observed vs Recalled Dietary Intake. J Acad Nutr Diet 2021;121(11):2233-41 e1. doi: 10.1016/j.jand.2021.06.009.

41.     Satija A, Yu E, Willett WC, et al. Understanding nutritional epidemiology and its role in policy. Adv Nutr 2015;6(1):5-18. doi: 10.3945/an.114.007492.

## Tables

**Table 1**. Usual nutrient intake difference between HEFI-2019 scores at the 90[th] percentile and the median HEFI-2019 score distribution in 4,089 adults aged 65 years or more from the CCHS 2015 - Nutrition.

| Nutrients[1] | Dietary Reference Intake group | | | | |
|---|---|---|---|---|---|
| | Adults, 65 y+ | Males, 65 to 70 y | Females, 65 to 70 y | Males, 71 y+ | Females, 71 y+ |
| Protein, g/d | 3 (-2, 8) | 2 (-4, 9) | 5 (-1, 11) | 2 (-4, 8) | 4 (-2, 10) |
| Calcium, mg/d | 4 (-50, 57) | -16 (-96, 65) | 21 (-47, 89) | -14 (-88, 60) | 25 (-38, 89) |
| Vitamin D, mcg/d | -0.5 (-1.0, 0.0) | -0.8 (-1.6, -0.1) | -0.1 (-0.7, 0.5) | -0.7 (-1.5, 0.0) | -0.1 (-0.7, 0.6) |
| Iron, mg/d | -0.2 (-1.0, 0.5) | -0.6 (-1.8, 0.6) | 0.2 (-0.7, 1.1) | -0.6 (-1.8, 0.6) | 0.3 (-0.5, 1.1) |
| Vitamin B12, mcg/d | -0.3 (-0.7, 0.1) | -0.4 (-0.9, 0.0) | -0.1 (-0.7, 0.4) | -0.4 (-0.8, 0.1) | -0.1 (-0.6, 0.4) |
| Vitamin B6, mg/d | 0.1 (0.0, 0.2) | 0.1 (0.0, 0.2) | 0.1 (0.0, 0.3) | 0.1 (0.0, 0.2) | 0.1 (0.0, 0.2) |
| Zinc, mg/d | 0.0 (-0.7, 0.7) | -0.4 (-1.5, 0.7) | 0.4 (-0.3, 1.2) | -0.3 (-1.3, 0.6) | 0.5 (-0.2, 1.2) |
| Folate, mcg/d | -13 (-39, 14) | -15 (-56, 26) | -6 (-36, 25) | -16 (-55, 22) | -2 (-31, 27) |
| Fibers, g/d | 3 (2, 5) | 4 (2, 7) | 3 (1, 5) | 4 (2, 6) | 3 (1, 5) |
| Magnesium, mg/d | 33 (14, 52) | 44 (14, 73) | 30 (6, 55) | 39 (13, 65) | 31 (9, 53) |
| Potassium, mg/d | 173 (17, 328) | 230 (6, 455) | 170 (-27, 368) | 214 (3, 424) | 178 (-7, 363) |
| Vitamin A, RAE/d | 16 (-37, 69) | 9 (-54, 71) | 26 (-54, 106) | 10 (-48, 68) | 26 (-42, 95) |

[1]Values are expected nutrient intake differences (95%CI) for a HEFI-2019 score at the 90th compared with the 50th percentile of the score distribution, and were estimated using linear regression models. Estimates reflect the expected nutrient intake difference when HEFI-2019 scores are increased to correspond to the 90th percentile of the score distribution, compared with the median HEFI-2019 score. All dietary intakes were modelled using the National Cancer Institute multivariate method (see Methods). 95%CI were estimated using 500 bootstrap weight replicates. CCHS, Canadian Community Health Survey; d, day; HEFI-2019, Healthy Eating Food Index-2019.

**Figure legends**

*Figure 1*: Distribution of total HEFI-2019 score, based on usual dietary intakes in 4,089 adults aged 65 years and more from the CCHS 2015 - Nutrition. Diamonds are means. Left and right whiskers indicate the 5th and 95th percentile, respectively. Dietary intakes were modelled using the National Cancer Institute multivariate method (see Methods). CCHS, Canadian Community Health Survey; DRI, Dietary Reference Intake.

*Figure 2*: Prevalence of inadequate nutrient intakes in 4,089 adults aged 65 years and older from the CCHS 2015 - Nutrition. Only nutrient intakes from foods were considered (i.e., excluding intakes from dietary supplement). Inadequate intakes are intakes below the age- and sex-specific Estimated Average Requirements (EAR). Potassium and fibers are not shown, because only Adequate Intakes (AI) values are available for these nutrients. All dietary intakes were modelled using the National Cancer Institute multivariate method (see Methods). 95%CI were estimated using 500 bootstrap weight replicates. CCHS, Canadian Community Health Survey.

*Figure 3*: Linear regression of nutrient intake on the total HEFI-2019 score in 4,089 adults aged 65 years or more from the CCHS 2015 - Nutrition. A positive relationship indicates that greater HEFI-2019 scores are associated with greater nutrient intake, and inversely. For visualization purpose, data points are from a random sample of 200 respondents selected proportionally to sampling weights. All dietary intakes were modelled using the National Cancer Institute multivariate method (see Methods). 95%CI were estimated using 500 bootstrap weight replicates. CCHS, Canadian Community Health Survey; HEFI-2019, Healthy Eating Food Index-2019.

*Figure 4*: Prevalence of inadequate intakes and difference for HEFI-2019 scores at the 90th compared with the 50th percentile of the score distribution in 4,089 adults aged 65 years or more from the CCHS 2015 - Nutrition. Inadequate intakes are intakes below the age- and sex-specific Estimated Average Requirements (EAR). Potassium and fibers are not shown, because only Adequate Intakes (AI) values are available for these nutrients. All dietary intakes were modelled using the National Cancer Institute multivariate method (see Methods). 95%CI were estimated using 500 bootstrap weight replicates. CCHS, Canadian Community Health Survey; HEFI-2019, Healthy Eating Food Index-2019.

Age and sex group

HEFI-2019 score based on usual intakes (/80), points