Predicting hearing help-seeking: Which features are important for a psychological

profiling module of a hearing mHealth application?

Giulia Angonese[1,2], Mareike Buhl[1,3,5], Inka Kuhlmann[1,2], Birger Kollmeier[1,3,4] and Andrea

Hildebrandt[1,2,4]

[1] Cluster of Excellence 'Hearing4all', Oldenburg, Germany;

[2] Department of Psychology, Psychological Methods and Statistics, Carl von Ossietzky

Universität Oldenburg, Germany;

[3] Department of Medical Physics and Acoustics, Carl von Ossietzky Universität Oldenburg,

Germany;

[4] Research Center Neurosensory Science, Carl von Ossietzky Universität Oldenburg,

Germany

[5] Institut Pasteur, Université Paris Cité, Inserm, Institut de l'Audition, F-75012 Paris, France

Corresponding authors:

Giulia Angonese and Andrea Hildebrandt, Department of Psychology, Carl von Ossietzky

Universität Oldenburg, Ammerländer Heerstraße 114-118, 26129 Oldenburg, Germany

Email: giulia.angonese@uni-oldenburg.de; andrea.hildebrandt@uni-oldenburg.de

19   **Abstract**

Introduction: Mobile health care solutions such as a Virtual Hearing Clinic (VHC) can

improve quality, accessibility and equity of health services, fostering early rehabilitation for

hearing loss. A VHC might be the first contact with a hearing diagnostic service and should

motivate users with hearing loss to seek professional help. A psychological profiling module

would secure more efficient treatment recommendations to promote help-seeking, after the

diagnostic use of the VHC.

Methods: N=185 (106 females) non-aided older individuals (M_{age}=63.8, SD_{age}=6.6)

with subjective hearing loss participated in a large ecological momentary assessment study.

We collected cross-sectional and longitudinal data on several psychological- and hearing-

related features that were previously found to predict help-seeking. Readiness to seek help was

assessed as outcome variable at study-end and after two months. Participants were classified

into help-seekers and non-seekers with supervised machine learning algorithms (Random

Forest, Naïve Bayes and Support Vector Machine). The most relevant features for prediction

were identified with feature importance analysis.

Results: The algorithms correctly predicted action to seek help at study-end in 66 to

70% of cases, reaching 75% classification accuracy at follow-up. Among the most important

features for classifications were the degree of hearing loss and its perceived consequences in

daily life, attitude towards hearing aids, personality traits like neuroticism and

conscientiousness.

Conclusion: This study contributes to the identification of individual characteristics that

predict help-seeking in older individuals with self-perceived hearing loss. Suggestions for the

implementation of a psychological profiling algorithm and for targeted recommendations in a

VHC are derived.

Keywords: Hearing loss, mobile health, help-seeking, ecological momentary

assessment, supervised classification, feature importance.

Introduction

Hearing enables people to experience their surroundings and to communicate with

others. Thus, hearing difficulties can have a strong impact on individuals' quality of life.

Hearing loss (HL) is one of the most common chronic diseases worldwide and it affects 20.3%

of the world's population. More than 60% of individuals with HL are older than 50 years of

age, with the principal cause being age-related HL (1). Untreated hearing difficulties have been

associated with lower self-rated health, depression, and anxiety, in addition to physical and

cognitive decline, dementia, and hospitalization in the older population (2–4). The primary

rehabilitative strategy for individuals with moderate to severe HL is the use of hearing aids

(HA), which increases activity levels, general health, quality of life (5) and decreases social

isolation and depressive symptoms (6) by supporting hearing ability and communication

efficacy. Despite the positive effects of hearing rehabilitation, the prevalence of HA use is still

limited to about 25% of the hearing impaired population (2, 4, 7, 8). Moreover, there is an

average delay of nine years between the time a person first acknowledges hearing difficulties

until the first contact with a hearing-health professional (9).

Developing easily accessible and affordable mobile health solutions in audiology could

promote broader and faster access to diagnosis and health services, fostering an early

rehabilitation and reducing the impact of HL on the individual. Before such mobile health

applications are implemented, there is need for basic research on potential components of a

Virtual Hearing Clinic (VHC) that could provide audiological self-test options and HA

algorithms for self-fitting. Through a VHC, users could receive specific treatment

recommendations depending on their hearing performance, including suggestions to visit a

hearing acoustician or ENT (Ear, Nose and Throat) physician, if indicated. In addition, users

could be given the possibility to experience listening conditions of a HA and its potential

benefits. However, even though professional support might be recommended, a fair amount of

users might still be reluctant to seek help. This would call for a careful characterization of "non-

help-seekers", namely individuals who might need more convincing incentives via a VHC

setting in order to take action. A VHC might often represent the first contact with a hearing

diagnostic service and should therefore motivate individuals in need (especially non-help-

seekers) to pursue elaborate professional help, in order to optimize the impact on the population.

It follows that the assessment of an individual with hearing difficulties should go beyond

simply quantifying HL and should consider individual characteristics that have been shown to

influence the readiness of individuals to seek professional help (4). Moreover, such information

collected by a VHC could later help clinicians in performing an adequate and personalized

counseling.

Acknowledgement and acceptance of hearing difficulties and their impact on everyday

life have been discussed as the most important predictive features of HA uptake (5, 6, 10, 11).

HL is frequently perceived as part of the natural ageing process and other health issues are

prioritized for treatment (4, 12). Even when HL is identified, individuals might reject the use

of a HA due to expected costs, stigma and negative stereotypes (5, 7, 10). However, a positive

attitude towards HA (13), high expectations to benefit from them (11) and perceived self-

efficacy in their daily management (14, 15) were shown to promote help-seeking, as well as

later HA uptake. Further relevant predictors are personal attitudes, beliefs, and personality traits.

Individuals who are more prone to seek help and successfully uptake a HA show higher internal

locus of control (2, 14) self-efficacy (14, 15), and agreeableness, as well as lower neuroticism

and openness (2). Altogether such individual characteristics refer to a general self-confidence

in the ability to cope with critical situations, good acceptance of others' suggestions and

recommendations, as well as less susceptibility to shame and embarrassment and less analytical

thinking.

95   The present study aims at identifying the most important predictive features for help-

96 seeking aiming to design a psychological profiling module in a VHC. Such module will

97 categorize help-seekers versus non-seekers and ultimately inform the design of personalized

98 treatment recommendations. For this purpose, data from a large number of questionnaires and

99 tests related to different psychological- and hearing-related individual characteristics, together

100 with multiple assessments of a hearing screening test were collected in a mobile study that

101 simulated a VHC. To target potential users of a future VHC, the study was addressed to

102 individuals with subjectively perceived hearing difficulties who had not yet been compensated

103 with HA. We used machine learning algorithms to predict readiness to seek professional help,

104 as assessed at the end of the study and after two months.

105   The following research questions guided our study design and analyses:

106   RQ1.  Which machine learning model can best predict help-seeking and

107     categorize individuals into help-seekers versus non-seekers?

108   RQ2.  Which psychological- and hearing-related features are most relevant to

109     profile individuals and tell help-seekers versus non-seekers apart?

110   RQ3.  How can feature importance measures inform the design of targeted

111     recommendations for users of a future VHC?

113          **Methods**

114 **Study overview**

115   We used the potential of Ecological Momentary Assessment (EMA, (16)), a

116 measurement approach to collect data in the everyday environment of the individual

117 ("Ecological") and to assess recent or immediate experiences and states ("Momentary"). The

118 study was conducted entirely online on personal mobile phones of the participants to

119 approximate the experience of dealing with a VHC mobile app. Communications with the

participants took place via email and SMS. The total assessment time of eight hours was

distributed across the working days of three consecutive weeks, with an overall daily testing

time of approximately 30 minutes. The first week (baseline assessment) included one

measurement per day, which could be performed at any preferred time. The second and third

week (EMA) included two measurement time-points per day (morning and evening) of

approximately 15 minutes each. Participants were remunerated with 10 Euros per hour. After

two months from the end of the study, participants were invited to complete a voluntary follow-

up online questionnaire. The study plan and data management have been approved by the

Research Ethics Committee of the Carl von Ossietzky Universität of Oldenburg. We, as single

researchers and as a team, are committed to and supporting Open Science practices. We have

therefore published a preprint of the manuscript on medrxiv.org in February 2023 and share

data analysis scripts via Zenodo (https://doi.org/10.5281/zenodo.7635920). The data will be

shared with interested researchers upon request, since the dataset will be used for further

projects and cannot be shared at the current time point.

Participants

Older adults above 50 years of age were recruited between September 2021 and August 2022

through the online platform Ebay's minijob announcements, the university intranet and via

mailing list services of several German universities' Guest-Audience and Senior-University

programs. Inclusions criteria were: subjective reports of hearing difficulties in daily life,

ownership of and the ability to use a smartphone, and good command of the German language.

The exclusion criterion was the use of hearing aids. Prior to enrolment, participants provided

written informed consent. The final dataset included $N_1 = 185$ participants, 106 female and 79

male (0 diverse), aged between 47 and 82 years, with $M_{age} = 63.1$ and $SD_{age} = 6.5$ (see Figure

1). One participant was below 50 years of age (47) but was nevertheless included in the final

sample, given that this value only slightly deviated from the planned age threshold. A

descriptive summary of participants' socio-demographic characteristics is provided in Table 1.

Figure 1. Participants' inclusion flow chart. A total number of 438 individuals demonstrated interest in the study and 192 participants were finally enrolled. Among the 186 individuals who successfully completed the study, one participant was excluded from the final data analysis due to the absence of hearing-test data. Out of the N_1=185 participants who completed the study, N_2=131 additionally participated to the follow-up questionnaire.

149 **Table 1.** Main socio-demographic characteristics of the participants.

Characteristic		n (% of total)
Age group (years)	47-60	70 (37.8)
	61-70	83 (44.9)
	70-82	32 (17.3)
Sex	female	106 (57.3)
Duration of hearing difficulties (years)	0 - 1	54 (29.2)
	2 - 5	89 (48.1)
	6 - 10	23 (12.4)
	10 – 21	19 (10.3)
Presence of Tinnitus		65 (35.1)
Previous doctor consultation for hearing difficulties		89 (48.1)
Presence of visual problems		134 (72.4)
Occupation status	employed	67 (36.2)
Monthly income (Euros)	< 1.500	62 (33.5)
	1.500 - 2.500	65 (35.1)
	2.500 – 4.000	42 (22.7)
	>4.000	16 (8.7)
Residential environment	countryside	2 (1.1)

	little town	29 (15.7)
	suburbs	48 (25.9)
	city	106 (57.3)
Self-estimated noise level at home	low	67 (36.2)
	moderate	113 (61.1)
	high	5 (2.7)

Note: These data were acquired during the baseline assessment through a self-report questionnaire.

Procedure

EMA tools. The data collection was carried out with the online, open-source software

formr (17). Here, different questionnaires and tests (see section Assessment) were combined

as surveys in a run that reproduced the desired design and could be assessed by users through

a specific link. A Customer Communication Platform (www.twilio.com) helped to send the

study link to the participants through daily text message reminders. The use of REST API

(Representational State Transfer Application Programming Interface) allows for data exchange

between platforms in a consistent format and in the respect of privacy and security. With the

individualized link received via SMS, participants could perform the study on their own

smartphone's browser. Seven participants performed the study on their computers since they

experienced technical difficulties with their smartphones.

During baseline assessment, cross-sectional data from a comprehensive set of 25

questionnaires and tests was collected. The questionnaires and tests were distributed on five

consecutive days in order to maximise study compliance and avoid priming effects on different

questionnaires. For a detailed list of all assessment tools and their references we refer to the

Supplementary Material 1 (Features and assessment tools). The following is a brief overview

of the assessed features.

Assessment of hearing-related features. Firstly, the assessment included self-reported

hearing difficulties and perceived consequences of HL, such as effects on social life

participation. Additionally, attitudes towards HA were evaluated with questionnaires on HA

expectations and stigma. Trait sensitivity to perceived noise was measured as a personality facet which was shown to be related to affect and neurosensory processing (18). We further assessed individuals' sound preferences profiles (19), aiming to gather more information about sound sensitivity and hearing habits. Finally, hearing health literacy was evaluated, since the ability to search, find and understand information related to hearing health was shown to influence self-management of HL (14).

Assessment of psychological features. As outlined in the introduction, personality traits (the Big Five (20)), locus of control and self-efficacy were shown to be associated with help-seeking and HA use, hence these features were included in the baseline assessment. Anxiety and depression were also measured, given their frequently demonstrated associations with HL (4,6), together with loneliness, which is seen as consequence of untreated HL (3). We further assessed optimism and sensory processing sensitivity, which refers to an individual's disposition to perceive and process stimuli (including auditory ones) more intensely than the average population (21). Perceived stress was also measured, since high levels of stress that are related to daily life, work or social situations may boost help-seeking behaviours (3). General health was also assessed given its predictive role for different steps of the HA uptake path (4, 6). The belief that HA are associated with old age and infirmity is often a barrier to HA uptake and use (4), therefore attitude towards ageing was assessed as well. For completeness, we also measured cognitive abilities (crystallized and fluid intelligence), despite discordant findings on associations between cognition and HA uptake (4, 6, 13). Lastly, participants were requested to complete a comprehensive questionnaire on socio-demographics (see Table 1).

Daily assessment of affect and hearing. Participants performed the daily measures in the morning, possibly right after waking up, and in the evening before going to bed. At each time, affect and hearing performance were assessed. This micro-longitudinal assessment

accounts for potential daily fluctuations of hearing performance and affect, which might

depend on particular daily events and states. The affect questionnaire included 14 items in line

with the Circumplex Model of Affect (Russel, 1980 (22)). Eight items were related to negative

affect and six to positive affect (23). The items were displayed in a randomized order at each

presentation and respondents had to indicate on a seven-point scale how much the specified

mood applied to them (1 = does not apply; 7 = applies fully). The items are listed in the

Supplementary Material 2 (Daily assessments). The affect questionnaire was presented before

and after the hearing test in order to assess mood at baseline and after receiving feedback on

the hearing test, respectively.

Hearing performance was assessed with the Digit Triplets Test (DTT) (23, 24) of the

Hörzentrum Oldenburg gGmbH (www.hz-ol.de). This screening instrument was designed to

measure speech intelligibility in noise by means of the Speech Recognition Threshold (SRT),

which indicates the Signal-to-Noise Ratio (SNR, difference between speech and noise level) at

which the participant reaches 50% speech intelligibility. Participants were invited to familiarize

with the test before starting the longitudinal assessment. They were instructed to perform the

test with the use of their personal headphones, but three participants reported technical

difficulties with their headphones and used loudspeakers instead. After completing the hearing

test, participants received a feedback on their performance in the form of a traffic-light colour,

where green indicated good (SRT < -7.1 dB SNR), yellow indicated intermediate (-7.1 >= SRT

< -5.1 dB SNR), and red reflected poor performance (SRT >= -5.1 dB SNR) (25,26). The

supplementary material provides detailed information on the hearing test and its feedback

(Supplementary Material 2 – Daily assessments; Supplementary Material 3 – Hearing test

feedback).

Outcome measures

At the end of the study, participants were asked to report on planned actions to seek

professional help for their perceived hearing difficulties, their motivation to seek help and the

source of this motivation. This information was also assessed at the beginning of the study.

These variables, as retrieved at the end of the study, were chosen as outcome measures for the

supervised machine learning (see below). The distribution of participants ($N_1 = 185$) along the

outcome classes considered is summarized in Table 2.

Action to seek help. Intention to seek professional help of was assessed with the

question: "Given the feedbacks of this study regarding your hearing performance, have you

made an appointment with one of the following doctors or a hearing care professional, or are

you planning to do so?" (followed by a list of hearing professionals). This variable was used to

create two outcome classes:

▪ participants ready to take action, who were planning to seek professional help

233   in the near future or had already taken an appointment (*action* class, $n_{11} = 64$);

234   ▪ participants not ready to take action, who did not plan to consult a hearing health

235   professional in the future (*no action* class, $n_{12} = 121$).

Action and motivation to seek help. A second outcome measure was taken into

account to further differentiate the *no action* class, in order to provide further insight for the

design of targeted recommendations in a VHC. Information on readiness to take action was

combined with the reported motivation to seek help at the end of the study. Motivation was

assessed through the question: "How motivated are you at the moment to seek help regarding

your hearing problems?" (1 = not motivated at all, 7 = very strongly motivated). The answer

spectrum was binarized by means of median split to create the following outcome classes:

▪ participants not ready to take action with high motivation (*no action & high*

244   *motivation* class, $n_{13} = 47$), who might particularly benefit from personalised

245   and tailored recommendations;

246   ■ participants not ready to take action who reported low motivation to seek help

247   (*no action & low motivation* class, $n_{14} = 74$);

248   ■ participants ready to take action, regardless of their motivation level (*action*

249   class, $n_{15} = 64$). This class was not further divided with respect to motivation,

250   since this would not result in different recommendations in a VHC. Moreover,

251   data exploration revealed that only seven individuals out of 64 in this category

252   reported low motivation.

Action to seek help at a follow-up. Two months after the end of the study, participants

who agreed to be contacted for a follow-up received a further questionnaire (in formr). The

short survey consisted of two multiple-choice questions and it was completed by 71% ($N_2 =$

131) of the participants. Individuals were asked to report again on their action to seek

professional help. In addition, they were asked to indicate whether the study participation

improved their awareness towards hearing difficulties. Only the first question was considered

as further outcome measure. The answers (given as four multiple-choice answers) were

binarized to achieve a class allocation comparable to the first outcome measure:

■ participants who reported to have completed an appointment with a hearing

262   professional; participants who had an appointment fixed, yet not completed;

263   participants who were planning to seek professional help in the near future

264   (*action at follow-up* class, $n_{21} = 52$);

265   ■ participants who did not plan to consult a hearing health professional (*no action*

266   *at follow-up* class, $n_{22} = 79$).

268   **Table 2**. Absolute class-wise frequencies of observations across the three outcomes considered.

		action at follow-up	no action at follow-up	no follow-up data	
action		33	12	19	64
no action	**high motivation**	12	22	13	47

low motivation	7	45	22	74
	52	79	54	

Note: The three outcomes considered are: action to seek help (N_1=185); action and motivation to seek help (N_1=185); action to seek help at follow-up (N_2=131). Additionally, the table provides an overview of those participants who did not complete the follow-up questionnaire.

Data pre-processing

Data analysis was performed with the R Software for Statistical Computing (27). Raw data from all questionnaires was imported from the online platform formr to R environment using the dedicated package formr (28). For each questionnaire or test presented at baseline, global scores were computed and considered as features. If both global scores and scale scores were available for a given assessment tool, only the scale scores were maintained if considered differentially relevant for the outcome. Hearing test results were received via E-Mail from the Hörzentrum Oldenburg and imported in R as .eml files. The performance feedback category (green, yellow, red) was additionally extracted and stored for each raw SRT result. Due to the particular implementation of the study in formr, participants could perform the hearing-test more than once at each measurement time-point. Whenever this happened, only the last SRT result at a given time-point was kept for analysis. This led to a removal of 3.9% of the raw SRT results. The longitudinal data on daily mood and hearing performance was summarized into the following features: individual mean SRT and its variability across all measurements, percentage of feedback type received -with green feedback (good performance) taken as reference category-, individual mean and variability of positive and negative affect at pre-test and at post-test (after receiving the hearing test feedback). The summarized longitudinal data was merged with the cross-sectional data, resulting in a wide-format data frame including 88 features.

Missing data occurred for the longitudinal measures of hearing performance and affect. A complete set of 20 SRT results were collected for 43.8% of participants, while at least 15

SRT results were obtained in 95.1% of cases. Only three participants provided less than 10

SRT results, with one participant completing only four measurements. Missing hearing data at

a specific time-point was considered Not Available (NA). Where an SRT result was missing,

the respective feedback and measures of affect at post-test were missing as well (NA). By

visual inspection of the individual SRT distributions, some specific outlier patterns were

identified. 14 participants showed a much larger SRT result at the first measurement, which

qualified as outlier following the interquartile range rule. These large SRT values (indicting

poor performance) were considered to be caused by misunderstanding of the hearing test

instructions and were therefore set to NA. The respective feedback category and measures of

affect where however not set to NA. This is because despite the unreliable SRT value,

participant's mood could still have been affected by the feedback received. With respect to

daily affect measures, two participants presented no data at measures of post-test affect, such

that summary measures could not be computed. In these cases, mean imputation technique was

applied: the sample mean and sample variance for negative and positive affect at post-test were

imputed to replace missing values.

Machine learning

The data was fed into three machine learning algorithms for supervised classification

along the outcome labels *action* versus *no-action*. We chose Naïve Bayes (NB), Random

Forests (RF) and Support Vector Machines (SVM) among other classifiers to cover a wide

range of model complexity (from simple models like NB, to more complex and non-linear ones

like SVM). We used Cross-Validation (CV) to train and subsequently test the model. The

algorithms have been implemented on R using the mlr package (29), following the approaches

described in (30) and (31). Given the presence of three different outcome measures, the same

analysis steps were carried out in parallel for each outcome, with a slight difference in the input

features included in the analysis. For the first outcome (*action to seek help*), data on motivation

at pre-study was kept in the features space, while motivation to seek help at the end of the study

was removed. Differently, for the second outcome (*action and motivation to seek help*), all data

on motivation at pre- and post-study was removed from the feature space. Implementation

details of the three algorithms are summarized next.

Naïve Bayes classifier (NB). This algorithm uses Bayes' rule to predict the probability

of an observation to belong to one of the outcome classes given its discriminant function values.

This probabilistic model takes continuous and categorical features into account and assumes

that they are independent (30). In the present implementation, after training the algorithm,

repeated 10-fold CV was used to evaluate the model's performance. A step-wise approach was

used to select the appropriate number of CV repetitions necessary to achieve accurate and

stable performance estimates (50, 100, 150 and 300 CV repetitions).

Random Forest (RF). Tree-based methods use recursive binary splitting to stratify the

features' space in smaller, non-overlapping regions used for classification. Trees are easy to

interpret but they lack of predictive power, since they tend to overfit the training data. RF can

be used to improve prediction accuracy (29, 31). This algorithm requires to tune a set of

hyperparameters which control the learning process and are selected (or tuned) by the algorithm

to obtain best performance. The following hyperparameters were considered:

▪ Ntree: number of trees to include. This value is usually fixed at a

computationally reasonable value rather than tuned (30). Ntree was set to 800.

▪ Mtry: the number of features to randomly sample at each iteration. A popular

value is given by $\sqrt{p}$ (where p = the number of predictors) (33). Different search

spaces were explored, with Mtry ranging between [1, 15].

▪ Nodesize: minimum number of cases to be included in a leaf. Different search

spaces were explored, with Nodesize ranging between [1, 20].

Tuning the algorithm and finding the best hyperparameter combination requires to

define an optimization algorithm, or search strategy, and evaluation method. We used grid-

search with 10-fold CV resampling. To evaluate model performance, nested CV was applied.

In this approach, an inner loop tunes the hyperparameters and an outer loop evaluates a wrapped

learner, which comprises the classification task, the learner type (RF) and the hyperparameter

tuning process. Here a 5-fold CV was applied as outer resample strategy.

Support Vector Machine (SVM). The SVM algorithm iteratively identifies a

hyperplane that separates labelled classes also in case of non-linear data distributions. For

multiclass problems, several models are built and then summarized for prediction. SVMs are

computationally expensive, but tend to perform very well on a variety of tasks conducted on

non-linearly separable classes. Additionally, the algorithm has the advantage to make no

assumptions on the features' distributional properties (29, 31). Similarly to RF, SVM requires

hyperparameter tuning. The following hyperparameters were considered:

▪ Kernel: the type of the kernel function that the algorithm uses to add further

dimensions to the features space and identify the separating hyperplane.

Polynomial, radial and sigmoid functions were included in the search space (30).

▪ Degree: the shape of the decision boundary in case of a polynomial kernel. The

search space was limited to [1, 5] to avoid the risk of overfitting (30).

▪ Cost: a penalty for having cases fall inside the class-separating margin. It is

recommended to tune both cost and gamma (see next point) on the logarithmic

scale (34), and a popular search space for cost is $[2^{-5}, 2^{15}]$ (35).

▪ Gamma: the influence of each case on the hyperplane. This hyperparameter

search space was set to the popular range $[2^{-15}, 2^{3}]$ (35).

Nested CV was used as previously described for RF. An inner resampling loop (with

10-fold CV) was applied for hyperparameter tuning and an outer loop (with 5-fold CV) for

performance evaluation.

Classification performance measures. The algorithms were evaluated in terms of

prediction accuracy on the test set which indicated the overall proportion of cases correctly

classified by the model as compared to the observed outcome. However, class imbalance in the

sample can negatively impact prediction accuracy, reducing its informativeness as performance

measure (35, 36). Matthews Correlation Coefficient (MCC) (yardstick package (38)) was

additionally taken into account to evaluate model performance. This coefficient improves over

accuracy measures in case of imbalanced datasets (36, 38) and can take values from -1 to 1,

with 1 indicating perfect prediction, 0 chance prediction and -1 inverse prediction. In addition

we computed the confusion matrix (`calculateConfusionMatrix()`, mlr package). Its

output provides the absolute number and the proportion of correct model predictions and

misclassifications for each outcome class. For binary outcomes, the confusion matrix allows to

estimate model sensitivity (i.e. accurately identifying seekers) and specificity (i.e. accurately

identifying non-seekers). In the present study, obtaining high specificity is of particular interest

in the context of a VHC. Indeed, individuals with HL who are not prone to seek help are the

main target population for tailored treatment recommendations and counselling.

Feature importance

After identifying the best performing machine learning algorithm, feature importance

was considered. Each feature receives a coefficient of importance that indicates its contribution

to model performance, regardless of the type of relationship (direction and linearity) between

the feature and the outcome. In RF, feature importance is model-dependent and it indicates

how much the feature contributes in reducing node impurity. Importance values were retrieved

by the function `getFeatureImportance()` (mlr package) applied on the RF model

trained with the tuned hyperparameters. These importance results have the advantage of being

inherent to the model and closely tied to its performance (40). Conversely, there are no model-

specific importance metrics available for NB and SVM. For these algorithms, the importance

value assigned to each feature corresponds to the area under the ROC (Receiver Operating

Characteristic) curve, which is computed from sensitivity and specificity measures (40). The

function `varImp()` from the caret package (41) was applied on the model trained with the

function `train()` (caret package), after ensuring comparable performance with the same

model previously trained in the mlr package.

Information on feature importance was used to identify which features are mostly

relevant for telling apart individuals depending on help-seeking intentions. No statistical

criterion exists to determine which importance value threshold should be used to retrieve

relevant features. Hence, three threshold values were inspected (the first 10, 15 and 20 features

in their importance ranking order) and evaluated in terms of predictive accuracy and

interpretability. Classification accuracy of these three feature sets were assessed on the

outcome data obtained at follow-up. For this analysis, the dataset was reduced to $N_2 = 131$

participants who completed the follow-up questionnaire. The important features were fed into

the best performing machine learning algorithm from the previous step.

409   **Results**

Classification performance

Predicting the action to seek help. A summary of the model-specific classification

accuracy for the first outcome (*action to seek help*) is provided in Table 3. The three algorithms

show similar overall performance accuracy estimates on the test set, correctly classifying about

67% to 70% of the cases in the full dataset ($N_1 = 185$). RF was the best performing algorithm

with an accuracy of 70.3% and an MCC = .282, indicating that the model's prediction improves

to about 30% over chance. By inspecting the confusion matrix, we observed that RF shows

high specificity, correctly classifying 91% of the cases belonging to the *no-action* class. The

NB classifier (10-fold CV repeated 50 times) showed the best sensitivity compared with the

competing algorithms, with 51% cases in the *action* class being correctly classified. RF and

NB were selected for feature importance analyses, given the good predictive performance and

high specificity of the RF, as well as highest sensitivity of the NB.

Predicting the action and motivation to seek help. The second part of Table 3

summarizes the classification performance with respect to the second outcome (*action and*

motivation to seek help), which contains three classes (see above). RF provided the highest

accuracy with 54.6% and MCC=.295, as compared to NB (10-fold CV repeated 100 times) and

SVM. However, the confusion matrix revealed that none of the three models was able to

adequately tell apart individuals within the *no action* class who differ with respect to high

versus low motivation. All algorithms can only correctly classify 9 to 25% of cases in the *no*

action & high motivation class. Potentially, an improvement in classification accuracy could

be achieved with a larger dataset in which the classes are better balanced, and with a more

reliable and elaborated measure of the participant's motivation to seek help. In view of these

limitations, the second outcome will not be considered for feature importance analysis.

Table 3. Model-specific overall performance (aggregated accuracy on the test set and MCC) and class-specific classification
accuracy (within class proportion of correct classifications) for the two outcomes measured at study-end.

			Action to seek help			
Model	Hyperparameters		Overall performance measures		Class-specific classification accuracy	
	space	tuned	Test accuracy	MCC	action (n_{11}=64)	no action (n_{12}=121)
NB			.66	.26	.51	.75
RF	ntree 800	800	.70	.23	.31	.91
	mtry [5,15]	13				
	nodesize [1,5]	1				
SVM	kernel	radial	.67	.23	.39	.82
	degree [1,5]	4				
	cost [2^{-5}, 2^{15}]	.01				
	gamma [2^{-15}, 2^3]	69.1				

	Action and motivation to seek help							
Model		Hyperparameters		Overall performance measures		Class-specific classification accuracy		
		space	tuned	Test accuracy	MCC	action $(n_{15}=64)$	no action & low motivation $(n_{14}=74)$	no action & high motivation $(n_{13}=47)$
NB				.495	.218	.47	.67	.25
RF	ntree	800	800	.546	.295	.55	.80	.09
	mtry	[8,10]	9					
	nodesize	[3,15]	2					
SVM	kernel		sigmoid	.492	.204	.53	.76	.02
	degree	[1,5]	1					
	cost	$[2^{-5}, 2^{15}]$	323					
	gamma	$[2^{-15}, 2^{3}]$	3.05×10^{-5}					

Note: results are based on the full data set (N_1=185). For RF and SVM the table additionally shows the hyperparameter search space used in the resampling procedure and the tuned values used for model training. The NB models selected were 10-fold CV repeated 50 and 100 times for the first and second outcome, respectively. (NB: Naïve Bayes classifier; RF: Random Forest; SVM: Support Vector Machine; MCC: Matthews Correlation Coefficient)

Feature importance

Feature importance was analysed based on the RF and NB algorithms predicting the first outcome, *action to seek help* at study-end on the full data set ($N_1 = 185$). For both models, features were first ranked in decreasing order according to their importance values. Three sets of features among the most important ones were taken into account for subsequent analysis:

- the top-10 features indicated by the two models, resulted in a total of 14 best features;

- the top-15 features indicated by the two models, resulted in a total of 19 best features;

- the top-20 features indicated by the two models, resulted in a total of 28 best features.

Figure 2 shows all 28 features with their importance ranking values originating from the RF and NB models. Next, the three sets of features were evaluated for their predictive performance and classification accuracy on the reduced dataset of $N_2 = 131$ participants who completed the follow-up questionnaire. NB and RF were trained on the three different feature sets for

predicting the action to seek help at follow-up. Results are summarized in Table 4. They show

that all feature sets provide good predictive performance and that the NB algorithm

outperforms RF, with an overall accuracy ranging between 73.2% to 75.0% and an MCC

between .426 and .469. Class-specific classification accuracy is comparable between NB and

RF, with the action class correctly classified in 52 to 63% of the cases and the no-action class

in 82 to 86% of cases. Figure 3 additionally provides an overview of model-specific relative

confusion matrices for action to seek help at study end and at follow-up.

Figure 2. Ranking of the most important features to predict *action to seek help* at study-end. Importance ranking are shown for the 20 most important features for the two models (NB and RF) trained on the full dataset (N_1=185), as previously summarized in Table 3. A total of 28 features are arranged on the *y*-axis with respect to their average ranking between the two models. (HL: hearing loss; HA: hearing aids; *M*: mean; *SD*: standard deviation; NB: Naïve Bayes; RF: Random Forest)

465 **Table 4.** Model-specific overall performance (aggregated accuracy on the test set and MCC) and class specific classification
466 accuracy (proportion of correct classifications) for the outcome measured at follow-up.

Action to seek help at follow-up

	Model	Hyperparameters		Overall performance measures		Class-specific classification accuracy	
			space	Test accuracy	MCC	action (n_{21}=52)	no action (n_{22}=79)
Top-10 features (*total 14*)	NB			.74	.44	.55	.86
	RF	ntree mtry nodesize	800 [1,4] [1,5]	.73	.43	.58	.84

Features	Model	Hyperparameter					
Top-15 features (*total 19*)	NB			.73	.43	.58	.83
	RF	ntree mtry nodesize	800 [1,10] [1,5]	.73	.41	.56	.84
Top-20 features (*total 28*)	NB			.75	.47	.63	.83
	RF	ntree mtry nodesize	800 [1,5] [1,5]	.70	.36	.52	.82

Note: Results are based on the reduced follow-up dataset (N_2=131). The table shows the results for three sets of features (10, 15 and 20 most important features for the two models). For NB 10-fold CV repeated 50 times was used. For RF the table additionally shows the hyperparameter search space used in the resampling procedure. (NB: Naïve Bayes classifier; RF: Random Forest; SVM: Support Vector Machine; MCC: Matthews Correlation Coefficient)

Figure 3. Visualization of the model-specific relative confusion matrices for the outcomes action to seek help, measured at study end (N_1 = 185) and action to seek help at follow-up (N_2 = 131) considering the set of top-10 most important features. Each plot illustrates the class-specific relative frequencies for each observed and predicted class, and provides a more immediate overview of each model's sensitivity (TP) and specificity (TN). (TP: True positive; TN: True Negative; FP: False Positive; FN: False Negative; NB: Naïve Bayes classifier; RF: Random Forest; SVM: Support Vector Machine)

The following is a brief description of 14 features ranked among the most important (previously identified as the top-10 features for the two models) in the prediction of *action to seek help at follow-up*:

- *Motivation to seek help* and source of this motivation at the beginning of the study;

- Individual's attitude and expectations towards HA, including: *general attitude towards HA*; *expectations towards HA*, as measured by the global score of the Expected Consequences of Hearing-Aid Ownership (ECHO) questionnaire (42), which assesses positive and negative expectations towards HA, expected services and costs, and assumptions about change in the personal image in case of HA use; and *stigma towards HA*, as measured by the Denial of Hearing Loss scale of the Attitudes towards Loss of Hearing Questionnaire (ALHQ–3.0) (43), which assesses acceptance of hearing aids and acknowledgement of HL;

- *Hearing performance* (mean SRT and its variability in the DTT) and percentage of negative feedback received (indicating poor performance);

- Perceived consequences of HL, including: *emotional consequences of HL*, as measured by the corresponding subscale of the Hearing Handicap Inventory (HHIE/A) questionnaire (40); and *self-reported HL*, as measured by the Qualities of Hearing subscale of the Speech, Spatial and Qualities of Hearing (SSQ) questionnaire (42), which addresses recognition, perceived clarity and naturalness of everyday sounds, as well as listening effort experienced in different hearing contexts;

- *High sensory-sensitivity personality*, as assessed through the Ease of Excitation subscale of the High Sensitive Person Scale (HSPS) questionnaire (44), which assesses emotional reactivity to physiological stimuli;

- Reported *physical health*, measured by the corresponding items of the Short-Form Health Survey-12 (SF-12) questionnaire (45).

501   Among the top-20 features, further features that were assigned lower importance values

502   but provide further insights toward targeted counselling in a VHC are:

503   ■ *Neuroticism*, which refers to a predisposition of experiencing negative emotions,

504   and *conscientiousness*, which relates to being proactive, organized and

505   methodical (2). These personality traits were assessed through the

506   corresponding items of the NEO Five-Factor Inventory (NEO-FFI)

507   questionnaire (46);

508   ■ *Monthly income*, which was categorized with three cut-off values (less than

509   1.500 Euros, between 1.500 and 2.500 Euros, between 2.500 and 4.000 Euros

510   and above 4.000 Euros);

511   ■ *Social consequences of HL*, as measured by the corresponding subscale of the

512   Hearing Handicap Inventory (HHIE/A) questionnaire (40); and the number of

513   people included in the individual's *social networks*, measured by the respective

514   index of the Social Network Index (47).

516   **Discussion**

The present study contributes to the identification of individuals' hearing-related and

psychological features that predict the readiness to seek professional help for hearing loss (HL).

Cross-sectional and longitudinal data have been collected in a large ecological momentary

assessment study. Potential users of a future Virtual Hearing Clinic (VHC), namely individuals

with subjective hearing difficulties, were classified into help-seekers and non-seekers by means

of supervised machine learning algorithms. Feature importance analyses helped to derive

relevant individual characteristics to be used in a prospective psychological profiling algorithm

of help seeking and support the design of a targeted recommendations module in a VHC.

Which machine learning model can best predict help-seeking?

Three machine learning classifiers correctly predicted *action to seek help* at study-end

in 66 to 70% of cases, clearly improving over chance prediction. This is a promising result

considering the complexity of the prediction outcome. As discussed earlier, several individual

factors can influence the decision to pursue hearing health care services and there can be

discrepancy between the intention and the upcoming concrete action. When predicting *action*

to seek help at follow-up with the selected important features, the performance of random forest

and naïve bayes models improved up to 75%, despite the smaller dataset ($N_2 = 131$). Measuring

help-seeking two months after the study-end provided a more valid measure of the participants'

behaviour. For example, among the participants who were categorized as non-seekers at study-

end, seven reported at follow-up to have completed an appointment with a hearing professional

and 12 were willing to do so in the near future. A more precise and objective outcome measure,

along with a larger dataset will contribute to further increase classification accuracy. In this

direction, future studies could benefit from the adoption of a theoretical model of health-related

behaviour change (48, 49). A theoretical conceptualization of the different stages and factors

that lead to behaviour change will help in defining precise outcome measures. Another factor

that might have limited overall classification performance is the type of predictive features

included, which were mostly self-reports rather than objectively measured traits. Momentary

assessments of such traits will need to be considered in the future in order to minimize self-

report biases. Note that multicollinearity as a potential statistical limitation was ruled out (the

correlation plots are available in the Supplementary Material 4). Random forest revealed high

accuracy in identifying *no action* both at study-end and at follow-up. Accurate identification

of non-seekers is the most relevant performance outcome in a VHC to design targeted

recommendations. Indeed, the envisioned profiling algorithm should be a system with high

specificity that motivates and promotes help-seeking, especially in those cases where users

would not spontaneously take action.

Which psychological- and hearing-related features are mostly relevant to profile a non-help-seeker?

Hearing performance appears to be one of the most important features to predict help-seeking. The association between degree of hearing loss and help-seeking, as well as with hearing aids (HA) uptake, is well established in literature (4, 9–11). The present results additionally highlight – for the first time in the literature – the predictive role of intra-individual fluctuations of hearing performance, emphasizing the need to move beyond the traditional view of considering hearing a stable neurosensory process (50). The implementation of repeated daily measurements of hearing performance through ecological momentary tests provides further insight on the impact of HL on the individual's everyday life. In line with this, feature importance findings emphasize the relevance of self-reports on the consequences of HL. The assessment should consider self-reported listening effort in different contexts, as well as perceived handicap and potential socio-emotional consequences of HL. Indeed, individuals who report greater negative impact of HL in their life are more prone to seek help and later uptake HA (9, 13). The individuals' self-awareness of HL can be validated or improved by providing repeated feedback on hearing performance in a VHC. As occurred at the follow-up survey, 85% of participants (out of $N_2 = 131$) reported increased awareness for their hearing abilities after receiving repeated feedback during the study.

Audiological factors alone are insufficient to predict help-seeking behaviour and the present results highlight the relevance of psychological characteristics. Individuals characterized by high sensitivity to sensory stimuli (21) and emotionally instable personality traits seem to perceive increased psychological discomfort following HL, even in presence of effective HA treatment (2). Furthermore, it appears that individuals who are proactive and methodical (i.e. with high conscientious trait values) are inclined to plan preventive medical consultations and to show more compliance to treatment recommendations. A further important

predictive feature for help-seeking is physical health. Literature suggests that individuals who

report poor health are more prone to uptake a HA (6) in order to improve their well-being.

However, other findings revealed individuals with better self-reported health as more likely to

undertake the first steps towards help-seeking (4, 13). Finally, according to present results and

previous findings (11, 13), investigating stigma, attitude and expectations towards HA informs

on individuals' readiness to seek help and to later uptake a HA. Stigma and negative stereotypes

related to HA may deter individuals from seeking help and can represent a barrier to HA use

(7, 10). An additional negative expectation, and potential obstacle to seeking help, is the

anticipated economic burden. Income was one of the most important features predicting help-

seeking in the present study. Indeed, individuals with higher socioeconomic status (9) and

higher income or pension earnings (3, 13) are more likely to initiate the process towards HA

uptake.

How can feature importance measures inform the design of targeted recommendations

for users of a future VHC?

Individuals' attitudes towards HA and self-recognition of HL not only are important

predictive features, but could additionally be considered as intervention target to motivate and

promote access to hearing care services, where needed. On the one hand, users profiled as

decided help-seekers after repeated use of a VHC could receive simple and straightforward

indications regarding hearing care needed. Those among them who should uptake a hearing

device (given their audiological outcome), could benefit from additional information on

available hearing-care services and professionals, in order to facilitate faster HA adoption rates.

On the other hand, users in need of HA who are profiled as non-seekers should obtain more

elaborated, targeted recommendations, which could act as an intervention on modifiable

predictive features like self-recognition of HL and attitude towards HA. Non-seekers could

receive a more detailed feedback on their hearing performance to improve awareness on their

hearing deficit. To promote positive attitude towards HA, information on the large range of

devices available, as well as example successful peer cases, could be provided. Knowledge on

accessible financial support for HA by insurance companies could additionally promote HA

uptake, given the predictive role of income and socio-economic status. Furthermore, an

implemented HA simulator in the VHC could offer possibilities to experience improved

listening conditions and improve positive expectations towards a HA. Elaborated information

on HA technologies, like the benefits of noise control and noise reduction algorithms, could

promote help-seeking by fostering knowledge in individuals who are more sensitive to

environmental noise (high sensory sensitivity trait).

The effectiveness of such recommendations could be increased through targeting or

tailoring communication. Targeted messages are designed for a specific population segment,

while tailored communication is individualized to the individual and was shown to be the most

effective in promoting health behaviour changes (48). Indeed, messages that are congruent to

the personality traits of the audience are more positively evaluated, persuasive and interesting

(51). The predictive role of personality traits such as neuroticism and conscientiousness for

help-seeking behaviour must be considered for efficient communication, both in the context of

a VHC and in clinical counselling. Individuals with high neuroticism trait are more susceptible

to perceived disease (52) and are drawn to action by motives of safety and security (51).

Therefore, they might be more persuaded from recommendations that highlight the positive

effects of HA and of an early intervention. Individuals characterized by low conscientious

personality require more support to improve adherence to treatment recommendations (53),

and would need more guidance in planning the subsequent steps towards help-seeking. For

example, a VHC would prompt these individuals regularly and remind them on the next step

to be taken.

Conclusion

This research provides initial knowledge towards a selection of tests and questionnaires

to be included in a psychological profiling algorithm to predict readiness to seek professional

help in individuals with self-reported hearing difficulties, in the context of a VHC.

Complementing the audiological assessments with a psychological profiling algorithm will

enable a VHC to capture a comprehensive picture of the user and deliver targeted and efficient

treatment recommendations depending on those profiles. The benefits of a psychological

assessment of the individual with HL might also extend to other applications within a VHC.

Future studies might explore potential relationships between psychological traits and, among

others HA fitting preferences and endurance in the fine-tuning process towards an optimal

aiding solution, openness to try new, elaborated technical solutions, preference for particular

VHC usability features. To conclude, a VHC as an easily accessible and affordable mobile

diagnostic tool could facilitate faster access to hearing-care services and subsequent earlier

intervention, where needed, to pursue the long-term goal of achieving "hearing for all".

Declarations

Conflicting interests: The Authors declare that there is no conflict of interest.

Funding: This research has been funded by the Deutsche Forschungsgemeinschaft

(DFG, German Research Foundation) under Germany's Excellence Strategy – EXC 2177/1 -

Project ID 390895286.

Ethical approval: The study protocol was approved by the Research Ethics Committee

of the Carl von Ossietzky Universität of Oldenburg (08.09.2021, Drs.EK/2020/020-01).

Guarantor: AH.

Contributorship: GA, AH and MB conceptualized the study and were involved in

protocol development, study design and data analysis. IK was involved in study design and

data analysis. BK contributed to study conceptualization and obtained funding. GA was

responsible for participants' recruitment, data collection and wrote the first draft of the

manuscript. All authors reviewed and edited the manuscript and approved the final version of

the manuscript.

Acknowledgements: We thank all the participants who took part in this study and the

Universities that helped us with recruitment: University of Oldenburg, Gasthörstudium;

Goethe-Universität Frankfurt am Main, Universität des 3. Lebensalters; Freie Universität

Berlin, GasthörerCard Programm; University of Köln, Gasthörer- und Seniorenstudium;

University of Kassel, Gasthörendenprogramm; University of Bielefeld, Studieren ab 50.

Preliminary results of this research were presented on symposia and conferences of the

"Hearing4All" Cluster of Excellence, on the VCCA Conference 2022 and the DGPs

Conference 2022. A preprint of the manuscript was published on medrxiv.org in February 2023.

Analyses scripts are available at https://doi.org/10.5281/zenodo.7635920. The data is available

upon request to the corresponding authors.

References

1. Haile LM, Kamenov K, Briant PS, Orji AU, Steinmetz JD, Abdoli A, et al. Hearing loss prevalence and years lived with disability, 1990–2019: findings from the Global Burden of Disease Study 2019. The Lancet. 2021 Mar 13;397(10278):996–1009.

2. Cox RM, Alexander GC, Gray GA. Who wants a hearing aid? Personality profiles of hearing aid seekers. Ear Hear. 2005;26(1):12–26.

3. Saunders GH, Chisolm TH, Wallhagen MI. Older adults and hearing help-seeking behaviors. Am J Audiol. 2012;21(2):331–7.

4. Sawyer CS, Armitage CJ, Munro KJ, Singh G, Dawes PD. Biopsychosocial Classification of Hearing Health Seeking in Adults Aged over 50 Years in England. Ear Hear. 2020;1215–25.

5. Fisher DE, Li CM, Hoffman HJ, Chiu MS, Themann CL, Petersen H, et al. Sex-specific predictors of hearing-aid use in older persons: The age, gene/environment susceptibility-Reykjavik study. Int J Audiol. 2015 Sep 2;54(9):634–41.

6. Nixon G, Sarant J, Tomlin D, Dowell R. Hearing aid uptake, benefit, and use: The impact of hearing, cognition, and personal factors. J Speech Lang Hear Res. 2021;64(2):651–63.

7. Southall K, Gagné JP, Jennings MB. Stigma : A negative and a positive influence on help-seeking for adults with acquired hearing loss. Int J Audiol. 2010;49:804–14.

8. Yong M, Willink A, McMahon C, McPherson B, Nieman CL, Reed NS, et al. Access to adults' hearing aids: Policies and technologies used in eight countries. Bull World Health Organ. 2019;97(10):699–710.

9. Simpson AN, Matthews LJ, Cassarly C, Dubno JR. Time from Hearing Aid Candidacy to Hearing Aid Adoption: A Longitudinal Cohort Study. Ear Hear. 2019 May 1;40(3):468–76.

10. Jenstad L, Moon J. Systematic Review of Barriers and Facilitators to Hearing Aid Uptake in Older Adults. Audiol Res. 2011 Mar 23;1(1):91–6.

11. Vestergaard Knudsen L, Öberg M, Nielsen C, Naylor G, Kramer SE. Factors Influencing Help Seeking, Hearing Aid Uptake, Hearing Aid Use and Satisfaction With Hearing Aids: A Review of the Literature. Trends Amplif. 2010;14(3):127–54.

12. Schneider JM, Gopinath B, McMahon CM, Britt HC, Harrison CM, Usherwood T, et al. Role of general practitioners in managing age-related hearing loss. Med J Aust. 2010 Jan 1;192(1):20–3.

13. Meyer C, Hickson L, Lovelock K, Lampert M, Khan A. An investigation of factors that influence help-seeking for hearing impairment in older adults. Int J Audiol. 2014;53(S1):S3–17.

14. Convery E, Hickson L, Meyer C, Keidser G. Predictors of hearing loss self-management in older adults. Disabil Rehabil. 2019;41(17):2026–35.

15. Meyer C, Hickson L, Fletcher A. Identifying the barriers and facilitators to optimal
hearing aid self-efficacy. Int J Audiol [Internet]. 2014;53(S1). Available from:
https://pubmed.ncbi.nlm.nih.gov/24447235/

16. Smyth J, Stone A. Ecological Momentary Assessment Research in Behavioral medicine.
J Happiness Stud. 2003;4(1):35–52.

17. Arslan RC, Walther MP, Tata CS. formr: A study framework allowing for automated
feedback generation and complex longitudinal experience-sampling studies using R.
Behav Res Methods 2019 521. 2019 Apr 1;52(1):376–87.

18. Senese VP, Ruotolo F, Ruggiero G, Iachini T. The italian version of the weinstein noise
sensitivity scale: Measurement invariance across age, sex, and context. Eur J Psychol
Assess. 2012;28(2):118–24.

19. Meis M, Huber R, Fischer RL, Schulte M, Spilski J, Meister H. Development and
psychometric properties of the sound preference and hearing habits questionnaire (SP-
HHQ). Int J Audiol. 2018;57(0):S118–29.

20. Goldberg LR. An alternative 'description of personality': The Big-Five factor structure. J
Pers Soc Psychol. 1990;59:1216–29.

21. Aron EN, Aron A. Sensory-Processing Sensitivity and Its Relation to Introversion and
Emotionality. J Pers Soc Psychol. 1997;73(2):345–68.

22. Posner J, Russell JA, Peterson BS. The circumplex model of affect: An integrative
approach to affective neuroscience, cognitive development, and psychopathology. Dev
Psychopathol. 2005 Jul;17(3):715.

23. Scott SB, Sliwinski MJ, Zawadzki M, Stawski RS, Kim J, Marcusson-Clavertz D, et al. A
Coordinated Analysis of Variance in Affect in Daily Life. Assessment. 2020 Dec
1;27(8):1683–98.

24. Smits C, Kapteyn T, Houtgast T. Development and validation of an automatic speech-in-
noise screening test by telephone. Int J Audiol. 2004 Feb 1;43:15–28.

25. Buschermöhle M, Wagener KC, Berg D, Meis M, Kollmeier B. The German Digits
Triplets Test (Part I): Implementations for Telephone, Internet and Mobile Devices.
Audiol Acoust Für Audiol. 2014;54(4):139–45.

26. Smits C, Merkus P, Houtgast T. How we do it: The Dutch functional hearing-screening
tests by telephone and internet. Clin Otolaryngol. 2006 Oct;31(5):436–40.

27. R Core Team. R: A Language and Environment for Statistical Computing [Internet].
Vienna, Austria; 2019. Available from: https://www.r-project.org/

28. Arslan RC. Formr.Org Survey Framework Utility R Package [Internet]. Zenodo; 2018.
Available from: https://zenodo.org/record/594253

29. Bischl B, Lang M, Kotthoff L, Schiffner J, Richter J, Studerus E, et al. mlr: Machine
Learning in R. J Mach Learn Res. 2017;17(170):1–5.

30. Rhys HI. Machine Learning with R, the tidyverse, and mlr. Manning Publications Co.;
2020.

31. Becker M, Binder M, Bischl B, Foss N, Kotthoff L, Lang M. mlr3book [Internet]. 2022.
Available from: https://mlr3book.mlr-org.com

32. James G, Witten D, Hastie T, Tibshirani R. An Introduction to Statistical Learning
[Internet]. Design. New York, NY: Springer New York; 2013. 618 p. (Springer Texts in
Statistics; vol. 103). Available from: http://books.google.com/books?id=9tv0taI8l6YC

33. Probst P, Bischl B. Tunability: Importance of Hyperparameters of Machine Learning
Algorithms [Internet]. Vol. 20, Journal of Machine Learning Research. 2019 p. 1–32.
Available from: http://jmlr.org/papers/v20/18-444.html.

34. Becker M, Ullmann T, Lang M, Bischl B, Richter J, Binder M. mlr3gallery: Practical
Tuning Series - Tune a Support Vector Machine [Internet]. 2021. Available from:
https://mlr3gallery.mlr-org.com/posts/2021-03-09-practical-tuning-series-tune-a-support-
vector-machine/

35. Hsu CW, Chang CC, Lin CJ. A Practical Guide to Support Vector Classification. 2016;
Available from: http://www.csie.ntu.edu.tw/~cjlin

36. Ali A, Mariyam Shamsuddin S, Ralescu A, Ralescu AL. Classification with class
imbalance problem: A review. Classif Int J Adv Soft Compu Appl [Internet]. 2013;5(3).
Available from: https://www.researchgate.net/publication/288228469

37. Delgado R, Tibau XA. Why Cohen's Kappa should be avoided as performance measure
in classification. PLOS ONE. 2019 set;14(9):e0222916.

38. Kuhn M, Vaughan D, Hvitfeldt E, RStudio. yardstick: Tidy Characterizations of Model
Performance [Internet]. 2022. Available from: https://CRAN.R-
project.org/package=yardstick

39. Jurman G, Riccadonna S, Furlanello C. A Comparison of MCC and CEN Error Measures
in Multi-Class Prediction. PLOS ONE. 2012 ago;7(8):e41882.

40. Kuhn M. The caret Package [Internet]. 2019. Available from:
https://topepo.github.io/caret/index.html

41. Kuhn M. caret: Classification and Regression Training [Internet]. 2020. Available from:
https://CRAN.R-project.org/package=caret

42. Kiessling J, Grugel L, Meister H, Meis M. Übertragung der Fragebögen SADL , ECHO
und SSQ ins Deutsche und deren Evaluation. Z Für Audiol. 2011;50(1):6–16.

43. Saunders GH, Cienkowski KM, Forsline A, Fausti S. Normative data for the Attitudes
towards Loss of Hearing Questionnaire. J Am Acad Audiol. 2005;16(9):637–52.

44. Konrad S, Herzberg PY. Psychometric properties and validation of a German high
Sensitive person scale (HSPS-G). Eur J Psychol Assess. 2019;35(3):364–78.

45. Drixler K, Morfeld M, Glaesmer H, Brähler E, Wirtz MA. Validierung der Messung
gesundheitsbezogener Lebensqualität mittels des Short-Form-Health-Survey-12 (SF-12
Version 2.0) in einer deutschen Normstichprobe. Z Psychosom Med Psychother.
2020;66(3):272–86.

46. Borkenau P, Ostendorf F. NEO-Fünf-Faktoren-Inventar (NEO-FFI) nach Costa und
McCrae: Handanweisung. Göttingen: Hogrefe.; 1993.

47. Cohen S. Social Ties and Susceptibility to the Common Cold. JAMA J Am Med Assoc.
1997 Jun 25;277(24):1940.

48. Noar SM, Benac CN, Harris MS. Does tailoring matter? Meta-analytic review of tailored
print health behavior change interventions. Psychol Bull. 2007;133(4):673–93.

49. Davis R, Campbell R, Hildon Z, Hobbs L, Michie S. Theories of behaviour and
behaviour change across the social and behavioural sciences: a scoping review. Health
Psychol Rev. 2015 Aug 7;9(3):323–44.

50. Kuhlmann I, Angonese G, Thiel C, Kollmeier B, Hildebrandt A. Are there good days and
bad days for hearing? – Quantifying intraindividual hearing variability. In revision.

51. Hirsh JB, Kang SK, Bodenhausen GV. Personalized Persuasion: Tailoring Persuasive
Appeals to Recipients' Personality Traits. Psychol Sci. 2012 Jun 1;23(6):578–81.

52. Ferguson E. Personality is of central concern to understand health: towards a theoretical
model for health psychology. Health Psychol Rev. 2013 May 1;7(sup1):S32–70.

53. Molloy GJ, O'Carroll RE, Ferguson E. Conscientiousness and Medication Adherence: A
Meta-analysis. Ann Behav Med. 2014 Feb 1;47(1):92–101.

797