

1 *Title:* Hypertension among persons living with HIV — Zambia, 2021; A cross-sectional study of a national  
2 electronic health record system

3

4 *Authors:* Jonas Z. Hines<sup>1</sup>, Jose Tomas Prieto<sup>2</sup>, Megumi Itoh<sup>1</sup>, Sombo Fwoloshi<sup>3,4</sup>, Khozya D. Zyambo<sup>3</sup>,  
5 Dalila Zachary<sup>1</sup>, Cecilia Chitambala<sup>1</sup>, Peter A. Minchella<sup>1</sup>, Lloyd B. Mulenga<sup>3</sup>, Simon Agolory<sup>1</sup>

6

7 *Affiliations:*

8 1. U.S. Centers for Disease Control and Prevention, Lusaka, Zambia

9 2. Palantir Technologies, Paris, France

10 3. Ministry of Health, Lusaka, Zambia

11 4. University of Zambia, School of Medicine

12

13 *Keywords:* Hypertension; Prevalence; HIV; Epidemiology; Non-communicable diseases; Zambia; Africa

14

15 *Corresponding author:*

16 Jonas Hines

17 U.S. Centers for Disease Control & Prevention (CDC) Zambia

18 351 Independence Avenue, Woodlands, Lusaka, Zambia

19 Phone: +260-211-257515

20 Email: [jhines1@cdc.gov](mailto:jhines1@cdc.gov)

21

22

23 **Abstract** (*word count: 337*)

24 *Background*

25 Hypertension is a major risk factor for cardiovascular disease, which is a common cause of death in  
26 Zambia. Data on hypertension prevalence in Zambia are scarce and limited to specific geographic areas  
27 and/or populations. We measured hypertension prevalence among persons living with HIV (PLHIV) in  
28 Zambia using a national electronic health record (EHR) system.

29

30 *Methods*

31 We did a cross-sectional study of hypertension prevalence among PLHIV aged  $\geq 18$  years in Zambia  
32 during 2021. Data were extracted from the SmartCare EHR, which covers  $\sim 90\%$  of PLHIV on treatment in  
33 Zambia. PLHIV with  $\geq 2$  recorded blood pressure (BP) readings in 2021 were included. Hypertension was  
34 defined as  $\geq 2$  elevated BP readings (i.e., systolic BP of  $\geq 140$  mmHg or diastolic BP of  $\geq 90$  mmHg) during  
35 2021 and/or on anti-hypertensive medication recorded in their EHR in the past five years. Multivariable  
36 logistic regression was used to assess associations between hypertension and independent variables.

37

38 *Results*

39 Among 750,098 PLHIV aged  $\geq 18$  years with  $\geq 2$  visits in SmartCare during 2021, 101,363 (13.5%) had  $\geq 2$   
40 blood pressure readings recorded in their EHR. Among these PLHIV, 14.7% (95% confidence interval [CI]:  
41 14.5-14.9) had hypertension during 2021. Only 8.9% of PLHIV with hypertension had an anti-  
42 hypertensive medication recorded in their EHR. The odds of hypertension were greater in older age  
43 groups compared to PLHIV aged 18-29 years (adjusted odds ratio [aOR] for 30-44 years: 2.6 [95% CI: 2.4-  
44 2.9]; aOR for 45-49 years: 6.4 [95% CI: 5.8-7.0]; aOR for  $\geq 60$  years: 14.5 [95% CI: 13.1-16.1]), urban areas  
45 (aOR: 1.9 [95% CI: 1.8-2.1]), and persons prescribed ART for  $\geq 6$ -month at a time (aOR: 1.1 [95% CI: 1.0-  
46 1.2]).

47

48 *Discussion*

49 Hypertension was common among a cohort of PLHIV in Zambia, with few having documentation of  
50 being on antihypertensive treatment. Most PLHIV were excluded from the analysis because of missing  
51 BP measurements in their EHR. Strengthening integrated management of non-communicable diseases in  
52 ART clinics might help to diagnose and treat hypertension in Zambia. Data completeness needs to be  
53 improved to routinely capture cardiovascular disease risk factors, including blood pressure readings  
54 consistently for PHLIV in their EHRs.

55

56 **Article** (*word count: 2,636*)

57 *Background*

58 Non-communicable diseases are common causes of death in sub-Saharan African countries (1). In  
59 Zambia, a country in southern Africa, cerebrovascular accidents (CVAs) and ischemic heart disease are  
60 among most common causes of death (2,3). With widespread availability of antiretroviral therapy (ART),  
61 non-communicable diseases are increasingly a major cause of morbidity among persons living with HIV  
62 (PLHIV) (4,5), including in Zambia (6,7). Hypertension is a known major risk factor for CVA and other  
63 cardiovascular diseases. Although country-specific data on prevalence of hypertension in sub-Saharan  
64 African countries is limited, one meta-analysis estimated prevalence to be 16% (8).

65  
66 PLHIV are at increased risk for noncommunicable diseases including cardiovascular disease (4,9–11). This  
67 risk might be attributed to effects of HIV (i.e., chronic inflammation) and/or side effects (e.g., metabolic  
68 syndrome, renal disease) of some ART (12,13). Some studies indicate PLHIV have a higher hypertension  
69 prevalence than people without HIV (14,15), with roughly 20-25% of PLHIV globally estimated to have  
70 hypertension (5,16,17). However, in sub-Saharan Africa, where most PLHIV reside, hypertension  
71 prevalence is not well-characterized at national levels and it is unclear if the prevalence of hypertension  
72 differs between PLHIV compared to people without HIV (15,18–20).

73  
74 Data on hypertension prevalence in Zambia are limited. In the 2018 Zambia Demographic and Health  
75 Survey, self-reported hypertension prevalence among women was 8.8% (this information was not  
76 ascertained from male participants) (21). Studies that objectively measure blood pressure have been  
77 limited geographically and/or conducted in special populations (7,22–26). For instance, in one study  
78 among PLHIV in Lusaka, hypertension prevalence was 6.4%, of which only one-half of persons were  
79 aware of their hypertension diagnosis (23). Notably, approximately one-quarter of PLHIV with

80 hypertension in that study had suffered a major cardiovascular event, including CVA or myocardial  
81 infarction.

82

83 Thus, current estimates of hypertension among PLHIV in Zambia give an incomplete picture. In this  
84 study, we sought to measure the proportion of PLHIV with hypertension using data routinely captured in  
85 a national electronic health record (EHR) system, SmartCare EHR.

86

### 87 *Methods*

88 We conducted a cross-sectional study of hypertension among PLHIV aged  $\geq 18$  years in Zambia from  
89 January to December 2021 (the last full calendar year of data available). We analyzed data from the  
90 SmartCare EHR, which was introduced for the HIV program in the early 2000s and has been scaled-up  
91 nationally since then. SmartCare EHR supports clinical care by providing patients with their longitudinal  
92 health record at any facility operating the EHR. As of 2021, SmartCare EHR was in use in  $\sim 1,500$  Zambian  
93 health facilities that provide care for approximately 90% of PLHIV on ART in Zambia.

94

95 Digitized SmartCare EHR data from health facilities are routinely consolidated and de-duplicated at the  
96 district and provincial levels, transported to Zambia MOH headquarters in Lusaka, and stored in  
97 Zambia's National Data Warehouse. All patient interactions (including clinical, pharmacy, and laboratory  
98 visits) at health facilities utilizing SmartCare EHR are recorded and data for most patients are entered  
99 into the system in real-time. In cases where health facilities record data on paper forms for retroactive  
100 data entry into SmartCare EHR, the process is completed prior to consolidation. Data from inpatient care  
101 are not captured in SmartCare EHR.

102

103 De-identified demographic data, clinical information, and pharmacy records were extracted from  
104 Zambia's SmartCare EHR system. Data were extracted for demographic characteristics, past medical  
105 history, medications, blood pressure measurements, height and weight, and laboratory data (CD4+  
106 count, HIV viral load, and creatinine). Data are cleaned upon ingestion in Foundry (Palantir Technologies,  
107 Paris, France) by casting laboratory tests results values and physical measurements to numeric format.  
108 For some variables (i.e., non-HIV past medical history, diet, amount of physical activity, smoking history,  
109 blood cholesterol levels, and blood glucose) data were either completely missing or too sparse to  
110 analyze (i.e., <1% complete). Only PLHIV aged  $\geq 18$  years with  $\geq 2$  clinical visits in 2021 were eligible for  
111 inclusion in the study (in Zambia, BP is measured at each clinical visit but not necessarily at other patient  
112 interactions captured in SmartCare EHR [e.g., lab check, pharmacy pick-up]).

113

114 We defined elevated blood pressure as a systolic blood pressure reading of  $\geq 140$  mmHg or diastolic  
115 blood pressure readings of  $\geq 90$  mmHg. We defined hypertension as having  $\geq 2$  systolic blood pressure  
116 (SBP) readings of  $\geq 140$  mmHg or  $\geq 2$  diastolic blood pressure (DBP) readings of  $\geq 90$  mmHg (27) during  
117 2021, or any PLHIV prescribed an antihypertensive medication in SmartCare EHR (including amloride,  
118 amlodipine, atenolol, carvedilol, enalapril, furosemide, hydralazine, hydrochlorothiazide, losartan,  
119 methyldopa, metoprolol, nifedipine, spironolactone, telmisartan, and valsartan) in the past five years.  
120 Because past medical history was not well captured in the EHR, we could not include persons with a  
121 historical hypertension diagnosis regardless of BP measurement readings during 2021. Among PLHIV  
122 with hypertension, we defined grade 2 hypertension as  $\geq 1$  reading with systolic blood pressure  $\geq 160$   
123 mmHg or diastolic blood pressure  $\geq 100$  mmHg and hypertensive urgency as  $\geq 1$  reading with systolic  
124 blood pressure  $\geq 180$  mmHg or diastolic blood pressure  $\geq 110$  mmHg (27).

125

126 Hypertension was measured only among PLHIV with  $\geq 2$  blood pressure readings during 2021 because  
127 the hypertension case definition required two blood pressure readings. 95% confidence intervals (CIs)  
128 were calculated using the Clopper-Pearson exact method in R using epiR package (28). Bivariable logistic  
129 regression was used to measure the association between hypertension and independent variables. We  
130 conducted a multivariable logistic regression with variables with  $\leq 10\%$  missingness (i.e., sex, age group,  
131 province, urban/rural designation, years on ART, current ART regimen, prescription length, and most  
132 recent viral load).

133  
134 We also conducted an additional age- and sex- adjusted analysis to investigate the relationships  
135 between hypertension and kidney function. Specifically, we assessed the relationship between elevated  
136 creatinine (i.e., glomerular filtration rate  $< 60$  mL/min/1.73m<sup>2</sup>) in the past year and hypertension. This  
137 was a separate analysis because creatinine data were too sparse to include in the multivariable analysis.

138  
139 Lastly, we also assessed the association between integrase-inhibitor-containing ART regimens and being  
140 overweight or having obesity, because of prior association between metabolic syndrome and this  
141 medication class. This was relevant to the study objectives given there is an association between  
142 metabolic syndrome and cardiovascular disease. Being overweight or having obesity and hypertension  
143 were the only two components of metabolic syndrome we could measure from SmartCare EHR; the  
144 other components (high triglycerides, low high-density lipoprotein, and elevated fasting glucose) were  
145 too sparsely captured. This analysis was age- and sex-adjusted.

146

147

148 *Results*

149 Among 1,299,263 active PLHIV in SmartCare EHR during 2021, there were 750,098 (57.7%) persons aged  
150  $\geq 18$  years that had  $\geq 2$  clinical visits in 2021 (Figure 1). Of these, 101,363 (13.5%) had  $\geq 2$  blood pressure  
151 readings recorded and were included in the analysis.

152  
153 The complete cohort differed from the analytic cohort for all variables, although absolute differences for  
154 key variables like sex and age were minor (e.g., 5.5% were aged  $\geq 60$  years in the complete cohort  
155 compared to 6.6% in analytic cohort) (Table 1). Health facilities with direct electronic data entry at the  
156 point-of-care, which were concentrated in Lusaka and Southern Provinces, had greater blood pressure  
157 data completeness than health facilities where data was captured on paper and retrospectively entered  
158 into SmartCare EHR (22.5% versus 5.3% captured  $\geq 2$  BP readings, respectively).

159  
160 Among PLHIV in the analytic dataset, the mean age was 41.6 years (standard deviation  $\pm 11.4$  years;  
161 range 18-104 years) and 64.7% were females. The median time on ART was 5.0 years (interquartile  
162 range: 2.0–9.0 years) and 95.7% of PLHIV were on dolutegravir-based ART regimens at most recent visit.

163  
164 During 2021, 35.0% of PLHIV had  $\geq 1$  elevated blood pressure reading. During 2021, 14.7% (95% CI: 14.5-  
165 14.9) of PLHIV had hypertension (Table 2). The proportion of PLHIV with hypertension increased with  
166 increasing age, from 4.3% among PLHIV aged 18-29 years, to 10.1% among PLHIV aged 30-44 years, to  
167 21.6% among PLHIV aged 45-59 years, and 37.4% for among PLHIV aged  $\geq 60$  years. Among PLHIV with  
168 hypertension, 60.7% had grade 2 hypertension and 27.0% had hypertensive urgency.

169  
170 Overall, 2.0% of PLHIV had an anti-hypertensive medication recorded in their EHR in the past five years;  
171 8.9% of PLHIV with two or more readings of SBP  $\geq 140$  mmHg or DBP  $\geq 90$  mmHg had an anti-  
172 hypertensive medication recorded in their EHR. Among any PLHIV with an anti-hypertensive medication



173 recorded, 85.9% had  $\geq 1$  reading with SBP  $\geq 140$  mmHg or DBP  $\geq 90$  mmHg and 60.2% had two elevated  
174 readings (i.e., were still hypertensive).

175

176 In the multivariable model, the odds of hypertension were greater for older age groups, PLHIV residing  
177 in urban areas and certain provinces, and PLHIV prescribed ART for  $\geq 6$ -month at a time (Table 2).

178 Although dolutegravir-based regimens were associated with higher odds of hypertension compared to  
179 efavirenz-based regimens in the bivariable analysis, there was no difference after adjustment in the  
180 multivariable model. PLHIV who were overweight or had obesity had greater odds of hypertension than  
181 normal weight PLHIV (although data missingness precluded inclusion of this characteristic in the  
182 multivariable model).

183

184 PLHIV with hypertension had greater odds of having an elevated creatinine (Table 3). Lastly, PLHIV on  
185 dolutegravir-based regimens had higher odds of being overweight or having obesity compared to  
186 persons on other regimens (adjusted OR: 1.16 [95% CI: 1.03–1.32]).

187

## 188 *Discussion*

189 Hypertension was common among PLHIV in Zambia. Many PLHIV with hypertension had dangerously  
190 high blood pressure readings putting them at elevated risk for cardiovascular disease including CVAs and  
191 acute cardiovascular events. This finding might explain why cardiovascular disease are among the most  
192 common causes of death among PLHIV in Zambia (6,29,30). Few PLHIV with hypertension had  
193 documentation of being on antihypertensive treatment and among those that were, most did not have  
194 their blood pressure under control (as demonstrated by elevated blood pressure readings in these  
195 patients). Other studies in Zambia indicate suboptimal levels of hypertension treatment and control  
196 (21,23), which is similar to other countries in Africa (20,31,32). Integrating noncommunicable disease

197 care into routine HIV care might increase prevention, diagnosis, and management of hypertension in  
198 Zambia, potentially reducing cardiovascular disease-related morbidity and mortality (33). Although most  
199 PLHIV were excluded from the study because of missing BP data, to our knowledge, this analysis is the  
200 largest cohort study of hypertension among PLHIV.

201  
202 That many hypertensive PLHIV still had elevated blood pressure readings despite antihypertensive  
203 treatment is demonstrative of the challenge of controlling hypertension even when treated (20,31,32);  
204 nevertheless this finding warrants action, with a focus on strategies to increase treatment of PLHIV with  
205 existing hypertension in Zambia and measures to prevent hypertension among those without it. Being  
206 older and overweight are established risk factors for hypertension, including among PLHIV (11,20). The  
207 observed geographic patterns of hypertension could be related to regional differences in diet (i.e., urban  
208 dwelling PLHIV consuming higher sodium diets or greater pollution exposure than those in rural areas)  
209 and/or be an artifact of poor data quality (i.e., highest proportion of PLHIV with hypertension observed  
210 in provinces that also had better data completeness).

211  
212 Some reports link ART use with increased prevalence of hypertension among PLHIV in Africa (7,34,35).  
213 Although we could not assess this relationship in this study that was confined to PLHIV on ART, longer  
214 ART duration was not associated with hypertension, which is in contrast to findings from several other  
215 studies from countries in Africa (17,36). Some types of ART have been associated with hypertension  
216 (37), but in this analysis there were no associations between ART regimens and hypertension in the  
217 multivariable model.

218  
219 The association of elevated BMI for PLHIV on dolutegravir-based regimens in this analysis could signify  
220 metabolic syndrome among these persons, with a potential side effect of integrase inhibitors (38); we

221 were not able to analyze glucose or lipid measurements to confirm this hypothesis. However, because  
222 many PLHIV in Zambia were transitioned to dolutegravir-based regimens from different ART regimens  
223 (i.e., efavirenz-based regimens) in the recent past, the elevated BMI among participants could also pre-  
224 date their transition to dolutegravir-based regimens so a different study design (i.e., cohort study) is  
225 warranted to further explore this potential signal. Nevertheless, the superior HIV viral load control and  
226 lower risk of HIV treatment failure make dolutegravir the preferred regimen in Zambia and other  
227 countries with generalized HIV epidemics (39).

228  
229 The study had several limitations. Most importantly, blood pressure data completeness was very low,  
230 with only approximately one-eighth of PLHIV in the dataset being analyzed. Despite this limitation, this is  
231 one of the largest studies of hypertension reported from sub-Saharan Africa to date and provides the  
232 first national level study of hypertension in Zambia. However, the estimate is not nationally  
233 representative and, furthermore, only represents an estimate among PLHIV in care who were captured  
234 by the SmartCare EHR. Comparison to non-HIV-infected persons was not possible, but as SmartCare EHR  
235 is integrated into other care settings in Zambia, this will become possible. For some variables, high  
236 amounts of missingness precluded their inclusion in the multivariable model and, furthermore, some  
237 important variables (e.g., non-HIV past medical history) were not available. Additionally, past medical  
238 history was not captured in the dataset, so the proportion of PLHIV with diagnosed hypertension could  
239 not be assessed. Furthermore, very few records had a blood pressure medication documented. This  
240 could reflect low levels of hypertension treatment, limited availability of sphygmomanometers, or could  
241 result from data entry omissions at the point-of-care. If antihypertensive medications were not  
242 consistently recorded, then the true prevalence of hypertension among PLHIV in Zambia is likely higher.  
243 Next, only blood pressure readings occurring over a one-year period were assessed. This approach  
244 reduced the likelihood that two elevated blood pressure measurements were separated by long periods

245 of time, but also led to the exclusion of blood pressure measurements that occurred outside of the  
246 defined period potentially affecting the hypertension estimates. Lastly, the EHR does not capture  
247 inpatient data, so information on consequences of uncontrolled hypertension (e.g., stroke or myocardial  
248 infarction) were not available.

249  
250 This analysis points to a need to improve hypertension management for PLHIV in Zambia. Some of the  
251 existing practices that rely on referring patients with elevated BPs to the outpatient department (i.e.,  
252 urgent care) for further evaluation might result in patient attrition, missing opportunities to adequately  
253 manage this common comorbidity among PLHIV in Zambia. Integrated primary care models for PLHIV  
254 have better outcomes for non-communicable disease management (40), and can even result in better  
255 viral suppression (41). ART clinics in Zambia might benefit from instituting integrated management of  
256 noncommunicable diseases, including life-style modification, anti-hypertensive therapy with appropriate  
257 treatment intensification, and medication adherence assessments for hypertension (42). An integrated  
258 primary care model for conditions like hypertension is possible in countries like Zambia and can improve  
259 patient outcomes (32,33). With integration, multi-month dispensing is a promising approach for co-  
260 management of HIV and hypertension (43).

261  
262 Routinely monitoring for hypertension, along with documenting other health measures (e.g., diet,  
263 smoking status, BMI) in the EHR would not only allow medical providers to determine client's  
264 cardiovascular risk for CVAs or other cardiovascular events and target treatment, but also might provide  
265 data needed to identify specific health facilities or clinicians that would benefit from educational  
266 interventions in management of these risk factors and conditions. EHRs like SmartCare are promising  
267 data sources for noncommunicable diseases surveillance given their reach, routine use in clinical setting,  
268 and richness of information. For this to be successful, data completeness needs to be improved to

269 routinely capture cardiovascular disease risk factors, including blood pressure readings consistently for  
270 PHLIV in their EHRs.

271

272 Hypertension was common among PLHIV in Zambia and many persons might not be adequately  
273 diagnosed or treated. It is important for ART clinic providers to consider hypertension among PLHIV and  
274 institute strategies to manage it appropriately. This will require adequate capacitation of the Zambian  
275 health workforce to recognize and manage hypertension. Additionally, care models that integrate  
276 hypertension (and other NCDs) management into ART clinics are promising strategies to improve care.  
277 EHRs might be used to routinely track program implementation at little additional data collection effort,  
278 and can be adapted into noncommunicable diseases surveillance systems. Addressing hypertension and  
279 other noncommunicable diseases will be important to reducing morbidity and mortality of PLHIV in  
280 Zambia.

281

282 *Ethics statement:* The study protocol was approved by the ERES Converge IRB in Lusaka, Zambia; it was  
283 also reviewed in accordance with CDC human research protection procedures and was determined to be  
284 research, but CDC investigators did not interact with human subjects or have access to identifiable data  
285 or specimens for research purposes. all methods were carried out in accordance with relevant guidelines  
286 and regulations. This project met requirements for waiver of informed consent documentation, which  
287 was granted by ERES Converge IRB in Zambia.

288

289 *Authorship Disclaimer*

290 The findings and conclusions in this report are those of the author(s) and do not necessarily represent  
291 the official position of the funding agencies

292

293 *PEPFAR Funding Acknowledgment*

294 This research has been supported by the President's Emergency Plan for AIDS Relief (PEPFAR) through  
295 the Centers for Disease Control and Prevention (CDC) through a cooperative agreement with the Zambia  
296 Ministry of Health and a cooperative agreement with Palantir Technologies

297

298 *Conflict of interest statement*

299 The authors have no conflicts to disclose

300

301

302 **References**

- 303 1. Abbatfati C, Abbas KM, Abbasi-Kangevari M, Abd-Allah F, Abdelalim A, Abdollahi M, et al. Global  
304 burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic  
305 analysis for the Global Burden of Disease Study 2019. *Lancet*. 2020;396(10258).
- 306 2. Institute for Health Metrics and Evaluation. Zambia: What causes the most deaths? [Internet].  
307 [cited 2020 Nov 17]. Available from: <http://www.healthdata.org/zambia>
- 308 3. Mapoma CC, Munkombwe B, Mwango C, Bwalya BB, Kalindi A, Gona NP. Application of verbal  
309 autopsy in routine civil registration in Lusaka District of Zambia. *BMC Health Serv Res*.  
310 2021;21(1):1–11.
- 311 4. Shah AS V, Stelzle D, Lee KK, Beck EJ, Alam S, Clifford S, et al. Global Burden of Atherosclerotic  
312 Cardiovascular Disease in People Living With HIV: Systematic Review and Meta-Analysis.  
313 *Circulation*. 2018 Sep;138(11):1100–12.
- 314 5. Patel P, Rose CE, Collins PY, Nuche-Berenguer B, Sahasrabuddhe V V, Peprah E, et al.  
315 Noncommunicable diseases among HIV-infected persons in low-income and middle-income  
316 countries: a systematic review and meta-analysis. *AIDS*. 2018 Jul;32 Suppl 1(Suppl 1):S5–20.
- 317 6. Cheelo M. Top causes of Mortality among Community Deaths in Zambia, 2020-2021. In: Annual  
318 National HIV/TB/COVID-19/Hepataitis B Technical Conference. Lusaka, Zambia; 2021.
- 319 7. Musekwa R, Hamooya BM, Koethe JR, Nzala S, Masenga SK. Prevalence and correlates of  
320 hypertension in HIV-positive adults from the Livingstone Central Hospital, Zambia. *Pan Afr Med J*.  
321 2021;39:237.
- 322 8. Twagirumukiza M, De Bacquer D, Kips JG, de Backer G, Stichele R Vander, Van Bortel LM. Current  
323 and projected prevalence of arterial hypertension in sub-Saharan Africa by sex, age and habitat:  
324 an estimate from population studies. *J Hypertens*. 2011 Jul;29(7):1243–52.
- 325 9. Benjamin LA, Corbett EL, Connor MD, Mzinganjira H, Kampondeni S, Choko A, et al. HIV,

- 326 antiretroviral treatment, hypertension, and stroke in Malawian adults: A case-control study.  
327 Neurology. 2016 Jan;86(4):324–33.
- 328 10. Walker RW, Jusabani A, Aris E, Gray WK, Unwin N, Swai M, et al. Stroke risk factors in an incident  
329 population in urban and rural Tanzania: a prospective, community-based, case-control study.  
330 Lancet Glob Heal. 2013 Nov;1(5):e282-8.
- 331 11. Ake JA, Polyak CS, Crowell TA, Kiweewa F, Semwogerere M, Maganga L, et al. Noninfectious  
332 Comorbidity in the African Cohort Study. Clin Infect Dis. 2019;69(4).
- 333 12. Hsue PY, Deeks SG, Hunt PW. Immunologic basis of cardiovascular disease in HIV-infected adults.  
334 J Infect Dis. 2012 Jun;205 Suppl(Suppl 3):S375-82.
- 335 13. Friis-Møller N, Sabin CA, Weber R, d'Arminio Monforte A, El-Sadr WM, Reiss P, et al. Combination  
336 antiretroviral therapy and the risk of myocardial infarction. N Engl J Med. 2003  
337 Nov;349(21):1993–2003.
- 338 14. Gallant J, Hsue PY, Shreay S, Meyer N. Comorbidities among US Patients with Prevalent HIV  
339 Infection - A Trend Analysis. In: Journal of Infectious Diseases. 2017.
- 340 15. Davis K, Perez-Guzman P, Hoyer A, Brinks R, Gregg E, Althoff KN, et al. Association between HIV  
341 infection and hypertension: a global systematic review and meta-analysis of cross-sectional  
342 studies. BMC Med. 2021;19(1).
- 343 16. Xu Y, Chen X, Wang K. Global prevalence of hypertension among people living with HIV: a  
344 systematic review and meta-analysis. J Am Soc Hypertens. 2017 Aug;11(8):530–40.
- 345 17. Bigna JJ, Ndoadoumgue AL, Nansseu JR, Tochie JN, Nyaga UF, Nkeck JR, et al. Global burden of  
346 hypertension among people living with HIV in the era of increased life expectancy: a systematic  
347 review and meta-analysis. J Hypertens. 2020 Sep;38(9):1659–68.
- 348 18. Dillon DG, Gurdasani D, Riha J, Ekoru K, Asiki G, Mayanja BN, et al. Association of HIV and ART  
349 with cardiometabolic traits in sub-Saharan Africa: a systematic review and meta-analysis. Int J



- 350 Epidemiol. 2013 Dec;42(6):1754–71.
- 351 19. Mogaka JN, Sharma M, Temu T, Masyuko S, Kinuthia J, Osoti A, et al. Prevalence and factors  
352 associated with hypertension among adults with and without HIV in Western Kenya. PLoS One.  
353 2022;17(1):e0262400.
- 354 20. Kwarisiima D, Balzer L, Heller D, Kotwani P, Chamie G, Clark T, et al. Population-Based  
355 Assessment of Hypertension Epidemiology and Risk Factors among HIV-Positive and General  
356 Populations in Rural Uganda. PLoS One. 2016;11(5):e0156309.
- 357 21. Zambia Statistics Agency, Ministry of Health, University Teaching Hospital Virology laboratory,  
358 ICF. Zambia Demographic and Health Survey 2018. 2019;540.
- 359 22. Yan LD, Chi BH, Sindano N, Bosomprah S, Stringer JS, Chilengi R. Prevalence of hypertension and  
360 its treatment among adults presenting to primary health clinics in rural Zambia: analysis of an  
361 observational database. BMC Public Health. 2015 Sep;15:933.
- 362 23. Bauer S, Wa Mwanza M, Chilengi R, Holmes CB, Zyambo Z, Furrer H, et al. Awareness and  
363 management of elevated blood pressure among human immunodeficiency virus-infected adults  
364 receiving antiretroviral therapy in urban Zambia: a call to action. Glob Health Action.  
365 2017;10(1):1359923.
- 366 24. Rush KL, Goma FM, Barker JA, Ollivier RA, Ferrier MS, Singini D. Hypertension prevalence and risk  
367 factors in rural and urban Zambian adults in western province: a cross-sectional study. Pan Afr  
368 Med J. 2018;30:97.
- 369 25. Mulenga D, Siziya S, Rudatsikira E, Mukonka VM, Babaniyi O, Songolo P, et al. District specific  
370 correlates for hypertension in Kaoma and Kasama rural districts of Zambia. Rural Remote Health.  
371 2013;13(3):2345.
- 372 26. Goma FM, Nzala SH, Babaniyi O, Songolo P, Zyaambo C, Rudatsikira E, et al. Prevalence of  
373 hypertension and its correlates in Lusaka urban district of Zambia : a population based survey. Int

- 374 Arch Med [Internet]. 2011;4(1):34. Available from: <http://www.intarchmed.com/content/4/1/34>
- 375 27. Unger T, Borghi C, Charchar F, Khan NA, Poulter NR, Prabhakaran D, et al. 2020 International  
376 Society of Hypertension Global Hypertension Practice Guidelines. *Hypertension*. 2020;75(6).
- 377 28. Stevenson M, Sergeant E, Nunes T, Heuer C, Marshall J, Sanchez J, et al. epiR: Tools for the  
378 Analysis of Epidemiological Data [Internet]. 2021. Available from: [https://cran.r-](https://cran.r-project.org/package=epiR)  
379 [project.org/package=epiR](https://cran.r-project.org/package=epiR)
- 380 29. Nutakki A, Chomba M, Chishimba L, Zimba S, Gottesman RF, Bahouth MN, et al. Risk factors and  
381 outcomes of hospitalized stroke patients in Lusaka, Zambia. *J Neurol Sci*. 2021 May;424:117404.
- 382 30. Kapombe P, Cheelo M, Kamalongo K, Tally L, Stoops E, Mwangi C, et al. Most common causes of  
383 death in Zambia by HIV status, 2020-2021. In: *AIDS 2022*. Montreal, Canada; 2022.
- 384 31. Muddu M, Tusubira AK, Sharma SK, Akiteng AR, Ssinabulya I, Schwartz JI. Integrated  
385 Hypertension and HIV Care Cascades in an HIV Treatment Program in Eastern Uganda: A  
386 Retrospective Cohort Study. *J Acquir Immune Defic Syndr*. 2019 Aug;81(5):552–61.
- 387 32. Heller DJ, Balzer LB, Kazi D, Charlebois ED, Kwarisiima D, Mwangwa F, et al. Hypertension testing  
388 and treatment in Uganda and Kenya through the SEARCH study: An implementation fidelity and  
389 outcome evaluation. *PLoS One*. 2020;15(1):e0222801.
- 390 33. Hickey MD, Ayieko J, Owaraganise A, Sim N, Balzer LB, Kabami J, et al. Effect of a patient-  
391 centered hypertension delivery strategy on all-cause mortality: Secondary analysis of SEARCH, a  
392 community-randomized trial in rural Kenya and Uganda. *PLoS Med*. 2021 Sep;18(9):e1003803.
- 393 34. Nduka CU, Stranges S, Sarki AM, Kimani PK, Uthman OA. Evidence of increased blood pressure  
394 and hypertension risk among people living with HIV on antiretroviral therapy: a systematic  
395 review with meta-analysis. *J Hum Hypertens*. 2016 Jun;30(6):355–62.
- 396 35. Lubega G, Mayanja B, Lutaakome J, Abaasa A, Thomson R, Lindan C. Prevalence and factors  
397 associated with hypertension among people living with HIV/AIDS on antiretroviral therapy in


- 398 Uganda. *Pan Afr Med J.* 2021;38:216.
- 399 36. Seaberg EC, Muñoz A, Lu M, Detels R, Margolick JB, Riddler SA, et al. Association between highly  
400 active antiretroviral therapy and hypertension in a large cohort of men followed from 1984 to  
401 2003. *AIDS.* 2005 Jun;19(9):953–60.
- 402 37. van Zoest RA, Wit FW, Kooij KW, van der Valk M, Schouten J, Kootstra NA, et al. Higher  
403 prevalence of hypertension in HIV-1-infected patients on combination antiretroviral therapy is  
404 associated with changes in body composition and prior stavudine exposure. *Clin Infect Dis.*  
405 2021;63(2).
- 406 38. Hamooya BM, Mulenga LB, Masenga SK, Fwemba I, Chirwa L, Siwingwa M, et al. Metabolic  
407 syndrome in Zambian adults with human immunodeficiency virus on antiretroviral therapy:  
408 Prevalence and associated factors. *Medicine (Baltimore).* 2021 Apr;100(14):e25236.
- 409 39. Phillips AN, Venter F, Havlir D, Pozniak A, Kuritzkes D, Wensing A, et al. Risks and benefits of  
410 dolutegravir-based antiretroviral drug regimens in sub-Saharan Africa: a modelling study. *Lancet*  
411 *HIV [Internet].* 2019 Feb 1;6(2):e116–27. Available from: [https://doi.org/10.1016/S2352-](https://doi.org/10.1016/S2352-3018(18)30317-5)  
412 [3018\(18\)30317-5](https://doi.org/10.1016/S2352-3018(18)30317-5)
- 413 40. Appenheimer A Ben, Bokhour B, McInnes DK, Richardson KK, Thurman AL, Beck BF, et al. Should  
414 Human Immunodeficiency Virus Specialty Clinics Treat Patients With Hypertension or Refer to  
415 Primary Care? An Analysis of Treatment Outcomes. *Open forum Infect Dis.* 2017;4(1):ofx005.
- 416 41. Hoang T, Goetz MB, Yano EM, Rossman B, Anaya HD, Knapp H, et al. The impact of integrated HIV  
417 care on patient health outcomes. *Med Care.* 2009 May;47(5):560–7.
- 418 42. Topp SM, Chipukuma JM, Chiko MM, Matongo E, Bolton-Moore C, Reid SE. Integrating HIV  
419 treatment with primary care outpatient services: Opportunities and challenges from a scaled-up  
420 model in Zambia. *Health Policy Plan.* 2013;28(4).
- 421 43. Kimera ID, Namugenyi C, Schwartz JI, Musimbaggo DJ, Ssenyonjo R, Atukunda P, et al. Integrated

422 multi-month dispensing of antihypertensive and antiretroviral therapy to sustain hypertension

423 and HIV control. *J Hum Hypertens.* 2022 Mar;1–7.

424

425



426 **Figures and Tables**

427

428 *Figure 1.* Sample size flow diagram for analysis of persons living with HIV with hypertension in Zambia,  
429 2021

430

431

432 \* This is the population which was eligible for inclusion in the analysis

433

434

435  
436 *Table 1.* Comparison of the variable distribution and completeness in the entire and analytic datasets for  
437 hypertension among persons living with HIV — Zambia, 2021

Variable	Entire dataset, n (%) (N=1,236,040)	Analytic dataset, n (%) (N=101,363)	p-value	
<i>Sex</i>				
Female	781,829 (63.3)	65,570 (64.7)	<0.01	
Male	454,211 (36.7)	35,793 (35.3)		
<i>Age group</i>				
18-29 years	232,099 (18.8)	16,191 (16.0)	<0.01	
30-44 years	585,455 (47.4)	45,850 (45.2)		
45-59 years	350,092 (28.3)	32,681 (32.2)		
≥60 years	68,394 (5.5)	6,641 (6.6)		
<i>Province</i>				
Central	125,704 (10.2)	2,813 (2.8)	<0.01	
Copperbelt	238,609 (19.3)	7,197 (7.1)		
Eastern	105,595 (8.5)	3,672 (3.6)		
Luapula	55,168 (4.5)	54.0 (0.1)		
Lusaka	343,981 (27.8)	45,414 (44.8)		
Muchinga	32,978 (2.7)	512 (0.5)		
Northern	52,014 (4.2)	58.0 (0.1)		
Northwestern	36,016 (2.9)	1,893 (1.9)		
Southern	128,299 (10.4)	35,082 (34.6)		
Western	91,156 (7.4)	4,120 (4.1)		
Missing	26,520 (2.1)	548 (0.5)		
<i>Urban/rural designation</i>				
Rural	387,472 (31.3)	18,454 (18.2)		<0.01
Urban	736,921 (59.6)	78,123 (77.1)		
Missing	111,647 (9.0)	4,786 (4.7)		
<i>Years on ART</i>				
0-1	340,014 (27.5)	19,087 (18.8)	<0.01	
2-4	419,400 (33.9)	28,182 (27.8)		
5-9	301,575 (24.4)	32,408 (32.0)		
≥10	175,051 (14.2)	21,686 (21.4)		
<i>Current ART regimen</i>				
Efavirenz-based	49,083 (4.0)	1,423 (1.4)	<0.01	
Dolutegravir-based	1,137,837 (92.1)	97,049 (95.7)		
Both efavirenz and dolutegravir listed	3,291 (0.3)	133 (0.1)		
Other	45,829 (3.7)	2,758 (2.7)		
<i>Most recent ART prescription length</i>				
<3 months	141,776 (11.5)	6,275 (6.2)	<0.01	
3-5 months	373,549 (30.2)	27,839 (27.5)		
6+ months	720,706 (58.3)	67,249 (66.3)		

Missing	9.00 (0.0)	0 (0.0)	
<i>Body mass index (kg/m<sup>2</sup>)</i>			
Normal (18.5-24.9)	336,236 (27.2)	51,056 (50.4)	<0.01
Low (<18.5)	59,860 (4.8)	8,776 (8.7)	
Overweight (25.0-29.9)	107,253 (8.7)	19,668 (19.4)	
Obesity (≥30.0)	53,674 (4.3)	10,585 (10.4)	
Missing	679,017 (54.9)	11,278 (11.1)	
<i>Initial CD4+ count (cells/mm<sup>3</sup>)</i>			
0-200	62,431 (5.1)	7,694 (7.6)	<0.01
201-350	61,544 (5.0)	7,195 (7.1)	
>350	107,473 (8.7)	12,847 (12.7)	
Missing	100,4592 (81.3)	73,627 (72.6)	
<i>Most recent CD4+ count (cells/mm<sup>3</sup>)*</i>			
0-200	59,590 (4.8)	6,994 (6.9)	<0.01
201-350	101,297 (8.2)	12,567 (12.4)	
>350	369,859 (29.9)	49,690 (49.0)	
Missing	705,294 (57.1)	32,112 (31.7)	
<i>Most recent viral load (copies/mL)<sup>†</sup></i>			
<1000	975,952 (79.0)	92,115 (90.9)	<0.01
1,000-9,999	16,881 (1.4)	1,494 (1.5)	
≥10,000	26,338 (2.1)	2,595 (2.6)	
Missing	216,869 (17.5)	5,159 (5.1)	
<i>Most recent creatinine<sup>‡</sup></i>			
Normal	36,457 (2.9)	8,892 (8.8)	<0.01
High	5,829 (0.5)	1,254 (1.2)	
Missing	119,3754 (96.6)	91,217 (90.0)	
<i>Data capture at point-of-care</i>			
Paper-based with retrospective input	495,498 (40.1)	73,406 (72.4)	<0.01
Direct electronic input	552,109 (44.7)	17,197 (17.0)	
Missing	188,433 (15.2)	10,760 (10.6)	

Variables are based on data from the most recent patient interaction in the electronic health record

\* 38.9% of recent CD4+ count measurements were from 2020 or 2021

† 98.4% of recent viral load measurements were from 2020 or 2021

‡ Elevated creatinine defined as ≥115 μmol/L in men and ≥98 μmol/L in women

ART: antiretroviral therapy

438

439

440 *Table 2. Hypertension prevalence and odds ratios of hypertension among persons living with HIV —*  
 441 *Zambia, 2021 (N=101,363)\**

	Prevalence, %	OR (95% CI)	aOR (95% CI) <sup>†</sup>
<i>Overall</i>	14.7		
<i>Sex</i>			
Female	13.7	Referent	Referent
Male	16.4	1.23 (1.19-1.28)	1 (0.96-1.04)
<i>Age group</i>			
18-29 years	4.3	Referent	Referent
30-44 years	10.1	2.53 (2.33-2.75)	2.61 (2.39-2.85)
45-59 years	21.6	6.19 (5.71-6.71)	6.36 (5.81-6.96)
≥60 years	37.4	13.42 (12.26-14.7)	14.54 (13.14-16.09)
<i>Province</i>			
Central	11.8	Referent	Referent
Copperbelt	18.1	1.65 (1.45-1.88)	1.25 (1.08-1.44)
Eastern	10.3	0.87 (0.74-1.01)	0.62 (0.5-0.78)
Luapula	5.6	0.44 (0.14-1.42)	0.4 (0.09-1.67)
Lusaka	15.4	1.37 (1.22-1.54)	1.11 (0.97-1.26)
Muchinga	3.7	0.29 (0.18-0.46)	0.37 (0.21-0.63)
Northern	10.3	0.87 (0.37-2.03)	1.23 (0.42-3.58)
Northwestern	12.4	1.06 (0.88-1.26)	0.92 (0.76-1.11)
Southern	14.6	1.28 (1.14-1.44)	0.97 (0.85-1.1)
Western	8.9	0.73 (0.62-0.85)	0.78 (0.66-0.92)
<i>Urban/rural designation</i>			
Rural	9.0	Referent	Referent
Urban	16.0	1.92 (1.82-2.03)	1.93 (1.81-2.05)
<i>Years on ART</i>			
0-1	10.8	Referent	Referent
2-4	12.8	1.21 (1.14-1.28)	1.04 (0.97-1.11)
5-9	15.0	1.46 (1.38-1.54)	1.02 (0.95-1.08)
≥10	20.1	2.09 (1.97-2.21)	1.07 (1.00-1.14)
<i>ART regimen<sup>‡</sup></i>			
Efavirenz-based	11.5	Referent	Referent
Dolutegravir-based	14.8	1.33 (1.13-1.57)	1.15 (0.96-1.38)
Other	13.7	1.22 (1.00-1.48)	1.00 (0.80-1.24)
<i>Most recent ART prescription length</i>			
<3 months	11.3	Referent	Referent
3-5 months	12.6	1.14 (1.04-1.24)	0.98 (0.89-1.09)
≥6 months	15.9	1.48 (1.36-1.60)	1.11 (1.01-1.22)
<i>Body mass index (kg/m<sup>2</sup>)</i>			
Normal (18.5-24.9)	11.6	Referent	
Low (<18.5)	8.6	0.72 (0.66-0.78)	
Overweight (25-29.9)	19.9	1.9 (1.82-1.99)	



Obesity ( $\geq 30.0$ )	28.7	3.08 (2.93-3.24)	
<i>Initial CD4+ count (cells/mm<sup>3</sup>)</i>			
0-200	15.2	Referent	
201-350	15.0	0.98 (0.9-1.07)	
>350	13.6	0.87 (0.81-0.95)	
<i>Most recent CD4+ count (cells/mm<sup>3</sup>)</i>			
0-200	15.2	Referent	
201-350	17.0	1.15 (1.06-1.24)	
>350	16.2	1.09 (1.01-1.16)	
<i>Most recent viral load count (copies/mL)</i>			
<1000	15.1	Referent	Referent
1,000-9,999	11.5	0.73 (0.62-0.86)	0.96 (0.81-1.14)
$\geq 10,000$	10.9	0.69 (0.61-0.78)	1.02 (0.89-1.17)
<i>Most recent creatinine <sup>¶</sup></i>			
Normal creatinine	17.3	Referent	
Elevated creatinine	28.1	1.88 (1.64-2.15)	

\* Hypertension defined as  $\geq 2$  systolic blood pressure readings of  $\geq 140$  mmHg or  $\geq 2$  diastolic blood pressure readings of  $\geq 90$  mmHg among persons with  $\geq 2$  clinical visits during 2021

<sup>†</sup> Adjusted for sex, age group, province, urban/rural designation, ART regimen, years on ART, script length, and body mass index

<sup>‡</sup> The most recently listed ART regimen in SmartCare electronic health record. If a regimen listed both dolutegravir and efavirenz, then it was excluded from the analysis (n= 128)

<sup>¶</sup> Elevated creatinine defined as  $\geq 115$   $\mu\text{mol/L}$  in men and  $\geq 98$   $\mu\text{mol/L}$  in women, corresponding to glomerular filtration rate of  $< 60$  mL/min/1.73m<sup>2</sup>

aOR: adjusted odds ratio; ART: antiretroviral therapy; CI: confidence interval; OR: odds ratio

443 *Table 3. Prevalence and odds ratios of elevated creatine among persons living with HIV — Zambia, 2021*  
 444 *(N=10,146)\**

	Prevalence, %	OR (95% CI)	aOR (95% CI) <sup>†</sup>
Overall	12.4		
Blood pressure			
Not hypertensive <sup>‡</sup>	10.9	Referent	Referent
Hypertensive <sup>‡</sup>	18.7	1.88 (1.64-2.15)	1.36 (1.18-1.57)
Sex			
Female	11.1	Referent	Referent
Male	14.4	1.35 (1.2-1.52)	1.19 (1.06-1.35)
Age Group			
18-29 years	4.8	Referent	Referent
30-44 years	10.2	2.28 (1.8-2.89)	2.21 (1.75-2.81)
45-59 years	15.7	3.72 (2.95-4.69)	3.38 (2.67-4.28)
≥60 years	27.0	7.42 (5.69-9.68)	6.37 (4.85-8.37)

\* Elevated creatinine defined as ≥115 μmol/L in men and ≥98 μmol/L in women, corresponding to glomerular filtration rate of <60 mL/min/1.73m<sup>2</sup>

<sup>†</sup> Adjusted for presence/absence of hypertension, sex, and age group

<sup>‡</sup> Hypertension defined as as having ≥2 systolic blood pressure readings of ≥140 mmHg or ≥2 diastolic blood pressure readings of ≥90 mmHg during the study period

aOR: adjusted odds ratio; CI: confidence interval; OR: odds ratio

445