Title

NeuroRehabilitation OnLine (NROL): Description of a multidisciplinary group

telerehabilitation innovation for stroke and neurological conditions using the TIDieR

checklist

Authors

Suzanne Ackerley[1,2] (sackerley@uclan.ac.uk), Neil Wilson[1] (nwilson9@uclan.ac.uk), Paul

Boland[1] (pboland2@uclan.ac.uk), Rosemary Peel[3] (rosemary.peel@mbht.nhs.uk , Louise

Connell[1,2] (laconnell@uclan.ac.uk).

[1]University of Central Lancashire, Preston, Lancashire, UK

[2] East Lancashire Hospitals NHS Trust, Burnley, Lancashire, UK

[3]University Hospitals of Morecambe Bay NHS Foundation Trust

Corresponding author

Louise Connell, School of Sport and Health Sciences, University of Central Lancashire, UK.

Email: laconnell@uclan.ac.uk

ABSTRACT

Objective

Providing recommended amounts of rehabilitation for stroke and neurological patients is

challenging. Telerehabilitation is viable for delivering rehabilitation and an acceptable

adjunct to in-person therapy. NeuroRehabilitation OnLine (NROL) was developed as a pilot

and subsequently operationalised as a regional innovation embedded across four National

Health Service (NHS) Trusts. The aim of this paper is to describe the NROL innovation to

assist future implementation and replication efforts.

Methods

The Template for Intervention Description and Replication (TIDieR) checklist, with guidance

from the TIDieR-Telehealth extension, was used to describe NROL. The description was

developed collaboratively by clinical academics, therapists, managers, and researchers.

Updated Consolidated Framework for Implementation Research domains were used to

describe the context in which the innovation was delivered.

Results

NROL delivers online group-based real-time neurorehabilitation with technology assistance

provided, incorporating multidisciplinary targeted therapy and peer support. Procedures,

materials and structure are detailed to demonstrate how NROL is embedded within a

healthcare system. NROL uses existing NHS therapy workforce alongside dedicated NROL

roles, including an essential technology support role. Selection of NROL groups is dependent

on patient need. The NROL innovation is modified over time in response to patient and staff

feedback. NROL described here is situated within a regional stroke and neurorehabilitation

network in North-West England, aligns with local and national strategies, and capitalises on

an existing clinical-academic partnership.

Conclusion

This comprehensive description of a regional NROL innovation, and clarification of core

components, should facilitate other healthcare settings to adapt and implement NROL for

their context. Continuous evaluation alongside implementation will ensure maximal impact

for neurorehabilitation.

STRENGTHS AND LIMITATIONS

• A systematic development process was undertaken using the TIDieR checklist, and

TIDieR-Telehealth extension, allowing a principled method for clear and user-friendly

description of a complex intervention to assist future implementation and

replication efforts.

• Stakeholder engagement was integral to the collaborative development of the NROL

innovation and its description, ensuring informed decision-making representing

multiple perspectives.

• The TIDieR checklist provides a succinct description of an intervention at a point in

time but may not capture the implementation complexities and evolutions.

However, the TIDieR checklist allows reporting on modifications and tailoring which

can help mitigate these limitations.

• The TIDieR checklist does not include an item on context, which is important to fully

understand interventions, hence in this article context it is deliberatively reported to

help situate the innovation.

Abstract word count: 258

Manuscript word count: 3966

BACKGROUND

Despite a wealth of evidence that greater amounts of rehabilitation can improve

outcomes(1-4) providing stroke and brain injury patients with the recommended amount of

therapy remains a clinical and workforce challenge(5). Increasing access to and opportunity

for therapy is a critical step to addressing shortfalls in therapy intensity and ideally needs to

be feasible with limited workforce resources.

Telerehabilitation, the provision of rehabilitation remotely via telecommunication devices,

has developed as a viable complement to in-person therapy and offers one possible solution

to help mitigate challenges. It can deliver conventional in-person therapies online with

equivalent outcomes(5-10) and with similar attendance levels and acceptability to

patients(11). Patients and staff report advantages in terms of saving time, energy and

travel(12-14). Although not a new approach, rapid expanded adoption of telerehabilitation

in routine practice was spurred by the COVID-19 pandemic(15-18), with further encouraging

data collected during this time(14,19-21).   In the UK and Ireland, clinical guidelines for

stroke rehabilitation now recommend the provision of therapist monitored, patient-centred

remote rehabilitation programmes alongside conventional in-person therapy(22).

A group-based real-time telerehabilitation innovation for patients with acquired brain injury

was developed and piloted in London, UK, entitled NROL (NeuroRehabilitation OnLine). This

standalone version of NROL demonstrated positive impacts on patient-reported

outcomes(20). NROL was subsequently adapted and operationalised within the UK National

Health Service (NHS) by embedding a recurring six-week multidisciplinary group-based

telerehabilitation innovation into an existing service. Implementation with concurrent

evaluation at a single NHS Trust level yielded positive results, demonstrating favourable

patient and staff opinion, sustainable staffing and resource efficiencies with good clinical

outcomes for patients(14). NROL was then expanded into a regional innovation involving

four NHS Trusts in the North-West of England aligning to the NHS new Integrated Care

System structure to work collaboratively across regions.   It remains part of a hybrid

approach to service delivery to complement in-person rehabilitation. NROL is acknowledged

as an exemplar initiative for delivering remote rehabilitation(22).

Despite the successful development of NROL, as yet it has only been described in its

standalone version(20) with no descriptive detail available for the current regional, multi-

site model. The Template for Intervention Description and Replication (TIDieR) checklist

specifies criteria for the reporting of complex interventions(23). This article describes the

NROL innovation developed for regional use within an existing healthcare service using the

TIDieR checklist guidance, with the aim of assisting future implementation and replication

efforts.

METHODS

This article describes NROL using the 12-items of the TIDieR checklist(23) and incorporates

guidance from the TIDieR-Telehealth extension(24). Checklist details were developed

iteratively and collaboratively, with input from staff involved in NROL implementation and

evaluation including clinical academics, therapists, managers, researchers and a patient

volunteer. Initially, individual TIDieR checklists describing each NROL group in a high level of

detail were produced. These informed development of NROL staff manual and standard

operating procedure documents. Finally, we abstracted from these documents to develop a

superordinate TIDieR checklist, aimed at describing NROL as a delivery method within an

existing healthcare system for application at a regional level.

Given the extensive interplay between an innovation and context, the settings in which

NROL was implemented are described according to the domains of the updated

Consolidated Framework for Implementation Research (CFIR)(25).

Patient and public involvement

NROL has been improved by partnership, with patients, carers and staff feedback shaping

the intervention described. The NROL patient volunteer co-produced individual TIDieR

checklists.

RESULTS

Item 1: Intervention (innovation) name

NeuroRehabilitation OnLine (NROL)

Item 2: Why

The purpose of NROL is to increase access and opportunity to neurorehabilitation for adult

patients actively receiving community stroke and neurorehabilitation services within an

existing healthcare system. NROL utilises an online delivery platform offering potential

advantages to staff and patients in terms of saving time, energy and travel, and enabling

delivery of more therapy to more people using an existing therapy workforce(14). Readily

accessible technology assistance to learn how to use the required technology and help

overcome barriers is important for optimising inclusivity and meeting the needs of this

patient population(26-28).

NROL delivers group-based therapy. Group rehabilitation formats can help maximise patient

contacts and staffing efficiencies whilst capitalising on the psychosocial benefits of peer

support opportunities(20, 29). Peer-based interventions are viewed positively by those with

acquired brain injury and neurological conditions(30, 31), and peer support elements are

recommended for interventions in stroke(32). Favourable evidence supports the use of

group therapy for the aspects of neurorehabilitation included within NROL(29, 33, 34).

NROL was embedded within an existing system (NHS), rather than being standalone, to

support a sustainable service delivery model that could augment rehabilitation provision.

The innovation aligns with strategic priorities at a local and national level. For example, NHS

England recommend using data and digital technologies, making the most effective use of

resources and establishing collaborative systems, as encompassed by NHS priorities and the

National Stroke Service Model(35, 36).

This TIDieR checklist describes NROL encompassing a regional approach involving four NHS

Trusts in the North-West of England. Cross-organisational working should allow a critical

mass of patients for groups where impairment incidence is relatively low (i.e., dysphasia

group) and a collective use of workforce. Collaborative working should also enable a

community of practice and shared learning.

Item 3: What (materials) and Item 4: What (procedures)

The materials and the procedures for NROL were developed for remote delivery. They will

be described together for the entry, delivery and exit of the 6-week NROL programme and

are depicted in the NROL process chart (Figure 1). A secure 'NROL hub' collaboration

platform (in Microsoft (MS) Teams) was created as a repository for shared NROL

information and resources. Key documents include a 'NROL standard operating procedure

(SOP)' and relevant organisational and governance approvals (e.g., Data Protection Impact

Assessment). NROL was branded with a logo used widely on staff and patient-facing

documentation, and t-shirts worn by the NROL team to increase visibility and establish a

sense of a cohesive team.

Insert Figure 1 here

NROL entry materials and procedures

Patients under the care of stroke and neurorehabilitation services with adequate access to

computer hardware and connectivity are eligible for NROL. The NROL staff manual provides

guidance to help therapy team members identify suitable patients. This living (continually

edited and updated) 'NROL staff manual' provides an overview of NROL, group information

and eligibility criteria, and referral process detail (see Supplementary file). Therapy team

members discuss NROL with their patient and gain verbal consent for their participation in

specified groups. A 'NROL patient leaflet' can be given to patients to complement verbal

information. Eligible, consenting patients are referred to NROL using a 'NROL referral form'.

Therapy team members submit the NROL referral form to the NROL hub. NROL support staff

enter referral form information onto a 'NROL database' (MS Excel), complete timetabling

and document technology assistance.

Prior to programme start, a NROL staff member contacts new patients to administer 'NROL

outcome measures', currently including the EQ-5D-5L for measuring health-related quality

of life(37), the Stroke Self-Efficacy Questionnaire (SSEQ) for measuring self-efficacy(38)

(modified for use with other neurological conditions), and the Patient Specific Functional

Scale (PSFS) for measuring activity performance(39). A NROL technology support staff

member completes a technology set-up with each patient to ensure they have the correct

equipment (e.g., Information technology (IT) hardware and software (MS Teams, email),

connectivity, group-specific requirements) and ability to access NROL, and a suitable home

environment to take part in their specified groups. A 'NROL technology support guide' can

be provided to patients for written guidance. Patients admitted to physical groups are

provided with a 'NROL physical group guide' describing what they need, and how to set up

their environment and webcam to safely perform exercises. Patients receive a 'NROL entry

email' detailing their personalised NROL 6-week programme, which is also sent to the

referrer. Relevant group invites are sent to patients and staff.

NROL delivery materials and procedures

Patients join NROL groups according to their personalised programme. Current groups are

described in the NROL staff manual (Supplementary file). Targeted therapy groups can

include 'physical' groups involving exercise or upper limb rehabilitation led by physiotherapy

and occupational therapy staff, and 'talking' groups including cognitive rehabilitation, living

well, fatigue management or communication therapy, led by occupational therapy,

psychology and speech and language therapy staff. Many of the groups are jointly led by

more than one discipline. Targeted therapy groups encompass interactive, educational, and

practical elements. In addition, community groups are offered to all patients and include an

introduction to NROL (Meet and Greet) and an optional weekly peer support group (Café

NROL). Patients are sent a prompt (i.e., text, email) in the first two weeks of their

programme to assist attendance. Technology assistance is available to patients and staff

during each NROL session. Throughout the programme if a patient is expected in a group

but does not join, the NROL technology support staff member makes contact to offer

assistance.

NROL is delivered online (MS Teams) on existing staff devices. NROL group facilitators and

the NROL technology support staff member require webcam and microphone-enabled

devices and a stable internet connection to deliver NROL groups. A headset can assist clear

communication. For physical groups, a large screen television with a portable microphone is

used for monitoring of patients whilst performing exercises. Session content is developed by

group facilitators and based on evidence-based practice guidelines, and may include

discussions, demonstrations, and shared presentations (MS PowerPoint). Group facilitators

and the NROL technology support staff member use the NROL database during sessions for

accessing patient details, timetabling information and for recording attendance. Access to a

telephone is required in case of adverse events. Group facilitators are required to ensure

clinical notes are entered for all patients in attendance at sessions. Weekly NROL staff

meetings are held to update and discuss NROL delivery and group facilitators meet

separately as required.

Patients require internet connectivity, and a webcam and microphone enabled device

appropriate to their groups (e.g., patients in physical groups require a large screen device),

with MS teams installed. Patients participating in physical groups require equipment for

enabling stability when performing exercises, such as a chair, frame or table. Specific

equipment may be requested for functional task practice and patients are advised to wear

comfortable footwear and clothing and have access to water or a suitable drink. Patients

participating in talking groups are advised to have a pen and paper for making notes or

practicing cognitive exercises. All patients should have a telephone in case of technology

issues or adverse events and should wear their pendant alarm if they have one.

NROL exit materials and procedures

On programme completion, patients receive a 'NROL exit email' with summary information

and resources, which is also sent to the referrer. NROL outcome measures are repeated,

and patients also complete a patient satisfaction survey.

Evaluation

Evaluation occurs alongside NROL delivery and includes analysis of service and patient level

information. Data (quantitative and qualitative) are sourced from the NROL database, and

outcome measures, and patient satisfaction surveys responses, and relevant analysis is

undertaken by the NROL leads. Data is summarised and feedback at multiple clinical levels

(delivery, organisational, regional) and to academic audiences.

Staff training

Meetings are held to discuss NROL evaluation findings. Shared learning sessions are run to

optimise learnings across groups. General NROL information is shared with staff across

trusts by NROL staff, including NROL leads and nominated NROL champions, during team

and service meetings, and via emails and posts on the NROL hub.

New staff are introduced to NROL during orientation. Staff can observe NROL sessions for

experiential learning, particularly when training to become a group facilitator. Staff training

and guidance is available as required and provided remotely by NROL leadership, support or

NROL therapy staff depending on training need. This includes training from technology

support and a 'NROL staff technology guide' to upskill staff and improve confidence.

Item 5: Who provided

NROL is provided by therapy staff from stroke and neurorehabilitation teams at partnered

NHS Trusts. This is supported by a wider NROL team encompassed within a clinical-academic

partnership. NROL uses an existing NHS workforce alongside dedicated NROL roles for the

operational and administrative oversight, and technology support.

Leadership

NROL leadership is managed through a clinical-academic partnership team consisting of

senior NHS Trust management staff, academic project staff, and a dedicated NROL

operational lead.

NROL therapy staff

NROL delivery is provided by NROL therapy staff comprised of existing NHS therapists and

therapy assistants, with student involvement. NROL therapy staff represent physiotherapy,

occupational therapy, speech and language therapy and psychology disciplines at a range of

seniority. Online groups are staffed by a minimum of two group facilitators (therapists

and/or therapy assistants).

Support staff

NROL delivery is facilitated by support staff. A dedicated NROL technology support staff

member provides technology assistance to patients and therapy staff. A NROL administrator

assists with processing referrals and patient correspondence.

Volunteers

Volunteers are previous NROL patients or members of the stroke and neurorehabilitation

community who support the running of community groups under guidance from the NROL

therapy team.

Item 6: How (mode of delivery)

Mode of delivery overlaps with the materials and procedures (see items 3 and 4). In brief,

NROL is a service delivery model providing group-based multidisciplinary rehabilitation as an

adjunct to in-person therapy. Sessions and meetings are conducted remotely (MS Teams).

NROL is embedded within an existing healthcare system (NHS) and makes use of existing

infrastructure (IT hardware and software).

Item 7: Where

For NROL delivery, group facilitators and the NROL technology support staff member attend

group sessions from a private, well-lit, quiet workspace. This involves real-time online co-

delivery from facilitators based at different geographical locations and NHS trusts. A room

with adequate space is required for demonstrating exercises in physical groups. Patients

take part in NROL groups from their own home (or suitable alternative) and also require a

quiet, well-lit, private room that is appropriate to the needs of the group.

Item 8: When and how much

NROL delivery is structured into recurring 6-week 'NROL' blocks. Blocks facilitate integration

of NROL into the existing service as they account for the need for patient flow.

NROL groups run during the working week (Monday to Friday). A NROL introductory group

(Meet and Greet) runs at the start of each block. All further group sessions are scheduled for

60 minutes, with most groups run weekly. The NROL timetable is modified over time

according to patient need and staffing capacity. Sessions can be rescheduled due to

unforeseen circumstances or bank holidays.

The maximum number of patients within a group is determined by the group facilitators and

decided pragmatically to ensure the best experience for patients and staff (see NROL staff

manual for more detail). In the event of too few or too many referrals for a group, deferral

to the next block or the running additional groups are considered.

Item 9: Tailoring

Tailoring is required at patient, group and block levels.

There is flexibility regarding when NROL is integrated within a patient's rehabilitation, and at

present optimal timing is unknown. Patients joining NROL are referred to appropriate

targeted therapy groups and a personalised programme created. Clinical reasoning should

determine how NROL fits with a patient's overall rehabilitation package of care. Patients can

attend more than one NROL block if clinically indicated and providing they remain under the

active care of their stroke or neurorehabilitation team.

The needs of patients in groups will be nuanced over time requiring a responsive approach.

All referrals are screened by group facilitators to ensure session content is tailored for the

current needs. NROL delivery should be considerate that language used is appropriate for

patients with a variety of neurological diagnoses (sudden, intermittent and progressive

conditions) and chronicities.

The structure of blocks, in terms of groups offered and frequency, actively considers patient

and workforce availability. The groups provided in each block is a balance between patient

demand and staff capacity. Some groups run all blocks, whilst others run intermittently.

There is an attempt to minimise groups running over holiday periods. Resourceful use of

available workforce is encouraged and the remote nature of NROL enables staff to deliver

NROL whilst working when from home or requiring work adjustments. Student involvement

can be incorporated to support groups.

Item 10: Modifications

This article describes the NROL innovation developed for regional use. It was modified from

a standalone version (20) and expanded from a project implemented at a single NHS trust

(14). The core components retained include provision of online real-time

neurorehabilitation with technology assistance, incorporating multidisciplinary targeted

therapy and community groups whilst embodying peer support. Adaptations for integration

within an existing healthcare system included inclusion of stroke and other neurological

conditions with varying chronicity, delivering as an adjunct to complement existing

rehabilitation, use of existing workforce and being run as recurring 6-week NROL blocks to

manage patient flow.

NROL continues to be modified over time. New groups are created to meet patient need or

to capitalise on changing staff skill sets. Groups can also be deferred, discontinued or

replaced based on patient and staff feedback.

Item 11: How well planned & Item 12: How well delivered (including fidelity)

Communication, resources and technology assistance are provided to optimise NROL entry

and participation (See *Items 3 & 4)*. Service data are obtained and reviewed to monitor

performance.

A detailed mixed-methods evaluation of NROL within a single trust is available(14).

Implementation and evaluation of the regional NROL programme in the North-West of

England is ongoing. This multi-trust regional innovation has to date delivered 4 x 6-week

NROL blocks, with 1082 patient contacts over 172 sessions provided to the 126 patients

referred to NROL. Withdrawal prior to programme start (17%) and drop-out (15%) are

reasonably low. The median number of patients participating in a NROL block is 32

(range=29-36). Ten different groups are currently offered across a range of therapies, with

typically nine groups running per block. Therapy staffing efficiencies (ratio 0.4) are

demonstrated compared to one-to-one delivery (ratio 1.0) and currently over 40 staff across

the North-West are involved with group delivery. Overall NROL patient attendance is good

(70%), with mechanisms in place to facilitate attendance (e.g., patients receive text/email

prompts in the first two weeks of their programme). Attendance to the weekly peer support

group (Café NROL) is optional however this opportunity is taken up by many patients (58%).

NROL outcome measure evaluation indicates improvements for many patients in health-

related quality of life and activity performance over the time of their NROL programme,

acknowledging that NROL is only one part of their overall rehabilitation. Overall patient

satisfaction is high, with 99% of patients reporting they would recommend NROL to another

person.

Context

The NROL innovation detailed in this article is delivered within a context that can be

described using the domains of inner and outer settings(25).

The inner setting is defined as the four NHS Trusts that are situated within the Lancashire

and South Cumbria Integrated Stroke and Neurorehabilitation Delivery Network that

provide community-based stroke and neurorehabilitation care for the region. The region has

a population of 1.8 million and covers a large geographical footprint with urban and rural

settlements, and ethnic diversity. Deprivation and poor health affect many, with differences

in life expectancy and quality of life varying significantly(40). In some neighbourhoods,

healthy life expectancy is 46.5 years.

Aligning with national strategies and policies, the Lancashire and South Cumbria region has

a vision to work collaboratively to share skills and expertise, data and best practice across

Trusts(41). A challenge is that Trusts have varied service remits (stroke, neurological or both)

and infrastructure (e.g. physical, staffing levels, technology systems, governance processes).

With regards to the outer setting, the impetus for starting NROL was the global pandemic,

which also influenced sociocultural values of staff and patients to increase the worthiness

and openness to use of remote technology. NROL also aligned with wider policy and

strategies(42-44), benefitted from an already established clinical-academic partnership

between the NHS and a university, and had funding from an external entity(45).

DISCUSSION

NROL utilises an online delivery platform, with dedicated technology assistance, to provide multidisciplinary real-time group therapy to patients actively receiving community stroke and neurorehabilitation. Telerehabilitation is recognised as having a vital role to play in future healthcare delivery(46) but as yet there are limited details of how to do this(47). To address this need, the TIDieR checklist is used as a structured method to provide a comprehensive description of a regional NROL innovation embedded within a healthcare system. The actual process of completing the TIDieR was time intensive but did provide the impetus for the team to clearly describe NROL, agreeing the core components. Consideration was given to the necessary balance of information to ensure comprehensive detail but attempting not to overwhelm. Further documentation is available as a supplementary file and by contacting the corresponding author. This description details an exemplar of telerehabilitation using NROL delivered in one region, and it is acknowledged that NROL will likely need adapting for other areas given the significant interaction between an innovation and its context. This article should facilitate other healthcare settings to adapt and implement NROL for their context.

Optimal adaptation of an innovation requires an understanding of the core components that cannot be changed versus the adaptable periphery that can be changed(25, 48, 49). The core components of NROL include provision of online real-time neurorehabilitation with technology assistance, incorporating multidisciplinary targeted therapy and community groups whilst embodying peer support. It is proposed that these core components should be consistently implemented for NROL but that the processes to achieve them are adapted

to fit local conditions. Examples include the use of MS Teams to deliver groups but other

online platforms are available; use of recurring 6-week NROL blocks but other timings may

suit other services; the number and types of targeted therapy groups will need to reflect

workforce capability and capacity. It is known innovations that have adaptability are more

likely to be used in clinical practice(50, 51). Further examination of the adaptive

components of NROL will help discern how it can be upscaled for use in a variety of

contexts.

Context is everything(52). A limitation of the TIDieR checklist is that it does not include an

item on context. Arguably to understand the innovation fully and guide future adaptations

an understanding of the context is required. This is because innovations are inextricably

linked to the context in which they are delivered, and achieving a good fit between these is

important to ensure the innovation works as intended(53). In this article, context is

deliberatively reported to help situate the innovation. NROL did take resources, time and

effort to implement as a regional initiative and details on the implementation will be

reported elsewhere. The need for resources to enable implementation is not

unexpected(49, 54, 55) and current healthcare systems are often not set-up to facilitate this

upfront effort. Influential contextual factors included leadership buy-in and commitment, a

clinical-academic partnership and fit with the local and broader strategic landscape.

Inevitably NROL will continue to evolve. The TIDieR checklist does allow for reporting on

modifications and tailoring. To date, the checklist has been primarily used for reporting

interventions in trials(56, 57) and there are limited examples of its use for adapted

interventions over time. This article documents current NROL delivery and captures its

retrospective modifications and tailoring. Going forward, transparent reporting of new

iterations of NROL together with descriptions of their context should be undertaken to aid

comparisons.

CONCLUSION

A hybrid approach incorporating telerehabilitation, to complement in-person therapy, is

required for a future-proof service that follows policy and guidelines. This comprehensive

description of a regional NROL innovation gives an example of successful implementation

within an existing healthcare system. It provides a platform for others to reduce duplication

of effort and help facilitate the use of telerehabilitation in clinical practice.   Adapted

versions of NROL are expected when implementing in different contexts. Transparent

reporting and continuous evaluation alongside NROL implementation are encouraged and

will ensure maximal impact for neurorehabilitation delivery.

 FIGURE LEGEND

Figure 1

NROL process chart. The key NROL procedures and staff and patient materials are outlined

over four stages; Referral, Entry, Delivery and Exit, with the staff involved indicated. This

chart can be used by the wider stroke and neurorehabilitation teams to help plan and

organise participation for suitable patients and understand the resources available and

needed from both a staff and patient perspective to support this. MS, Microsoft; IT,

information technology.

Availability of data/ resources

Digital copies of NROL materials are available on request by contacting Professor Louise

Connell (laconnell@uclan.ac.uk).

Funding

This work was supported by a generous donation from the charity SameYou (Charity

number 1170102). The funders had no role in writing the manuscript.

Competing interests

The authors declare that they have no competing interests.

Author contributions

SA and LC conceived the study. All authors (SA, LC, NW, PB, RP) were involved in the

planning and conduct of the work described in the paper. SA, LC and NW drafted the initial

manuscript. All authors were involved in revising the manuscript. All authors have given final

approval of the version to be published and agree to be accountable for all aspects of the

work.

Acknowledgements:

We would like to express our gratitude to the individuals and services that contribute to

NROL, and have contributed to this work, from East Lancashire Hospitals NHS Trust,

University Hospitals of Morecambe Bay NHS Foundation Trust, Blackpool Teaching Hospitals

NHS Foundation Trust and Lancashire and South Cumbria NHS Foundation Trust. Special

448 thanks go to Ian Grimshaw for providing technological support for patients and staff, as well

449 as supporting administrative and organisational operations. We extend our appreciation to

450 the UCL N-ROL team for their sharing of expertise and experience of online delivery and

451 valuable discussions. Many thanks to the SameYou team for their generous fundraising and

452 support which have made NROL possible.

REFERENCES

REFERENCES

1.   Lohse KR, Lang CE, Boyd LA. Is more better? Using metadata to explore dose-response
relationships in stroke rehabilitation. Stroke. 2014;45(7):2053-8.10.1161/strokeaha.114.004695
2.   French B, Thomas LH, Leathley MJ, Sutton CJ, McAdam J, Forster A, et al. Repetitive Task
Training for Improving Functional Ability After Stroke. Stroke. 2009;40(4):e98-
e9.10.1161/strokeaha.108.519553
3.   Schneider EJ, Lannin NA, Ada L, Schmidt J. Increasing the amount of usual rehabilitation
improves activity after stroke: a systematic review. Journal of Physiotherapy. 2016;62(4):182-
7.https://doi.org/10.1016/j.jphys.2016.08.006
4.   Kwakkel G, Van Peppen R, Wagenaar RC, Wood Dauphinee S, Richards C, Ashburn A, et al.
Effects of Augmented Exercise Therapy Time After Stroke. Stroke. 2004;35(11):2529-
39.10.1161/01.str.0000143153.76460.7d
5.   Sentinel Stroke National Audit Programme. The Road to Recovery. The Ninth SSNAP Annual
Report. [Internet] 2022. Available from:
https://www.strokeaudit.org/Documents/National/Clinical/Apr2021Mar2022/Apr2021Mar2022-
AnnualReport.aspx [accessed 30 March 2023]
6.   Cramer SC, Dodakian L, Le V, McKenzie A, See J, Augsburger R, et al. A Feasibility Study of
Expanded Home-Based Telerehabilitation After Stroke. Front Neurol.
2020;11:611453.10.3389/fneur.2020.611453
7.   Laver KE, Adey-Wakeling Z, Crotty M, Lannin NA, George S, Sherrington C. Telerehabilitation
services for stroke. Cochrane Database Syst Rev.
2020;1(1):Cd010255.10.1002/14651858.CD010255.pub3
8.   Stroke Association. A New Era for Stroke. [Internet] 2016. Available from:
https://www.stroke.org.uk/sites/default/files/anefs_report_web.pdf [accessed 30 March 2023]
9.   Parrott D, Ibarra S. Randomized Controlled Trial Comparing a Telemedicine Brain Injury
Coping Skills (BICS) group intervention to traditional in-person BICS for Brain Injury Patients and
Caregivers. Archives of Physical Medicine and Rehabilitation.
2021;102(10):e15.https://doi.org/10.1016/j.apmr.2021.07.434
10.   Khan F, Amatya B, Kesselring J, Galea M. Telerehabilitation for persons with multiple
sclerosis. Cochrane Database Syst Rev. 2015(4):CD010508.10.1002/14651858.CD010508.pub2
11.   Rietdijk R, Power E, Attard M, Togher L. Acceptability of telehealth-delivered rehabilitation:
Experiences and perspectives of people with traumatic brain injury and their carers. Journal of
Telemedicine and Telecare. 2022;28(2):122-34.10.1177/1357633x20923824
12.   Hargreaves DL, D. Stroke, GIRFT Programme National Specialty Report. [Internet] 2022.
Available from: https://gmnisdn.org.uk/wp-content/uploads/2022/04/Stroke-GiRFT-report-Apr-
22.pdf) [accessed
13.   Chartered Society of Physiotherapy. National Evaluation of Remote Physiotherapy Services:
the findings. [Internet] 2022. Available from: https://www.csp.org.uk/professional-
clinical/professional-guidance/remote-consultations/national-evaluation-remote-0 [accessed 30
March 2023]
14.   Ackerley S, Wilson N, Boland P, Reed J, Connell L. Implementation of neurological group-
based telerehabilitation within existing healthcare: a mixed methods evaluation. BMC Health
Services Research journal. Under review.
15.   Prvu Bettger J, Resnik LJ. Telerehabilitation in the Age of COVID-19: An Opportunity for
Learning Health System Research. Phys Ther. 2020;100(11):1913-6.10.1093/ptj/pzaa151
16.   Wosik J, Fudim M, Cameron B, Gellad ZF, Cho A, Phinney D, et al. Telehealth transformation:
COVID-19 and the rise of virtual care. Journal of the American Medical Informatics Association.
2020;27(6):957-62.10.1093/jamia/ocaa067

17.   Taylor A, Caffery L, Gesesew H, King A, Bassal A-r, Ford K, et al. How Australian Health Care
Services Adapted to Telehealth During the COVID-19 Pandemic: A Survey of Telehealth Professionals.
Frontiers in Public Health. 2021;9.10.3389/fpubh.2021.648009
18.   Ross MH, Nelson M, Parravicini V, Weight M, Tyrrell R, Hartley N, et al. Staff perspectives on
the key elements to successful rapid uptake of telerehabilitation in medium-sized public hospital
physiotherapy departments. Physiotherapy Research International. 2022.10.1002/pri.1991
19.   Yang CL, Waterson S, Eng JJ. Implementation and Evaluation of the Virtual Graded Repetitive
Arm Supplementary Program (GRASP) for Individuals With Stroke During the COVID-19 Pandemic
and Beyond. Phys Ther. 2021;101(6).10.1093/ptj/pzab083
20.   Beare B, Doogan CE, Douglass-Kirk P, Leff AP, Ward N. Neuro-Rehabilitation OnLine (N-ROL):
description and evaluation of a group-based telerehabilitation programme for acquired brain injury.
Journal of Neurology, Neurosurgery and Psychiatry. 2021;92(12):1354.10.1136/jnnp-2021-326809
21.   Buckingham S, Anil K, Demain S, Gunn H, Jones RB, Kent B, et al. Telerehabilitation for
people with physical disabilities and movement impairment: development and evaluation of an
online toolkit for practitioners and patients. Disability and Rehabilitation. 2022:1-
8.10.1080/09638288.2022.2074549
22.   Royal College of Physicians. National Clinical Guideline for Stroke for the United Kingdom
and Ireland. [Internet] 2022. Available from: https://uploads-
ssl.webflow.com/62c3d8070eda8516a8ba9475/6380a7e6a9a3db745a8e8315_2023%20National%2
0Clinical%20Guideline%20for%20Stroke%20consultation%20document.pdf [accessed 30 March
2023]
23.   Hoffmann TC, Glasziou PP, Boutron I, Milne R, Perera R, Moher D, et al. Better reporting of
interventions: template for intervention description and replication (TIDieR) checklist and guide.
BMJ. 2014;348(mar07 3):g1687-g.10.1136/bmj.g1687
24.   Rhon DI, Fritz JM, Kerns RD, McGeary DD, Coleman BC, Farrokhi S, et al. TIDieR-telehealth:
precision in reporting of telehealth interventions used in clinical trials - unique considerations for the
Template for the Intervention Description and Replication (TIDieR) checklist. BMC Medical Research
Methodology. 2022;22(1).10.1186/s12874-022-01640-7
25.   Damschroder LJ, Reardon CM, Widerquist MAO, Lowery J. The updated Consolidated
Framework for Implementation Research based on user feedback. Implementation Science.
2022;17(1).10.1186/s13012-022-01245-0
26.   Mace RA, Mattos MK, Vranceanu A-M. Older adults can use technology: why healthcare
professionals must overcome ageism in digital health. Translational Behavioral Medicine.
2022;12(12):1102-5.10.1093/tbm/ibac070
27.   Tsai H-YS, Shillair R, Cotten SR. Social Support and "Playing Around". Journal of Applied
Gerontology. 2017;36(1):29-55.10.1177/0733464815609440
28.   Doorley JD, Mace RA, Popok PJ, Grunberg VA, Ragnhildstveit A, Vranceanu AM. Feasibility
Randomized Controlled Trial of a Mind-Body Activity Program for Older Adults With Chronic Pain and
Cognitive Decline: The Virtual "Active Brains" Study. Gerontologist. 2022;62(7):1082-
94.10.1093/geront/gnab135
29.   Miller MJ, Pak SS, Keller DR, Barnes DE. Evaluation of Pragmatic Telehealth Physical Therapy
Implementation During the COVID-19 Pandemic. Phys Ther. 2021;101(1).10.1093/ptj/pzaa193
30.   Aterman S, Ghahari S, Kessler D. Characteristics of peer-based interventions for individuals
with neurological conditions: a scoping review. Disability and Rehabilitation. 2022:1-
32.10.1080/09638288.2022.2028911
31.   Hughes R, Fleming P, Henshall L. Peer support groups after acquired brain injury: a
systematic review. Brain Injury. 2020;34(7):847-56.10.1080/02699052.2020.1762002
32.   Clark E, Maccrosain A, Ward NS, Jones F. The key features and role of peer support within
group self-management interventions for stroke? A systematic review. Disability and Rehabilitation.
2020;42(3):307-16.10.1080/09638288.2018.1498544

33.   Fleming V, Brownsett S, Krason A, Maegli MA, Coley-Fisher H, Ong Y-H, et al. Efficacy of
spoken word comprehension therapy in patients with chronic aphasia: a cross-over randomised
controlled trial with structural imaging. Journal of Neurology, Neurosurgery & Psychiatry.
2021;92(4):418-24.10.1136/jnnp-2020-324256
34.   Ward NS, Brander F, Kelly K. Intensive upper limb neurorehabilitation in chronic stroke:
outcomes from the Queen Square programme. Journal of Neurology, Neurosurgery & Psychiatry.
2019;90(5):498-506.10.1136/jnnp-2018-319954
35.   NHS England. National Stroke Service Model: Integrated Stroke Delivery Networks. [Internet]
2021. Available from: https://www.england.nhs.uk/wp-content/uploads/2021/05/stroke-service-
model-may-2021.pdf [accessed 30 March 2023]
36.   Stroke Association. Shaping stroke research to rebuild lives. [Internet] 2021. Available from:
https://www.stroke.org.uk/sites/default/files/research/stroke_priority_setting_partnership_full_rep
ort.pdf [accessed 30 March 2023]
37.   EuroQol Research Foundation. EQ-5D-3L User Guide. [Internet] 2018. Available from:
https://euroqol.org/publications/user-guides [accessed 30 March 2023]
38.   Riazi A, Aspden T, Jones F. Stroke Self-efficacy Questionnaire: a Rasch-refined measure of
confidence post stroke. J Rehabil Med. 2014;46(5):406-12.10.2340/16501977-1789
39.   Stratford P, Gill C, Westaway M, Binkley J. Assessing Disability and Change on Individual
Patients: A Report of a Patient Specific Measure. Physiotherapy Canada. 1995;47(4):258-
63.10.3138/ptc.47.4.258
40.   Lancashire Independent Economic Review. Deep Dive: Health,Wealth & Wellbeing.
[Internet] 2021. Available from: https://www.lancashireier.org/wp-
content/uploads/2021/12/LIER_Health_Wealth_and_Wellbeing_2021_v1.pdf [accessed 30 March
2023]
41.   Lancashire and South Cumbria Provider Collaborative. Our vision. [Internet] 2022. Available
from: https://lscprovidercollaborative.nhs.uk/about-us/our-vision [accessed 16 February 2023]
42.   World Health A. Digital health. Geneva: World Health Organization, 2018 2018.
https://apps.who.int/iris/handle/10665/279505
43.   World Health Assembly. eHealth. Geneva: World Health Organization, 2005 2005.
https://apps.who.int/iris/handle/10665/20378
44.   NHS England and NHS Inprovement. 2022/23 priorities and operational planning guidance.
[Internet] 2022. Available from: https://www.england.nhs.uk/wp-
content/uploads/2022/02/20211223-B1160-2022-23-priorities-and-operational-planning-guidance-
v3.2.pdf [accessed 30 March 2023]
45.   SameYou. NROL. [Internet] 2022. Available from: https://www.sameyou.org/nrol [accessed
16 February 2023]
46.   NHS Health Education England. The Topol Review. [Internet] 2023. Available from:
https://topol.hee.nhs.uk/the-topol-review/ [accessed 16 February 2023]
47.   English C, Ceravolo MG, Dorsch S, Drummond A, Gandhi DB, Halliday Green J, et al.
Telehealth for rehabilitation and recovery after stroke: State of the evidence and future directions.
International Journal of Stroke. 2022;17(5):487-93.10.1177/17474930211062480
48.   Haynes A, Brennan S, Redman S, Williamson A, Gallego G, Butow P. Figuring out fidelity: a
worked example of the methods used to identify, critique and revise the essential elements of a
contextualised intervention in health policy agencies. Implementation Science.
2015;11(1).10.1186/s13012-016-0378-6
49.   Greenhalgh T, Papoutsi C. Spreading and scaling up innovation and improvement. BMJ.
2019:l2068.10.1136/bmj.l2068
50.   Rogers E. Diffusion of innovations: 5th Edition. New York: Free Press; 2003.
51.   Gustafson DH, Sainfort F, Eichler M, Adams L, Bisognano M, Steudel H. Developing and
Testing a Model to Predict Outcomes of Organizational Change. Health Services Research.
2003;38(2):751-76.10.1111/1475-6773.00143

52.   Bate P. The Health Foundation: Perspectives on context. [Internet] 2014. Available from:
https://www.health.org.uk/sites/default/files/PerspectivesOnContextBateContextIsEverything.pdf
[accessed 30 March 2023]
53.   Hawe P. Lessons from Complex Interventions to Improve Health. Annual Review of Public
Health. 2015;36(1):307-23.10.1146/annurev-publhealth-031912-114421
54.   Standing C, Standing S, McDermott M-L, Gururajan R, Kiani Mavi R. The Paradoxes of
Telehealth: a Review of the Literature 2000-2015. Systems Research and Behavioral Science.
2018;35(1):90-101.10.1002/sres.2442
55.   Signal N, Martin T, Leys A, Maloney R, Bright F. Implementation of telerehabilitation in
response to COVID-19: Lessons learnt from neurorehabilitation clinical practice and education. New
Zealand Journal of Physiotherapy. 2020;48:117-26.10.15619/NZJP/48.3.03
56.   Tume LN, Blackwood B, McAuley DF, Morris K, Peters MJ, Jordan J, et al. Using the TIDieR
checklist to describe the intervention of the Sedation and Weaning in Children(SANDWICH) trial.
Nursing in Critical Care. 2022.10.1111/nicc.12810
57.   Carlsson H, Rosén B, Björkman A, Pessah-Rasmussen H, Brogårdh C. SENSory re-learning of
the UPPer limb (SENSUPP) after stroke: development and description of a novel intervention using
the TIDieR checklist. Trials. 2021;22(1).10.1186/s13063-021-05375-6