**Title**

NeuroRehabilitation OnLine (NROL): Description of a multidisciplinary group

telerehabilitation innovation for stroke and neurological conditions using the TIDieR

checklist

**Authors**

Suzanne Ackerley[1,2] (sackerley@uclan.ac.uk), Neil Wilson[1] (nwilson9@uclan.ac.uk), Paul

Boland[1] (pboland2@uclan.ac.uk), Rosemary Peel[3] (rosemary.peel@mbht.nhs.uk , Louise

Connell[1,2] (laconnell@uclan.ac.uk).

[1]University of Central Lancashire, Preston, Lancashire, UK

[2] East Lancashire Hospitals NHS Trust, Burnley, Lancashire, UK

[3] University Hospitals of Morecambe Bay NHS Foundation Trust

**Corresponding author**

Louise Connell, School of Sport and Health Sciences, University of Central Lancashire, UK. E-

mail: laconnell@uclan.ac.uk

**Abstract**

Background

Providing recommended amounts of rehabilitation for stroke and neurological patients is

challenging. Telerehabilitation is viable for delivering rehabilitation and an acceptable adjunct

to in-person therapy. NeuroRehabilitation OnLine (NROL) was developed as a pilot and

subsequently operationalised as a regional innovation embedded across four National Health

Service (NHS) Trusts.

Objective

To describe the NROL innovation to assist future implementation and replication efforts.

Methods

The Template for Intervention Description and Replication (TIDieR) checklist, with guidance

from the TIDieR-Telehealth extension, was used to describe NROL. The description was

developed collaboratively by clinical academics, therapists, managers, and researchers.

Updated Consolidated Framework for Implementation Research domains were used to

describe the context in which the innovation was delivered.

Results

NROL delivers online group-based real-time neurorehabilitation with technology assistance

provided, incorporating multidisciplinary targeted therapy and peer support. Procedures,

materials and structure are detailed to demonstrate how NROL is embedded within a

healthcare system. NROL uses existing NHS therapy workforce alongside dedicated NROL

roles, including an essential technology support role. Selection of NROL groups is dependent

on patient need. The NROL innovation is modified over time in response to patient and staff

feedback. NROL described here is situated within a regional stroke and neurorehabilitation

network in North-West England, aligns with local and national strategies, and capitalises on

an existing clinical-academic partnership.

Conclusion

This comprehensive description of a regional NROL innovation, and clarification of core

components, should facilitate other healthcare settings to adapt and implement NROL for

their context. Continuous evaluation alongside implementation will ensure maximal impact

for neurorehabilitation.

**Background**

Despite a wealth of evidence that greater amounts of rehabilitation can improve outcomes (1-4) providing stroke and brain injury patients with the recommended amount of therapy remains a clinical and workforce challenge (5). Increasing access to and opportunity for therapy is a critical step to addressing shortfalls in therapy intensity and ideally needs to be feasible with limited workforce resources.

Telerehabilitation, the provision of rehabilitation remotely via telecommunication devices, has developed as a viable complement to in-person therapy and offers one possible solution to help mitigate challenges. It can deliver conventional in-person therapies online with equivalent outcomes (5-10) and with similar attendance levels and acceptability to patients (11). Patients and staff report advantages in terms of saving time, energy and travel (12-14). Although not a new approach, rapid expanded adoption of telerehabilitation in routine practice was spurred by the COVID-19 pandemic (15-18), with further encouraging data collected during this time (14, 19-21). In the UK and Ireland, clinical guidelines for stroke rehabilitation now recommend the provision of therapist monitored, patient-centred remote rehabilitation programmes alongside conventional in-person therapy (22).

A group-based real-time telerehabilitation innovation for patients with acquired brain injury was developed and piloted in London, UK, entitled NROL (NeuroRehabilitation OnLine). This standalone version of NROL demonstrated positive impacts on patient-reported outcomes (20). NROL was subsequently adapted and operationalised within the UK National Health Service (NHS) by embedding a recurring six-week multidisciplinary group-based

telerehabilitation innovation into an existing service. Implementation with concurrent

evaluation at a single NHS Trust level yielded positive results, demonstrating favourable

patient and staff opinion, sustainable staffing and resource efficiencies with good clinical

outcomes for patients (14). NROL was then expanded into a regional innovation involving four

NHS Trusts in the North-West of England aligning to the NHS new Integrated Care System

structure to work collaboratively across regions.  It remains part of a hybrid approach to

service delivery to complement in-person rehabilitation. NROL is acknowledged as an

exemplar initiative for delivering remote rehabilitation (22).

Despite the successful development of NROL, as yet it has only been described in its

standalone version (20) with no descriptive detail available for the current regional, multi-site

model. The Template for Intervention Description and Replication (TIDieR) checklist specifies

criteria for the reporting of complex interventions (23). This article describes the NROL

innovation developed for regional use within an existing healthcare service using the TIDieR

checklist guidance, with the aim of assisting future implementation and replication efforts.

**Methods**

This article describes NROL using the 12-items of the TIDieR checklist (23) and incorporates

guidance from the TIDieR-Telehealth extension, providing telehealth-specific criteria for

effective description of telehealth interventions (24). Checklist details were developed

iteratively and collaboratively, with input from staff involved in NROL implementation and

evaluation including clinical academics, therapists, managers, and researchers. Initially,

individual TIDieR checklists describing each NROL group in a high level of detail were

produced. These informed development of NROL staff manual and standard operating

procedure documents. Finally, we abstracted from these documents to develop a

superordinate TIDieR checklist, aimed at describing NROL as a delivery method within an

existing healthcare system for application at a regional level.

Given the extensive interplay between an innovation and context, the settings in which NROL

was implemented are described according to the domains of the updated Consolidated

Framework for Implementation Research (CFIR) (25).

**Results**

*Item 1: Intervention (innovation) name*

NeuroRehabilitation OnLine (NROL)

*Item 2: Why*

The purpose of NROL is to increase access and opportunity to neurorehabilitation for adult

patients actively receiving community stroke and neurorehabilitation services within an

existing healthcare system. NROL utilises an online delivery platform offering potential

advantages to staff and patients in terms of saving time, energy and travel, and enabling

delivery of more therapy to more people using an existing therapy workforce (14). Readily

accessible technology assistance to learn how to use the required technology and help

overcome barriers is important for optimising inclusivity and meeting the needs of this patient

population (26-28).

NROL delivers group-based therapy. Group rehabilitation formats can help maximise patient

contacts and staffing efficiencies whilst capitalising on the psychosocial benefits of peer

support opportunities (20, 29). Peer-based interventions are viewed positively by those with

acquired brain injury and neurological conditions (30, 31), and peer support elements are

recommended for interventions in stroke (32). Favourable evidence supports the use of group

therapy for the aspects of neurorehabilitation included within NROL (29, 33, 34).

NROL was embedded within an existing system (NHS), rather than being standalone, to

support a sustainable service delivery model that could augment rehabilitation provision. The

innovation aligns with strategic priorities at a local and national level. For example, NHS

England recommend using data and digital technologies, making the most effective use of

resources and establishing collaborative systems, as encompassed by NHS priorities and the

National Stroke Service Model (35, 36).

This TIDieR checklist describes NROL encompassing a regional approach involving four NHS

Trusts in the North-West of England. Cross-organisational working should allow a critical mass

of patients for groups where impairment incidence is relatively low (i.e., dysphasia group) and

a collective use of workforce.  Collaborative working should also enable a community of

practice and shared learning.

***Item 3: What (materials) and Item 4: What (procedures)***

The materials and the procedures for NROL were developed for remote delivery. They will be

described together for the entry, delivery and exit of the 6-week NROL programme and are

depicted in Figure 1. A secure 'NROL hub' collaboration platform (in MS Teams) was created

as a repository for shared NROL information and resources. Key documents include a 'NROL

standard operating procedure (SOP)' and relevant organisational and governance approvals

(e.g., Data Protection Impact Assessment). NROL was branded with a logo used widely on staff

and patient-facing documentation, and t-shirts worn by NROL group facilitators and

technology support to increase visibility and establish a sense of a cohesive team.

**Figure 1**

| | Procedure | Staff Material | Patient Material |
|---|---|---|---|

**Referral: Stroke & Neurorehabilitation team**

**Patient identified for NROL**
Therapist identifies patient need and ensures they meet eligibility criteria for NROL and relevant groups

NROL staff manual

NROL patient leaflet

**Consent gained**
Therapist discusses NROL and suitable groups with the patient, confirms IT hardware and internet connectivity access, and gains verbal consent

NROL referral form

**Referral submission**
Therapists completes referral form including group-specific outcome measures and uploads to NROL hub

NROL hub (Collaboration platform)

IT hardware webcam/microphone enabled, with internet connectivity

**Entry: NROL lead & support staff**

**Referral processing**
NROL support enter referral information onto database and perform group timetabling with lead oversight

NROL database

NROL outcome measures

IT software (MS Teams, Email)

NROL technology support guide

**Patient set-up**
Completion of outcome measures and technology set-up

NROL physical group guide

NROL entry email

**Patient programmes**
Patients emailed their NROL programmes and relevant group invites emailed to patients and staff

NROL group invites

**Delivery: NROL therapy & support staff**

Clinical notes

**Meet and Greet**

Talking

Physical

Community

Weekly MDT Meetings

NROL regional SOP

Telephone/pendant alarm

IT hardware, software, internet connectivity, patient data system, group-specific equipment & telephone

Physical: environmental & therapy equipment as detailed by group facilitators

NROL staff technology guide

NROL database

Talking: Pen and paper

**Exit: NROL Support staff**

**Patient exit**
Completion of outcome measure & patient satisfaction survey
Patients sent exit email with group specific summary information and resources

NROL outcome measures

NROL exit email

NROL patient satisfaction survey

Group specific resources

**NROL entry materials and procedures**

Patients under the care of stroke and neurorehabilitation services with adequate access to

computer hardware and connectivity are eligible for entry to NROL. The NROL staff manual

provides guidance to help therapy team members identify suitable patients. This living

(continually edited and updated) 'NROL staff manual' provides an overview of NROL, group

information and eligibility criteria, and referral process detail (see Supplementary file).

Therapy team members discuss NROL with their patient and gain verbal consent for their

participation in specified groups. A 'NROL patient leaflet' can be given to patients to

complement verbal information. Eligible, consenting patients are referred to NROL using a

'NROL referral form'.

Therapy team members submit the NROL referral form to the NROL hub. NROL support staff

enter referral form information onto a 'NROL database' (MS Excel), a document used to

collate patient demographic and service information, complete timetabling and document

technology assistance.

Prior to programme start, a NROL staff member contacts new patients to administer 'NROL

outcome measures', currently including the EQ-5D-5L for measuring health-related quality of

life (37), the Stroke Self-Efficacy Questionnaire (SSEQ) for measuring self-efficacy (38)

(modified for use with other neurological conditions), and the Patient Specific Functional

Scale (PSFS) for measuring activity performance (39). A NROL technology support staff

member completes a technology set-up with each patient to ensure they have the correct

equipment (e.g., IT hardware and software (MS Teams, email), connectivity, group-specific

requirements) and a suitable home environment to take part in their specified groups. A

'NROL technology support guide' can be provided to patients for written guidance. Patients

admitted to physical groups are provided with a 'NROL physical group guide' describing what

they need, and how to set up their environment and webcam to safely perform exercises.

Patients receive a 'NROL entry email' detailing their personalised NROL 6-week programme,

which is also sent to the referrer. Relevant group invites are sent to patients and staff.

**NROL delivery materials and procedures**

Patients join NROL groups according to their personalised programme. Current groups are

described in the NROL staff manual (Supplementary file). Targeted therapy groups can include

'physical' groups involving exercise or upper limb rehabilitation led by physiotherapy and

occupational therapy staff, and 'talking' groups including cognitive rehabilitation, living well,

fatigue management or communication therapy, led by occupational therapy, psychology and

speech and language therapy staff. Many of the groups are jointly led by more than one

discipline. Targeted therapy groups encompass interactive, educational, and practical

elements. In addition, community groups are offered to all patients and include an

introduction to NROL (Meet and Greet) and an optional weekly peer support group (Café

NROL). Patients are sent a prompt (i.e., text, email) in the first two weeks of their programme

to assist attendance. Technology assistance is available to patients and staff during each NROL

session. Throughout the programme if a patient is expected in a group but does not join, the

NROL technology support staff member makes contact to offer assistance.

NROL is delivered online (MS Teams video conferencing application) on existing staff devices.

NROL group facilitators and the NROL technology support staff member require webcam and

microphone-enabled devices and a stable (preferably wired) internet connection to deliver

NROL groups. A headset can assist clear communication. For physical groups, a large screen

television with a portable microphone is used for monitoring of patients whilst performing

exercises. Session content is developed by group facilitators and based on evidence-based

practice guidelines, and may include discussions, demonstrations, and shared presentations

(MS Powerpoint). Group facilitators and the NROL technology support staff member use the

NROL database during sessions for accessing patient details, timetabling information and for

recording attendance. Access to a telephone is required in case of adverse events. Group

facilitators are required to ensure clinical notes are entered for all patients in attendance at

sessions. Weekly NROL staff meetings are held to update and discuss NROL delivery and group

facilitators meet separately as required.

Patients require internet connectivity, and a webcam and microphone enabled device

appropriate to their groups (e.g., patients in physical groups require a large screen device),

with MS teams installed. Patients participating in physical groups require equipment for

enabling stability when performing exercises, such as a chair, frame or table. Specific

equipment may be requested for functional task practice and patients are advised to wear

comfortable footwear and clothing and have access to water or a suitable drink. Patients

participating in talking groups are advised to have a pen and paper for making notes or

practicing cognitive exercises. All patients should have a telephone in case of technology

issues or adverse events and should wear their pendant alarm if they have one.

**NROL exit materials and procedures**

On programme completion, patients receive a 'NROL exit email' with summary information

and resources, which is also sent to the referrer. NROL outcome measures are repeated, and

patients also complete a patient satisfaction survey.

**Evaluation**

Evaluation occurs alongside NROL delivery and includes analysis of service and patient level

information. Data (quantitative and qualitative) are sourced from the NROL database, and

outcome measures, and patient satisfaction surveys responses, and relevant analysis is

undertaken by the NROL leads. Data is summarised and feedback at multiple clinical levels

(delivery, organisational, regional) and to academic audiences.

**Staff training**

Meetings are held to discuss NROL evaluation findings. Shared learning sessions are run to

optimise learnings across groups. General NROL information is shared with staff across trusts

by NROL staff, including NROL leads and nominated NROL champions, during team and

service meetings, and via emails and posts on the NROL hub.

New staff are introduced to NROL during orientation. Staff can observe NROL sessions for

experiential learning, particularly when training to become a group facilitator. Staff training

and guidance is available as required and provided remotely by NROL leadership, support or

NROL therapy staff depending on training need. This includes training from technology

support and a 'NROL staff technology guide' to upskill staff and improve confidence.

*Item 5: Who provided*

NROL is provided by therapy staff from stroke and neurorehabilitation teams at partnered

NHS Trusts. This is supported by a wider NROL team encompassed within a clinical-academic

partnership. NROL uses an existing NHS workforce alongside dedicated NROL roles for the

operational and administrative oversight, and technology support.

**Leadership**

NROL leadership is managed through a clinical-academic partnership team consisting of

senior NHS Trust management staff, academic project staff, and a dedicated NROL

operational lead.

**NROL therapy staff**

NROL delivery is provided by NROL therapy staff comprised of existing NHS therapists, therapy

assistants and students. NROL therapy staff represent physiotherapy, occupational therapy,

speech and language therapy and psychology disciplines at a range of seniority. Online groups

are staffed by a minimum of two group facilitators (therapists and/or therapy assistants).

**Support staff**

NROL delivery is facilitated by support staff. A dedicated NROL technology support staff

member provides technology assistance to patients and therapy staff. A NROL administrator

assists with processing referrals and patient correspondence.

**Volunteers**

Volunteers are previous NROL patients or members of the stroke and neurorehabilitation

community who support the running of community groups under guidance from the NROL

therapy team.

**Item 6: How (mode of delivery)**

Mode of delivery overlaps with the materials and procedures (see items 3 and 4). In brief, NROL is a service delivery model providing group-based rehabilitation as an adjunct to in-person therapy. Sessions and meetings are conducted remotely (MS Teams). NROL is embedded within an existing healthcare system (NHS) and makes use of existing infrastructure (IT hardware and software).

**Item 7: Where**

For NROL delivery, group facilitators and the NROL technology support staff member attend group sessions from a private, well-lit, quiet workspace. This involves real-time online co-delivery from facilitators based at different geographical locations and NHS trusts. A room with adequate space is required for demonstrating exercises in physical groups. Patients take part in NROL groups from their own home (or suitable alternative) and also require a quiet, well-lit, private room that is appropriate to the needs of the group.

**Item 8: When and how much**

NROL delivery is structured into recurring 6-week 'NROL' blocks. Blocks facilitate integration of NROL into the existing service as they account for the need for patient flow.

NROL groups run during the working week (Monday to Friday). A NROL introductory group (Meet and Greet) runs at the start of each block. All further group sessions are scheduled for 60 minutes, with most groups run weekly. The NROL timetable is modified over time

according to patient need and staffing capacity. Sessions can be rescheduled due to

unforeseen circumstances or bank holidays.

The maximum number of patients within a group is determined by the group facilitators and

decided pragmatically to ensure the best experience for patients and staff (see NROL staff

manual for more detail).  In the event of too few or too many referrals for a group, deferral

to the next block or the running additional groups are considered.

***Item 9: Tailoring***

Tailoring is required at patient, group and block levels.

There is flexibility regarding when NROL is integrated within a patient's rehabilitation, and at

present optimal timing is unknown. Patients joining NROL are referred to appropriate

targeted therapy groups and a personalised programme created. Clinical reasoning should

determine how NROL fits with a patient's in-person overall rehabilitation package of care.

Patients can attend more than one NROL block if clinically indicated and the patients remains

under the active care of their stroke or neurorehabilitation team.

The needs of patients in groups will be nuanced over time requiring a responsive approach.

All referrals are screened by group facilitators to ensure session content is tailored for the

current needs. NROL delivery should be considerate that language used is appropriate for

patients with a variety of neurological diagnoses (sudden, intermittent and progressive

conditions) and chronicities.

The structure of blocks, in terms of groups offered and frequency, actively considers

workforce availability.  The groups provided in each block is a balance between patient

demand and staff capacity. Some groups run all blocks, whilst others run intermittently. There

is an attempt to minimise groups running over holiday periods. Resourceful use of available

workforce is encouraged and the remote nature of NROL enables staff to deliver NROL whilst

working when from home or requiring work adjustments.  Student involvement can be

incorporated to support groups.

***Item 10: Modifications***

This article describes the NROL innovation developed for regional use. It was modified from

a standalone version (20) and expanded from a project implemented at a single NHS trust

(14).  The core components retained include provision of online real-time neurorehabilitation

with technology assistance, incorporating multidisciplinary targeted therapy and community

groups whilst embodying peer support.  Adaptations for integration within an existing

healthcare system included inclusion of stroke and other neurological conditions with varying

chronicity, delivering as an adjunct to complement existing rehabilitation, use of existing

workforce and being run as recurring 6-week NROL blocks to manage patient flow.

NROL continues to be modified over time. New groups are created to meet patient need or

to capitalise on changing staff skill sets. Groups can also be deferred, discontinued or replaced

based on patient and staff feedback.

***Item 11: How well planned & Item 12: How well delivered (including fidelity)***

Communication, resources and technology assistance are provided to optimise NROL entry

and participation (See *Items 3 & 4)*. Service data are obtained and reviewed to monitor

performance.

A detailed mixed-methods evaluation of NROL within a single trust is available *(Ackerley et al*

*23).*  Implementation and evaluation of the regional NROL programme in the North-West of

England is ongoing. This multi-trust regional innovation has to date delivered 4 x 6-week NROL

blocks, with 1082 patient contacts over 172 sessions provided to the 126 patients referred to

NROL. Withdrawal prior to programme start (17%) and drop-out are reasonably low (15%).

The median number of patients participating in a NROL block is 32 (range = 29-36). Ten

different groups are currently offered across a range of therapies, with typically nine groups

running per block. Therapy staffing efficiencies (ratio 0.4) are demonstrated compared to

one-to-one delivery (ratio 1.0) and currently over 40 staff across the North-West are involved

with group delivery. Overall NROL patient attendance is good (70%), with mechanisms in

place to facilitate attendance (e.g., patients receive text/email prompts in the first two weeks

of their programme). Attendance to the weekly peer support group (Café NROL) is optional

however this opportunity is taken up by many patients (58%). Patient NROL outcome measure

evaluation indicates improvements for many patients in health-related quality of life and

activity performance over the time of their NROL programme, acknowledging that NROL is

only one part of their overall rehabilitation. Overall patient satisfaction is high, with 99% of

patients reporting they would recommend NROL to another person.

***Context***

The NROL innovation detailed in this article is delivered within a context that can be described

using the domains of inner and outer settings (25).

The inner setting is defined as the four NHS Trusts that are situated within the Lancashire and

South Cumbria Integrated Stroke and Neurorehabilitation Delivery Network that provide

community-based stroke and neurorehabilitation care for the region. The region has a

population of 1.8 million and covers a large geographical footprint with urban and rural

settlements, and ethnic diversity. Deprivation and poor health affect many, with differences

in life expectancy and quality of life varying significantly (40).  In some neighbourhoods,

healthy life expectancy is 46.5 years.

Aligning with national strategies and policies, the Lancashire and South Cumbria region has a

vision to work collaboratively to share skills and expertise, data and best practice across Trusts

(41). A challenge is that Trusts have varied service remits (stroke, neurological or both) and

infrastructure (e.g. physical, staffing levels, technology systems, governance processes).

With regards to the outer setting, the impetus for starting NROL was the global pandemic,

which also influenced sociocultural values of staff and patients to increase the worthiness and

openness to use of remote technology. NROL also aligned with wider policy and strategies

(42-44), benefitted from an already established clinical-academic partnership between the

NHS and a university, and had funding from an external entity (45).

**Discussion**

NROL utilises an online delivery platform, with dedicated technology assistance, to provide

multidisciplinary real-time group therapy to patients actively receiving community stroke and

neurorehabilitation. Telerehabilitation is recognised as having a vital role to play in future

healthcare delivery (46) but as yet there are limited details of how to do this (47).  To address

this need, the TIDieR checklist is used as a structured method to provide a comprehensive

description of a regional NROL innovation embedded within a healthcare system. The actual

process of completing the TIDieR was time intensive but did provide the impetus for the team

to clearly describe NROL, agreeing the core components. Consideration was given to the

necessary balance of information to ensure comprehensive detail but attempting not to

overwhelm. Further documentation is available as supplementary files and by contacting the

corresponding author. This description details an exemplar of telerehabilitation using NROL

delivered in one region, and it is acknowledged that NROL will likely need adapting for other

areas given the significant interaction between an innovation and its context. This article

should facilitate other healthcare settings to adapt and implement NROL for their context.

Optimal adaptation of an innovation requires an understanding of the core components that

cannot be changed versus the adaptable periphery that can be changed (48, 49). The core

components of NROL include provision of online real-time neurorehabilitation with

technology assistance, incorporating multidisciplinary targeted therapy and community

groups whilst embodying peer support. It is proposed that these core components should be

consistently implemented for NROL but that the processes to achieve them are adapted to fit

local conditions. Examples include the use of MS Teams to deliver groups but other online

platforms are available; use of recurring 6-week NROL blocks but other timings may suit other

services; the number and types of targeted therapy groups will need to reflect workforce

capability and capacity. It is known innovations that have adaptability are more likely to be

used in clinical practice (50, 51).  Further examination of the adaptive components of NROL

will help discern how it can be upscaled for use in a variety of contexts.

Context is everything (52).  A limitation of the TIDieR checklist is that it does not include an

item on context.  Arguably to understand the innovation fully and guide future adaptations

an understanding of the context is required. This is because innovations are inextricably linked

to the context in which they are delivered, and achieving a good fit between these is

important to ensure the innovation works as intended (53). In this article, context is

deliberatively reported to help situate the innovation.  NROL did take resources, time and

effort to implement as a regional initiative and details on the implementation will be reported

elsewhere. The need for resources to enable implementation is not unexpected (49, 54, 55)

and current healthcare systems are often not set-up to facilitate this upfront effort. Influential

contextual factors included leadership buy-in and commitment, a clinical-academic

partnership and fit with the local and broader strategic landscape.

Inevitably NROL will continue to evolve. The TIDieR checklist does allow for reporting on

modifications. To date, the checklist has been primarily used for reporting interventions in

trials (56, 57) and there are limited examples of its use for adapted interventions over time.

This article documents current NROL delivery and captures its retrospective modifications.

Going forward, transparent reporting of new iterations of NROL together with descriptions of

their context should be undertaken to aid comparisons.

**Conclusion**

A hybrid approach incorporating telerehabilitation, to complement in-person therapy, is

required for a future-proof service that follows policy and guidelines. This comprehensive

description of a regional NROL innovation gives an example of successful implementation

within an existing healthcare system. It provides a platform for others to reduce duplication

of effort and help facilitate the use of telerehabilitation in clinical practice. Adapted versions

of NROL are expected when implementing in different contexts. Transparent reporting and

continuous evaluation alongside NROL implementation are encouraged and will ensure

maximal impact for neurorehabilitation delivery.

Availability of data/ resources

Digital copies of NROL materials are available on request by contacting Professor Louise

Connell (laconnell@uclan.ac.uk).

*Competing interests:*

The authors declare that they have no competing interests.

*Funding:*

NROL was supported by a generous donation from the charity SameYou (Charity number 1170102).

The funders had no role in the design of the study and collection, analysis, and interpretation of data

or in writing the manuscript.

## References

**References**

1.      Lohse KR, Lang CE, Boyd LA. Is more better? Using metadata to explore dose-response
relationships in stroke rehabilitation. Stroke. 2014;45(7):2053-8.
2.      French B, Thomas LH, Leathley MJ, Sutton CJ, McAdam J, Forster A, et al. Repetitive Task
Training for Improving Functional Ability After Stroke. Stroke. 2009;40(4):e98-e9.
3.      Schneider EJ, Lannin NA, Ada L, Schmidt J. Increasing the amount of usual rehabilitation
improves activity after stroke: a systematic review. Journal of Physiotherapy. 2016;62(4):182-7.
4.      Kwakkel G, Van Peppen R, Wagenaar RC, Wood Dauphinee S, Richards C, Ashburn A, et al.
Effects of Augmented Exercise Therapy Time After Stroke. Stroke. 2004;35(11):2529-39.
5.      Sentinel Stroke National Audit Programme. The Road to Recovery. The Ninth SSNAP Annual
Report 2022 [Available from:
https://www.strokeaudit.org/Documents/National/Clinical/Apr2021Mar2022/Apr2021Mar2022-
AnnualReport.aspx.
6.      Cramer SC, Dodakian L, Le V, McKenzie A, See J, Augsburger R, et al. A Feasibility Study of
Expanded Home-Based Telerehabilitation After Stroke. Front Neurol. 2020;11:611453.
7.      Laver KE, Adey-Wakeling Z, Crotty M, Lannin NA, George S, Sherrington C. Telerehabilitation
services for stroke. Cochrane Database Syst Rev. 2020;1(1):Cd010255.
8.      Stroke Association. A New Era for Stroke2016. Available from:
https://www.stroke.org.uk/sites/default/files/anefs_report_web.pdf.
9.      Parrott D, Ibarra S. Randomized Controlled Trial Comparing a Telemedicine Brain Injury
Coping Skills (BICS) group intervention to traditional in-person BICS for Brain Injury Patients and
Caregivers. Archives of Physical Medicine and Rehabilitation. 2021;102(10):e15.
10.     Khan F, Amatya B, Kesselring J, Galea M. Telerehabilitation for persons with multiple
sclerosis. Cochrane Database Syst Rev. 2015(4):CD010508.
11.     Rietdijk R, Power E, Attard M, Togher L. Acceptability of telehealth-delivered rehabilitation:
Experiences and perspectives of people with traumatic brain injury and their carers. Journal of
Telemedicine and Telecare. 2022;28(2):122-34.
12.     Hargreaves DL, D. Stroke, GIRFT Programme National Specialty Report2022. Available from:
https://gmnisdn.org.uk/wp-content/uploads/2022/04/Stroke-GiRFT-report-Apr-22.pdf)
13.     Chartered Society of Physiotherapy. National Evaluation of Remote Physiotherapy Services:
the findings 2022 [Available from: https://www.csp.org.uk/professional-clinical/professional-
guidance/remote-consultations/national-evaluation-remote-0.
14.     Ackerley S, Wilson N, Boland P, Reed J, Connell L. Implementation of neurological group-
based telerehabilitation within existing healthcare: a mixed methods evaluation. BMC Health
Services Research journal. Under review.
15.     Prvu Bettger J, Resnik LJ. Telerehabilitation in the Age of COVID-19: An Opportunity for
Learning Health System Research. Phys Ther. 2020;100(11):1913-6.
16.     Wosik J, Fudim M, Cameron B, Gellad ZF, Cho A, Phinney D, et al. Telehealth transformation:
COVID-19 and the rise of virtual care. Journal of the American Medical Informatics Association.
2020;27(6):957-62.
17.     Taylor A, Caffery L, Gesesew H, King A, Bassal A-r, Ford K, et al. How Australian Health Care
Services Adapted to Telehealth During the COVID-19 Pandemic: A Survey of Telehealth Professionals.
Frontiers in Public Health. 2021;9.
18.     Ross MH, Nelson M, Parravicini V, Weight M, Tyrrell R, Hartley N, et al. Staff perspectives on
the key elements to successful rapid uptake of telerehabilitation in medium-sized public hospital
physiotherapy departments. Physiotherapy Research International. 2022.
19.     Yang CL, Waterson S, Eng JJ. Implementation and Evaluation of the Virtual Graded Repetitive
Arm Supplementary Program (GRASP) for Individuals With Stroke During the COVID-19 Pandemic
and Beyond. Phys Ther. 2021;101(6).

20.    Beare B, Doogan CE, Douglass-Kirk P, Leff AP, Ward N. Neuro-Rehabilitation OnLine (N-ROL):
description and evaluation of a group-based telerehabilitation programme for acquired brain injury.
Journal of Neurology, Neurosurgery and Psychiatry. 2021;92(12):1354.
21.    Buckingham S, Anil K, Demain S, Gunn H, Jones RB, Kent B, et al. Telerehabilitation for
people with physical disabilities and movement impairment: development and evaluation of an
online toolkit for practitioners and patients. Disability and Rehabilitation. 2022:1-8.
22.    Royal College of Physicians. National Clinical Guideline for Stroke for the United Kingdom
and Ireland.2022. Available from: https://uploads-
ssl.webflow.com/62c3d8070eda8516a8ba9475/6380a7e6a9a3db745a8e8315_2023%20National%2
0Clinical%20Guideline%20for%20Stroke%20consultation%20document.pdf.
23.    Hoffmann TC, Glasziou PP, Boutron I, Milne R, Perera R, Moher D, et al. Better reporting of
interventions: template for intervention description and replication (TIDieR) checklist and guide.
BMJ. 2014;348(mar07 3):g1687-g.
24.    Rhon DI, Fritz JM, Kerns RD, McGeary DD, Coleman BC, Farrokhi S, et al. TIDieR-telehealth:
precision in reporting of telehealth interventions used in clinical trials - unique considerations for the
Template for the Intervention Description and Replication (TIDieR) checklist. BMC Medical Research
Methodology. 2022;22(1).
25.    Damschroder LJ, Reardon CM, Widerquist MAO, Lowery J. The updated Consolidated
Framework for Implementation Research based on user feedback. Implementation Science.
2022;17(1).
26.    Mace RA, Mattos MK, Vranceanu A-M. Older adults can use technology: why healthcare
professionals must overcome ageism in digital health. Translational Behavioral Medicine.
2022;12(12):1102-5.
27.    Tsai H-YS, Shillair R, Cotten SR. Social Support and "Playing Around". Journal of Applied
Gerontology. 2017;36(1):29-55.
28.    Doorley JD, Mace RA, Popok PJ, Grunberg VA, Ragnhildstveit A, Vranceanu AM. Feasibility
Randomized Controlled Trial of a Mind-Body Activity Program for Older Adults With Chronic Pain and
Cognitive Decline: The Virtual "Active Brains" Study. Gerontologist. 2022;62(7):1082-94.
29.    Miller MJ, Pak SS, Keller DR, Barnes DE. Evaluation of Pragmatic Telehealth Physical Therapy
Implementation During the COVID-19 Pandemic. Phys Ther. 2021;101(1).
30.    Aterman S, Ghahari S, Kessler D. Characteristics of peer-based interventions for individuals
with neurological conditions: a scoping review. Disability and Rehabilitation. 2022:1-32.
31.    Hughes R, Fleming P, Henshall L. Peer support groups after acquired brain injury: a
systematic review. Brain Injury. 2020;34(7):847-56.
32.    Clark E, Maccrosain A, Ward NS, Jones F. The key features and role of peer support within
group self-management interventions for stroke? A systematic review. Disability and Rehabilitation.
2020;42(3):307-16.
33.    Fleming V, Brownsett S, Krason A, Maegli MA, Coley-Fisher H, Ong Y-H, et al. Efficacy of
spoken word comprehension therapy in patients with chronic aphasia: a cross-over randomised
controlled trial with structural imaging. Journal of Neurology, Neurosurgery & Psychiatry.
2021;92(4):418-24.
34.    Ward NS, Brander F, Kelly K. Intensive upper limb neurorehabilitation in chronic stroke:
outcomes from the Queen Square programme. Journal of Neurology, Neurosurgery & Psychiatry.
2019;90(5):498-506.
35.    NHS England. National Stroke Service Model: Integrated Stroke Delivery Networks2021.
Available from: https://www.england.nhs.uk/wp-content/uploads/2021/05/stroke-service-model-
may-2021.pdf.
36.    Stroke Association. Shaping stroke research to rebuild lives2021. Available from:
https://www.stroke.org.uk/sites/default/files/research/stroke_priority_setting_partnership_full_rep
ort.pdf.

37.    EuroQol Research Foundation. EQ-5D-3L User Guide 2018 [Available from:
https://euroqol.org/publications/user-guides.
38.    Riazi A, Aspden T, Jones F. Stroke Self-efficacy Questionnaire: a Rasch-refined measure of
confidence post stroke. J Rehabil Med. 2014;46(5):406-12.
39.    Stratford P, Gill C, Westaway M, Binkley J. Assessing Disability and Change on Individual
Patients: A Report of a Patient Specific Measure. Physiotherapy Canada. 1995;47(4):258-63.
40.    Lancashire Independent Economic Review. Deep Dive: Health,Wealth & Wellbeing2021.
Available from: https://www.lancashireier.org/wp-
content/uploads/2021/12/LIER_Health_Wealth_and_Wellbeing_2021_v1.pdf.
41.    Lancashire and South Cumbria Provider Collaborative. Our vision 2022 [16 February 2023].
Available from: https://lscprovidercollaborative.nhs.uk/about-us/our-vision
42.    World Health A. Digital health. Geneva: World Health Organization; 2018.
43.    World Health Assembly. eHealth. Geneva: World Health Organization; 2005.
44.    NHS England and NHS Inprovement. 2022/23 priorities and operational planning
guidance2022. Available from: https://www.england.nhs.uk/wp-
content/uploads/2022/02/20211223-B1160-2022-23-priorities-and-operational-planning-guidance-
v3.2.pdf
45.    SameYou. NROL 2022 [16 February 2023]. Available from: https://www.sameyou.org/nrol.
46.    NHS Health Education England. The Topol Review 2023 [16 February 2023]. Available from:
https://topol.hee.nhs.uk/the-topol-review/
47.    English C, Ceravolo MG, Dorsch S, Drummond A, Gandhi DB, Halliday Green J, et al.
Telehealth for rehabilitation and recovery after stroke: State of the evidence and future directions.
International Journal of Stroke. 2022;17(5):487-93.
48.    Haynes A, Brennan S, Redman S, Williamson A, Gallego G, Butow P. Figuring out fidelity: a
worked example of the methods used to identify, critique and revise the essential elements of a
contextualised intervention in health policy agencies. Implementation Science. 2015;11(1).
49.    Greenhalgh T, Papoutsi C. Spreading and scaling up innovation and improvement. BMJ.
2019:l2068.
50.    Rogers E. Diffusion of innovations: 5th Edition. New York: Free Press; 2003.
51.    Gustafson DH, Sainfort F, Eichler M, Adams L, Bisognano M, Steudel H. Developing and
Testing a Model to Predict Outcomes of Organizational Change. Health Services Research.
2003;38(2):751-76.
52.    Bate P. The Health Foundation: Perspectives on context. Available from:
https://www.health.org.uk/sites/default/files/PerspectivesOnContextBateContextIsEverything.pdf.
53.    Hawe P. Lessons from Complex Interventions to Improve Health. Annual Review of Public
Health. 2015;36(1):307-23.
54.    Standing C, Standing S, McDermott M-L, Gururajan R, Kiani Mavi R. The Paradoxes of
Telehealth: a Review of the Literature 2000-2015. Systems Research and Behavioral Science.
2018;35(1):90-101.
55.    Signal N, Martin T, Leys A, Maloney R, Bright F. Implementation of telerehabilitation in
response to COVID-19: Lessons learnt from neurorehabilitation clinical practice and education. New
Zealand Journal of Physiotherapy. 2020;48:117-26.
56.    Tume LN, Blackwood B, McAuley DF, Morris K, Peters MJ, Jordan J, et al. Using the TIDieR
checklist to describe the intervention of the Sedation and Weaning in Children( SANDWICH) trial.
Nursing in Critical Care. 2022.
57.    Carlsson H, Rosén B, Björkman A, Pessah-Rasmussen H, Brogårdh C. SENSory re-learning of
the UPPer limb (SENSUPP) after stroke: development and description of a novel intervention using
the TIDieR checklist. Trials. 2021;22(1).