Stress in Autism (STREAM): A Study Protocol on the Role of Circadian Activity, Sleep Quality and Sensory Hyper-Reactivity

Christine M. Falter-Wagner[1*], Clara C. Gernert[1*], Valdas Noreika[2,3], Barbara Jachs[3], Nazia Jassim[4,5], Kathryn Gibbs[6], Joaquim Streicher[3], Hannah Betts[3], Tristan A. Bekinschtein[3]

[1] Department of Psychiatry and Psychotherapy, Medical Faculty, Ludwig-Maximilians-University, Munich, Germany

[2] Department of Psychology, School of Biological and Behavioural Sciences, Queen Mary University of London, London, United Kingdom

[3] Consciousness and Cognition Lab, Department of Psychology, University of Cambridge, Cambridge, United Kingdom

[4] Autism Research Centre, Department of Psychiatry, University of Cambridge, Cambridge, United Kingdom

[5] Prediction and Learning Lab, Department of Psychology, University of Cambridge, Cambridge, United Kingdom

[6] Independent researcher

[*] Co-first authorship

Correspondence concerning this manuscript should be addressed to:
Christine Falter-Wagner
Department of Psychiatry and Psychotherapy
Medical Faculty, LMU Munich
Nussbaumstr. 7, 80336 Munich, Germany
Email: christine.falter@med.uni-muenchen.de

Tristan Bekinschtein
Consciousness and Cognition Lab
Department of Psychology
University of Cambridge
Cambridge, CB2 3EB, United Kingdom (UK)
Email: tb419@cam.ac.uk

Abstract

This proof-of-concept study protocol describes a novel approach towards investigating

interrelationships between stress indices, sleep and circadian activation patterns, and sensory

sensitivities in autism. We will adopt an individualised approach, collecting 17 days of data per person

from 20 participants with and 20 without autism. Participants' sleep will be tracked by using wearable

EEG headbands and a sleep diary. Diurnal monitoring of heart rate and electrodermal activity via

wearables will serve as proxies of stress and will be time-locked to subjective experience traces collected

throughout the day. These traces quantify the experience dimensions such as stress and sensory

sensitivities (e.g., sensory overload). Capturing the dynamics of subjective experience phase-locked to

neural and physiological proxies between and within individuals can potentially tackle some of the most

pressing issues in autism: sleep problems, sensory overload and stress, and will ultimately lead to an

individualised prognostic approach to mental health in autism.

1. Introduction

Due to the high prevalence of anxiety, depression, and suicidality among autistic individuals, mental health is a pressing theme and one of the top priorities in the field of autism research (1-2). Stress is a key player for mental health in the framework of the currently prevalent stress-vulnerability model of mental health problems. Importantly, stress in the life of an autistic person is likely to be complex and autism-specific, thus necessitating first-hand accounts. Indeed, the value of first-hand experience accounts by people with autism has been proclaimed already several decades ago (3) and yet, case studies or qualitative and nuanced assessment of experience is limited, and rarely linked to neurocognitive mechanisms. In the planned study, we adopt a quantitative approach to capture complex individual experiences by temporal experience tracing (TET). Next, we link these subjective, quantifiable data to neural and physiological measurements. The resulting rich neurophenomenological dataset of each participant will be analysed to capture the complex interplay between experiences and physiological dynamics.

Subjective impressions of people have been assessed primarily with Likert scales and questionnaires or with qualitative approaches via interviews. The scales are a rough tool to get information about a person's experience, the qualitative approach is time-consuming, requires expertise, and in practice, the outcome is difficult to compare between participants. To account for this, we developed a more fine-grained method to capture relevant dimensions of experience (4). Once these dimensions of experience are defined (typically 10-14 dimensions; e.g., attention, boredom, effort, social anxiety, rumination), people draw and trace in time the intensity of each dimension, allowing subjective aspects of experience to be treated as an objective measure of the stream of consciousness. These transformed dimensions of experience into time series can be then analysed in parity and combination with daytime and sleep-related activity, heartbeat and electrodermal activity data from wristbands, as well as with portable brain measures via EEG headbands. To directly tap into real-life experience of stress, the quantified subjective experiences will be related to individual self-measurements of several stress-related neurobehavioural factors. One major source of stress in autism has been suggested to be sensory-processing difficulties (5-7). Beyond autism it is known that sensory disturbances are increased because of sleep difficulties and circadian abnormalities, for instance, diurnal fluctuations of sensory processing partially depend on

the individual chronotype (e.g., 8-9). Indeed, sleep difficulty is also known to be a very common

problem in autism (e.g., 10-14) with 85.6 % of autistic individuals reporting at least one sleep-related

concern (13). Furthermore, a positive relationship has been shown between sleep issues and autism

symptom severity (15). In particular, recent large meta-analytic findings showed decreased sleep

efficiency and a higher proportion of light sleep in people with autism spectrum disorder (ASD) across

the lifespan (16-18). Importantly, poor sleep has been shown to correlate with psychological issues,

adaptive functioning and physical health in people with ASD (19). Likewise, individuals with autism

report circadian abnormalities (20).

Taking these findings together, we hypothesise a vicious circle between disturbed sleep and circadian

irregularities increasing sensory issues that lead to distress, which then further aggravates sleep

problems. Indeed, there are hints of daytime stress and hypothalamic-pituitary-adrenal axis

dysregulation in association with both sleep problems and autism (21-22). In a review of early evidence

on the effects of meditation on autism, it has been shown that meditative activities might lead to

improved relaxation, reduced clinical symptoms and better quality of life (23). Given meditation lends

itself as an activity from which subjective experience dynamics can be extracted, performing a daily

meditation session and reporting experiences afterwards over a period of time might be a good option

to capture individuals' experiences in a specific task.

Thus, in the first step, the current study aims at measuring individual interrelationship patterns of stress,

sensory reactivity, circadian activity, and sleep quality in autism. Importantly, the planned study will

build on an individualised approach to collecting rich longitudinal data sets from individuals over a

period of time in an ecological setting (i.e., fully home-based assessment). Our long-term goal with the

data collected in this study is to set the foundation for the development of individualised prediction of

episodes of sensory sensitivity changes and distress, based on e.g., sleep parameters, for people with

autism as a prevention strategy for mental health comorbidities.

Finally, honouring the relevance of a participatory approach to autism research, the study protocol was

developed iteratively incorporating feedback on feasibility, intelligibility, and relevance from volunteer

pilot participants with and without autism. Adjustments and advancements to the design, instructions,

schedule of testing and content (e.g., dimensions tested with TET) were made in the course of this

participatory process.

2. Materials and Methods

The study PRE.2021.091, named as 'Circadian and homeostatic effects on stress and daytime

neurophenomenology in autism spectrum conditions', was approved by the Cambridge Psychology

Research Ethics Committee on the 15th of March 2022 (see Approval Letter). Participants consent is

informed and written.

2.1 Aim, design and set up of the study

The main aims of this proof-of-concept study are to a) quantify detailed physiological and

phenomenological data to be able to explore relevant and meaningful aspects of stress and experience

in autistic individuals in everyday life; b) analyse whole-night sleep data, measured subjectively and

objectively over several days, in adults with and without ASD for characterization and comparison; c)

explore daytime parameters of the autonomic nervous system within and between both groups; d)

investigate whether participants' traces of daily meditation can be clustered in common patterns of

experience; e) explore if night or meditation neural and physiological features can be used as classifiers

for data-driven experience clusters, f) determine the differential predictive power of daytime stress,

sensory hypersensitivities, and physical activity on participants' sleep behaviour and vice versa (i.e.

concept of vicious circle), g) investigate whether we can develop an individualised data-driven

prediction model of distress and sensory hypersensitivity episodes for autistic individuals. The general

gist of the data acquisition protocol is based on remotely collecting data at participants' homes, using

non-invasive technological devices in their daily life. This includes the use of a custom-built app

programmed in *Unity* using the programming language *C#*, an EEG headband (*Dreem*) and a wristband

(E4, *Empatica*). We will provide full equipment, including a smartphone where required, for the time

of participation.

Initially, demographic data will be collected on paper. Then participants will be asked to fill out a series

of questionnaires via the app and stored them in the data server at the University. Standard

questionnaires for both groups of participants, Control Group (CG) the and ASD group, will comprise

the following: *Beck Depression Inventory* (BDI-II) (24), the short version of the *Autism Spectrum*

Quotient (AQ-10) (25), the short version of the *Sensory Perception Quotient* (SPQ-10) (26), the

Pittsburgh Sleep Quality Index (PSQI) (27), the *State-Trait Anxiety Inventory* with focus on items of

trait anxiety (STAI (Trait)) (28), the *Perceived Stress Questionnaire* (PSQ) (29) and the *Culture Fair*

Intelligence Test (CFT 20-R) (30). The participant is assigned a number for the study, the key for which

will be stored separately from the study data. The pseudonymised number is used as an identifier for the

clinical questionnaires. During at least 17 days, participants will be asked to fill out diaries regarding

their sleep quality and their activities throughout each day. Both diaries are included in the app. They

will also be asked to use the TET tool in the app to describe the dynamics of their daily experiences

(e.g., tracing the level of stress, see Figure 3). Participants' sleep will be tracked using EEG-headbands

(*Dreem*) during the night. Diurnal monitoring of heart rate (HR), heart rate variability (HRV) and skin

conductance (SC) via a wristband (E4, *Empatica*) will serve as objective, physiological parameters of

sympathetic and parasympathetic activity. In Phase I, baseline data will be gathered for three days.

In Phase II, participants will be asked to additionally perform a meditation session once a day for 14

days at all, lasting 20 minutes. For an overview of the recording schedule, see Figure 1. An audio guide

helps the participants with the structure of the meditation task, which is separated into a breathing and

body scanning part, each one lasting 10 minutes. After the meditation session participants fill out the

meditation TET tool in the app to report about the dynamics of their experiences made whilst meditating

(e.g., tracing the level of rumination, see Figure 4). On the final day, participants will fill out the BDI-

II (24) referring to depressive symptoms in the past week to compare participants' depressive

symptomatology before and during participating in our study.

2.2 Sample size, inclusion, and exclusion criteria

In this proof-of-concept study we are aiming to recruit at least 40 adults (20 CG/20 ASD) (31) for

analyses at the group and single participant individual. However, the strength lays in the weeks

repetitions per participant that allows for nested statistical models that lend robustness and nuance to the

questions (32). Inclusion criteria for both groups of participants will comprise the following: age

between 18 and 65 years, normal or corrected-to-normal vision, IQ>70 in the CFT 20-R (30) and verbal

in order to understand instructions and be able to ask questions. Autistic individuals should have

received a medical diagnosis of autism with the code 6A02.0 (ICD-11), F84.5 (ICD-10), or F84.0 (ICD-

10). Further psychiatric comorbidities (ICD-10 F codes) in the ASD group are no exclusion criteria in

general but will be collected within demographic data collection. No diagnosed psychiatric (ICD-10 F

codes) or acute neurological disorder should be present in the control group (CG). Medication, whether

taken regularly or on demand, is also not an exclusion criterion for participation, but will be reported in

the diary.

2.3 Recruitment

Autistic individuals will be recruited in Germany and the UK through the outpatient department for

autism spectrum disorders at the LMU Munich, the Cambridge Autism Research Centre (CARD), and

through autism-related network partners (e.g., *Autistica* in the UK and *WGAS* in Germany). The CG will

additionally be recruited using the SONA System, an online platform for research participant

recruitment.

2.4 Measures and Procedure

Participants will undergo training in the use of the equipment and the app either online or in

person. Participants' consent form and demographic data will be collected on paper beforehand.

Afterwards, participants will receive an anonymized identifier to log into an app. At the beginning of

Phase I participants will fill out several standard questionnaires first (see Table 1). All questionnaires

are included in the app and need to be answered only once, except from the BDI-II (24), which needs to

be answered for a second time at the end of the study.

EEG data (sleep and meditation) and physiological data (HR and SC) as objective parameters are

recorded for at least 17 days. Several neural markers will be extracted from participants' EEG data

recorded while sleeping (see Table 2) or meditating (see Table 3). For an overview of the headband's

electrodes and channels, please see Figure 5. Physiological parameters, like HR and SC, will be assessed

during the day by using a wristband (E4, Empatica) (see Tables 4-5 and Figure 1).

Beside from questionnaires and diaries (i.e., sleep and activity diary) the app we have created also

contains the TET tool for reporting about daily and meditation experiences (see Table 6). In both TET

assessments participants have the option to define a personalised dimension at the end.

Table 1

Questionnaires: Subjective Markers

	Questionnaires	
Marker	Start of Study	End of Study
Depressive Symptoms	BDI-II (Beck et al., 1996)	BDI-II (Beck et al., 1996)
Autistic Traits	AQ-10 (Allison et al., 2012)	
Sensory Perception	SPQ-10 (Greenberg et al., 2018)	
Sleep Quality and Habits	PSQI (Buysse et al., 1989)	
Trait Anxiety	STAI (Trait) (Spielberger et al., 1983)	
Experience of Stress	PSQ (Fliege et al., 2009)	
IQ testing	CFT 20-R (Weiß et al., 2006)	

158 *Note*. Seven questionnaires need to be filled out once at the start of the study. Participants are asked to

159 fill out the BDI-II (Beck et al., 1996) for a second time at the end of the study.

161 **Table 2**

162 *Sleep: Neural Parameters*

Abbreviation	Marker	Unit
TST	Total sleep time	Minutes
SOL	Sleep onset latency	Minutes
MAI	Micro arousal index	Number
WASO	Wake after sleep onset (WASO)	Minutes
N1	Stage N1 duration	Minutes
N2	Stage N2 duration	Minutes
N3	Stage N3 duration	Minutes
REM	Stage REM duration	Minutes
NREM	Stage NREM duration	Minutes
N1_percentage	Proportion of TST in stage N1	Ratio (%)
N2_percentage	Proportion of TST in stage N2	Ratio (%)
N3_percentage	Proportion of TST in stage N3	Ratio (%)
REM_percentage	Proportion of TST in stage REM	Ratio (%)
NREM_percentage	Proportion of TST in stage NREM	Ratio (%)
AW	Awakenings	Number
TIB	Total time in bed	Minutes
SE	Sleep Efficiency = TST/TIB	Ratio (%)

163 *Note.* Neural parameters which are suggested to be extracted from whole-night sleep EEG data.

166   **Table 3**

167   *Meditation: Neural Parameters*

Abbreviation	Channel	Note
Ch1	Fz-O1	Frontal to occipital
Ch3	Fz-F7	Frontal
Ch4	F8-F7	Frontal interhemispheric
Ch5	F7-O1	Frontal to occipital
Ch7	Fz-F8	Frontal
Abbreviation	Marker	Note
δ	Delta power (1-3 Hz)	Spectral parameter
Θ	Theta power (4-7 Hz)	Spectral parameter
α	Alpha power (8-12 Hz)	Spectral parameter
β	Beta power (13-30 Hz)	Spectral parameter
γ	Gamma power (31-48 Hz)	Spectral parameter
SE	Spectral entropy	Spectral parameter
MSF	Median spectral frequency	Spectral parameter
FAA	Frontal alpha asymmetry	Spectral parameter
PE- α	Permutation entropy in alpha	Information theory
PE- Θ	Permutation entropy in theta	Information theory

168   *Note.* EEG channels and neural parameters which are proposed to use for analysis and classification.

169   EEG data are measured with the *Dreem* headband.

171   **Table 4**

172   *Physiological Parameters: Heart Rate and Heart Rate Variability*

1. Heart Rate Measures		
Parameter	Description	Unit
HR	Mean heart rate	Beats per minute (bpm)
HR_Range	Difference between the highest and lowest heart rate	Beats per minute (bpm)
SNS	Sympathetic activity index	Index
PNS	Parasympathetic activity index	Index
Stress	Stress index	Index
2. Heart Rate Variability (HRV): Time-Domain Measures		
Parameter	Description	Unit
SDNN	Standard deviation of NN intervals	Milliseconds (ms)
SDNN_Index	Standard deviation of the average NN intervals for each 5 min segment	Milliseconds (ms)
pNN50	Percentage of successive RR intervals that differ more than 50 ms	Percentage (%)
3. Heart Rate Variability (HRV): Frequency-Domain Measures		
Parameter	Description	Unit
LF_power	Relative power of the low-frequency band (0.04-0.15 Hz)	Percentage (%)
HF_power	Relative power of the high-frequency band (0.15-0.4 Hz)	Percentage (%)
LF/HF	Ratio of low-frequency to high-frequency power	Ratio (%)

173   *Note.* Several heart rate and heart rate variability (HRV) measures get extracted from the wristband's

174   (E4, *Empatica*) blood volume pressure data.

176 **Table 5**

177 *Physiological Parameters: Skin Conductance*

1. Skin Conductance (SC)		
Parameter	Description	Unit
SC_Mean	Mean value of skin conductance	Micro Siemens (uS)
SC_Range	Difference between the highest and lowest point in SC	Micro Siemens (uS)
SC_AUC	Area under curve (AUC) for the SC signal	Index
2. SC: Phasic Activity		
Parameter	Description	Unit
Phasic_Mean	Mean value of the phasic activity signal	Micro Siemens (uS)
Phasic_Std	Standard deviation of the phasic activity signal	Micro Siemens (uS)
Phasic_Range	Difference between the highest and lowest point in phasic activity	Micro Siemens (uS)
Phasic_Peak	Number of peaks in phasic activity	Index
Peak_Amplitude	Average of peaks' amplitude	Micro Siemens (uS)
Phasic_AUC	Area under curve for the phasic activity signal	Index
Phasic_AUC_Percentage	Proportion of Phasic_AUC in SC_AUC	Ratio (%)
3. SC: Tonic Activity		
Parameter	Description	Unit
Tonic_Mean	Mean value of tonic activity	Micro Siemens (uS)
Tonic_Range	Difference between the highest and lowest point in tonic activity	Micro Siemens (uS)
Tonic_AUC	Area under curve for the tonic activity signal	Index
Tonic_AUC_Percentage	Proportion of Tonic_AUC in SC_AUC	Ratio (%)
Phasic_Tonic_AUC	Ratio of Phasic_AUC and Tonic_AUC	Ratio (%)

178 *Note.* Measured levels of skin conductance (SC) gets decomposed into phasic and tonic activity. For all

179 three variables several parameters will be calculated.

180 **Table 6**

181 *Dimensions for TET*

Number	Daily Experiences	Meditation
1	Meta-Awareness	Meta-Awareness
2	Wakefulness	Wakefulness
3	Boredom	Boredom
4	Sensory Avoidance	Effort
5	Social Avoidance	Source
6	Physical Tension	Attention
7	Scenario Anxiety	Body
8	Rumination	Sensory Seeking
9	Stress	Physical Tension
10	Pain	Scenario Anxiety
11	Personalised Dimension	Rumination
12		Stress
13		Personalised Dimension

Note. Ten dimensions are defined for TET referring to participants' daily experiences. 12 dimensions

are described for the TET after meditation. For both tasks, participants can add another personalised

dimension.

2.5 Data management

The experimental data will be stored on a server hosted by the Clinical School Computing Service from

Addenbrooke's Hospital (Cambridge, UK). The EEG recordings will be stored temporarily in a cloud

operated by the EEG provider *Rhythm* by *Dreem*. Wristband recordings are stored by *Empatica* and

hosted by Amazon Web Services located in the US. All data will be moved daily to the Universities

appropriate servers. All collected raw data, logged with a number code, will be stored permanently in a

backup in the High-Performance Computing (HPC) at the University of Cambridge.

Personally identifiable information will be kept separately from the experimental data and will not be

provided to individuals beyond the research team. Experimental data will not contain personally

identifiable information. Participants will receive an anonymized identifier for the EEG headband and

the wristband. The code for decrypting the personal data is only available to the database administrator

and is locked on a local computer that is not connected to the network. Clinical data will be stored in

the Clinical School Computing Service separated from the physiological and behavioural data to ensure

compartmentalization and data privacy. All EEG and behavioural data storage will comply with the

requirements of the Data Protection Act. All data will be stored in a secure system for at least 10 years.

2.6 Risk management

Dreem headbands are registered as an FDA Class II Medical Device in the USA (*Dreem*, n.d.). Both

devices, the wristband and the headband, collect data non-invasively and have a consumer CE marking.

For more specific safety instructions participants will first have to meet the inclusion criteria. Second,

participants will have to sign a consent form before starting the study, which highlights the most

important safety instructions and warnings when using the devices, amongst other information. These

safety considerations refer to the manual and the instruction sheet of the producers. Third, all

participants will undergo operator training to learn how to use the devices. Even though there are no

actual associated risks for participants, this approach is to ensure correct data collection and a

comfortable use of the devices. Fourth, participants will be asked to watch an instruction video produced

by the research group explaining how to wear the devices. The video will be included in the app for easy

access. Fifth, participants can always get in touch with members of the research team via email or phone.

Finally, it will be made sure that participants understand that they can stop data collection and study

participation at any time without giving a reason.

We considered three main risks in our study design and address them as follows:

A) Recruitment, dropout, missing data

Participants can interrupt the study protocol for a short break (e.g., due to acute illness) and can define the time when performing the daily meditation flexibly according to their normal daily schedule. The study protocol of 17 days can be extended when data collection was unsuccessful in more than two days and the participant is willing to extend data acquisition. The experimenter will monitor the data quality. If data quality is unsatisfactory, the researcher will guide the participant to mitigate any issues. Furthermore, nested statistical models are used for analysis which are robust regarding missing data. Participants will be encouraged to record data continuously over several days, as this will allow us to understand potential effects of a cumulative sleep-deficit on daytime sensory processing and subjective experiences of stress. Data collection will be stopped if participants are unable to start data collection within two weeks of receiving the devices or if the break between data collection days exceeds one week.

B) Equipment malfunction

We will keep spare devices and send participants replacements if needed. For participants who do not have a compatible smartphone we will offer a smartphone with the installed app on it for the time when participating in the study. Additionally, we created a website version, where participants can anonymously answer all questionnaires online.

C) Participants' digital literacy

Participants may lack digital skills to use the equipment needed to perform the study. We will test for digital skills and task comprehension at the start of the experiment. We only embark participants on the study who demonstrate sufficient digital skills and sufficient skills using the wearables correctly to avoid too high burden for those participants without digital literacy skills. At the same time, participants will be supported by the experimenters throughout the study (e.g., via virtual meetings).

Any risk from our equipment is managed by safety checks on computers and the devices directly after and before the start of each new participant, following the respective Standard Operating Procedures. It should be emphasised that EEG is an entirely passive monitoring measurement that detects tiny electrical

currents generated by the brain's own activity. These signals are then amplified and recorded by a

computer. The wristband from *Empatica* gathers physiological data non-invasively by LED lights and

SC electrodes.

2.7 Data analysis plan

Given the high prevalence of sleep disorders (33) in autism, we assume significant differences in

subjective ratings and measures of sleep quality when analysing participants' sleep diaries (17).

H1) We expect autistic individuals to report longer sleep onset latency, more night awakenings, lower

sleep duration and lower sleep efficiency in the sleep diary compared to individuals of the control group.

H1.1) We hypothesise that objective parameters, referring to whole-night EEG data (for outcome

variables, see Table 2), are in line with participants' subjective reports.

H2) We expect differences between autistic individuals and control individuals in their EEG sleep

patterns (17-18).

H3) We hypothesise that autistic individuals show higher sympathetic activity and stress indices

throughout the day, lower heart rate variability and more phasic activity in their SC time series compared

to the CG.

We aim to clarify the differential predictive power of daytime stress and physical activity, anxiety,

depression, and autism traits on participants' sleep behaviour. In the next step we want to focus on

participants' TET reports and their potential as predictive markers for past and upcoming sleep

disturbances.

H4) We predict that autistic individuals will show higher absolute values and a higher variance in the

graphs of the TET tool for daily dimensions when reporting about dimensions that are linked to autism

specific behaviour (i.e., social avoidance, sensory avoidance).

H5) For the meditation task, we expect an association in both groups (ASD/CG) between peri-meditation

EEG neural dynamics and self-reported meditation-induced changes in stress levels.

H6) We hypothesise an association between meditation EEG neural activity and the following night's sleep quality.

H7) We expect the TET parameters of stress, reported in the daily and meditation TET tasks, to be associated and also related to the following night's self-reported and objectively measured sleep quality.

Exploratorily, EEG features of the meditation session will be used for classification of data-driven experience clusters for meditation, while sleep EEG features will be used for classification of psychophysiological clusters of participants' daytime autonomous nervous system activity level (like in 32).

2.8 Timeline of the study

During 2021-2022 we designed the study project and refined the analysis pipeline by incorporating input from autistic advisors (Autism Research Centre, University of Cambridge). Additionally, an app was programmed for digitalisation of the TET tool (see Figure 3 and 4). All diaries and questionnaires, except from the demographics, were included in the app to make data acquisition easier and more flexible. So far, we have collected pilot data from 6 participants (3 ASD/ 3 CG). We are aiming to recruit participants in the UK and Germany over at least the next 24 months. Each participant will be asked to collect data for 17 days in a row. In parallel to recruitment, we will refine our analysis pipeline by using already collected pilot data.

3. Discussion

This study addresses several common co-occurring conditions in ASD: sleep disorders, sensory hypersensitivity, and stress. The study protocol described herein includes several aspects of novelty in the field of autism research concerning the design and research methods. First, subjective experiences are quantified on an individual level and related to psychophysiological and neurophysiological (i.e., objective) measurements. Second, data are collected over a prolonged period of time allowing for analysis of individual patterns of interrelation between circadian activities and sleep, sensory hypersensitivity, and stress. Third, the meaningful dimensions of the experience during the day and

during the period of meditation were discussed and agreed with autistic individuals, including autistic

researchers. For that we expect to not only make the study relevant clinically but more importantly, to

learn more about the individual aspects of stress experience in autism. And fourth, participants will be

assessed during their daily lives at home using wearables and digital measurements that enable us to

apply lab-level control within the ecological setting of people's real-life experience and thereby increase

relevance.

3.1 Limitations of the study design

Aside from the methodological advantages and the importance of the addressed research questions in

the field of autism, the study design has limitations. First of all, a personalised approach over a longer

period implies that it is more difficult to collect data from a large sample size. There might be a sampling

bias as we gather data from people who are able to cope with the wearable equipment. Nevertheless, we

will learn important and new aspects about the interrelation of individual stress experience, sensory

hypersensitivity as well as circadian and sleep irregularities by focusing on personalised patterns gained

from rich individual longitudinal data sets (34). Additionally, we aim to investigate the potential of

objective and subjective outcome variables for classification of clusters within and between individuals

with and without ASD. Questions about the directionality of effects and etiological pathways remain a

target for future prospective studies and are outside the scope of the current study design. Given the

heterogeneity in the autism phenotype and with respect to constantly changing environmental factors,

beside from individual, time, and context dependent factors (e.g., unknown coping strategies,

comorbidities, social support), we might not derive a general pattern across all participants. Still, we

expect to be able to assess individual interrelationships that can inform about factors relevant for

prediction of stress and sensory overload. In this respect, we account for several potential cofactors in

our study by using different diaries and addressing several dimensions of experiences, besides giving

participants extra space to report any irregularity of the day and their state.

As TET is a method in which participants graph intensity of their experiences retrospectively, this might

lead to memory biases. Recall biases are not specific for autistic individuals and could likewise occur in

control participants and for the traditional alternatives such as rating scales. Additionally, we advise participants to draw their experiences directly after performing the meditation session to reduce the recall bias in this task. We are convinced though that TET offers a great opportunity to get more insight into the dynamics of subjective experiences, something that cannot be derived from standard rating scales (32).

As two wearables, a wristband and a headband, are used for data acquisition in this research project we are aware that data quality might be lower due to motion artefacts and lower sampling frequencies compared to clinical assessments, where devices such as an electrocardiogram (ECG) for patients' HR or polysomnography (PSG) for assessing sleep quality are used. Additionally, the *Dreem* headband has a scalp coverage limited to a few frontal and occipital electrodes, which yields EEG data with low spatial resolution. Despite of these limitations, as we want to capture naturalistic behaviour, experiences, and physiological responses of individuals in their daily and relevant environment, it is crucial to use non-lab based and non-invasive data acquisition methods with an uncomplicated handling. Wearables offer this possibility and flexibility (35). We address upcoming issues in data quality by giving participants regular feedback about the quality of their recordings and discuss individual adjustments in the process of data acquisition, in addition to adapted pre-processing methods when analysing raw data.

3.2 Dissemination plans

The results of the study will be presented at relevant conferences including those, which have a broad audience such as individuals with autism and their families as well as practitioners in the field, who might be interested in the patterns of interrelation between sleep, sensory hyper- or hyposensitivity, and stress. Furthermore, the use and validation of the method using quantified experiences and its direct physiological counterparts is a strength in itself and is applicable to any group of participants, hence relevant outside the autism and stress research. Thus, preference will be given to more general psychology journals for publications resulting from the current study.

3.3 Handling of amendments to the study, including termination

As we have already gathered pilot data and have refined the study protocol in a participatory research

collaboration over the last year, we are now in the position to run a full proof-of-concept study. Any

unforeseen amendment to the study protocol will be first addressed to the University ethics committee.

After their approval, changes will be discussed with the participants directly. As we have created the

app for this study project, changes can be easily implemented by our research teams. Participants will

be informed that they are free to withdraw their consent at any time without giving a reason for

termination. The experimenter will stay in close contact with the participants to note any upcoming

technical or other issues early and make adjustments where required on an individual level. Additionally,

it is always possible for participants to interrupt the study schedule for some days (e.g., because of

illness) and continue later.

4. Conclusion

The current study protocol will deliver insights into individual patterns of objective and subjective stress

experienced by individuals with autism and relate it to their circadian activities and sleep and any

occurring episodes of sensory hypersensitivity. The results will help tailor future interventions to tackle

stress reduction for better mental health outcomes in the autistic population.

Authors' contributions

All authors contributed to the design of the study protocol.

Data availability statement

This article does not contain personal data.

Acknowledgements

The authors were supported by a Cambridge-LMU Strategic Partnership Grant (granted to TB, VN and

CFW).

5. References

1. Cassidy S, Bradley P, Robinson J, Allison C, McHugh M, Baron-Cohen S. Suicidal ideation and

suicide plans or attempts in adults with Asperger's syndrome attending a specialist diagnostic clinic: a

clinical cohort study. The Lancet Psychiatry [Internet]. 2014 Jul;1(2):142–7. Available from:

https://www.thelancet.com/journals/lanpsy/article/PIIS2215-0366(14)70248-2/fulltext

2. Gatfield O, Mangan C, Haar T, Kinniburgh A, Rodger S. 2016 Autism Research Priorities Survey

Report FINAL REPORT [Internet]. 2016 [cited 2023 Feb 15]. Available from:

https://www.autismcrc.com.au/sites/default/files/inline-files/Research%20priorities%20survey%20-

%20Final%20report.pdf

3. Cesaroni L, Garber M. Exploring the experience of autism through firsthand accounts. Journal of

Autism and Developmental Disorders. 1991 Sep;21(3):303–13.

4. Jachs B, Camino Garcia M, Canales-Johnson A, Bekinschtein TA. Drawing the experience dynamics

of meditation. bioRxiv. 2022 Mar 7; Available from: https://doi.org/10.1101/2022.03.04.482237

5. Green SA, Ben-Sasson A. Anxiety Disorders and Sensory Over-Responsivity in Children with Autism

Spectrum Disorders: Is There a Causal Relationship? Journal of Autism and Developmental Disorders.

2010 Apr 10;40(12):1495–504. Available from: https://doi.org/10.1007/s10803-010-1007-x

6. Reynolds S, Lane SJ, Thacker L. Sensory Processing, Physiological Stress, and Sleep Behaviors in

Children with and without Autism Spectrum Disorders. OTJR: Occupation, Participation and Health.

2011 May 20;32(1):246–57. Available from: https://doi.org/10.3928/15394492-20110513-02

7. Wigham S, Rodgers J, South M, McConachie H, Freeston M. The Interplay Between Sensory

Processing Abnormalities, Intolerance of Uncertainty, Anxiety and Restricted and Repetitive

Behaviours in Autism Spectrum Disorder. Journal of Autism and Developmental Disorders. 2014 Sep

27;45(4):943–52. Available from: https://doi.org/10.1007/s10803-014-2248-x

8. Engel-Yeger B, Shochat T. The Relationship between Sensory Processing Patterns and Sleep Quality

in Healthy Adults. Canadian Journal of Occupational Therapy. 2012 Jun;79(3):134–41. Available from:

https://doi.org/10.2182/cjot.2012.79.3.2

9. Vasak M, Williamson J, Garden J, Zwicker JG. Sensory Processing and Sleep in Typically

Developing Infants and Toddlers. American Journal of Occupational Therapy. 2015 May

29;69(4):6904220040p1. Available from: https://doi.org/10.5014/ajot.2015.015891

10. Elrod MG, Hood BS. Sleep Differences Among Children With Autism Spectrum Disorders and

Typically Developing Peers. Journal of Developmental & Behavioral Pediatrics. 2015 Apr;36(3):166–

77. Available from: https://doi.org/10.1097/dbp.0000000000000140

11. Hohn VD, de Veld DMJ, Mataw KJS, van Someren EJW, Begeer S. Insomnia Severity in Adults

with Autism Spectrum Disorder is Associated with sensory Hyper-Reactivity and Social Skill

Impairment. Journal of Autism and Developmental Disorders. 2019 Feb 9;49(5):2146–55.

12. Lugo J, Fadeuilhe C, Gisbert L, Setien I, Delgado M, Corrales M, et al. Sleep in adults with autism

spectrum disorder and attention deficit/hyperactivity disorder: A systematic review and meta-analysis.

European Neuropsychopharmacology. 2020 Jul.

Available from: https://doi.org/10.1016/j.euroneuro.2020.07.004

13. Singer EV, Niarchou M, Maxwell-Horn A, Hucks D, Johnston R, Sutcliffe JS, et al. Characterizing

sleep disorders in an autism-specific collection of electronic health records. Sleep Medicine [Internet].

2022 Apr 1 [cited 2022 Apr 9];92:88–95. Available from: https://doi.org/10.1016/j.sleep.2022.03.009

14. Weir E, Allison C, Ong KK, Baron-Cohen S. An investigation of the diet, exercise, sleep, BMI, and

health outcomes of autistic adults. Molecular Autism. 2021 May 8;12(1).

15. Schreck KA, Mulick JA, Smith AF. Sleep problems as possible predictors of intensified symptoms

of autism. Research in Developmental Disabilities. 2004 Feb;25(1):57–66. Available from:

https://doi.org/10.1016/j.ridd.2003.04.007

16. Chen H, Yang T, Chen J, Chen L, Dai Y, Zhang J, et al. Sleep problems in children with autism

spectrum disorder: a multicenter survey. BMC Psychiatry. 2021 Aug 16;21(1). Available from:

https://doi.org/10.1186%2Fs12888-021-03405-w

17. Díaz-Román A, Zhang J, Delorme R, Beggiato A, Cortese S. Sleep in youth with autism spectrum

disorders: systematic review and meta-analysis of subjective and objective studies. Evidence Based

Mental Health. 2018 Oct 25;21(4):146–54. Available from: https://doi.org/10.1136/ebmental-2018-

300037

18. Morgan B, Nageye F, Masi G, Cortese S. Sleep in Adults with Autism Spectrum Disorder: A

Systematic Review and Meta-Analysis of subjective and objective studies. Sleep Medicine. 2019 Aug.

Available from: https://doi.org/10.1016/j.sleep.2019.07.019

19. Han E, Scior K, Avramides K, Crane L. A Systematic Review on Autistic people's Experiences of

Stigma and Coping Strategies. Autism Research: Official Journal of the International Society for Autism

Research [Internet]. 2021 Dec 8;15(1). Available from: https://pubmed.ncbi.nlm.nih.gov/34881514/

20. Pinato L, Galina Spilla CS, Markus RP, Cruz-Machado S da S. Dysregulation of circadian rhythms

in autism spectrum disorders. Current pharmaceutical design [Internet]. 2019 [cited 2019 Nov

15];10.2174/1381612825666191102170450.

Available from: https://www.ncbi.nlm.nih.gov/pubmed/31682208

21. Corbett BA, Mendoza S, Wegelin JA, Carmean V, Levine S. Variable cortisol circadian rhythms in

children with autism and anticipatory stress. Journal of Psychiatry & Neuroscience : JPN [Internet].

2008   May   [cited   2023   Feb   15];33(3):227.   Available   from:

http://www.ncbi.nlm.nih.gov/pmc/articles/pmc2441887/

22. Gernert C, Falkai P, Falter-Wagner CM. The Generalized Adaptation Account of Autism. Frontiers

in Neuroscience. 2020 Oct 6;14. Available from: https://doi.org/10.3389/fnins.2020.534218

23. Sequeira S, Ahmed M. Meditation as a Potential Therapy for Autism: A Review. Autism Research

and Treatment. 2012;2012:1–11. Available from: https://doi.org/10.1155/2012/835847

24. Beck AT, Steer RA, Brown G. Beck Depression Inventory–II. PsycTESTS Dataset. 1996.

Available from: https://psycnet.apa.org/doi/10.1037/t00742-000

25. Allison C, Auyeung B, Baron-Cohen S. Toward Brief "Red Flags" for Autism Screening: The Short

Autism Spectrum Quotient and the Short Quantitative Checklist in 1,000 Cases and 3,000 Controls.

Journal of the American Academy of Child & Adolescent Psychiatry. 2012 Feb;51(2):202-212.e7.

Available from: https://doi.org/10.1016/j.jaac.2011.11.003

26. Greenberg DM, Warrier V, Allison C, Baron-Cohen S. Testing the Empathizing–Systemizing theory

of sex differences and the Extreme Male Brain theory of autism in half a million people. Proceedings of

the National Academy of Sciences. 2018 Nov 12;115(48):12152–7.

27. Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh sleep quality index: A

new instrument for psychiatric practice and research. Psychiatry Research [Internet]. 1989

May;28(2):193–213. Available from: https://pubmed.ncbi.nlm.nih.gov/2748771/

28. Spielberger CD, Gorsuch RL, Lushene R, Vagg PR, Jacobs GA. Manual for the State-Trait Anxiety

Inventory (Form Y1-Y2). Palo Alto, CA: Consulting Psychologists Press; 1983.

29. Fliege H, Rose M, Arck P, Levenstein S, Klapp BF. PSQ - Perceived Stress Questionnaire.

www.psycharchives.org [Internet]. 2009 Jan 1 [cited 2023 Feb 15]; Available from:

https://www.psycharchives.org/en/item/018f244d-58d8-4105-aeb2-b800443c6009

30. Weiß RH. CFT 20-R: Revision. Goettingen: Hogrefe; 2006.

31. Fraenkel JR, Wallen NE, Hyun HH. How to design and evaluate research in education. New York:

Mcgraw-Hill Humanities/Social Sciences/Languages; 2012.

32. Jachs B. The Neurophenomenology of Meditative States: Introducing Temporal Experience Tracing

to Capture Subjective Experience States and their Neural Correlates [Internet].

www.repository.cam.ac.uk.   2022   [cited   2023   Feb   15].   Available   from:

https://www.repository.cam.ac.uk/handle/1810/332709

33. Petruzzelli MG, Matera E, Giambersio D, Marzulli L, Gabellone A, Legrottaglie AR, et al.

Subjective and Electroencephalographic Sleep Parameters in Children and Adolescents with Autism

Spectrum Disorder: A Systematic Review. Journal of Clinical Medicine. 2021 Aug 30;10(17):3893.

Available from: https://doi.org/10.3390/jcm10173893

34. Backman CL, Harris SR. CASE STUDIES, SINGLE-SUBJECT RESEARCH, AND N OF 1

RANDOMIZED TRIALS. American Journal of Physical Medicine & Rehabilitation. 1999

Mar;78(2):170–6. Available from: https://doi.org/10.1097/00002060-199903000-00022

35. Arnal PJ, Thorey V, Debellemaniere E, Ballard ME, Bou Hernandez A, Guillot A, et al. The Dreem

Headband compared to polysomnography for electroencephalographic signal acquisition and sleep

staging. Sleep [Internet]. 2020 May 20 [cited 2021 May 29];43(11). Available from:

https://academic.oup.com/sleep/article/43/11/zsaa097/5841249

6. Figures

Figure 1

Overview of the Daily Schedule (Phase II)

Note. The figure illustrates an example of a daily routine in the context of the study (Phase II). A) This

timeseries shows which tasks are included, ranging from one night-sleep EEG recording to the next

night. In the morning participants fill out a sleep diary, wear the wristband for assessing physiological

data during the whole day, perform a meditation session and report about their experiences afterwards,

fill out an activity diary and use the temporal experiences tracing (TET) tool for reporting experiences

made during the day in the evening by using an app. B) This timeseries shows the belonging time points

for performing a task.

Figure 2

Screenshots from the App: Overview

A)   B)   C)

528 *Note*. A) The figure presents the menu page of the app. B) First, several questionnaires need to be filled

529 out at the beginning of the study. C) A daily checklist is included in the app which helps the participants

530 to track their progress day per day.

531 **Figure 3**

532 *Screenshots from the App: Daily Experiences and TET*

A) B)

534 *Note.* A) Tracing the dynamics of daily experiences at the end of the day is one task in the project. In

535 this example participants are asked to trace their feelings of 'Stress' throughout the day. B) Example of

536 an answer for this question. Participants can draw the graph directly in the app.

537   **Figure 4**

538   *Screenshots from the App: Meditation and TET*

A)   B)

Note. A) After a daily meditation session, participants report their subjective experiences made while

meditating. This example shows the item 'Rumination'. B) Example of an answer for this dimension.

The grey lines in the graph correspond to a gong, coming every 5 minutes in the audio, which serves as

an aid to anchor the memory of participants' experiences. The meditation lasts 20 minutes in total.

Figure 5

EEG Channels

Note. The bipolar channel montage of the portable EEG headband *(Dreem).* Adapted and reprinted with

permission from *The Neurophenomenology of Meditative States: Introducing Temporal Experience*

Tracing to capture Subjective Experience States and their Neural Correlates (PhD Thesis), by B. Jachs,

2021.