Research article: Myocardial Infarction; Early Prognostic Instrumental & Laboratory Markers: Single Cross Sectional Analysis

Running tittle: Current cardiopathologies management advances

Authors' list: Basheer Abdullah Marzoog[1,*], Natalia Igorevna Bloshkina[1], Yuli Vadimovna Maltseva[1], Vera Sergeevna Gromova [2], Anastasia Olegvona Evlakhova [3], Elena Ivanovna Gorshinina[1]

[1] National Research Mordovia State University. Address: Bolshevitskaya Street, 68, Saransk, Rep. Mordovia, Russia. Postal address: Russia, Mordovia republic, Saransk, Bolshevitskaya Street, 31, 430005.

[2] Immanuel Kant Baltic Federal University, Russia, Kaliningrad, Alexander Nevsky Street, 14. ORCID 0000-0002-8707-8876

[3] Saint Petersburg State Pediatric Medical University, , Russia, Litovskaya St., 2, St. Petersburg, 194100

[*]Corresponding author: Basheer Abdullah Marzoog, undergraduate medical school student at National Research Mordovia State University (marzug@mail.ru, +79969602820). Address: Bolshevitskaya Street, 68, Saransk, Rep. Mordovia, 430005. Postal address: Mordovia republic, Saransk, Bolshevitskaya Street, 31. ORCID: 0000-0001-5507-2413, Scopus ID: 57486338800

Competing interests: No competing interests regarding the publication.

All authors read and approved the final version of the manuscript

Abstract

Background: Cardiovascular events remain one of the frequently reported causes of death globally.

Aims: Assess the early laboratory and instrumental changes in systemic manner.

Objectives: Patients with fresh myocardial infarction that has been occurred in less than 30 days before hospitalization in the rehabilitation hospital.

Materials and methods: A single large scale retrospective cohort sectional study for patient with fresh myocardial infarction for the period of 2014-2019, data collected from Mordovia Rehabilitation Hospital. The study involved 154 patients and analyzed 76 parameters for each patient. The t-test, Pearson correlation coefficient, and ROC test has been used. For statistical analysis used Statistica program.

Results: The sample included 154 patients with history of fresh myocardial infarction. Of 154. 51(33.11%) female and 103 (66.88 %) males. chronic heart failure (CHF), which is seen in 42 (27.27273 %) patients, 101 (65.58442 %) did not have a CHF, and 11 (7.14286 %) missing data. Diabetes mellitus (DM) has been observed in 28 (18.18182 %), 118 (76.62338 %) without DM, Hypertension has been seen in 118 (76.62338 %), 28 (18.18182%) did not have hypertension, chronic kidney disease (CKD) has been seen in 11 (7.14%) patients, 131 (85.06494 %) did not have CKD, Chronic obstructive pulmonary disease (COPD) existed in 14 (9.09091%), 132 (85.71429 %) did not have COPD. Post myocardial infarction (MI) CHF has been seen in 108 (70.12987%), 42 (27.27273 %) did not have CHF. Post MI arrythmia seen in 50 (32.47%) patients, 99 (64.28571%) did not have arrythmia. Early post MI complication such as aneurysm has been seen in 12 (7.79221%), 138 (89.61039 %) did not have aneurysm, Dressler syndrome; pericarditis has been seen in 4 (2.59740%) patients, pneumonitis seen in 1 (0.64935%) patient, and pleuritis have not been seen.

Conclusions: Systemic manifestations include kidney function impairment or development of new kidney disease. Of these changes include increase in the aorta basement diameter, increase in the right ventricle wall thickness, decrease in the level of serum red blood cell, serum potassium level, and serum calcium level, with increase in the serum myoglobulin.

Others: Females are more frequently affected with post MI urinary tract infection.

Keywords: Myocardial infarction; Cardiology; Myocardiopathies; Myocardiocyte; Angiogenesis; Regeneration

Introduction

Myocardial infarction (MI) and it is associated sequalae remains the most frequently reported cause of mortality and morbidity worldwide [1]. The development of cardiovascular events is associated with several modified and non-modified risk factors [2]. The lifestyle changes have huge impact on the overall health of the cardiovascular system [3, 4]. Unfortunately, sedentary life style is seen in majority of myocardial infarction victims which is associated with development of several predisposition factors for the cardiovascular events such as dyslipidemia, insulin resistance, and central obesity or visceral obesity [5–7]. Cumulatively, these early signs of cardiovascular homeostasis disturbance are ignored to provoke the progression of dyslipidemia and insulin resistance to reach enough level to induce the emergence of cardiovascular event such as MI [8, 9].

Assessment of the early post-MI changes is of huge impact on the further therapy plan and the prognosis of MI victims. One of the early specific markers of myocardial infarction is elevation in the cardiac lactate dehydrogenase 1 and troponins including troponin I and T. However, systemic changes are rarely assessed in early pot-MI patients [10]. The role of these changes remains not clear in the determination of the prognosis of MI patients and the potential risk of development of complications such as renal failure. In the current retrospective analysis, we covered the potential role of these systemic and the regularly checked cardiac markers in the prognosis determination and prognosis of the patients.

Materials and methods

A retrospective cohort analysis for 155 patients for the period 2014-2019. The data has been collected from the republican rehabilitation hospital. The consent of the patients has been taken for gathering, analyzing and publication the material and any associated photos or graphs. The missing data has been treated as absent and removed from the analysis. All the measurement units are according to the local units used in the laboratories. There was no access to the 2020–2022 year data because of the COVID-19 issues. Pleuritis was not seen in patients and been excluded from the complications report. The presented t test results are all statistically significant, and other non-mentioned results are statistically not significant at $p<0.05$. the leukocytes and the red blood cell level is always mentioned without the $*10^9$ or $*10^{12}$, respectively. We could not assess the risk hazard for the estimated parameters and the survival rate, limitations of the study.

T-test, Pearson correlation coefficient, and ROC analysis have been used using Statistica 12 program (StatSoft, Inc. (2014). STATISTICA (data analysis software system), version 12. www.statsoft.com.).

Results

The sample included 154 patients with history of fresh myocardial infarction. Of 154. 51(33.11%) female and 103 (66.88 %) males. In the sample, 131 (85.06494 %) lives in the city, 17 (11.03896 %) lives in the village, and 5 (3.24675%) lives in the town.

Majority of the patient had a concomitant disease prior to their myocardial infarction. 132 (85.71429%) had a concomitant disease, 18 (11.68831%) did not have a concomitant disease, and 4 (2.59740 %) missing data. Of the concomitant diseases is chronic heart failure (CHF), which is seen in 42 (27.27273 %) patients, 101 (65.58442 %) did not have a CHF, and 11 (7.14286 %) missing data. Diabetes mellitus (DM) has been observed in 28 (18,18182 %), 118 (76.62338 %) without DM, and 8 (5.19481 %) missing data. Hypertension has been seen in 118 (76.62338 %), 28 (18.18182%) did not have hypertension, and 8 (5.19481%) missing data. Chronic kidney disease (CKD) has been seen in 11 (7.14%) patients, 131 (85.06494 %) did not have CKD, and 12 (7.79221 %) missing data. Chronic obstructive pulmonary disease (COPD) existed in 14 (9.09091%), 132 (85.71429 %) did not have COPD, and 8 (5.19481 %) missing data. Disrupted lipid profile was present in 21 (13.63636 %), 121 (78.57143 %) did not have dyslipidemia, and 12 (7.79221 %) missing data. Varicose vein is coexisted in 3 (1.94805%), 137 (88.96104%) did not have varicose vein, and 14 (9.09091%) missing data.

In terms of post myocardial infarction complications, 145 (94.15584 %) patients had complication, 5 (3.24675%) did not have complication, and 4 (2.59740%) missing data. Post myocardial infarction (MI) CHF has been seen in 108 (70.12987%), 42 (27.27273 %) did not have CHF, and 4 (2.59740 %) missing data. Post MI arrythmia seen in 50 (32.47%) patients, 99 (64.28571%) did not have arrythmia, 5 (3.24675%) missing data. Early post MI complication such as aneurysm has been seen in 12 (7.79221%), 138 (89.61039 %) did not have aneurysm, and 4 (2.59740%) missing data. Late post MI complications including Dressler syndrome including the pericarditis has been seen in 4 (2.59740%) patients, 146 (94.80519%) did not have pericarditis, and 4 (2.59740 %) missing data. Additional components of the Dressler syndrome include pneumonitis, which is seen in 1 (0.64935%) patient, 148 (96.10390%) did not have pneumonitis, and 5 (3.24675 %) missing data. Whereas, the third component of the Dressler syndrome, pleuritis, have not been seen, therefore, it has not been reported.

Interestingly, 43 (27.92208%) patients had re-infarct, 107 (69.48052 %) were for the first time, and 4 (2.59740 %) missing data. Assessment of the activity level of tissue-type plasminogen activator (t-PA) through d-dimer level in the serum, it was elevated in 7

(4.54545%) patients, 5 (3.24675%) normal range, other 142 (92.20779%) did not measure their d-dimer level.

The primary descriptive statistics demonstrated in (*Table*) 1.

Table 1: The primary descriptive statistics of the sample. Abbreviations: EF; ejection fraction, EDV; end diastolic volume, LV; left ventricle, LA; left atrium, RV; right ventricle, RA; right atrium, Hb; hemoglobin, RBC; red blood cell, ESR; erythrocytes sedimentation rate, CKP-MB; creatinine phosphokinase-MB fraction, LDH; lactate dehydrogenase, ALT; alanine transaminase, AST; aspartate transaminase, LDL; low density lipoprotein, TAG; triacyl glycerides, APTT; activated prothrombin time, PT; prothrombin time.

Variable	Descriptive Statistics					
	Valid N	% Valid obs.	Mean	Median	Minimum	Maximum
Hospitalization days	153	99.35065	14.53	15.000	0.0000	24
Age (years)	152	98.70130	61.64	62.500	30.0000	86
Body weight (Kg)	31	20.12987	82.26	85.000	62.0000	112
Height (cm)	31	20.12987	170.58	172.000	155.0000	185
BMI (Kg/m^2)	31	20.12987	28.56	27.888	21.7993	41
HR (beat/mint)	148	96.10390	79.82	78.000	53.0000	130
SBP (mmHg)	148	96.10390	139.57	140.000	60.0000	260
DBP (mmHg)	148	96.10390	84.05	80.000	40.0000	130
Aorta diameter (mm)	144	93.50649	35.21	35.000	20.0000	46
EF (%)	142	92.20779	54.55	55.000	25.0000	80
EDV (ml)	113	73.37662	134.30	129.000	62.0000	234
LV (mm)	132	85.71429	52.35	51.150	10.0000	89
LA (mm)	133	86.36364	40.09	40.000	28.0000	62
RV (mm)	144	93.50649	26.75	26.250	5.0000	44
RA (mm)	130	84.41558	35.73	36.000	24.0000	49
Hb	143	92.85714	137.10	138.000	85.0000	170
Leukocytes (/L)	141	91.55844	8.82	8.300	2.8000	18
RBC (/L)	118	76.62338	5.12	4.505	2.6000	24
Lymphocytes (%)	130	84.41558	25.51	25.500	2.0000	54
Thrombocytes (/L)	111	72.07792	206.76	200.000	83.0000	506
ESR (mm/hr)	138	89.61039	19.04	15.500	2.0000	60

Variable	Descriptive Statistics					
	Valid N	% Valid obs.	Mean	Median	Minimum	Maximum
Ketones (mmol/l)	1	0.64935	0.50	0.500	0.5000	1
Urine proteins (mmol/l)	137	88.96104	5.27	0.000	0.0000	150
Leukocytes (microscopically)	142	92.20779	2.64	1.000	0.0000	50
Epithelial cells (microscopically)	136	88.31169	1.86	1.000	0.0000	20
Troponin I (ng/ml)	126	81.81818	4.74	1.205	0.0000	33
Myoglobulin (ng/ml)	3	1.94805	111.30	71.000	63.0000	200
CKP-MB 1 (U/l)	129	83.76623	97.01	29.000	6.0000	2881
CKP-ÌÂ 2 (U/l)	71	46.10390	79.26	42.000	6.0000	508
LDH 1 (U/l)	108	70.12987	652.30	475.500	30.0000	3900
LDH 2 (U/l)	56	36.36364	21640.80	492.000	97.0000	1177
INR (Sec)	125	81.16883	1.25	1.000	0.8000	18
Total bilirubin (mmol/l)	108	70.12987	16.53	17.680	4.4200	36
ALT (mmol/l)	140	90.90909	42.65	28.000	1.1900	361
AST (mmol/l)	140	90.90909	55.57	30.000	5.1000	423
Total cholesterol (mmol/l)	141	91.55844	5.23	5.100	2.4000	9
LDL (mmol/l)	138	89.61039	23.20	3.240	0.9200	2766
TAG (mmol/l)	140	90.90909	3.65	1.565	0.2700	138
Na (mmol/l)	122	79.22078	139.76	142.750	1.7600	151
K (mmol/l)	126	81.81818	6.02	4.060	1.0000	92
Creatinine (µmol/L)	142	92.20779	81.56	84.000	0.0000	176
Ca (mmol/l)	115	74.67532	1.69	1.180	0.2500	38
Urea (mmol/l)	139	90.25974	7.79	6.700	2.5000	93
Glucose (mmol/l)	135	87.66234	6.86	6.100	4.0000	19
APTT (Sec)	74	48.05195	27.31	25.500	15.0000	76
Fibrinogen (ng/ml)	117	75.97403	3386.85	3500.000	300.0000	7500

	Descriptive Statistics					
Variable	Valid N	% Valid obs.	Mean	Median	Minimum	Maximum
PT (Sec)	1	0.64935	94.00	94.000	94.0000	94

Males are affected in myocardial infarction earlier in comparing to female, the cut-off point of age is 67 years old. (*Figure 1*)

In male group the mean age is 59.76 years old and in female group the mean age is 65.46 years old, which is statistically significant difference, t value 3.30972, p<0.001170. Moreover, mean height of the male is 172.57 cm, and in female 164.88 cm, which is statistically significant difference, t value -3.26225, p< 0.002829. On echocardiogram, in male group, the mean diameter of the aorta is 36.23 mm, in female group, the mean diameter of the aorta is 33.04 mm, which is statistically significant difference, t value -4.24952, p<0.000039. In male group, the mean right atrium wall thickness on echocardiogram equal to 36.26 mm, in female group equal to 34.56 mm, which is statistically significant difference, t value -2.06826, p< 0.040630. Mean hemoglobin level in male was 140.26 and in female was 130.66, which is statistically significant difference, t value -3.71265, p<0.000295. In urine analysis, in male group, mean number of leukocytes in urine was 1.52 under microscope and in female group was 4.91, which is statistically significant difference, t value 3.11688, p< 0.002218. Moreover, in urine analysis, number of the epithelial cells in the male 1.36 under microscope and in female group 2.91, which is statistically significant difference, t value 3.12012, p< 0.002214. In male group, the mean myoglobulin level was 67.00 ng/ml and in female was 199.90 ng/ml, which is statistically significant difference, t value 19.18246, p< 0.033158.

Patients with concomitant diseases have a mean heart rate (HR) equal to 80.82 beat/mint and patients without concomitant disease have mean HR equal to 72.39 beat/mint, which is statistically significant difference, t value 2.05541, p< 0.041633. Additionally, patients with concomitant diseases have a mean right ventricle wall thickness of 27.13 mm and patients without concomitant disease have a mean right ventricle wall thickness of 23.85 mm, which is statistically significant difference, t value 2.79969, p< 0.005833. Moreover, patients with concomitant disease have a mean protein in urine 3.18 mg/l, and the patients without concomitant disease had a mean level of urine protein 21.25 mg/l, which is statistically significant difference, t value -3.10080, p< 0.002353. Interestingly, patients with concomitant disease had lower calcium level a mean equal to 1.31 mmol/l and the patients without

concomitant disease had a mean calcium level 3.84 mmol/l, which is statistically significant difference, t value -2.67394, p< 0.008608. (*Figure 2*)

Patients with CHF have a mean thrombocytes level equal to 185.26 /l, whereas, patients without CHF have a mean thrombocytes level 216.24 /l, which is statistically significant difference, t value - 2.15343, p< 0.033573.

Mean glucose level in diabetic patients equal to 9.93 mmol/l and in non-diabetic patients equal to 6.13 mmol/l, which is statistically significant difference, t value -7.9724, p< 0.0001. Coexistence of DM in patients increases the mean left ventricle wall thickness to equal 55.60 mm, whereas, in patients without DM, the mean left ventricle wall thickness to equal 51.42 mm, which is statistically significant difference, t value -2.5388, p< 0.012341. Mean erythrocytes sedimentation rate (ESR) in DM patients equal to 23.60 mm/hr, and equal to 17.43 mm/hr in non-diabetic patients, which is statistically significant difference, t value -2.2236, p< 0.027887. Mean creatinine phosphokinase-MB fraction is dramatically higher in DM patients 233.36 U/L and lower in non-diabetic patients 66.05 U/l, which is statistically significant difference, t value --2.6267, p< 0.009726. Diabetic patients experience high level of triacyl glyceride (TAG) 2.17 mmol/l, whereas, non-diabetic patients had a lower TAG level 1.66 mmol/l, which is statistically significant difference, t value -2.5633, p< 0.011481.

Hypertensive patients have a low mean level of red blood cells (RBC) level 4.55 /l in compare to non-hypertensive patients 7.7, which is statistically significant difference, t value -4.63276, p< 0.000010. In hypertensive patients, the mean LDH 1 is 608.21 U/l, INR is 1.05, sodium is 141.70 mmo/l, potassium is 4.04 mmol/l, and calcium is 1.14 mmol/l, whereas, in non-hypertensive patients, the mean level of LDH 1 is 908.8 U/l, INR is 2.1, sodium is 130.6 mmol/l, potassium is 14.8 mmol/l, calcium is 3.9 mmol/l, which is statistically significant difference; t value -2.07903, -3.04226, 2.51546, -3.62102, -3.19601; p< 0.040123, 0.002890, 0.013268, 0.000432, 0.001828, respectively.

CKD patients have high mean troponin I level equal to 22.70 ng/ml in compare to non-CKD patients 3.72 ng/ml, which is statistically significant difference, t value -6.84999, p< 0.0001. Also, CKD patients have higher mean level of CPK-MB fraction equal to 475.61 U/l in compare to non-CKD patients equal to 78.38 U/l, which is statistically significant difference, t value -3.63297, p< 0.000415. Mean creatinine level is higher in CKD patients 109.81 µmol/L comparing to non-CKD patients, which is statistically significant difference, t value -3.63297, p< 0.000415.

Interestingly, COPD patients have a mean left ventricle wall thickness of 48.46 mm, which is less than non-COPD patients 52.73 mm, which is statistically significant difference, t value

1.99093, p< 0.048653. However, COPD patients have a mean right atrium wall thickness of 39.55 mm in compare to non-COPD patients 35.41 mm, which is statistically significant difference, t value -3.06986, p< 0.002632. Moreover, COPD patients have a mean RBC level equal to 145.29 in compare to non-COPD patients 136.39, which is statistically significant difference, t value -2.11854, p< 0.035948.

Patients with dyslipidemia have mean troponin I 9.4 ng/ml, whereas, patients without dyslipidemia have mean troponin I 3.64 ng/ml, which is statistically significant difference, t value -2.99785, p< 0.003316. Additionally, patients with dyslipidemia have mean total bilirubin 19.3 mmol/l, and 15.72 mmol/l in patients without dyslipidemia, which is statistically significant difference, t value -2.12114, p< 0.036358. Moreover, patients with dyslipidemia have a mean sodium serum level 132.3 mmol/l and a mean serum potassium level 8.4 mmol/l, whereas, in patients without dyslipidemia, the mean sodium serum level 142.24 mmol/l and a mean serum potassium level 4.07 mmol/l, which is statistically significant difference; t value 2.74875, -2.09792; p< 0.006966, 0.038083.

The mean leukocytes level in patients with varicose veins is 12.07 /l, whereas, in patients without varicose veins, the mean leukocytes level is 8.71 /l, which is statistically significant difference; t value -1.99832, p< 0.047801. The mean RBC level in patients with varicose veins is 9.87 /l, whereas, in patients without varicose veins, the mean RBC level is 4.56 /l, which is statistically significant difference; t value -6.88122, p< 0.000000. The mean ESR level in patients with varicose veins is 37.67 mm/hr, whereas, in patients without varicose veins, the mean ESR level is 18.25 mm/hr, which is statistically significant difference; t value -2.64654, p< 0.009161.

In terms of complications, patients with complications have been hospitalized for 14.65 days, whereas, patients without complications hospitalized for 11.40 days, which is statistically significant difference; t value 2.05492, p< 0.041657. In patients with complications, the mean aorta diameter 35.38 mm, whereas, in patients without complications, the mean aorta diameter 29.25 mm, which is statistically significant difference; t value 2.77454, p< 0.006278. In patients with complications, the mean right ventricle wall thickness 26.89 mm, whereas, in patients without complications, the mean right ventricle wall thickness 21.25mm, which is statistically significant difference; t value 2.44283, p< 0.015809. In patients with complications, the mean RBC level 4.87 /l, whereas, in patients without complications, the mean RBC level 12.40 /l, which is statistically significant difference; t value -5.37900, p< 0.0001. In patients with complications, in general urine analysis, microscopically, the mean urea proteins level 4.30, whereas, in patients without complications, the mean urea proteins

level 50.00, which is statistically significant difference; t value -3.61867, p< 0.000419. In patients with complications, the mean potassium level 5.37 mmol/l and mean calcium level 1.58 mmol/l, whereas, in patients without complications, the mean potassium level 32.99 mmol/l and mean calcium level is 5.99 mmol/l, which is statistically significant difference; t value -3.75574, -2.05930; p< 0.000265, 0.041784, respectively.

Patients with post-MI CHF have mean myoglobulin level 199.90 ng/ml, whereas, in patients without post-MI CHF, the mean myoglobulin level 67.00ng/ml, which is statistically significant difference; t value 19.18246, p< 0.033158.

Patients with history of previous infarct have creatinine 90.48 µmol/l in compare to patients without reinfarction 78.01 µmol/l, which is statistically significant difference; t value 2.26506, p< 0.025056.

The mean age of patients with arrythmia is 65.58 years old, whereas, the mean age of patients without arrythmia is 59.68 years old, which is statistically significant difference; t value 3.32256, p< 0.001129. The mean height of patients with arrythmia is 173.67 cm, whereas, the mean height of patients without arrythmia is 168.61 cm, which is statistically significant difference; t value 2.23386, p< 0.034675. The mean ejection fraction (EF) of patients with arrythmia is 50.40 %, whereas, the mean EF of patients without arrythmia is 56.43 %, which is statistically significant difference; t value -3.81906, p< 0.000202. The mean left atrium wall thickness of patients with arrythmia is 42.14 mm, whereas, the mean left atrium wall thickness of patients without arrythmia is 39.12 mm, which is statistically significant difference; t value 3.00570, p< 0.003189. The mean right atrium wall thickness of patients with arrythmia is 37.36 mm, whereas, the mean right atrium wall thickness of patients without arrythmia is 34.95 mm, which is statistically significant difference; t value 2.96567, p< 0.003614. The mean CKP-MB of patients with arrythmia is 181.82 U/l, whereas, the mean CKP-MB of patients without arrythmia is 56.68 U/l, which is statistically significant difference; t value 2.34423, p< 0.020643. The mean lactate dehydrogenase (LDH) serum level of patients with arrythmia is 888.37 U/l, whereas, the mean lactate dehydrogenase serum level of patients without arrythmia is 533.49 U/l, which is statistically significant difference; t value 3.16430, p< 0.002039. The mean aspartate transaminase serum level of patients with arrythmia is 71.53 mmol/l, whereas, the mean lactate dehydrogenase serum level of patients without arrythmia is 47.38 mmol/l, which is statistically significant difference; t value 2.23750, p< 0.026893.

The mean EF of patients with aneurysm is 43.00 %, whereas, the mean EF of patients without aneurysm is 55.59 %, which is statistically significant difference; t value 4.57958, p<

0.000010. The mean leukocytes level in patients with aneurysm is 9.70 /l, whereas, the mean leukocytes of patients without aneurysm is 2.11 /l, which is statistically significant difference; t value -3.83162, p< 0.000192. The mean epithelial cell level in general urine analysis in patients with aneurysm is 5.40, whereas, the mean epithelial cell level in patients without aneurysm is 1.58, which is statistically significant difference; t value -4.41313, p< 0.000021.

The mean thickness of the left atrium of patients with pericarditis is 45.5mm, whereas, the mean thickness of the left atrium of patients without pericarditis is 39.90 mm, which is statistically significant difference; t value -2.01849, p< 0.045598. The mean creatinine level of patients with pericarditis is 52.5 µmol/L, whereas, the mean creatinine level of patients without pericarditis is 82.39 µmol/L, which is statistically significant difference; t value 1.98986, p< 0.048568.

Patients with pneumonitis have mean aorta diameter of 20.00 mm, whereas, the mean aorta diameter of patients without pneumonitis is 35.32 mm, which is statistically significant difference; t value 3.55083, p< 0.000524. Patients with pneumonitis have mean serum leukocytes level of 14.80 /l, whereas, the mean serum leukocytes level of patients without pneumonitis is 8.70 /l, which is statistically significant difference; t value -2.11170, p< 0.036527. Patients with pneumonitis have mean ESR level of 48.00 mm/hr, whereas, the mean ESR level of patients without pneumonitis is 18.47 mm/he, which is statistically significant difference; t value -2.36894, p< 0.019267. Moreover, patients with pneumonitis have mean urine leukocyte level of 22.00, whereas, the mean urine leukocyte level of patients without pneumonitis is 2.53, which is statistically significant difference; t value -3.15623, p< 0.001963. The mean CKP-MB in patients with pneumonitis 1164.00 U/l, whereas, the mean CKP-MB of patients without pneumonitis is 89.60 U/l, which is statistically significant difference; t value -3.92161, p< 0.000144.

The mean aorta diameter, EF, EDV, troponin I, CKP-MB, APTT in patients with elevated troponin T is 28.00 mm, 36.00 %, 62.00 ml, 33.00 ng/ml, 2881.00 U/l, and 49.10 sec, respectively. Whereas, the mean aorta diameter, EF, EDV, troponin I, CKP-MB, APTT in patients without elevation in troponin T is 35.31 mm, 54.68%, 134.95 ml, 4.51 ng/ml, 75.26 U/l, 27.01 sec, respectively, which is statistically significant difference; t value 2.3478, 2.0136, 2.0775, -3.6144, -19.3413, -2.5572, p< 0.020266, 0.045967, 0.040063, 0.000436, 0.000000, 0.012661, respectively.

The mean age of patients with elevated D-dimer level is 68.14 years old, and mean age of patients with normal D-dimer is 56.60 years old, which is statistically significant difference; t value 2.25820, p< 0.047513. The mean wall thickness of the right ventricle of patients with

elevated D-dimer level is 25.93 mm, and mean wall thickness of the right ventricle of patients with normal D-dimer is 29.40 mm, which is statistically significant difference; t value -2.35428, p< 0.040345.

Comparing the dependent variable shows no statistical differences. (*Figure 3*)

In terms of correlations, a full table shows the correlation between all the continuous variables is presented. (*Table 2*)

An interesting correlation between the LDL level and the diameter of the aorta basement has been seen. (*Figure 4*) It has been shown that when the left ventricle wall thickness increases the EF dramatically decrease. (*Figure 4*)

Discussion

Post-MI changes include early and late pathophysiological changes, early post-MI complications include ventricular rupture, aneurism, arrythmia, sudden cardiac arrest and death [11–13]. However, the late changes include pericarditis, pneumonitis, pleuritis, and post infarction cardiosclerosis and dyskinesia [14, 15]. Such manifestations are associated with some laboratory and instrumental changes. Assessment of these changes help prevention or slow down, or at least make a prognosis for the MI patients. Low level of thrombocytes in patients with concomitant CHF return to the use of anti-aggregatnts and anticoagulants for a long time.

High leukocytes level in diabetic patients endorses the hypothesis of chronic mild inflammatory syndrome in patients with metabolic syndrome [16–19]. The high TAG values in diabetic patients is due to the lipogenesis from high glucose serum levels and metabolic disorders associated with the chronic lipid peroxidation and biogenesis disorders [16, 20].

The dramatic increase in the wall thickness of patients with COPD is an indicator of a stable stage of pulmonary heart, which is associated with pulmonary hypertension and blood stagnation in the pulmonary circulation [21, 22]. Typically, these patients experience cough and shortness of breath on the background of distress syndrome, fluid accumulation in the pulmonary tissue, and impaired ability of the pneumocytes for gas exchange. As a compensation for the respiratory hypoxemia, the elevation in the levels of the erythropoietin and the further erythropoiesis elevation, which is demonstrated in the high level of Hb in COPD patients.

Leukocytosis is seen in patients with varicose vein is indicator of the persistent inflammation with elevation in the tissue hypoxia and further elevation in hypoxia inducing factor to stimulate the erythropoietin production and further increasing in the RBC level in patients

with varicose veins [23, 24]. This persistent inflammation is confirmed by the elevation in the ESR level in patients with varicose veins [25–28].

Post MI complications reported in 4/5 of the sample and these complications associated with some laboratory and instrumental changes. Suggesting that post MI laboratory changes are not limited to the local myocardial infarction changes to involves systemic manifestations such as kidney function impairment or development of new kidney disease. Of these changes include increase in the aorta basement diameter, increase in the right ventricle wall thickness, decrease in the level of RBC, serum potassium level, and serum calcium level, with increase in the serum myoglobulin.

Re-infarct worsens the kidney function and induces renal failure [29–32]. Aging is a risk factor for MI and arrythmia development. Arrythmia increases CPK-MB, LDH, AST, wall thickness of both atriums, and decrease EF. The higher the troponin level the less EF and the poorest further sequalae.

Early post-MI aneurysm significantly impairs the cardiac output and further reduces the systemic systolic blood pressure. These changes are associated with the reduction in the blood flow in to the renal afferent fibers, which are sensed by the juxtaglomerular apparatus that leads to activation the renin angiotensin aldosterone system with the anti-diuretic hormone. Therefore, reduction in the kidney function is seen in MI patients. Additionally, elevation of creatinine in aneurysm patients is typical for impaired kidney function particularly filtration due to hypoperfusion of the glomeruli.

Patients with pneumonitis are having a high leukocyte in serum and in the urine, high ESR, and high CKP-MB fraction. However, these patients experience small diameter of the aorta basement.

Conclusions

The early prognostic value of the instrumental and laboratory changes is complication dependent. With age increase the level of D-dimer and thrombocytes. The higher the troponin level the less EF and the poorest further sequalae. Systemic manifestations include kidney function impairment or development of new kidney disease. Of these changes include increase in the aorta basement diameter, increase in the right ventricle wall thickness, decrease in the level of serum red blood cell, serum potassium level, and serum calcium level, with increase in the serum myoglobulin.

List of abbreviations

CHF; chronic heart failure, DM; Diabetes mellitus, CKD; Chronic kidney disease, COPD; Chronic obstructive pulmonary disease, MI; myocardial infarction, HR; heart rate, ESR; erythrocytes sedimentation rate, RBC; red blood cells

Declarations

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

The study approved by the National Research Mordovia State University6 Russia, from "Ethics Committee Requirement N8/2 from 30.06.2022".

HUMAN AND ANIMAL RIGHTS

No animals were used in this research. All human research procedures followed were in accordance with the ethical standards of the committee responsible for human experimentation (institutional and national), and with the Helsinki Declaration of 1975, as revised in 2013.

CONSENT FOR PUBLICATION

Written informed consent was obtained from the participants for publication of study results and any accompanying images.

AVAILABILITY OF DATA AND MATERIALS

Not applicable.

FUNDING

None.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

Declared none.

References

[1]   Tsao, C. W.; Aday, A. W.; Almarzooq, Z. I.; Anderson, C. A. M.; Arora, P.; Avery, C. L.; Baker-Smith, C. M.; Beaton, A. Z.; Boehme, A. K.; Buxton, A. E.; et al. Heart Disease and Stroke Statistics-2023 Update: A Report From the American Heart Association. *Circulation*, **2023**. https://doi.org/10.1161/CIR.0000000000001123.

[2]   Yusuf, S.; Joseph, P.; Rangarajan, S.; Islam, S.; Mente, A.; Hystad, P.; Brauer, M.;

Kutty, V. R.; Gupta, R.; Wielgosz, A.; et al. Modifiable Risk Factors, Cardiovascular Disease and Mortality in 155,722 Individuals from 21 High-, Middle-, and Low-Income Countries. *Lancet (London, England)*, **2020**, *395* (10226), 795. https://doi.org/10.1016/S0140-6736(19)32008-2.

[3]   Ward, Z. J.; Bleich, S. N.; Cradock, A. L.; Barrett, J. L.; Giles, C. M.; Flax, C.; Long, M. W.; Gortmaker, S. L. Projected U.S. State-Level Prevalence of Adult Obesity and Severe Obesity. *N. Engl. J. Med.*, **2019**, *381* (25), 2440–2450. https://doi.org/10.1056/NEJMSA1909301.

[4]   Grundy, S. M. A Changing Paradigm for Prevention of Cardiovascular Disease: Emergence of the Metabolic Syndrome as a Multiplex Risk Factor. *Eur. Hear. J. Suppl.*, **2008**, *10* (suppl_B), B16–B23. https://doi.org/10.1093/EURHEARTJ/SUM044.

[5]   Abdullah Marzoog, B. Adaptive and Compensatory Mechanisms of the Cardiovascular System and Disease Risk Factors in Young Males and Females. *New Emirates Med. J.*, **2022**, *04*. https://doi.org/10.2174/04666221128110145.

[6]   Bays, H. E.; Taub, P. R.; Epstein, E.; Michos, E. D.; Ferraro, R. A.; Bailey, A. L.; Kelli, H. M.; Ferdinand, K. C.; Echols, M. R.; Weintraub, H.; et al. Ten Things to Know about Ten Cardiovascular Disease Risk Factors. *Am. J. Prev. Cardiol.*, **2021**, *5*, 100149. https://doi.org/10.1016/J.AJPC.2021.100149.

[7]   Powell-Wiley, T. M.; Poirier, P.; Burke, L. E.; Després, J. P.; Gordon-Larsen, P.; Lavie, C. J.; Lear, S. A.; Ndumele, C. E.; Neeland, I. J.; Sanders, P.; et al. Obesity and Cardiovascular Disease: A Scientific Statement From the American Heart Association. *Circulation*, **2021**, *143* (21), e984. https://doi.org/10.1161/CIR.0000000000000973.

[8]   Tian, X.; Chen, S.; Wang, P.; Xu, Q.; Zhang, Y.; Luo, Y.; Wu, S.; Wang, A. Insulin Resistance Mediates Obesity-Related Risk of Cardiovascular Disease: A Prospective Cohort Study. *Cardiovasc. Diabetol.*, **2022**, *21* (1). https://doi.org/10.1186/S12933-022-01729-9.

[9]   Lavie, C. J.; Ozemek, C.; Carbone, S.; Katzmarzyk, P. T.; Blair, S. N. Sedentary Behavior, Exercise, and Cardiovascular Health. *Circ. Res.*, **2019**, *124* (5), 799–815. https://doi.org/10.1161/CIRCRESAHA.118.312669.

[10]   Lorber, D. Importance of Cardiovascular Disease Risk Management in Patients with Type 2 Diabetes Mellitus. *Diabetes, Metab. Syndr. Obes.*, **2014**, *7*, 169–183. https://doi.org/10.2147/DMSO.S61438.

[11]   Damluji, A. A.; Van Diepen, S.; Katz, J. N.; Menon, V.; Tamis-Holland, J. E.; Bakitas,

M.; Cohen, M. G.; Balsam, L. B.; Chikwe, J. Mechanical Complications of Acute Myocardial Infarction: A Scientific Statement From the American Heart Association. *Circulation*, **2021**, *144* (2), E16–E35. https://doi.org/10.1161/CIR.0000000000000985.

[12] Konstam, M. A.; Kramer, D. G.; Patel, A. R.; Maron, M. S.; Udelson, J. E. Left Ventricular Remodeling in Heart Failure: Current Concepts in Clinical Significance and Assessment. *JACC Cardiovasc. Imaging*, **2011**, *4* (1), 98–108. https://doi.org/10.1016/J.JCMG.2010.10.008.

[13] Ibanez, B.; James, S.; Agewall, S.; Antunes, M. J.; Bucciarelli-Ducci, C.; Bueno, H.; Caforio, A. L. P.; Crea, F.; Goudevenos, J. A.; Halvorsen, S.; et al. 2017 ESC Guidelines for the Management of Acute Myocardial Infarction in Patients Presenting with ST-Segment Elevation. *Eur. Heart J.*, **2018**, *39* (2), 119–177. https://doi.org/10.1093/EURHEARTJ/EHX393.

[14] Marzoog, B. A.; Alexandrovich, K. O.; Nikolaevich, T. M.; Vladimirovich, K. S. Post-Coronary Artery Bypass Graft Complications; Potential Causes and Risk Factors. *medRxiv*, **2023**, 2022.12.29.22284005. https://doi.org/10.1101/2022.12.29.22284005.

[15] Elbadawi, A.; Elgendy, I. Y.; Mahmoud, K.; Barakat, A. F.; Mentias, A.; Mohamed, A. H.; Ogunbayo, G. O.; Megaly, M.; Saad, M.; Omer, M. A.; et al. Temporal Trends and Outcomes of Mechanical Complications in Patients With Acute Myocardial Infarction. *JACC Cardiovasc. Interv.*, **2019**, *12* (18), 1825–1836. https://doi.org/10.1016/J.JCIN.2019.04.039.

[16] Marzoog, B. Lipid Behavior in Metabolic Syndrome Pathophysiology. *Curr. Diabetes Rev.*, **2022**, *18* (6), e150921196497. https://doi.org/10.2174/1573399817666210915101321.

[17] Marzoog, B. A. Tree of Life; Endothelial Cell Physiopathology, the Good Guy Is a Partner in Crime! *Anat. Cell Biol.*, **2023**.

[18] Marzoog, B. A.; Vlasova, T. I. Beta-Cell Autophagy under the Scope of Hypoglycemic Drugs; Possible Mechanism as a Novel Therapeutic Target. *Obe. Metab.*, **2022**, *18* (4), 465–470. https://doi.org/10.14341/omet12778.

[19] Marzoog, B. A. The Metabolic Syndrome Puzzles; Possible Pathogenesis and Management. *Curr. Diabetes Rev.*, **2022**, *18*. https://doi.org/10.2174/1573399818666220429100411.

[20] Glowinski, J.; Glowinski, S. Generation of Reactive Oxygen Metabolites by the Varicose Vein Wall. *Eur. J. Vasc. Endovasc. Surg.*, **2002**, *23* (6), 550–555. https://doi.org/10.1053/EJVS.2002.1656.

[21]   Marzoog, B. A. Pulmonary Fibrosis; Risk Factors and Molecular Triggers, Insight for Neo Therapeutic Approach. *Curr. Respir. Med. Rev.*, **2022**, *18* (4), 257–264. https://doi.org/10.2174/1573398X18666220806124019.

[22]   Marzoog, B. A. Local Lung Fibroblast Autophagy in the Context of Lung Fibrosis Pathogenesis. *Curr. Respir. Med. Rev.*, **2023**, *19* (1), 6–11. https://doi.org/10.2174/1573398X19666221130141600.

[23]   Poredos, P.; Spirkoska, A.; Rucigaj, T.; Fareed, J.; Jezovnik, M. K. Do Blood Constituents in Varicose Veins Differ From the Systemic Blood Constituents? **2015**. https://doi.org/10.1016/j.ejvs.2015.04.031.

[24]   Lattimer, C. R.; Kalodiki, E.; Geroulakos, G.; Hoppensteadt, D.; Fareed, J. Are Inflammatory Biomarkers Increased in Varicose Vein Blood? *Clin. Appl. Thromb.*, **2016**, *22* (7), 656–664. https://doi.org/10.1177/1076029616645330/ASSET/IMAGES/LARGE/10.1177_1076 029616645330-FIG2.JPEG.

[25]   Bontekoe, E.; Brailovsky, Y.; Hoppensteadt, D.; Bontekoe, J.; Siddiqui, F.; Newman, J.; Iqbal, O.; Reed, T.; Fareed, J.; Darki, A. Upregulation of Inflammatory Cytokines in Pulmonary Embolism Using Biochip-Array Profiling. *Clin. Appl. Thromb.*, **2021**, *27*. https://doi.org/10.1177/10760296211013107.

[26]   Dzieciuchowicz, L.; Espinosa, G.; Páramo, J. A. Increased Levels of Metalloproteinase 10 and Hemostatic Markers in Patients with Noncomplicated Primary Varicose Veins. *Clin. Appl. Thromb.*, **2015**, *21* (7), 684–687. https://doi.org/10.1177/1076029613517167.

[27]   Gwozdzinski, L.; Pieniazek, A.; Bernasinska-Slomczewska, J.; Hikisz, P.; Gwozdzinski, K. Alterations in the Plasma and Red Blood Cell Properties in Patients with Varicose Vein: A Pilot Study. *Cardiol. Res. Pract.*, **2021**, *2021*. https://doi.org/10.1155/2021/5569961.

[28]   Jezovnik, M. K.; Fareed, J.; Poredos, P. Patients with a History of Idiopathic Deep Venous Thrombosis Have Long-Term Increased Levels of Inflammatory Markers and Markers of Endothelial Damage. *Clin. Appl. Thromb.*, **2017**, *23* (2), 124–131. https://doi.org/10.1177/1076029616670259.

[29]   Wang, C.; Pei, Y. Y.; Ma, Y. H.; Ma, X. L.; Liu, Z. W.; Zhu, J. H.; Li, C. S. Risk Factors for Acute Kidney Injury in Patients with Acute Myocardial Infarction. *Chin. Med. J. (Engl).*, **2019**, *132* (14), 1660. https://doi.org/10.1097/CM9.0000000000000293.

[30] Shacham, Y.; Leshem-Rubinow, E.; Steinvil, A.; Assa, E. Ben; Keren, G.; Roth, A.; Arbel, Y. Renal Impairment According to Acute Kidney Injury Network Criteria among ST Elevation Myocardial Infarction Patients Undergoing Primary Percutaneous Intervention: A Retrospective Observational Study. *Clin. Res. Cardiol.*, **2014**, *103* (7), 525–532. https://doi.org/10.1007/S00392-014-0680-8.

[31] Tsai, T. T.; Patel, U. D.; Chang, T. I.; Kennedy, K. F.; Masoudi, F. A.; Matheny, M. E.; Kosiborod, M.; Amin, A. P.; Messenger, J. C.; Rumsfeld, J. S.; et al. Contemporary Incidence, Predictors, and Outcomes of Acute Kidney Injury in Patients Undergoing Percutaneous Coronary Interventions: Insights from the NCDR Cath-PCI Registry. *JACC Cardiovasc. Interv.*, **2014**, *7* (1), 1–9. https://doi.org/10.1016/J.JCIN.2013.06.016.

[32] Maksimczuk, J.; Galas, A.; Krzesiński, P. What Promotes Acute Kidney Injury in Patients with Myocardial Infarction and Multivessel Coronary Artery Disease—Contrast Media, Hydration Status or Something Else? *Nutr. 2023, Vol. 15, Page 21*, **2022**, *15* (1), 21. https://doi.org/10.3390/NU15010021.

Figure legend

Figure 1: Cut-off point for the age of the patients divided in to male and female. Males are less than 67 years old. Whereas, female affected at more 67 years old.

Figure 2: Diagramed presentation of the concomitant and the post MI compilations. From below up to complications (CHF, DM, hypertension, CKD, COPD, dyslipidemia, varicose veins) belongs to concomitant diseases before the development of MI. Abbreviations: CHF; chronic heart failure, COPD; chronic obstructive pulmonary disease, CKD; chronic kidney disease, DM; diabetes mellites.

Figure 3: graphical presentation for the dynamical changes in the levels of LDH (lactate dehydrogenase) and CPK-MB (creatinine phosphokinase-MB fraction. Both LDH and CPK-

MB fraction measured by U/L. this analysis conducted between two days for all patients as a follow up after MI.

Figure 4: A direct correlation between LDL level and the aorta diameter. Additionally, an opposite correlation between the left ventricle (LV) wall thickness and the EF.

Variable	Spearman Rank Order Correlations (Acute MI in Total Post MI 5) MD pairwise deleted Marked correlations are significant at p <,05000		
	Date of infarct	Date of hosp.	Date of dischargment
Date of infarct	1,000000	0,998755	0,998662
Date of hosp.	0,998755	1,000000	0,999754
Date of dischargment	0,998662	0,999754	1,000000
Hospitalization days	-0,492819	-0,480996	-0,470104
Age	-0,014406	-0,017981	-0,017534
Body weight	-0,042648	-0,056196	-0,057004
Hieght	-0,171595	-0,175662	-0,173637
BMI	-0,050015	-0,060710	-0,063937
HR	-0,026789	-0,029367	-0,028830
SBP	0,086786	0,096680	0,094712
DBP	0,050813	0,050794	0,050194
Aorta diameter	-0,029993	-0,022536	-0,025733
EF	0,080422	0,083670	0,083384
EDV	-0,045009	-0,050675	-0,052638
LV	-0,001416	-0,023427	-0,022092
LA	0,123265	0,117261	0,118687
RV	0,224438	0,209813	0,208153
RA	0,130204	0,133860	0,135790
Hb	0,037855	0,024145	0,023418
leukocytes	0,142190	0,118443	0,118092
RBC	0,013550	-0,027649	-0,029574
lymphocytes	0,091797	0,095098	0,093676
Thrombocytes	-0,210074	-0,213754	-0,213379
ESR	-0,014440	-0,027477	-0,027431
Ketones			
Urine protiens	0,302814	0,302253	0,305086
leukocytes	-0,344187	-0,325607	-0,325124
epithelial cells	-0,404424	-0,392751	-0,392272
Troponin I	-0,082746	-0,087620	-0,089514
Myoglobulin	0,500000	0,500000	0,500000
CKP-MB 1	-0,161461	-0,146670	-0,146545
CKP-İÅ 2	-0,369494	-0,367680	-0,369441
LDH 1	0,030811	0,026900	0,027219
LDH 2	0,123665	0,119836	0,113804
INR	0,222657	0,210509	0,210587
Total bili	-0,070080	-0,093753	-0,091667
ALT	-0,025792	-0,033286	-0,036998
AST	0,033419	0,045471	0,045172
Total choles	-0,081606	-0,083591	-0,082453
LDL	-0,120629	-0,114472	-0,110837
TAG	0,075539	0,041802	0,040081
Na	0,295534	0,320231	0,323290
K	0,392714	0,347683	0,345315
Creatinin	-0,123623	-0,108991	-0,107730
Ca	0,752734	0,655580	0,652954
uria	0,173533	0,172153	0,172156
Glucose	0,198889	0,197077	0,199063
APTT	-0,219685	-0,213583	-0,214560
Fibrinogen	0,070849	0,061743	0,060104
PT			

Variable	Spearman Rank Order Correlations (Acute MI in Total Post MI 5) MD pairwise deleted Marked correlations are significant at p <,05000			
	Hospitalization days	Age	Body weight	Hieght
Date of infarct	-0,492819	-0,014406	-0,042648	-0,171595
Date of hosp.	-0,480996	-0,017981	-0,056196	-0,175662
Date of dischargment	-0,470104	-0,017534	-0,057004	-0,173637
Hospitalization days	1,000000	0,092495	-0,239856	0,084892
Age	0,092495	1,000000	-0,056415	0,005208
Body weight	-0,239856	-0,056415	1,000000	-0,178452
Hieght	0,084892	0,005208	-0,178452	1,000000
BMI	-0,336077	0,029871	0,898731	-0,536541
HR	0,045764	-0,060908	0,438935	-0,249103
SBP	-0,063265	-0,054161	0,136715	-0,045358
DBP	0,004338	-0,110726	0,041841	-0,011282
Aorta diameter	-0,056556	-0,058935	-0,108477	0,305970
EF	-0,041657	-0,145918	-0,120649	-0,248037
EDV	-0,066685	-0,098928	0,438158	0,294385
LV	0,021873	-0,076707	0,348287	0,273553
LA	-0,006042	0,177610	0,177460	0,185164
RV	-0,087129	0,037892	0,415737	-0,152526
RA	-0,018941	0,161934	0,180343	0,397011
Hb	-0,162796	-0,157068	0,085367	0,176164
leukocytes	0,078156	-0,021030	-0,100778	-0,041827
RBC	-0,185504	-0,230622	0,050290	0,136868
lymphocytes	-0,187678	-0,150462	-0,124957	-0,175052
Thrombocytes	-0,053971	-0,000641	0,182193	-0,059169
ESR	0,026856	0,200394	-0,132872	-0,297634
Ketones				
Urine protiens	-0,030714	-0,031736	-0,234565	0,200668
leukocytes	0,220460	0,072826	0,111436	-0,010736
epithelial cells	0,251582	0,047546	0,069070	0,060775
Troponin I	0,036854	0,170201	0,307384	0,038983
Myoglobulin	-0,500000			
CKP-MB 1	0,062096	0,120242	0,221078	0,154298
CKP-ÌÂ 2	-0,060476	0,003794	-0,462185	0,692409
LDH 1	0,075959	0,172099	0,461451	0,083129
LDH 2	-0,351822	-0,035860	-0,342356	0,882919
INR	-0,122041	0,088783	-0,087567	-0,115722
Total bili	0,083439	0,119390	0,142107	0,304978
ALT	-0,149559	0,006031	0,357989	-0,136986
AST	-0,089148	0,053983	0,255829	0,051121
Total choles	0,135370	0,189818	0,032294	-0,138274
LDL	0,159497	0,103719	0,214156	-0,089237
TAG	-0,018082	-0,089894	0,190830	-0,346317
Na	-0,074054	0,042710	0,099801	-0,086951
K	-0,119386	0,096715	0,262988	-0,486864
Creatinin	0,118681	0,196371	-0,007508	0,035929
Ca	-0,365114	-0,006410	0,283974	-0,338311
uria	-0,032561	0,189761	0,121533	-0,404216
Glucose	-0,002611	0,166185	0,654202	-0,048985
APTT	0,142006	0,049935	0,003153	-0,204839
Fibrinogen	0,014194	-0,101096	-0,200220	-0,219706
PT				

Variable	Spearman Rank Order Correlations (Acute MI in Total Post MI 5) MD pairwise deleted Marked correlations are significant at p <,05000				
	BMI	HR	SBP	DBP	Aorta diameter
Date of infarct	-0,050015	-0,026789	0,086786	0,050813	-0,029993
Date of hosp.	-0,060710	-0,029367	0,096680	0,050794	-0,022536
Date of dischargment	-0,063937	-0,028830	0,094712	0,050194	-0,025733
Hospitalization days	-0,336077	0,045764	-0,063265	0,004338	-0,056556
Age	0,029871	-0,060908	-0,054161	-0,110726	-0,058935
Body weight	0,898731	0,438935	0,136715	0,041841	-0,108477
Hieght	-0,536541	-0,249103	-0,045358	-0,011282	0,305970
BMI	1,000000	0,475827	0,233983	0,134203	-0,303926
HR	0,475827	1,000000	0,164042	0,193671	0,056941
SBP	0,233983	0,164042	1,000000	0,837480	0,062520
DBP	0,134203	0,193671	0,837480	1,000000	0,109910
Aorta diameter	-0,303926	0,056941	0,062520	0,109910	1,000000
EF	-0,032926	-0,178774	0,093906	0,042189	0,050386
EDV	0,321078	0,012821	0,129426	0,181506	0,126137
LV	0,062654	0,108568	0,078351	0,214669	0,217956
LA	-0,090691	-0,033390	0,127778	0,117089	0,220792
RV	0,343756	0,122876	0,107076	0,187414	0,339438
RA	-0,040683	0,009769	-0,000967	0,056850	0,222980
Hb	0,065826	0,031176	0,017701	0,064565	0,077202
leukocytes	-0,091010	-0,047458	-0,010262	0,043619	-0,101208
RBC	0,131963	-0,109265	0,008063	0,010929	-0,012348
lymphocytes	0,044404	0,126764	-0,023606	0,000627	0,038371
Thrombocytes	0,253130	0,129454	-0,104800	-0,053676	-0,208526
ESR	-0,000494	0,084674	-0,118841	-0,048738	-0,041583
Ketones					
Urine protiens	-0,238145	-0,171914	0,073485	0,051278	-0,084270
leukocytes	0,186406	-0,010679	0,051028	-0,026506	-0,196673
epithelial cells	0,114878	0,018641	0,042596	-0,026188	-0,137258
Troponin I	0,335063	-0,054693	-0,195500	-0,180407	-0,105639
Myoglobulin		1,000000	1,000000	0,866025	
CKP-MB 1	0,102362	0,072393	-0,149235	-0,148068	-0,134453
CKP-ÌÂ 2	-0,543938	0,019912	-0,021319	0,048399	0,081346
LDH 1	0,437790	-0,043696	-0,196905	-0,227802	-0,317959
LDH 2	-0,357143	-0,065697	-0,047346	-0,023404	-0,025929
INR	0,016422	-0,176321	-0,009411	-0,032399	0,012770
Total bili	0,013691	0,091752	-0,074120	-0,152479	0,105600
ALT	0,462577	0,168135	-0,014180	-0,079591	-0,081878
AST	0,191586	0,058756	-0,161133	-0,161825	0,016592
Total choles	0,167920	0,074696	0,130019	0,103124	-0,272978
LDL	0,287161	-0,019979	0,048523	0,051746	-0,290000
TAG	0,243163	-0,001737	0,159888	0,203202	-0,064514
Na	0,147481	0,017900	0,073128	0,023359	0,012202
K	0,402687	0,055393	0,000735	-0,018963	0,099267
Creatinin	-0,051752	-0,018731	-0,115542	-0,229538	-0,130249
Ca	0,422982	-0,077756	-0,065510	-0,085972	-0,005432
uria	0,221784	0,076878	-0,020419	-0,078703	-0,032316
Glucose	0,497591	0,077596	-0,030442	-0,022622	0,053932
APTT	0,145463	0,034608	-0,055021	-0,086966	-0,118503
Fibrinogen	-0,051268	-0,068684	0,014181	-0,081389	-0,066697
PT					

Variable	Spearman Rank Order Correlations (Acute MI in Total Post MI 5) MD pairwise deleted Marked correlations are significant at p <,05000					
	EF	EDV	LV	LA	RV	RA
Date of infarct	0,080422	-0,045009	-0,001416	0,123265	0,224438	0,130204
Date of hosp.	0,083670	-0,050675	-0,023427	0,117261	0,209813	0,133860
Date of dischargment	0,083384	-0,052638	-0,022092	0,118687	0,208153	0,135790
Hospitalization days	-0,041657	-0,066685	0,021873	-0,006042	-0,087129	-0,018941
Age	-0,145918	-0,098928	-0,076707	0,177610	0,037892	0,161934
Body weight	-0,120649	0,438158	0,348287	0,177460	0,415737	0,180343
Hieght	-0,248037	0,294385	0,273553	0,185164	-0,152526	0,397011
BMI	-0,032926	0,321078	0,062654	-0,090691	0,343756	-0,040683
HR	-0,178774	0,012821	0,108568	-0,033390	0,122876	0,009769
SBP	0,093906	0,129426	0,078351	0,127778	0,107076	-0,000967
DBP	0,042189	0,181506	0,214669	0,117089	0,187414	0,056850
Aorta diameter	0,050386	0,126137	0,217956	0,220792	0,339438	0,222980
EF	1,000000	-0,204007	-0,306233	-0,037623	0,037063	0,051806
EDV	-0,204007	1,000000	0,816552	0,427731	0,232208	0,153922
LV	-0,306233	0,816552	1,000000	0,373137	0,198793	0,180908
LA	-0,037623	0,427731	0,373137	1,000000	0,421956	0,553528
RV	0,037063	0,232208	0,198793	0,421956	1,000000	0,222865
RA	0,051806	0,153922	0,180908	0,553528	0,222865	1,000000
Hb	0,022881	0,107920	0,016682	0,074045	0,114354	0,011420
leukocytes	-0,181839	0,209526	0,089332	0,077030	0,139811	-0,037217
RBC	0,070459	0,014637	0,022377	-0,065412	0,136926	-0,022704
lymphocytes	0,019908	-0,086165	-0,038685	-0,084508	-0,061272	-0,150101
Thrombocytes	-0,007202	0,042597	-0,055031	-0,195668	-0,085269	-0,108218
ESR	-0,008205	0,068129	0,064889	-0,004561	-0,030127	0,084620
Ketones						
Urine protiens	0,162382	-0,145296	-0,080618	-0,139515	-0,001364	0,111536
leukocytes	-0,102500	0,026230	-0,042291	0,040654	-0,106300	-0,140144
epithelial cells	-0,153045	-0,042611	-0,066126	-0,099064	-0,243669	-0,234743
Troponin I	-0,053490	-0,054769	-0,119882	-0,106785	-0,094003	-0,032516
Myoglobulin	1,000000			-0,500000		
CKP-MB 1	-0,059120	0,042945	-0,080071	-0,114084	-0,232948	-0,029421
CKP-ÌÂ 2	-0,037202	-0,058452	-0,086273	0,007858	-0,009298	0,032700
LDH 1	-0,121081	0,078570	-0,081344	-0,087928	-0,057713	-0,049061
LDH 2	-0,131816	-0,114369	-0,150290	-0,053576	-0,082071	0,011156
INR	-0,041426	0,021536	-0,046075	0,055728	0,098168	0,162673
Total bili	-0,161993	0,154520	0,007623	0,182757	0,195401	0,054035
ALT	-0,068176	0,069499	0,009479	0,064334	0,083873	0,022768
AST	-0,100612	0,069440	0,068600	-0,049618	0,001424	0,077034
Total choles	0,001418	-0,010848	-0,034502	-0,022993	-0,008355	-0,160652
LDL	-0,003403	-0,023996	-0,070730	-0,008003	0,006907	-0,091096
TAG	0,135686	-0,012581	0,085614	0,071785	0,096231	-0,036389
Na	0,028201	-0,062749	0,045815	0,144556	-0,055881	0,181103
K	0,047078	0,058684	-0,002292	0,117294	0,285462	0,147145
Creatinin	-0,136330	-0,094480	-0,066634	-0,027856	-0,207834	-0,101998
Ca	0,020465	-0,047146	-0,021513	0,265162	0,180772	0,166642
uria	0,005515	-0,078861	-0,037700	0,189547	0,194250	0,101188
Glucose	-0,146818	0,043452	0,129861	0,175194	0,223505	0,171877
APTT	-0,201638	0,173079	0,040094	-0,129817	-0,000931	-0,139865
Fibrinogen	0,067573	-0,119314	-0,116229	0,006765	-0,007770	-0,072335
PT						

Variable	Spearman Rank Order Correlations (Acute MI in Total Post MI 5) MD pairwise deleted Marked correlations are significant at p <,05000				
	Hb	leukocytes	RBC	lymphocytes	Thrombocytes
Date of infarct	0,037855	0,142190	0,013550	0,091797	-0,210074
Date of hosp.	0,024145	0,118443	-0,027649	0,095098	-0,213754
Date of dischargment	0,023418	0,118092	-0,029574	0,093676	-0,213379
Hospitalization days	-0,162796	0,078156	-0,185504	-0,187678	-0,053971
Age	-0,157068	-0,021030	-0,230622	-0,150462	-0,000641
Body weight	0,085367	-0,100778	0,050290	-0,124957	0,182193
Hieght	0,176164	-0,041827	0,136868	-0,175052	-0,059169
BMI	0,065826	-0,091010	0,131963	0,044404	0,253130
HR	0,031176	-0,047458	-0,109265	0,126764	0,129454
SBP	0,017701	-0,010262	0,008063	-0,023606	-0,104800
DBP	0,064565	0,043619	0,010929	0,000627	-0,053676
Aorta diameter	0,077202	-0,101208	-0,012348	0,038371	-0,208526
EF	0,022881	-0,181839	0,070459	0,019908	-0,007202
EDV	0,107920	0,209526	0,014637	-0,086165	0,042597
LV	0,016682	0,089332	0,022377	-0,038685	-0,055031
LA	0,074045	0,077030	-0,065412	-0,084508	-0,195668
RV	0,114354	0,139811	0,136926	-0,061272	-0,085269
RA	0,011420	-0,037217	-0,022704	-0,150101	-0,108218
Hb	1,000000	0,268726	0,654935	0,087015	0,058451
leukocytes	0,268726	1,000000	0,109458	-0,407888	0,278365
RBC	0,654935	0,109458	1,000000	0,111199	0,164037
lymphocytes	0,087015	-0,407888	0,111199	1,000000	-0,163473
Thrombocytes	0,058451	0,278365	0,164037	-0,163473	1,000000
ESR	-0,424196	-0,053640	-0,344304	0,087865	0,071924
Ketones					
Urine protiens	-0,194071	0,143997	0,042784	-0,208550	-0,053757
leukocytes	-0,141955	0,072283	-0,037207	-0,139760	0,175763
epithelial cells	-0,036518	0,088763	-0,007425	-0,058215	0,092318
Troponin I	0,228199	0,273322	0,188032	-0,034776	0,242812
Myoglobulin	-0,500000	1,000000		0,500000	
CKP-MB 1	0,046591	0,118350	-0,082001	0,063300	0,164946
CKP-ÍÂ 2	0,116045	0,071282	-0,021656	-0,153597	0,205310
LDH 1	-0,070553	0,252933	0,001593	-0,061029	0,095874
LDH 2	0,254468	0,286870	0,126967	-0,211282	0,349153
INR	0,101077	0,187285	0,052727	-0,213453	0,200752
Total bili	0,215681	-0,040948	0,089829	0,074228	-0,220295
ALT	0,011558	0,067994	0,073779	-0,006331	0,035331
AST	-0,009187	0,072187	-0,080989	-0,038706	-0,083374
Total choles	0,066861	-0,025760	0,023385	0,111800	-0,011878
LDL	0,114681	-0,024549	0,091089	0,066237	0,051375
TAG	0,157777	0,203652	0,172810	0,157562	-0,017568
Na	0,081063	-0,068359	-0,027294	0,157465	-0,155198
K	0,218370	0,047010	0,187586	0,193137	-0,172012
Creatinin	-0,024685	0,068294	-0,137145	-0,124911	0,183592
Ca	0,194971	0,060749	0,269192	0,198846	-0,046922
uria	-0,211940	0,037870	-0,139107	-0,165130	-0,004645
Glucose	-0,008374	0,205235	0,090559	-0,125482	-0,062960
APTT	0,102761	0,104666	0,174735	0,011001	0,316938
Fibrinogen	-0,051919	-0,093511	0,020243	0,044111	-0,224921
PT					

Variable	Spearman Rank Order Correlations (Acute MI in Total Post MI 5) MD pairwise deleted Marked correlations are significant at p <,05000				
	ESR	Ketones	Urine protiens	leukocytes	epithelial cells
Date of infarct	-0,014440		0,302814	-0,344187	-0,404424
Date of hosp.	-0,027477		0,302253	-0,325607	-0,392751
Date of dischargment	-0,027431		0,305086	-0,325124	-0,392272
Hospitalization days	0,026856		-0,030714	0,220460	0,251582
Age	0,200394		-0,031736	0,072826	0,047546
Body weight	-0,132872		-0,234565	0,111436	0,069070
Hieght	-0,297634		0,200668	-0,010736	0,060775
BMI	-0,000494		-0,238145	0,186406	0,114878
HR	0,084674		-0,171914	-0,010679	0,018641
SBP	-0,118841		0,073485	0,051028	0,042596
DBP	-0,048738		0,051278	-0,026506	-0,026188
Aorta diameter	-0,041583		-0,084270	-0,196673	-0,137258
EF	-0,008205		0,162382	-0,102500	-0,153045
EDV	0,068129		-0,145296	0,026230	-0,042611
LV	0,064889		-0,080618	-0,042291	-0,066126
LA	-0,004561		-0,139515	0,040654	-0,099064
RV	-0,030127		-0,001364	-0,106300	-0,243669
RA	0,084620		0,111536	-0,140144	-0,234743
Hb	-0,424196		-0,194071	-0,141955	-0,036518
leukocytes	-0,053640		0,143997	0,072283	0,088763
RBC	-0,344304		0,042784	-0,037207	-0,007425
lymphocytes	0,087865		-0,208550	-0,139760	-0,058215
Thrombocytes	0,071924		-0,053757	0,175763	0,092318
ESR	1,000000		-0,031909	-0,021033	-0,089231
Ketones		1,000000			
Urine protiens	-0,031909		1,000000	0,037846	-0,035824
leukocytes	-0,021033		0,037846	1,000000	0,789820
epithelial cells	-0,089231		-0,035824	0,789820	1,000000
Troponin I	-0,039481		0,045506	0,071426	0,106241
Myoglobulin	0,500000		0,866025	0,500000	0,500000
CKP-MB 1	0,098663		-0,044192	0,172589	0,239036
CKP-ÌÂ 2	0,013043		-0,291265	0,219115	0,329329
LDH 1	0,047350		0,137175	0,128269	0,040671
LDH 2	-0,210693		0,046534	-0,120852	-0,084567
INR	-0,019873		0,145399	-0,010482	-0,113416
Total bili	-0,048101		0,032055	-0,140302	-0,083612
ALT	0,013972		-0,037007	-0,097826	-0,028856
AST	0,000510		0,091987	-0,132582	-0,041266
Total choles	-0,128001		-0,061160	0,106976	0,121417
LDL	-0,102900		-0,057283	0,212155	0,184569
TAG	0,038587		0,007271	-0,001248	0,020937
Na	-0,101750		-0,070218	-0,177667	-0,162685
K	-0,033521		-0,049361	-0,202077	-0,275099
Creatinin	-0,097531		0,030465	0,177259	0,158177
Ca	-0,090898		0,007566	-0,376993	-0,433478
uria	0,165846		0,190483	0,099496	0,036004
Glucose	0,021904		0,109950	0,067847	0,160787
APTT	-0,032073		-0,184248	0,223590	0,313909
Fibrinogen	-0,019113		0,032734	0,074379	0,100294
PT					

Variable	Spearman Rank Order Correlations (Acute MI in Total Post MI 5) MD pairwise deleted Marked correlations are significant at p <,05000				
	Troponin I	Myoglobulin	CKP-MB 1	CKP-ÌÂ 2	LDH 1
Date of infarct	-0,082746	0,50000	-0,161461	-0,369494	0,030811
Date of hosp.	-0,087620	0,50000	-0,146670	-0,367680	0,026900
Date of dischargment	-0,089514	0,50000	-0,146545	-0,369441	0,027219
Hospitalization days	0,036854	-0,50000	0,062096	-0,060476	0,075959
Age	0,170201		0,120242	0,003794	0,172099
Body weight	0,307384		0,221078	-0,462185	0,461451
Hieght	0,038983		0,154298	0,692409	0,083129
BMI	0,335063		0,102362	-0,543938	0,437790
HR	-0,054693	1,00000	0,072393	0,019912	-0,043696
SBP	-0,195500	1,00000	-0,149235	-0,021319	-0,196905
DBP	-0,180407	0,86603	-0,148068	0,048399	-0,227802
Aorta diameter	-0,105639		-0,134453	0,081346	-0,317959
EF	-0,053490	1,00000	-0,059120	-0,037202	-0,121081
EDV	-0,054769		0,042945	-0,058452	0,078570
LV	-0,119882		-0,080071	-0,086273	-0,081344
LA	-0,106785	-0,50000	-0,114084	0,007858	-0,087928
RV	-0,094003		-0,232948	-0,009298	-0,057713
RA	-0,032516		-0,029421	0,032700	-0,049061
Hb	0,228199	-0,50000	0,046591	0,116045	-0,070553
leukocytes	0,273322	1,00000	0,118350	0,071282	0,252933
RBC	0,188032		-0,082001	-0,021656	0,001593
lymphocytes	-0,034776	0,50000	0,063300	-0,153597	-0,061029
Thrombocytes	0,242812		0,164946	0,205310	0,095874
ESR	-0,039481	0,50000	0,098663	0,013043	0,047350
Ketones					
Urine protiens	0,045506	0,86603	-0,044192	-0,291265	0,137175
leukocytes	0,071426	0,50000	0,172589	0,219115	0,128269
epithelial cells	0,106241	0,50000	0,239036	0,329329	0,040671
Troponin I	1,000000		0,490390	0,265996	0,374820
Myoglobulin		1,00000			
CKP-MB 1	0,490390		1,000000	0,386079	0,576087
CKP-ÌÂ 2	0,265996		0,386079	1,000000	0,067851
LDH 1	0,374820		0,576087	0,067851	1,000000
LDH 2	0,319420		0,127235	0,437527	0,314912
INR	0,208372		0,057017	0,000653	0,141873
Total bili	0,166543		0,189142	0,018849	0,112833
ALT	0,130182		0,152500	0,057059	0,200843
AST	0,170510		0,194561	0,053218	0,241813
Total choles	0,124187	0,50000	-0,040978	-0,006981	0,082099
LDL	0,195942	0,50000	0,061061	0,116077	0,097868
TAG	0,075288	0,50000	-0,001916	-0,046766	0,003054
Na	0,073038	-1,00000	-0,145310	-0,202524	0,056657
K	0,111597	0,50000	0,007184	-0,162865	0,162139
Creatinin	0,177284	-1,00000	0,260587	-0,008614	0,226690
Ca	0,002295		-0,255494	-0,448271	-0,032190
uria	0,077832	0,50000	0,050858	-0,208204	0,114875
Glucose	-0,027007	1,00000	-0,062950	-0,082192	0,112243
APTT	0,126145		0,309909	0,423780	0,087702
Fibrinogen	-0,126763		0,019966	-0,141352	0,038093
PT					

Variable	Spearman Rank Order Correlations (Acute MI in Total Post MI 5) MD pairwise deleted Marked correlations are significant at p <,05000				
	LDH 2	INR	Total bili	ALT	AST
Date of infarct	0,123665	0,222657	-0,070080	-0,025792	0,033419
Date of hosp.	0,119836	0,210509	-0,093753	-0,033286	0,045471
Date of dischargment	0,113804	0,210587	-0,091667	-0,036998	0,045172
Hospitalization days	-0,351822	-0,122041	0,083439	-0,149559	-0,089148
Age	-0,035860	0,088783	0,119390	0,006031	0,053983
Body weight	-0,342356	-0,087567	0,142107	0,357989	0,255829
Hieght	0,882919	-0,115722	0,304978	-0,136986	0,051121
BMI	-0,357143	0,016422	0,013691	0,462577	0,191586
HR	-0,065697	-0,176321	0,091752	0,168135	0,058756
SBP	-0,047346	-0,009411	-0,074120	-0,014180	-0,161133
DBP	-0,023404	-0,032399	-0,152479	-0,079591	-0,161825
Aorta diameter	-0,025929	0,012770	0,105600	-0,081878	0,016592
EF	-0,131816	-0,041426	-0,161993	-0,068176	-0,100612
EDV	-0,114369	0,021536	0,154520	0,069499	0,069440
LV	-0,150290	-0,046075	0,007623	0,009479	0,068600
LA	-0,053576	0,055728	0,182757	0,064334	-0,049618
RV	-0,082071	0,098168	0,195401	0,083873	0,001424
RA	0,011156	0,162673	0,054035	0,022768	0,077034
Hb	0,254468	0,101077	0,215681	0,011558	-0,009187
leukocytes	0,286870	0,187285	-0,040948	0,067994	0,072187
RBC	0,126967	0,052727	0,089829	0,073779	-0,080989
lymphocytes	-0,211282	-0,213453	0,074228	-0,006331	-0,038706
Thrombocytes	0,349153	0,200752	-0,220295	0,035331	-0,083374
ESR	-0,210693	-0,019873	-0,048101	0,013972	0,000510
Ketones					
Urine protiens	0,046534	0,145399	0,032055	-0,037007	0,091987
leukocytes	-0,120852	-0,010482	-0,140302	-0,097826	-0,132582
epithelial cells	-0,084567	-0,113416	-0,083612	-0,028856	-0,041266
Troponin I	0,319420	0,208372	0,166543	0,130182	0,170510
Myoglobulin					
CKP-MB 1	0,127235	0,057017	0,189142	0,152500	0,194561
CKP-IÂ 2	0,437527	0,000653	0,018849	0,057059	0,053218
LDH 1	0,314912	0,141873	0,112833	0,200843	0,241813
LDH 2	1,000000	0,382124	0,002090	0,286460	0,447358
INR	0,382124	1,000000	0,107299	-0,044216	0,049075
Total bili	0,002090	0,107299	1,000000	0,175968	0,135787
ALT	0,286460	-0,044216	0,175968	1,000000	0,572879
AST	0,447358	0,049075	0,135787	0,572879	1,000000
Total choles	-0,049465	-0,111364	0,135346	0,068525	0,067669
LDL	-0,043774	-0,011290	0,158629	0,020542	0,064199
TAG	-0,033778	-0,095456	-0,039331	0,126804	0,040082
Na	0,126636	0,138965	-0,085883	-0,015813	0,130889
K	-0,091874	0,175417	0,134458	0,035404	0,062157
Creatinin	-0,170041	0,185665	0,069496	-0,021837	0,002576
Ca	-0,065550	0,229418	-0,076928	0,098510	-0,040195
uria	-0,156195	0,132454	0,048647	0,248900	0,155723
Glucose	0,089111	0,031761	-0,009808	0,246838	0,156518
APTT	0,197419	0,167639	0,073983	0,084182	-0,016993
Fibrinogen	-0,089828	-0,389201	0,105851	-0,003601	-0,044486
PT					

Variable	Spearman Rank Order Correlations (Acute MI in Total Post MI 5) MD pairwise deleted Marked correlations are significant at p <,05000				
	Total choles	LDL	TAG	Na	K
Date of infarct	-0,081606	-0,120629	0,075539	0,29553	0,392714
Date of hosp.	-0,083591	-0,114472	0,041802	0,32023	0,347683
Date of dischargment	-0,082453	-0,110837	0,040081	0,32329	0,345315
Hospitalization days	0,135370	0,159497	-0,018082	-0,07405	-0,119386
Age	0,189818	0,103719	-0,089894	0,04271	0,096715
Body weight	0,032294	0,214156	0,190830	0,09980	0,262988
Hieght	-0,138274	-0,089237	-0,346317	-0,08695	-0,486864
BMI	0,167920	0,287161	0,243163	0,14748	0,402687
HR	0,074696	-0,019979	-0,001737	0,01790	0,055393
SBP	0,130019	0,048523	0,159888	0,07313	0,000735
DBP	0,103124	0,051746	0,203202	0,02336	-0,018963
Aorta diameter	-0,272978	-0,290000	-0,064514	0,01220	0,099267
EF	0,001418	-0,003403	0,135686	0,02820	0,047078
EDV	-0,010848	-0,023996	-0,012581	-0,06275	0,058684
LV	-0,034502	-0,070730	0,085614	0,04582	-0,002292
LA	-0,022993	-0,008003	0,071785	0,14456	0,117294
RV	-0,008355	0,006907	0,096231	-0,05588	0,285462
RA	-0,160652	-0,091096	-0,036389	0,18110	0,147145
Hb	0,066861	0,114681	0,157777	0,08106	0,218370
leukocytes	-0,025760	-0,024549	0,203652	-0,06836	0,047010
RBC	0,023385	0,091089	0,172810	-0,02729	0,187586
lymphocytes	0,111800	0,066237	0,157562	0,15747	0,193137
Thrombocytes	-0,011878	0,051375	-0,017568	-0,15520	-0,172012
ESR	-0,128001	-0,102900	0,038587	-0,10175	-0,033521
Ketones					
Urine protiens	-0,061160	-0,057283	0,007271	-0,07022	-0,049361
leukocytes	0,106976	0,212155	-0,001248	-0,17767	-0,202077
epithelial cells	0,121417	0,184569	0,020937	-0,16269	-0,275099
Troponin I	0,124187	0,195942	0,075288	0,07304	0,111597
Myoglobulin	0,500000	0,500000	0,500000	-1,00000	0,500000
CKP-MB 1	-0,040978	0,061061	-0,001916	-0,14531	0,007184
CKP-ÌÂ 2	-0,006981	0,116077	-0,046766	-0,20252	-0,162865
LDH 1	0,082099	0,097868	0,003054	0,05666	0,162139
LDH 2	-0,049465	-0,043774	-0,033778	0,12664	-0,091874
INR	-0,111364	-0,011290	-0,095456	0,13897	0,175417
Total bili	0,135346	0,158629	-0,039331	-0,08588	0,134458
ALT	0,068525	0,020542	0,126804	-0,01581	0,035404
AST	0,067669	0,064199	0,040082	0,13089	0,062157
Total choles	1,000000	0,830689	0,417038	0,09591	-0,006258
LDL	0,830689	1,000000	0,319855	0,06534	-0,075583
TAG	0,417038	0,319855	1,000000	-0,00482	0,096150
Na	0,095906	0,065343	-0,004822	1,00000	0,349223
K	-0,006258	-0,075583	0,096150	0,34922	1,000000
Creatinin	0,083018	0,087098	-0,032333	-0,01335	-0,060077
Ca	-0,108344	-0,126906	0,066131	0,47663	0,575797
uria	0,046621	0,055896	0,027175	0,09747	0,132856
Glucose	0,048354	0,026693	0,161067	-0,03075	0,004122
APTT	0,110501	0,148558	-0,068772	-0,11314	0,049147
Fibrinogen	0,044268	-0,029297	0,160949	-0,09123	0,017443
PT					

Variable	Spearman Rank Order Correlations (Acute MI in Total Post MI 5) MD pairwise deleted Marked correlations are significant at p <,05000				
	Creatinin	Ca	uria	Glucose	APTT
Date of infarct	-0,12362	0,752734	0,173533	0,198889	-0,219685
Date of hosp.	-0,10899	0,655580	0,172153	0,197077	-0,213583
Date of dischargment	-0,10773	0,652954	0,172156	0,199063	-0,214560
Hospitalization days	0,11868	-0,365114	-0,032561	-0,002611	0,142006
Age	0,19637	-0,006410	0,189761	0,166185	0,049935
Body weight	-0,00751	0,283974	0,121533	0,654202	0,003153
Hieght	0,03593	-0,338311	-0,404216	-0,048985	-0,204839
BMI	-0,05175	0,422982	0,221784	0,497591	0,145463
HR	-0,01873	-0,077756	0,076878	0,077596	0,034608
SBP	-0,11554	-0,065510	-0,020419	-0,030442	-0,055021
DBP	-0,22954	-0,085972	-0,078703	-0,022622	-0,086966
Aorta diameter	-0,13025	-0,005432	-0,032316	0,053932	-0,118503
EF	-0,13633	0,020465	0,005515	-0,146818	-0,201638
EDV	-0,09448	-0,047146	-0,078861	0,043452	0,173079
LV	-0,06663	-0,021513	-0,037700	0,129861	0,040094
LA	-0,02786	0,265162	0,189547	0,175194	-0,129817
RV	-0,20783	0,180772	0,194250	0,223505	-0,000931
RA	-0,10200	0,166642	0,101188	0,171877	-0,139865
Hb	-0,02468	0,194971	-0,211940	-0,008374	0,102761
leukocytes	0,06829	0,060749	0,037870	0,205235	0,104666
RBC	-0,13715	0,269192	-0,139107	0,090559	0,174735
lymphocytes	-0,12491	0,198846	-0,165130	-0,125482	0,011001
Thrombocytes	0,18359	-0,046922	-0,004645	-0,062960	0,316938
ESR	-0,09753	-0,090898	0,165846	0,021904	-0,032073
Ketones					
Urine protiens	0,03047	0,007566	0,190483	0,109950	-0,184248
leukocytes	0,17726	-0,376993	0,099496	0,067847	0,223590
epithelial cells	0,15818	-0,433478	0,036004	0,160787	0,313909
Troponin I	0,17728	0,002295	0,077832	-0,027007	0,126145
Myoglobulin	-1,00000		0,500000	1,000000	
CKP-MB 1	0,26059	-0,255494	0,050858	-0,062950	0,309909
CKP-ÌÂ 2	-0,00861	-0,448271	-0,208204	-0,082192	0,423780
LDH 1	0,22669	-0,032190	0,114875	0,112243	0,087702
LDH 2	-0,17004	-0,065550	-0,156195	0,089111	0,197419
INR	0,18567	0,229418	0,132454	0,031761	0,167639
Total bili	0,06950	-0,076928	0,048647	-0,009808	0,073983
ALT	-0,02184	0,098510	0,248900	0,246838	0,084182
AST	0,00258	-0,040195	0,155723	0,156518	-0,016993
Total choles	0,08302	-0,108344	0,046621	0,048354	0,110501
LDL	0,08710	-0,126906	0,055896	0,026693	0,148558
TAG	-0,03233	0,066131	0,027175	0,161067	-0,068772
Na	-0,01335	0,476633	0,097467	-0,030751	-0,113135
K	-0,06008	0,575797	0,132856	0,004122	0,049147
Creatinin	1,00000	-0,021655	0,477372	0,009566	0,130883
Ca	-0,02165	1,000000	0,226440	0,114659	-0,213578
uria	0,47737	0,226440	1,000000	0,223992	-0,038725
Glucose	0,00957	0,114659	0,223992	1,000000	-0,091630
APTT	0,13088	-0,213578	-0,038725	-0,091630	1,000000
Fibrinogen	-0,03245	-0,074313	0,054461	0,016700	-0,101748
PT					

Variable	Spearman Rank Order Correlations (Acute MI in Total Post MI 5) MD pairwise deleted Marked correlations are significant at p <,05000	
	Fibrinogen	PT
Date of infarct	0,070849	
Date of hosp.	0,061743	
Date of dischargment	0,060104	
Hospitalization days	0,014194	
Age	-0,101096	
Body weight	-0,200220	
Hieght	-0,219706	
BMI	-0,051268	
HR	-0,068684	
SBP	0,014181	
DBP	-0,081389	
Aorta diameter	-0,066697	
EF	0,067573	
EDV	-0,119314	
LV	-0,116229	
LA	0,006765	
RV	-0,007770	
RA	-0,072335	
Hb	-0,051919	
leukocytes	-0,093511	
RBC	0,020243	
lymphocytes	0,044111	
Thrombocytes	-0,224921	
ESR	-0,019113	
Ketones		
Urine protiens	0,032734	
leukocytes	0,074379	
epithelial cells	0,100294	
Troponin I	-0,126763	
Myoglobulin		
CKP-MB 1	0,019966	
CKP-ÌÂ 2	-0,141352	
LDH 1	0,038093	
LDH 2	-0,089828	
INR	-0,389201	
Total bili	0,105851	
ALT	-0,003601	
AST	-0,044486	
Total choles	0,044268	
LDL	-0,029297	
TAG	0,160949	
Na	-0,091235	
K	0,017443	
Creatinin	-0,032451	
Ca	-0,074313	
uria	0,054461	
Glucose	0,016700	
APTT	-0,101748	
Fibrinogen	1,000000	
PT		1,000000

Variable	Correlations (Acute MI in Total Post MI 5) Marked correlations are significant at p < ,05000 N=30 (Casewise deletion of missing data)
	Body weight
BMI	0,895885