1 **Campus Sewage Water Surveillance based dynamics and infection trends of**

2 **SARS-CoV-2 variants during third wave of COVID-19 in Pune, India**

3 Vinita Malik[a,1], Vinay Rajput [a,b,1], Rinka Pramanik[a,1], Rachel Samson[a,b], Rakesh

4 kumar Yadav[a,b], Pradnya Kadam[c], Nikita Shah[c], Rutuja Sawant [c], Unnati Bhalerao[c],

5 Manisha Tupekar[c], Soumen Khan[c], Priyanki Shah [e,] LS Shashidhara[c,e], Sanjay

6 Kamble[d], Syed Dastager[a,b], Krishanpal Karmodiya[c], Mahesh Dharne[a,b*]

7 [a]National Collection of Industrial Microorganisms (NCIM), Biochemical Sciences

8 Division, CSIR-National Chemical Laboratory, Pune 411008, Maharashtra, India

9 [b]Academy of Scientific and Innovative Research (AcSIR), Ghaziabad 201002, Uttar

10 Pradesh, India

11 [c]Department of Biology, Indian Institute of Science Education and Research, Pune

12 411008, Maharashtra, India

13 [d]Chemical Engineering and Process Development Division (CEPD), CSIR-National

14 Chemical Laboratory, Pune 411008, Maharashtra, India

15 [e] The Pune Knowledge Cluster (PKC), Savitribai Phule Pune University, Pune 411007,

16 Maharashtra, India

17 [1] These authors contributed equally

18 Corresponding author: National Collection of Industrial Microorganisms (NCIM),

19 Biochemical Sciences Division, CSIR-National Chemical Laboratory, Pune 411008,

20 India. E-mail address: ms.dharne@ncl.res.in (M.S. Dharne).

Highlights

- The first study on campus sewage water for SARS-CoV-2 surveillance in India
- Early detection of Omicron VOC signals during early November 2021
- Sequencing revealed the presence of Omicron VOC fragments prior to clinical cases reported on campus
- Genomic analysis indicated transition of Delta to Omicron lineage in late December 2021 and potentially led to the third wave
- Combining qRT-PCR and sequencing could be useful for on-campus tracking of variants using wastewater surveillance

Abstract

The wastewater-based epidemiology (WBE) of SARS-CoV-2 is a quick and cost-effective method of tracking virus transmission. However, few studies reported on campus or in academic or residential settings worldwide. In this study, we demonstrated the WBE approach to detect, monitor, and evaluate genomic variants of SARS-CoV-2 fragments in a sewage treatment plant (STP) located on the campus of CSIR National Chemical Laboratory, Pune, India. Herein we describe the early warning capability of WBE, with viral load rise in campus sewage water up to 14 days before its clinical detection. This was supported further by a significant correlation between SARS-CoV-2 RNA concentration and clinically reported COVID-19 cases on campus. Additionally, we comprehended the probable targets missed by the quantitative qRT-PCR using amplicon-based sequencing due to low viral load. The analysis revealed the presence of signature mutations of the Omicron

(S:N679K, S:N764K, S:D796Y, N:P13L, ORF1a:T3255I, ORF1a:K856R,

ORF1a:P3395H, and N:S413R) before the lineage was first detected globally. Further,

we used Lineage decomposition (LCS) tool to detect the Variant of Concern

(VOC)/Variant of Interest (VOI) signals upto a month earlier in sewage water samples.

The analysis also indicated the transition of lineage from Delta to Omicron in late

Decemeber,2021. This is the first study in India highlighting the use of on-campus

STP to evaluate the local spread of SARS-CoV-2, which could aid in preventing

COVID-19 in academic institutes/universities. This study proves the usefulness of

WBE as an early warning system for detecting, tracking and tracing VOCs using the

sequencing approach. The current study could aid in taking critical decisions to tackle

the pandemic scenario on campus.

Keywords: SARS-CoV-2, wastewater-based epidemiology, Early warning, on-

campus surveillance, Next-generation sequencing.

1. Introduction

Following the detection of the first SARS-CoV-2 infection (Severe Acute Respiratory

Syndrome Coronavirus 2) in late December 2019, several variants of this virus have

been reported globally. The constantly changing genome of SARS-CoV-2 has

challenged healthcare systems worldwide (Zhu et al., 2020). SARS-CoV-2 had

infected over 44 million people and killed over 0.5 million people in India as of

November 1, 2022 (Worldometer 2022). Despite administrating more than two

million doses of the Covid-19 vaccine (https://www.mohfw.gov.in/), the government

is still taking action to speed up the immunization campaign to contain the virus.

However, the major obstacle for public health is the emergence of new variants of

SARS-CoV-2 due to rapid mutations in its genome. The World Health Organization

(WHO) has categorized SARS-CoV-2 variants into Variants of Interest (VOIs) and

Variants of Concern (VOCs) based on their characteristics and impact on the global

community. Monitoring the occurrence of the variants in the population is critical for

timely public health interventions. The Alpha (B.1.1.7),B.1.351 (Beta), P.1(Gamma),

B.1.617.2 (Delta), and B.1.1.529 (Omicron) are examples of notable VOCs of SARS-

CoV-2 (Brief, 2021; Fujino et al., 2021; Galloway et al., 2020; Tegally et al., 2021;

WHO, 2021). In October 2020, the Delta variant surfaced in India, resulting in the

devastating second COVID-19 outbreak in the summer of 2021. During the Delta

wave, India's average death rate of 3 percent doubled over only three months (Jha et

al., 2022). The Omicron variant caused the surge in COVID-19 infection, which was

reported first in South Africa in late November 2021(He et al., 2021). Infectiousness

and transmission rates for the Omicron variant were more than those of the earlier

lineages (Duong et al., 2022; Wolter et al., 2022). Omicron is reportedly highly

resistant to antibody-directed inactivation by vaccine-triggered sera and antibodies

from both vaccinated and SARS-CoV-2 infected individuals (Hoffman et al., 2022;

Tada et al., 2022; Wilhelm et al., 2021).

Primarily, genomic surveillance of SARS-CoV-2 is accomplished through

clinical sample sequencing (Grimaldi et al., 2022). It has been observed and widely

accepted that sequencing all clinically positive samples is not economically feasible

(Tyson et al., 2020). Even though SARS-CoV-2 is primarily a respiratory virus, its

genome can also be found in infected patients' respiratory tract, stool, and

urine(Wölfel et al., 2020). Both symptomatic and asymptomatic cases shed the virus

in their stool, eventually reaching sewage treatment plants (STPs) (Mizumoto et al.,

2020; Nishiura et al., 2020; Treibel et al., 2020; Wan et al., 2020). Based on the above

findings, implementing WBE is an inexpensive virus surveillance approach that does

not infringe on human rights (Carducci et al., 2020;O'Brien and Xagoraraki, 2019). In

the recent past, WBE has been demonstrated to be instrumental during the H1N1

influenza outbreak and in detecting the prevalence of Poliovirus, Norovirus, Hepatitis

A and E virus, adenovirus, and rotavirus (Ahmed et al., 2021; Hellmér et al., 2014;

Lago et al., 2003). SARS-CoV-2 RNA is detected and quantified in wastewater using

quantitative reverse transcription polymerase chain reaction (qRT-PCR) (Hillary et al.,

2020). However, the qRT-PCR-based method for detecting variants in SARS-CoV-2

positive samples is only practical for already identified variants and cannot be utilized

to identify novel variants (Dikdan et al., 2022; Harper et al., 2021; Shen et al., 2021).

Wastewater surveillance is one of the most effective complementary epidemiological

methods for detecting the presence and evolution of SARS-CoV-2 variants (Amman

et al., 2022). Wastewater has been used to track SARS-CoV-2 quite early in the

pandemic (Randazzo et al., 2020; Sherchan et al., 2020). Several researchers have

asserted that SARS-CoV-2 can be found in wastewater from all over the world (Markt

et al., 2022; Pecia et al., 2022; Wurtz et al., 2021; Wurtzer et al., 2021). WBE allows

the detection of mutations before clinical testing is done (Bar-Or et al., 2021;

Karthikeyen et al., 2022). The Centers for Disease Control and Prevention (CDC),

Atlanta, USA, has also acknowledged the use of environmental wastewater

surveillance techniques to track and identify circulating SARS-CoV-2 variants and

issued guidelines on its use in the community National Wastewater Surveillance

System, CDC, USA).

In this study, we present the applicability of WBE to test the presence of

SARS-CoV-2 and its variants in sewage water from a research campus-based STP in

India. We collected samples from the STP inlet of the CSIR-National Chemical

Laboratory (CSIR-NCL) campus in Pune, India. To determine the SARS-CoV-2

frequency and genetic diversity of variants in campus sewage water, all samples were

analyzed using high-throughput next-generation sequencing (NGS) technology. We

also demonstrated that sewage water-based genomic surveillance could be a

preventive measure for averting outbreaks and a reliable tool for identifying novel

variants that would otherwise escape from variant-specific qRT-PCR assays.

2. Material and Methods

2.1 Sample collection and downstream processing

CSIR-National Chemical Laboratory, Pune, India (18°32'31.2"N; 73°48'43.2"E) is a

federal research institution and home to a closed society of approximately 5,000

employees and students (Fig.1). Domestic water is treated via a sewerage line

connected to a Phytorid STP with a capacity of 0.15 million liters per day (MLD).

The sample collection included a grab-sampling method, executed following the CDC,

USA guidelines of COVID-19 wastewater sampling, and approved by the Institutional

Biosafety Committee (IBSC). Between November 2021 and April 2022, one litre of

each of the 38 samples was collected from the inlet in a sterile plastic container

between 09:00 am and 10:00 am. All the samples were processed immediately after

collection per CDC, USA guidelines of COVID-19 sewage water sampling. The

sampling bottle was wiped with 70% ethanol and promptly subjected to 1h

pasteurization at 60°C. Of the total 1L of the sample, an aliquot of 160 ml was used

for the concentration of the RNA fragments of SARS-CoV-2 using the polyethylene

glycol (PEG) precipitation method (Ahmed et al., 2020). Briefly, the sample was

centrifuged at 5000xg (5910 R, Eppendorf, Germany) for 10 minutes at 4°C to collect

the larger debris as a pellet. Filtration was done twice, first with Whatman qualitative

filter paper grade 1 with a diameter of 47 mm (Merck, USA), and then with 0.22 µm

PES filter membrane (diameter 47mm) using vacuum filtration assembly (Tarsons,

India). The resulting filtrate was added to a 250ml HDPE bottle (HiMedia, India),

containing 8% (w/v) PEG 8000 (HiMedia, India) and 1.7% of molecular biology

grade NaCl (w/v) (HiMedia, India). This mixture was incubated for 16 h at 4°C in a

rotating shaker water bath (Equitron medica, India) at 175 rpm. It was centrifuged at

10000xg for 1 h at 4°C to collect the viral pellet. The supernatant was removed, and

the pellet was resuspended in 560 µL of TE buffer (pH 7.0). One part of the

resuspended pellet (280 µL) was stored at -80°C for future use, and the remaining

(280 µL) was used to extract viral RNA. QiagenQIAmp Viral RNA Minikit (Qiagen,

Germany) was used for viral RNA extraction according to the manufacturer's

instructions. The RNA was eluted in 80 µL of elution buffer and stored at -80°C for

downstream analysis. After obtaining permission from the Institutional Biosafety

Committee (IBSC), the used materials were disposed off.

2.2 Molecular detection

The qRT-PCR approach was used for SARS-CoV-2 RNA detection in sewage water

samples. The experiment was conducted using an ICMR-approved GenePath

Diagnostics qRT-PCR kit (GenePath DxCoViDx One v2.1.1TK-Quantitative). It is a

one-tube multiplex qRT-PCR assay that includes E, N and RdRP genes of SARS-

CoV-2, with the RNase P gene serving as a nucleic acid isolation control. Each

sample was tested in triplicates, wherein the PCR cocktail for individual reaction

contained 9.25 µL of master mix, 0.75 µL of a primer-probe mix, and 5 µL of

template in a 15 µL total reaction volume. The thermal cycling protocol consisted of

pre-incubation at 37°C for 5 min, reverse transcription for 7 min at 52°C, RT

inactivation at 95°C for 3 min 30 sec and 40 amplification cycles of denaturation at

95°C for 5 sec, and extension at 58 °C for 35 sec. All qRT-PCR experiments were

conducted using an Applied Biosystems 7500 Fast Real-Time PCR system (USA).

Viral load in samples was quantified using an online Covid-19 Viral Load Calculation

Tool provided by the kit's manufacturer.

2.3 SARS-CoV-2 whole genome sequencing

All the samples were sequenced at the Next Generation Genomics facility at the

Indian Institute of Science Education and Research (IISER), Pune, India. Sample

libraries were prepared using Illumina Covid seq RUO kits (Illumina, USA). Post-

annealing with random hexamers, the first strand synthesis reaction was carried out,

followed by cDNA amplification in two separate multiplex PCR reactions, including

primers designed to cover the entire genome of SARS-CoV-2. Using the Illumina

PCR Indexes, the amplified products were further processed for tagmentation and

adapter ligation, followed by enrichment and clean-up of amplicons as per the

manufacturer's instructions. Pooled libraries were quantified using the Qubit 4.0

fluorometer (Invitrogen, USA). All the pools were normalized to a final concentration

of 4.0nM using the Illumina Pooling calculator. For sequencing, the final pooled

library was denatured, neutralized, and adjusted to a final concentration of 1.4pM, and

paired-end sequencing was performed using the Illumina NextSeq 550 sequencing

platform (Illumina, USA).

2.4 Bioinformatic analysis

A dedicated in-house bioinformatics pipeline was used for analyzing raw data to

identify the main circulating lineages in sewage water. In the first step, raw reads

were checked for quality and filtered using fastp (Chen et al., 2018). After that, the

reference genome of SARS-CoV-2 (MN908947.3) was used to align the filtered reads

by BWA Mem (Li and Durbin, 2009).

Basic alignment and coverage statistics were determined by SAM tools

coverage/bedcov (Li et al., 2009; Quinlan and Hall, 2010). iVar was used for single

nucleotide variant (SNV) calling using a minimum base quality score of 20, minimum

frequency 0.03, and minimum read depth of 10 (Grubaugh et al., 2019). To predict the

lineage composition, we used LCS- a mixture model for determining SARS-CoV-2

variant composition in environmental samples (Valieris et al., 2022). Scripting was

done in Python/Bash to manually check variant-associated mutations, followed by

data plotting in R Studio.

3. Results

3.1 SARS-CoV-2 quantification

The goals of our study were to (i) detect and quantify SARS-CoV-2 genetic material

in campus sewage water, (ii) monitor SARS-CoV-2 infection trend on campus, (iii)

monitor viral variants of SARS-CoV-2 through NGS, and (iv) implement sewage

water surveillance within the campus, incorporating it as a component of the

pandemic response system. Overall, 38 sewage water samples collected from the

campus STP inlet from November 2021 to April 2022 were investigated for the

occurrence of SARS-CoV-2 RNA. Samples were considered positive when any two

of three target genes for SARS-CoV-2 showed Ct value ≤ 35 as per kit instructions.

Nineteen samples were positive, with Ct values ranging between 29.23 and 34.52,

representing a 50% overall positive detection rate (Supplementary Table 1). The

remaining 19 samples were negative.

Further, to understand the transmission dynamics of SARS-CoV-2 in sewage water at

the community level, we calculated the viral load of each sample in copies/ml using

the calculation tool provided by the qRT-PCR kit manufacturer (Fig.2). The first

significant increase in viral load was observed in sample PR-41 (111.44 copies/ml)

collected on November 22, 2021 (Supplementary Table1), two weeks before the first

positive case was reported on campus. Viral load was later observed to increase

exponentially from January 03, 2022 (PR-50) and the highest viral load was detected

on January 20, 2022 in sample PR-55 (526 copies/mL). SARS-CoV-2 RNA copies

continued to be found from February 2022 to April 2022, with a relatively higher viral

load in sample PR-62 (229 copies/mL). A total of 184 positive cases of COVID-19

were reported between November 2021 and April 2022 among the students, staff, and

faculty, or their family members residing on the CSIR-NCL campus (Supplementary

Table 2). However, the first case of COVID-19 on campus was reported to authorities

on December 06, 2021. This correlates with our experimental data showing an

increase in viral load 14 days before its first reporting. Pearson's correlation for viral

load showed a significant correlation ($R^2=0.68$, p-value>$7.9e^{-11}$) with COVID-19

cases reported on campus (Fig.2B). Although no new COVID-19 cases were reported

on campus after February 07, 2022, the SARS-CoV-2 RNA sustained with few spikes

in the concentration of viral load during February 2022 to April 2022.

3.2 Overview of SARS-CoV-2 sequencing, detected nucleotide substitutions and

deletions

All 38 sewage water samples collected during the third wave of Omicron from

November 2021 to April 2022 were sequenced using the Illumina Next Seq 500

platform. Samples were grouped into positive (n=19) and negative (n=19) categories

based on SARS-CoV-2 qRT-PCR (Supplementary Table 1). Since the qRT-PCR

method uses primer pairs that target specific viral fragments and the SARS-CoV-2

virus is susceptible to degradation in wastewater, that would eventually make

capturing viral fragments with limited qRT-PCR primer pairs challenging. As a result,

when wastewater is processed for qRT-PCR-based detection, the probability

phenomenon may pose a challenge to the limited qRT-PCR primers. Currently, the

whole genome NGS protocols use multiple primers targeting the entire genome, thus

increasing the probability of detecting the minute amount of nucleic acid fragments in

sewage water. Therefore, we sequenced all the samples regardless of qRT-PCR

results to find the SARS-CoV-2 fragments and to understand the mutation profile of

SARS-CoV-2 better.

A total of 124.16 million reads were generated from sequencing, of which

26.46 million reads were mapped to the SARS-CoV-2 reference genome

(MN908947.3). Out of 38 sequenced samples, 27 showed mean genomic coverage

above 50% ranging between 99.81% and 57.2%. At the same time, 16 out of 27

samples showed mean genome coverage uniformity above 90% (Supplementary

Table1). Sequencing results of qRT-PCR positive (n=19) and negative samples (n=19)

showed significant evidence for the presence of COVID-19 genomic material in the

samples, supported by the linear regression analysis between viral concentration and

reference genome coverage (Pearson correlation coefficient: R^2 =0.18, p-value>0.003)

(Fig.2.C). Their reads mapped ranged between 1.7 million and 34 thousand (Fig. 3).

Overall, 2108 unique nucleotide substitutions (Supplementary Fig.1) were

found compared with the reference Wuhan genome of SARS-CoV-2 (MN908947.3).

Among the nucleotide substitutions found: 941 mutations were in ORF1a polyprotein;

519 in ORF1ab polyprotein; 88 in ORF3a protein; 323 in spike protein; 19 in

envelope protein; 48 in membrane protein; 10 in ORF6 protein; 19 in ORF7a protein;

15 in ORF7b protein; 17 in ORF8 protein; 101 in nucleocapsid phosphoprotein and 08

in ORF10 protein. A total of 119 deletions were detected in samples, 61 in ORF1a, 22

in ORF1ab, 6 in ORF3a, 22 in spike protein; 03 in membrane glycoprotein; 01 in

ORF7a protein; 01 in ORF7b protein; and 03 in nucleocapsid phosphoprotein.

3.3 Early detection of Omicron signature mutations

Sequencing data were analyzed and searched for the presence of signature mutations

of different lineages of SARS-CoV-2VOCs, Alpha (B.1.1.7), Beta (B.1.351), Gamma

(P.1), Delta (B.1.617.2) and Omicron (B.1.1.529) and VOIs, Kappa (B.1.617.1),

Epsilon (B.1.427, B.1.429), Eta (B.1.525), Iota (B.1.526) and Lambda (C.37) (Fig. 4).

A total of 5214 signature mutations were detected, representing 16 highly

transmissible variants of COVID-19. Samples collected from November to December

2021 were highly dominated by Delta lineage (Fig.5A). The transition of Omicron

lineage over delta lineage was observed in late December 2021, which is supported by

the detection of a higher number of Omicron-associated signature mutations from

January 2022 onwards (Fig. 5B), Omicron mutations ranging between 30-42

nucleotide substitutions per sample, followed by continuous detection till March 2022.

The Omicron lineage of SARS-CoV-2 contains 62 characteristic deletions and non-

synonymous nucleotide substitutions, including 30 substitutions on the spike region

(Dejnirattisai et al., 2022). We detected the first signal of Omicron in sample PR-39

collected on November 08, 2021. A total of 11 signature mutations were detected, 09

from the spike gene (S:P681H, S:T478K, S:D614G, S:H655Y, S:N679K, S:N764K,

S:L452R, S:D796Y, S:T478K;), 01 from N gene (N:P13L) and 01 from ORF1a gene

(ORF1a:K856R). Similarly, 05 markers were detected in sample PR-40 (S:L452R,

S:T478K, S:D614G, ORF1a:T3255I, ORF1a:P3395H), 06 in PR-41 (S:L452R,

S:T478K, S:D614G, N:P13L, ORF1a:T3255I, ORF1a:P3395H) and 04 in PR-42

(S:L452R, S:T478K, S:D614G, ORF1a:T3255I) (Supplementary Tables 3 and 4).

3.4 SARS-CoV-2 prevalence and lineage trends in sewage water

LCS was used to ascertain the relative frequencies of SARS-CoV-2 variants in

sewage water. A total of 426 lineages were detected, representing 16 highly

transmissible variants of SARS-CoV-2 (VOCs and VOIs). We found that the Delta

lineage dominated in (B.1.617.2) November 2021 (68%) and December 2021 (61%);

after that Kappa (10% and 18%), BA.3 (5% and 7%), BA.2 (3%), BA.1 (2% and 0.1%)

and others (8%) (Fig.5).The Omicron family lineages increased exponentially from

January 2022 onwards and fully replaced Delta lineage by February 2022. BA.2.X

was the most dominant lineage (53.6% and 55%), followed by BA.1 (17% and 4%),

BA.2 (2% and 12%), BA.2.12 (0% and 12%), BA.5 (1% and 9%) and Iota (3% and

0.4%) respectively in January and February 2022. Finally, the linear-mixed effect

model was employed to understand the lineage trend from November 2021 to April

2022. The Omicron lineages (BA.1, BA.2, BA.4, and BA.5) showed an increasing

trend, whereas the Delta lineage (B.1.617.2) showed a decreasing trend from

November 2021 to February 2022(Fig. 6). **Fig.6:** Linear mixed model based monthly

proportion of SARS-CoV-2 lineages/sub-lineages

Discussion

Wastewater surveillance provides information on the viral load of SARS-

CoV-2 and the introduction of new variants (VOI/VOC) in the community (Johnson

et al., 2022; Oloye et al., 2022). WBE can be highly prioritised in India, especially

with the emergence of new variants and limited COVID-19 testing. Our study demonstrates how genomic surveillance of sewage water from on-campus STP can be utilised for detection, monitoring, and evaluation of the spread of SARS-CoV-2, which effectively helps to take decisions for the protection against SARS-CoV-2 in academic institutes/universities as documented by Betancourt et al., 2021; Corchis-Scottet al., 2021; Gibas C et al., 2021; Kantor et al., 2022; Karthikeyan et al., 2021; Mangwana et al., 2022; Wang et al., 2022; and Vo et al., 2022 across the world. Several studies have revealed the detection of SARS-CoV-2 from wastewater in India (Arora et al., 2022; Dharmadhikari et al., 2022;Joshi et al., 2022). However, to the best of our knowledge, this is the first study from the country to use the WBE technique to prevent the spread of SARS-CoV-2 on an academic campus.

A higher viral load of SARS-CoV-2 is strongly linked to increased disease virulence, progression, and death rate (Fajnzylber et al., 2020; Kawasuji et al., 2022). Serosurveillance of COVID-19 in Pune city showed a rise in daily new cases from the last week of December, 2021 (Supplementary Fig.3). Whereas by the end of January, 2022 the pick of daily active and new cases were observed, which decreased by the end of March, 2022 (http://cms.unipune.ac.in/~bspujari/Covid19/Pune2/). The trend of daily active and new cases reported in the serosurveillance complements the clinical case reported on campus. We observed an increase in the viral load of SARS-CoV-2 in sewage water prior to the first clinical case reported on campus. Recent studies from the USA (Al-Falsity et al., 2022;Karthikeyan et al., 2021; Wu et al., 2022), Mexico(Padilla-Reyes et al., 2022); Greece (Galani et al., 2022), England (Martin et al., 2020; Morvan et al., 2022), South Africa (Johnson et al., 2022), Brazil (de Araújo et al., 2022); Japan (Malla et al., 2022), and India (Joshi et al., 2022) also revealed similar results which are consistent with our findings. The strong link

between SARS-CoV-2 viral load in sewage water and reports of newly confirmed COVID-19 cases per day suggests that these infected people contribute substantial viral load to the sewage water (Hasan et al., 2021). These findings demonstrate WBE's early warning capability (Kirby et al., 2022; Lastra et al., 2022; Wang et al., 2022).

In most locations, the epidemic caused by the Omicron variant had a lower impact on health than previous COVID-19 waves due to increased herd immunity and possibly diminished nature of SARS-CoV-2 infections (Nealon and Cowling, 2022). Nyberg and colleagues also found that the Omicron variant is less severe than the Delta lineage of SARS-CoV-2 in terms of mortality and hospitalization (Nyberg et al., 2022; Post and Lorenzo-Redondo, 2022). Although the viral load was higher and continuously detected in the on-campus STP sewage water, the reported COVID-19-positive cases were low. The aforementioned higher viral load could be correlated with the incidence of several asymptomatic COVID-19-positive individuals on campus who did not test themselves for the underlying condition. Another attribute could be the overestimation of the number of COVID-19-positive cases. Therefore, our study supports using the WBE approach as a realistic indicator for monitoring SARS-CoV-2 and future outbreaks.

The qRT-PCR method has been established as a standard for assessing viral nucleic acid from patients and has also been used for environmental studies with its utility in WBE studies (Ahmed et al., 2022; Johnson et al., 2022). Initially, researchers depended on qRT-PCR and Sanger sequencing to establish the existence of viral fragments of SARS-CoV-2 in wastewater. WBE studies are founded on SARS-CoV-2 fecal shedding (Crank et al., 2022; Li et al., 2022); however, the fecal

matter would be vastly diluted by the addition of household sewage water and rain,

establishing a probability framework for detecting viral fragments (Novoa et al., 2022;

Wade et al., 2022). We detected SARS-CoV-2 in 100% of qRT-PCR negative

samples, concluding that the amplicon sequencing protocol is much more sensitive in

detecting the SARS-CoV-2 fragments in sewage water than qRT-PCR. Therefore, our

study emphasizes the efforts of sequencing, irrespective of the qRT-PCR result, not

only for viral detection but the monitoring of newly emerging variants, novel

mutations, and evolutionary studies of SARS-CoV-2.

Variant tracking from wastewater cannot confirm the presence of a specific

variant because COVID-19 genetic material in wastewater is often fragmented

(Gregory et al., 2022; Wolfe et al., 2022); as a result, it poses a significant challenge

to detect all characteristics mutations of a particular variant linked to a single genome.

However, detecting multiple mutations associated with a variant or unique mutations

absent in others increases the confidence of the presence of a specific variant in

wastewater (Kirby et al., 2022; Zahmatkesh et al., 2022). Therefore, we also analyzed

all the samples manually using the iVar SNP calling pipeline to gain in-depth insights

into the mutation profile of low-sequenced samples. Surprisingly, we found Omicron

signals in early November 2021, whereas Omicron was discovered for the first time in

Botswana on November 11, 2021 (Rahmani and Rezaei, 2022). Later, WHO

designated variant B.1.1529 as a VOC on November 26, 2021. India reported the first

case of Omicron on December 02, 2021, in Karnataka state, followed by the first

detection of Omicron in Maharashtra state on December 04, 2021. However, we

detected several unique Omicron-associated mutations in our data in early November

2021 (S:N679K, S:N764K, S:D796Y, N:P13L, ORF1a:T3255I, ORF1a:K856R,

ORF1a:P3395H, N:S413R) which provides substantial evidence that Omicron variant

was likely present earlier on campus in India before it was first detected clinically

across the globe. Similar reports from the USA and France also provide strong

evidence for the presence of the Omicron variant in early November 2021 in

community wastewater before the first detection from the clinical sample (Ferré et al.,

2022; Kirby et al., 2022).

Wastewater testing captures contributions from all residents within a locality,

which provides a less skewed outcome of viral genetic diversity and public health

(Karthikeyan et al., 2021; Safford et al., 2022; Sims and Kasprzyk-Hordern, 2020).

Clinical testing has limitations such as providing data only on those who agree to be

tested, missing asymptomatic cases, resource allocation, and affordability concerns.

We used LCS, a statistical model for estimating SARS-CoV-2 variants composition in

pooled samples (Valieris et al., 2022). Our samples showed significant SARS-CoV-2

lineage diversity across months, including lineages that were not reported from

clinical samples, highlighting the discrepancies in clinical testing and sequencing. We

identified the shift around January and February 2022, defined by a sharp rise of

Omicron lineages (BA.1, BA.2, BA.3, BA.4, BA.5) and a decrease of previously

dominant lineages (Delta and Kappa).

Furthermore, BA.4 and BA.5 signals were detected in January and February

2022, two months earlier than their first report from a Telangana, India clinical

sample in late March 2022. Hence, the emergence of variants in wastewater indicates

that a sizeable number of people were infected with specific variants, contributing to

the total viral load. One limitation of our study is the lack of sequencing data of

clinically reported positive cases from campus, which could have aided in correlating

with the emerging and prevailing variants of SARS-CoV-2 circulating on the campus.

Throughout the surveillance period, we informed decision makers (COVID-

19 task force) about SARS-CoV-2 detection in sewage water of campus-STP

approximately 48 h after sample collection, which helped in policy-making decisions

to alert the community on campus. Last, our work illustrates the utility of WBE for

monitoring COVID-19 on campus, and similar settings could help manage SARS-

CoV-2 and other future emerging pathogens of concern.

4. Conclusions

This study concludes that the use of WBE at the community level would be an

effective means of preventing SARS-CoV-2 outbreaks. An increase in viral load in

the community detected early with the help qRT-PCR approach indicates its

importance as an alarming feature of WBE. The amplicon sequencing protocol can be

sensitive in detecting the fragments of SARS-CoV-2 in wastewater, which were failed

to get caught in qRT-PCR. It can also be concluded that to provide comprehensive

information about population infection dynamics; more emphasis should be on the

amplicon sequencing-based approaches to gain information on the emerging variants.

Overall, this work has shown that implementing real-time WBE on-campus and

integrating it with the response system would effectively prevent and contain future

outbreaks.

Data availability: All data produced in the present study are available upon

reasonable request to the authors. Clinical COVID-19 data was used from CMS

dashboard (http://cms.unipune.ac.in/~bspujari/Covid19/Pune2/)

Credit authorship contribution statement

Vinita Malik: Conceptualization, Data curation, Formal analysis, Investigation,

Methodology, Validation, Roles/Writing - original draft

Vinay Rajput: Data curation, Formal analysis, Investigation, Methodology, Software,

Visualization, Roles/Writing - original draft

Rinka Pramanik: Investigation, Methodology, Visualization, Roles/Writing -

original draft

Rachel Samson: Writing - review & editing

Rakeshkumar Yadav: Writing - review & editing

Pradnya Kadam: Methodology

Nikita Shah: Methodology

Rutuja Sawant: Methodology

Unnati Bhalerao: Methodology

Manisha Tupekar: Methodology

Soumen Khan: Methodology

Priyanki Shah: Funding acquisition, Project administration, Supervision

LS Shashidhara: Funding acquisition, Project administration, Supervision

Sanjay Kamble: Investigation, Formal analysis

Syed Dastager: Investigation, Formal analysis

Krishanpal Karmodiya: Formal analysis

Mahesh Dharne: Conceptualization, Funding acquisition, Investigation, Project

administration, Resources, Supervision, Writing - review & editing.

Declaration of competing interest: The authors declare that they have no known

competing financial interests or personal relationships that could have appeared to

influence the work reported in this paper.

Acknowledgments

This work is funded by a grant from Rockefeller Foundation, USA (Project code GAP

332926). The authors express their gratitude to the Director General, Council of

Scientific and Industrial Research (CSIR), New Delhi; Directors of CSIR-NCL,,

IISER-Pune and Pune Knowledge Cluster, for encouragement and support. Authors

are indebted to Prof. Aurnab Ghose (IISER-Pune for grant administration support and

discussions). Thanks to Dr. Narendra Kadoo (Chair, NCL Covid task force) and Dr.

Chetan Gadgil (CSIR-NCL) for their valuable suggestions. Thanks to the

Administrative office for providing on-campus case records and the Engineering and

Civil services section of CSIR-NCL for all the support during study. The manuscript

has been thoroughly checked for plagiarism content using iThenticate software.

References

Ahmed, W., Angel, N., Edson, J., Bibby, K., Bivins, A., O'Brien, J.W., Choi, P.M.,

Kitajima, M., Simpson, S.L., Li, J., 2020. First confirmed detection of SARS-

CoV-2 in untreated wastewater in Australia: a proof of concept for the

wastewater surveillance of COVID-19 in the community. Science of the Total

Environment 728, 138764.

Ahmed, W., Smith, W.J.M., Metcalfe, S., Jackson, G., Choi, P.M., Morrison, M.,

Field, D., Gyawali, P., Bivins, A., Bibby, K., 2022. Comparison of RT-qPCR

and RT-dPCR platforms for the trace detection of SARS-CoV-2 RNA in

wastewater. ACS ES&T Water.

Ahmed, W., Tscharke, B., Bertsch, P.M., Bibby, K., Bivins, A., Choi, P., Clarke, L.,

Dwyer, J., Edson, J., Nguyen, T.M.H., 2021. SARS-CoV-2 RNA monitoring in

wastewater as a potential early warning system for COVID-19 transmission in

the community: A temporal case study. Science of The Total Environment 761,

144216.

Al-Faliti, M., Kotlarz, N., McCall, C., Harris, A.R., Smith, A.L., Stadler, L.B., de los

Reyes, F.L.I.I.I., Delgado Vela, J., 2022. Comparing Rates of Change in SARS-

CoV-2 Wastewater Load and Clinical Cases in 19 Sewersheds Across Four

Major Metropolitan Areas in the United States. ACS ES&T Water.

https://doi.org/10.1021/acsestwater.2c00106

Amman, F., Markt, R., Endler, L., Hupfauf, S., Agerer, B., Schedl, A., Richter, L.,

Zechmeister, M., Bicher, M., Heiler, G., 2022. Viral variant-resolved wastewater

surveillance of SARS-CoV-2 at national scale. Nat Biotechnol 1–9.

Bar-Or, I., Weil, M., Indenbaum, V., Bucris, E., Bar-Ilan, D., Elul, M., Levi, N.,

Aguvaev, I., Cohen, Z., Shirazi, R., 2021. Detection of SARS-CoV-2 variants by

genomic analysis of wastewater samples in Israel. Science of The Total

Environment 789, 148002.

Chen, R.E., Zhang, X., Case, J.B., Winkler, E.S., Liu, Y., VanBlargan, L.A., Liu, J.,

Errico, J.M., Xie, X., Suryadevara, N., 2021. Resistance of SARS-CoV-2

variants to neutralization by monoclonal and serum-derived polyclonal

antibodies. Nat Med 27, 717–726.

Chen, S., Zhou, Y., Chen, Y., Gu, J., 2018. fastp: an ultra-fast all-in-one FASTQ

preprocessor. Bioinformatics 34, i884–i890.

Clark, A.E., Wang, Z., Cantarel, B., Kanchwala, M., Xing, C., Chen, L., Irwin, P., Xu,

Y., Oliver, D., Lee, F., 2021. Multiplex fragment analysis identifies SARS-CoV-

2 variants. MedRxiv.

Crank, K., Chen, W., Bivins, A., Lowry, S., Bibby, K., 2022. Contribution of SARS-

CoV-2 RNA shedding routes to RNA loads in wastewater. Science of The Total

Environment 806, 150376.

Crits-Christoph, A., Kantor, R.S., Olm, M.R., Whitney, O.N., Al-Shayeb, B., Lou,

Y.C., Flamholz, A., Kennedy, L.C., Greenwald, H., Hinkle, A., 2021. Genome

sequencing of sewage detects regionally prevalent SARS-CoV-2 variants. mBio

12, e02703-20.

de Araújo, J.C., Mota, V.T., Teodoro, A., Leal, C., Leroy, D., Madeira, C., Machado,

E.C., Dias, M.F., Souza, C.C., Coelho, G., Bressani, T., Morandi, T., Freitas,

G.T.O., Duarte, A., Perdigão, C., Tröger, F., Ayrimoraes, S., de Melo, M.C.,

Laguardia, F., Reis, M.T.P., Mota, C., Chernicharo, C.A.L., 2022. Long-term

monitoring of SARS-CoV-2 RNA in sewage samples from specific public places

and STPs to track COVID-19 spread and identify potential hotspots. Science of

The Total Environment 838, 155959.

https://doi.org/https://doi.org/10.1016/j.scitotenv.2022.155959

Dejnirattisai, W., Huo, J., Zhou, D., Zahradník, J., Supasa, P., Liu, C., Duyvesteyn,

H.M.E., Ginn, H.M., Mentzer, A.J., Tuekprakhon, A., 2022. SARS-CoV-2

Omicron-B. 1.1. 529 leads to widespread escape from neutralizing antibody

responses. Cell 185, 467–484.

Dharmadhikari, T., Rajput, V., Yadav, R., Boargaonkar, R., Patil, D., Kale, S.,

Kamble, S.P., Dastager, S.G., Dharne, M.S., 2022. High throughput sequencing

based direct detection of SARS-CoV-2 fragments in wastewater of Pune, West

India. Science of The Total Environment 807, 151038.

Duong, D.B., King, A.J., Grépin, K.A., Hsu, L.Y., Lim, J.F.Y., Phillips, C., Thai, T.T.,

Venkatachalam, I., Vogt, F., Yam, E.L.Y., 2022. Strengthening national

capacities for pandemic preparedness: a cross-country analysis of COVID-19

cases and deaths. Health Policy Plan 37, 55–64.

Fajnzylber, J., Regan, J., Coxen, K., Corry, H., Wong, C., Rosenthal, A., Worrall, D.,

Giguel, F., Piechocka-Trocha, A., Atyeo, C., 2020. SARS-CoV-2 viral load is

associated with increased disease severity and mortality. Nat Commun 11, 1–9.

Galani, A., Aalizadeh, R., Kostakis, M., Markou, A., Alygizakis, N., Lytras, T.,

Adamopoulos, P.G., Peccia, J., Thompson, D.C., Kontou, A., 2022. SARS-CoV-

2 wastewater surveillance data can predict hospitalizations and ICU admissions.

Science of The Total Environment 804, 150151.

Gregory, D.A., Trujillo, M., Rushford, C., Flury, A., Kannoly, S., San, K.M., Lyfoung,

D., Wiseman, R.W., Bromert, K., Zhou, M.-Y., 2022. Genetic Diversity and

Evolutionary Convergence of Cryptic SARS-CoV-2 Lineages Detected Via

Wastewater Sequencing. medRxiv.

Grimaldi, A., Panariello, F., Annunziata, P., Giuliano, T., Daniele, M., Pierri, B.,

Colantuono, C., Salvi, M., Bouché, V., Manfredi, A., 2022. Improved SARS-

CoV-2 sequencing surveillance allows the identification of new variants and

signatures in infected patients. Genome Med 14, 1–15.

Grubaugh, N.D., Gangavarapu, K., Quick, J., Matteson, N.L., de Jesus, J.G., Main,

B.J., Tan, A.L., Paul, L.M., Brackney, D.E., Grewal, S., 2019. An amplicon-

based sequencing framework for accurately measuring intrahost virus diversity

using PrimalSeq and iVar. Genome Biol 20, 1–19.

He, X., Hong, W., Pan, X., Lu, G., Wei, X., 2021. SARS‑CoV‑2 Omicron variant:

characteristics and prevention. MedComm (Beijing) 2, 838–845.

Hillary, L.S., Malham, S.K., McDonald, J.E., Jones, D.L., 2020. Wastewater and

public health: the potential of wastewater surveillance for monitoring COVID-19.

Curr Opin Environ Sci Health 17, 14–20.

Izquierdo-Lara, R., Elsinga, G., Heijnen, L., Munnink, B.B.O., Schapendonk, C.M.E.,

Nieuwenhuijse, D., Kon, M., Lu, L., Aarestrup, F.M., Lycett, S., 2021.

Monitoring SARS-CoV-2 circulation and diversity through community

wastewater sequencing, the Netherlands and Belgium. Emerg Infect Dis 27, 1405.

Jha, P., Deshmukh, Y., Tumbe, C., Suraweera, W., Bhowmick, A., Sharma, S.,

Novosad, P., Fu, S.H., Newcombe, L., Gelband, H., 2022. COVID mortality in

India: National survey data and health facility deaths. Science (1979) 375, 667–

671.

Johnson, R., Sharma, J.R., Ramharack, P., Mangwana, N., Kinnear, C., Viraragavan,

A., Glanzmann, B., Louw, J., Abdelatif, N., Reddy, T., 2022. Tracking the

circulating SARS-CoV-2 variant of concern in South Africa using wastewater-

based epidemiology. Sci Rep 12, 1–12.

Joshi, M., Kumar, M., Srivastava, V., Kumar, D., Rathore, D.S., Pandit, R., Graham,

D.W., Joshi, C.G., 2022. Genetic sequencing detected the SARS-CoV-2 delta

variant in wastewater a month prior to the first COVID-19 case in Ahmedabad

(India). Environmental Pollution 310, 119757.

Karthikeyan, S., Nguyen, A., McDonald, D., Zong, Y., Ronquillo, N., Ren, J., Zou, J.,

Farmer, S., Humphrey, G., Henderson, D., 2021. Rapid, large-scale wastewater

surveillance and automated reporting system enable early detection of nearly 85%

of COVID-19 cases on a university campus. mSystems 6, e00793-21.

Kawasuji, H., Morinaga, Y., Tani, H., Yoshida, Y., Takegoshi, Y., Kaneda, M., Murai,
Y., Kimoto, K., Ueno, A., Miyajima, Y., 2022. SARS☐CoV☐2 RNAemia with a
higher nasopharyngeal viral load is strongly associated with disease severity and
mortality in patients with COVID☐19. J Med Virol 94, 147–153.

Kirby, A.E., Welsh, R.M., Marsh, Z.A., Alexander, T.Y., Vugia, D.J., Boehm, A.B.,
Wolfe, M.K., White, B.J., Matzinger, S.R., Wheeler, A., 2022. Notes from the
field: early evidence of the SARS-CoV-2 B. 1.1. 529 (Omicron) variant in
community wastewater—United States, November–December 2021. Morbidity
and Mortality Weekly Report 71, 103.

Lastra, A., Botello, J., Pinilla, A., Urrutia, J.I., Canora, J., Sánchez, J., Fernández, P.,
Candel, F.J., Zapatero, A., Ortega, M., 2022. SARS-CoV-2 detection in
wastewater as an early warning indicator for COVID-19 pandemic. Madrid
region case study. Environ Res 203, 111852.

Li, H., Durbin, R., 2009. Fast and accurate short read alignment with Burrows–
Wheeler transform. bioinformatics 25, 1754–1760.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
Abecasis, G., Durbin, R., 2009. The sequence alignment/map format and
SAMtools. Bioinformatics 25, 2078–2079.

Li, X., Kulandaivelu, J., Guo, Y., Zhang, S., Shi, J., O'Brien, J., Arora, S., Kumar, M.,
Sherchan, S.P., Honda, R., 2022. SARS-CoV-2 shedding sources in wastewater
and implications for wastewater-based epidemiology. J Hazard Mater 432,
128667.

Malla, B., Thakali, O., Shrestha, S., Segawa, T., Kitajima, M., Haramoto, E., 2022.
Application of a high-throughput quantitative PCR system for simultaneous

monitoring of SARS-CoV-2 variants and other pathogenic viruses in wastewater.

Science of the Total Environment 853, 158659.

Mangwana, N., Archer, E., Muller, C.J.F., Preiser, W., Wolfaardt, G., Kasprzyk-

Hordern, B., Carstens, A., Brocker, L., Webster, C., McCarthy, D., 2022.

Sewage surveillance of SARS-CoV-2 at student campus residences in the

Western Cape, South Africa. Science of The Total Environment 851, 158028.

Markt, R., Endler, L., Amman, F., Schedl, A., Penz, T., Büchel-Marxer, M.,

Grünbacher, D., Mayr, M., Peer, E., Pedrazzini, M., 2022. Detection and

abundance of SARS-CoV-2 in wastewater in Liechtenstein, and the estimation of

prevalence and impact of the B. 1.1. 7 variant. J Water Health 20, 114–125.

Martin, J., Klapsa, D., Wilton, T., Zambon, M., Bentley, E., Bujaki, E., Fritzsche, M.,

Mate, R., Majumdar, M., 2020. Tracking SARS-CoV-2 in sewage: evidence of

changes in virus variant predominance during COVID-19 pandemic. Viruses 12,

1144.

Mizumoto, K., Kagaya, K., Zarebski, A., Chowell, G., 2020. Estimating the

asymptomatic proportion of coronavirus disease 2019 (COVID-19) cases on

board the Diamond Princess cruise ship, Yokohama, Japan, 2020.

Eurosurveillance 25, 2000180.

Morvan, M., Lojacomo, A., Souque, C., Wade, M., Hoffmann, T., Pouwels, K.,

Singer, A., Bunce, J., Engeli, A., Grimsley, J., 2022. Estimating SARS-CoV-2

prevalence from large-scale wastewater surveillance: insights from combined

analysis of 44 sites in England. International Journal of Infectious Diseases 116,

S24.

Nealon, J., Cowling, B.J., 2022. Omicron severity: milder but not mild. Lancet 399,

412.

Nishiura, H., Kobayashi, T., Miyama, T., Suzuki, A., Jung, S., Hayashi, K., Kinoshita,

R., Yang, Y., Yuan, B., Akhmetzhanov, A.R., 2020. Estimation of the

asymptomatic ratio of novel coronavirus infections (COVID-19). International

journal of infectious diseases 94, 154–155.

Novoa, B., Ríos-Castro, R., Otero-Muras, I., Gouveia, S., Cabo, A., Saco, A., Rey-

Campos, M., Pájaro, M., Fajar, N., Aranguren, R., 2022. Wastewater and marine

bioindicators surveillance to anticipate COVID-19 prevalence and to explore

SARS-CoV-2 diversity by next generation sequencing: One-year study. Science

of the Total Environment 833, 155140.

Nyberg, T., Ferguson, N.M., Nash, S.G., Webster, H.H., Flaxman, S., Andrews, N.,

Hinsley, W., Bernal, J.L., Kall, M., Bhatt, S., 2022. Comparative analysis of the

risks of hospitalisation and death associated with SARS-CoV-2 omicron (B. 1.1.

529) and delta (B. 1.617. 2) variants in England: a cohort study. The Lancet 399,

1303–1312.

Padilla-Reyes, D.A., Álvarez, M.M., Mora, A., Cervantes-Avilés, P.A., Kumar, M.,

Loge, F.J., Mahlknecht, J., 2022. Acquired insights from the long-term

surveillance of SARS-CoV-2 RNA for COVID-19 monitoring: The case of

Monterrey Metropolitan Area (Mexico). Environ Res 210, 112967.

https://doi.org/https://doi.org/10.1016/j.envres.2022.112967

Pérez-Cataluña, A., Chiner-Oms, Á., Cuevas-Ferrando, E., Díaz-Reolid, A., Falcó, I.,

Randazzo, W., Girón-Guzmán, I., Allende, A., Bracho, M.A., Comas, I., 2022.

Spatial and temporal distribution of SARS-CoV-2 diversity circulating in

wastewater. Water Res 211, 118007.

Post, L.A., Lorenzo-Redondo, R., 2022. Omicron: fewer adverse outcomes come with

new dangers. The Lancet 399, 1280–1281.

Quinlan, A.R., Hall, I.M., 2010. BEDTools: a flexible suite of utilities for comparing

genomic features. Bioinformatics 26, 841–842.

https://doi.org/10.1093/bioinformatics/btq033

Rahmani, S., Rezaei, N., 2022. SARS-CoV-2 Omicron (B. 1.1. 529) Variant: No

Time to Wait! Acta Bio Medica: Atenei Parmensis 93.

Randazzo, W., Cuevas-Ferrando, E., Sanjuán, R., Domingo-Calap, P., Sánchez, G.,

2020. Metropolitan wastewater analysis for COVID-19 epidemiological

surveillance. Int J Hyg Environ Health 230, 113621.

Safford, H.R., Shapiro, K., Bischel, H.N., 2022. Wastewater analysis can be a

powerful public health tool—if it's done sensibly. Proceedings of the National

Academy of Sciences 119, e2119600119.

Scott, L.C., Aubee, A., Babahaji, L., Vigil, K., Tims, S., Aw, T.G., 2021. Targeted

wastewater surveillance of SARS-CoV-2 on a university campus for COVID-19

outbreak detection and mitigation. Environ Res 200, 111374.

Sherchan, S.P., Shahin, S., Ward, L.M., Tandukar, S., Aw, T.G., Schmitz, B., Ahmed,

W., Kitajima, M., 2020. First detection of SARS-CoV-2 RNA in wastewater in

North America: a study in Louisiana, USA. Science of The Total Environment

743, 140621.

Sims, N., Kasprzyk-Hordern, B., 2020. Future perspectives of wastewater-based

epidemiology: monitoring infectious disease spread and resistance to the

community level. Environ Int 139, 105689.

Tada, T., Zhou, H., Dcosta, B.M., Samanovic, M.I., Chivukula, V., Herati, R.S.,

Hubbard, S.R., Mulligan, M.J., Landau, N.R., 2022. Increased resistance of

SARS-CoV-2 Omicron variant to neutralization by vaccine-elicited and

therapeutic antibodies. EBioMedicine 78, 103944.

Treibel, T.A., Manisty, C., Burton, M., McKnight, Á., Lambourne, J., Augusto, J.B.,

Couto-Parada, X., Cutino-Moguel, T., Noursadeghi, M., Moon, J.C., 2020.

COVID-19: PCR screening of asymptomatic health-care workers at London

hospital. The Lancet 395, 1608–1610.

Tyson, J.R., James, P., Stoddart, D., Sparks, N., Wickenhagen, A., Hall, G., Choi, J.H.,

Lapointe, H., Kamelian, K., Smith, A.D., 2020. Improvements to the ARTIC

multiplex PCR method for SARS-CoV-2 genome sequencing using nanopore.

BioRxiv.

Valieris, R., Drummond, R.D., Defelicibus, A., Dias-Neto, E., Rosales, R.A., Tojal da

Silva, I., 2022. A mixture model for determining SARS-Cov-2 variant

composition in pooled samples. Bioinformatics 38, 1809–1815.

Vo, V., Tillett, R.L., Chang, C.-L., Gerrity, D., Betancourt, W.Q., Oh, E.C., 2022.

SARS-CoV-2 variant detection at a university dormitory using wastewater

genomic tools. Science of The Total Environment 805, 149930.

Vogels, C.B.F., Breban, M.I., Ott, I.M., Alpert, T., Petrone, M.E., Watkins, A.E.,

Kalinich, C.C., Earnest, R., Rothman, J.E., Goes de Jesus, J., 2021. Multiplex

qPCR discriminates variants of concern to enhance global surveillance of SARS-

CoV-2. PLoS Biol 19, e3001236.

Wade, M.J., Jacomo, A. lo, Armenise, E., Brown, M.R., Bunce, J.T., Cameron, G.J.,

Fang, Z., Gilpin, D.F., Graham, D.W., Grimsley, J.M.S., 2022. Understanding

and managing uncertainty and variability for wastewater monitoring beyond the

pandemic: Lessons learned from the United Kingdom national COVID-19

surveillance programmes. J Hazard Mater 424, 127456.

Wan, Y., Shang, J., Graham, R., Baric, R.S., Li, F., 2020. Receptor recognition by the novel coronavirus from Wuhan: an analysis based on decade-long structural studies of SARS coronavirus. J Virol 94, e00127-20.

Wang, H., Miller, J.A., Verghese, M., Sibai, M., Solis, D., Mfuh, K.O., Jiang, B., Iwai, N., Mar, M., Huang, C., 2021. Multiplex SARS-CoV-2 genotyping PCR for population-level variant screening and epidemiologic surveillance. medRxiv.

Wang, Y., Liu, P., Zhang, H., Ibaraki, M., VanTassell, J., Geith, K., Cavallo, M., Kann, R., Saber, L., Kraft, C.S., 2022. Early warning of a COVID-19 surge on a university campus based on wastewater surveillance for SARS-CoV-2 at residence halls. Science of The Total Environment 821, 153291.

Wilhelm, A., Widera, M., Grikscheit, K., Toptan, T., Schenk, B., Pallas, C., Metzler, M., Kohmer, N., Hoehl, S., Helfritz, F.A., 2021. Reduced neutralization of SARS-CoV-2 Omicron variant by vaccine sera and monoclonal antibodies. MedRxiv.

Wolfe, M., Hughes, B., Duong, D., Chan-Herur, V., Wigginton, K.R., White, B.J., Boehm, A.B., 2022. Detection of SARS-CoV-2 Variants Mu, Beta, Gamma, Lambda, Delta, Alpha, and Omicron in Wastewater Settled Solids Using Mutation-Specific Assays Is Associated with Regional Detection of Variants in Clinical Samples. Appl Environ Microbiol 88, e00045-22.

Wölfel, R., Corman, V.M., Guggemos, W., Seilmaier, M., Zange, S., Müller, M.A., Niemeyer, D., Jones, T.C., Vollmar, P., Rothe, C., 2020. Author Correction: Virological assessment of hospitalized patients with COVID-2019. Nature 588, E35–E35.

Wolter, N., Jassat, W., Walaza, S., Welch, R., Moultrie, H., Groome, M., Amoako, D.G., Everatt, J., Bhiman, J.N., Scheepers, C., 2022. Early assessment of the

clinical severity of the SARS-CoV-2 omicron variant in South Africa: a data

linkage study. The Lancet 399, 437–446.

Wu, F., Xiao, A., Zhang, J., Moniz, K., Endo, N., Armas, F., Bonneau, R., Brown,

M.A., Bushman, M., Chai, P.R., 2022. SARS-CoV-2 RNA concentrations in

wastewater foreshadow dynamics and clinical presentation of new COVID-19

cases. Science of The Total Environment 805, 150121.

Wurtz, N., Penant, G., Jardot, P., Duclos, N., la Scola, B., 2021. Culture of SARS-

CoV-2 in a panel of laboratory cell lines, permissivity, and differences in growth

profile. European Journal of Clinical Microbiology & Infectious Diseases 40,

477–484.

Xagoraraki, I., O'Brien, E., 2020. Wastewater-based epidemiology for early detection

of viral outbreaks, in: Women in Water Quality. Springer, pp. 75–97.

Zahmatkesh, S., Sillanpaa, M., Rezakhani, Y., Wang, C., 2022. Review of Concerned

SARS-CoV-2 Variants Like Alpha (B. 1.1. 7), Beta (B. 1.351), Gamma (P. 1),

Delta (B. 1.617. 2), and Omicron (B. 1.1. 529), as well as Novel Methods for

Reducing and Inactivating SARS-CoV-2 Mutants in Wastewater Treatment

Facilities. Journal of Hazardous Materials Advances 100140.

Zhu, N., Zhang, D., Wang, W., Li, X., Yang, B., Song, J., Zhao, X., Huang, B., Shi,

W., Lu, R., 2020. A novel coronavirus from patients with pneumonia in China,

2019. New England journal of medicine.

Fig.1: Sampling site-campus STP (18°32'31.2"N; 73°48'43.2"E)

Fig.2: Analysis of SARS-CoV-2 viral load in WW against reported COVID-19 cases

on campus. A. SARS-CoV-2 viral load dynamics in sewage water and reported

COVID-19 cases on campus. B. Linear regression correlation plot of viral

concentration in wastewater with reported COVID-19 cases on-campus. (Pearson

correlation coefficient: $R^2 = 0.68$, p-value>$7.9e^{-11}$). C. Linear regression correlation

plot of viral concentration in wastewater with genome coverage (Pearson correlation

coefficient: $R^2 = 0.18$, p-value>0.003)

Fig. 3: Distribution of the number of reads mapped to the SARS-CoV-2 reference

genome per sample (MN908947.3)

Fig. 4: Detection of transition between SARS-CoV-2 VOC Delta and Omicron by

sewage water genomic surveillance.

Fig. 5: LCS analysis-based distribution of different lineages/sub-lineages of SARS-

CoV-2. **A**. Relative abundance in the percentage of the identified SARS-CoV-2

lineages/sub-lineages**. B**. Proportion of SARS-CoV-2 lineages from November 2021-

February, 2022.

Fig.6: Linear mixed model based monthly proportion of SARS-CoV-2 lineages/sub-

lineages

Fig.1: Sampling site-campus STP (18°32'31.2"N 73°48'43.2"E)

Fig.2 Analysis of SARS-CoV-2 viral load in WW against reported COVID-19 cases on campus. **A.** SARS-CoV-2 viral load dynamics in sewage water and reported COVID-19 cases on campus. **B.** Linear regression correlation plot of viral concentration in wastewater with reported COVID-19 cases on-campus. (Pearson correlation coefficient: R^2 =0.68, p-value>7.9e-11). **C.** Linear regression correlation plot of viral concentration in wastewater with genome coverage (Pearson correlation coefficient: R^2 =0.18, p-value>0.003)

Fig. 3: Distribution of the number of reads mapped to SARS-CoV-2 reference genome per sample (MN908947.3).

Fig. 4: Detection of transition between SARS-CoV-2 VOC Delta and Omicron by sewage water genomic surveillance

Samples chosen have more than 50% of genome coverage with the reference genome (MN908947.3).

Fig. 5: LCS analysis-based distribution of different lineages/sub-lineages of SARS-CoV-2. **A**. Relative abundance in the percentage of the identified SARS-CoV-2 lineages/sub-lineages. **B**. Proportion of SARS-CoV-2 lineages from November 2021-February, 2022.

Each box represents different variants and each line represents its trend.

Fig. 6: Linear mixed model based monthly proportion of SARS-CoV-2 lineages/sub-lineages.