**<u>TITLE PAGE</u>**

**1. Title**

- A Multi-layer Perceptron Neural Network for Predicting the Osteoporosis in Women Using Physical

Activity Factors

**2. Corresponding author:**

Jun-hee Kim (JH Kim), PhD, PT

Address: 1, Yeonsedae-gil, Maeji-ri, Heungeop-myeon, Wonju-si, Gangwon-do, 26493, Laboratory of

KEMA AI Research (KAIR), Department of Physical Therapy, College of Software and Digital

Healthcare Convergence, Yonsei University, Wonju, South Korea.

TEL: 82-33-760-2497 / FAX: 82-33-760-2496 / E-mail: move@yonsei.ac.kr

**ABSTRACT**

**Introduction**: Osteoporosis (OP) is a bone disease caused by a decrease in bone mineral density (BMD). OP is common in women because BMD gradually decreases after age 35. OP due to decreased BMD is highly likely to cause fatal traumatic injuries such as hip fracture. The purpose of this study was developed and evaluated a multi-layer perceptron neural network model that predicts OP using physical characteristics and activity factors of adult women over the age of 35 whose BMD begins to decline.

**Materials and Methods**: Data from KNHANES were used to develop a multi-layer perceptron model for predicting OP. Data preprocessing included variable selection and sample balancing, and LASSO was used for feature selection. The model used 5 hidden layers, dropout and batch normalization and was evaluated using evaluation scores such as accuracy and recall score calculated from a confusion matrix.

**Results**: Models were trained and evaluated to predict OP using selected features including age, quality of life index, weight, grip strength and average working hours per week. The model achieved 76.8% accuracy, 74.5% precision, 80.5% recall, 77.4% F1 score, and 74.8% ROC AUC.

**Conclusion**: A multi-layer perceptron neural network for predicting OP diagnosis using physical characteristics and activity factors in women aged 35 years or older showed relatively good performance. Since the selected variables can be easily measured through surveys, assessment tool, and digital hand dynamometer, this model will be useful for screening elderly women with OP or not in areas with poor medical facilities or difficult access.

**Keywords**: Osteoporosis, predictive model, multi-layer perceptron, deep learning, physical activity

## Introduction

Osteoporosis (OP) is a bone disease that occurs when bone mineral density (BMD) and bone mass decrease [1]. In other words, a decrease in BMD below a certain level is regarded as the onset of OP [1]. In women, rapid BMD and skeletal growth occur from menarche, when the secretion of sex steroids, including estrogen, increases [2]. BMD in adult women peaks between the ages of 25 and 35, after which women experience a gradual decline throughout their lives [2]. A decrease in BMD is prominent from menopause, when female hormones decrease, resulting in a higher rate of OP than in men [3, 4]. In particular, estrogen is thought to be important for maintaining BMD, and it has been reported that bone loss may occur as estrogen levels drop during menopause, resulting in a much higher rate of OP [5, 6].

Patients diagnosed with OP have a high risk of fracture due to trauma compared to normal people [7–9]. In particular, the femur, which makes up the hip joint, is a part where fractures commonly occur due to OP, and hip fracture can cause great difficulties in independent activities of daily life, and in severe cases, can be life threatening [10]. The 1-year mortality rate due to hip fracture increases by 2% every year, and in the case of women, the 10-year mortality rate after hip fracture is reported to be 16% [11]. In addition, once a fracture occurs, the risk of a second fracture increases more than twice, and when a second fracture occurs, the patient's mortality rate is much higher [12]. The mortality within 42 months after a second hip fracture was 57.2% [13]. Because the mortality rate of patients with hip fractures is high, it is important to predict or diagnose OP early to prevent hip fractures from occurring.

BMD is measured using dual energy X-ray absorptiometry (DXA) scans of the spine and hip to diagnose OP [14]. For postmenopausal women, BMD is determined by T-score compared to reference values of race- and sex-matched young adults [14]. However, one of the disadvantages of diagnosing OP through DXA scan is that measurement errors may occur due to surrounding soft tissue [15]. Therefore, recently, studies applying machine learning such as deep learning to predict OP with high performance or to predict OP with a simpler X-ray imaging technique have been actively conducted. Learning hip radiographs and clinical data through deep learning can improve diagnostic performance

for OP classification and prediction even without using the DXA scans [16]. Similarly, diagnosis

using deep learning based on lumbar radiographs had the potential to screen for OP and osteopenia

even without using DXA scans [17]. In addition, The CT image-based deep learning diagnosis method

showed higher accuracy and could output higher specificity or sensitivity for OP evaluation [18].

Deep learning models that predict OP using these medical diagnostic devices have been studied, but

deep learning models that predict OP according to data that can be acquired without professional

knowledge, such as physical characteristics or activity levels, have not been studied. Bone is an organ

that continuously undergoes remodeling processes of resorption and formation, and BMD is formed

by the balance of these processes [19]. According to the physical stress theory, bone can remain

constant in an environment given an appropriate amount of stress [20]. In other words, factors that

regulate stress on bones, such as physical activity, are important in controlling BMD [20]. Low

activity level was considered a risk factor for OP, and vigorous physical activity such as strength

training or aerobic exercise was recommended to prevent OP [21, 22]. However, despite the benefits

of moderate loads for BMD, too much activity due to overtraining at elite sports levels can have

negative consequences in BMD.[23] Moderate intensity physical activity may help prevent OP, but

excessive intensity physical activity may increase the risk of OP [24, 25]. Therefore, data related to

physical activity will be able to be trained through deep learning as factors for predicting OP.

The purpose of this study was to create a multi-layer perceptron neural network model that predicts

the diagnosis of OP in adult women with a high risk of OP using physical characteristics and physical

activity-related factors, and to evaluate the performance of the created multi-layer perceptron deep

learning model.

**88**  **Materials and Methods**

**89**  1.  Data Source

**90**  This study used data samples obtained from the 6th to 8th KNHANES (2015-2019), the national

**91**  statistics of the Korea Disease Control and Prevention Agency. From a total of 39,759 data samples,

**92**  data were extracted from women aged 35 years or older, the age at which bone density begins to

**93**  decline. Data samples with missing values were excluded, and 11,929 samples were finally selected.

**94**  The selected data samples were classified into 9915 data samples of normal subjects and 2014 data

**95**  samples of subjects diagnosed with OP (Figure 1). This study was approved by the Public Institutional

**96**  Review Board Designated by Ministry of Health and Welfare (approval number: P01-202303-01-003)

**97**

**98**  [Figure 1. Flowchart of the study]

**99**

**100**  2.  Data Preprocessing and Feature Selection

**101**  The physical characteristics and activity variables of the subjects were extracted from the data for

**102**  analysis. Age, weight, height, waist circumference, grip strength, body mass index, quality of life,

**103**  average working hours per week, walking days per week, and strength training days per week were

**104**  selected as continuous variables. Activity restriction, occupational activity, work-related moderate-

**105**  intensity physical activity, work-related moderate-intensity physical activity, leisure-related high-

**106**  intensity physical activity, leisure-related moderate-intensity physical activity, and weight change

**107**  were selected as categorical variables (Table 1).

**108**  To address potential imbalances in the data between diagnosed osteoporosis (OP) and normal

**109**  samples, a RandomUnderSampler function was used. This technique randomly selected an equivalent

**110**  number of normal samples to match the number of OP samples. Additionally, continuous variables

**111**  were scaled using the StandardScaler function, while categorical variables were transformed using

**112**  one-hot encoding methods. These steps were taken to ensure that the data was prepared appropriately

for subsequent analysis. To avoid overfitting due to many variables, LASSO rules were used to select

appropriate variables before generating the multi-layer perceptron model [26]. Alpha representing the

degree of regularization was set at 0.01. After the data was preprocessed, it was split into two sets:

train and test data. The ratio of the train data to the test data was 7:3. This division allowed for the

model to be trained on a portion of the data and then tested on a separate set to evaluate its

performance.

[Table 1. Description of Variables]

3.   Multi-layer Perceptron Neural Network

A multi-layer perceptron is a distributed information processing structure composed of nodes. Each

layer consists of one or more nodes. The input layer receives a signal from the outside and passes the

output to the hidden layer through a weighted connection. It is then computed in the hidden layer and

passed to the output layer to perform computations and generate predictions. The multi-layer

perceptron model created in this study consists of one input layer, one output layer, and five hidden

layers. The five layers consisted of 128, 64, 32, 16, and 8 nodes, respectively. The ReLU activation

function was applied to the 5 hidden layers, and the sigmoid activation function was applied to the

output layer to predict the OP. To prevent overfitting, the dropout ratio was set to 0.25 and batch

normalization was performed. For the gradient descent algorithm to find the minimum loss function,

Adam optimizer was used. The ratio of data split and used for validation while the model was being

trained was 0.2. In order to reduce the learning time for building the model, the EarlyStopping

function was used to terminate learning when the validation loss value did not improve for 20 epochs.

[Figure 2. Schematic diagram of multi-layer perceptron neural network]

4.   Model evaluation

The model's performance was assessed by generating a confusion matrix using the test data, which

classified the predicted class and the actual class as TP (true positive), FP (false positive), FN (false

negative), and TN (true negative). The accuracy, precision, recall, F1-score, and ROC AUC were

calculated from the confusion matrix to evaluate the model's overall performance.

**Results**

1.   Feature Selection

The variables selected using LASSO regularization were age, quality of life index, weight, grip

strength, and average working hours per week.

2.   Multi-layer Perceptron Neural Network Model Training

The multi-layer perceptron neural network model created using the selected features showed the

following accuracy and loss (Figure 3). Training of the model was completed at the 39th epoch. The

average accuracy of the top 5 models built in the process of model training was 77.34%.

[Figure 3. Training accuracy and loss curve]

3.   Model Performance

The accuracy score, which represents the number of correctly predicted data among the entire test

data by the model composed of multi-layer perceptron layer neural networks, was 76.8%. The

precision score for indicating actual OP among test data samples that were predicted to have OP was

74.5%. The recall score representing the prediction of having an OP diagnosis among test data

samples with an actual OP diagnosis was 80.5%. The F1 score, the harmonized mean of precision and

recall, was 77.4%. The ROC AUC score, which indicates how well the multi-layer perceptron neural

network model can classify each class, was 74.8%.

[Figure 4. Confusion matrix for osteoporosis prediction]

**Discussion**

In this study, a multi-layer perceptron neural network model was created using factors related to the

physical characteristics of women over the age of 35 whose bone density begins to decrease. The

factors selected to create the model included not only age, weight, and grip strength, but also the

quality of life and average working hours per week. All the evaluation indicators evaluating the

overall performance of the model created with the above factors exceeded 74%. Among the

evaluations, the recall score, which means the sensitivity of OP diagnosis, exceeded 80%.

The factors selected to model of this study may have been related to BMD, which is the criterion

for diagnosis of OP. Lee et al. [27] conducted a study on the relationship between BMD and age based

on the 4th KNHANES survey data between 2008 and 2009, similar to the subjects in the data sample

of this study. In this study, women's BMD was highest in the 10-19 year age group and declined at a

rate of 0.66 to 1.08% per year until age 80 years [27]. Additionally, moderate physical stress can

stimulate bone remodeling and increase density [28]. Therefore, factors related to physical stress, such

as muscle strength and weight, appear to be related to BMD as factors that can be controlled, unlike

factors such as age [28, 29]. Luo et al. [30] analyzed the relationship between grip strength and BMD

based on a sample of NHANES 2013–2014 survey data of Americans, adjusting for all other factors

that could affect BMD, such as age, body mass index, use of female hormones, and physical activity.

Grip strength has been shown to be associated with increased femoral neck and total lumbar spine

BMD in premenopausal and postmenopausal women [30]. Kim et al. [31] conducted a study on the

relationship between women's body weight and BMD.[31] In this study, they found a significant

difference in weight between the abnormal group with a BMD T-score of less than -1 and the normal

group with a T-score of -1 or greater. Additionally, a weak correlation existed between body weight

and T-score, with each 1 kg increase in weight reducing the risk of abnormal BMD by 3.7%.[31]

The EQ-5D used to measure the quality of life variable is a tool to evaluate health-related quality of

life at three levels for a total of five items: motor ability, self-management, daily life, pain and

discomfort, and anxiety and depression [32]. In other words, the quality of life variable through the

EQ-5D index can be seen as a variable that includes physical activity. The average working hours per

week selected in the model of this study may be seen as a factor that quantitatively reflects the degree

of physical activity [29, 33]. However, there is no study that directly conducted the relationship

between working hours and OP, and rather, the results of other studies on the relationship between OP

and work can refute this view. It has been hypothesized that endocrine disorders caused by night shifts

may indirectly affect the bone physiology of night shift workers [34]. In addition, excessive physical

activity may cause menstrual disorders in women, resulting in decreasing BMD [35]. Nevertheless,

epidemiological evidence suggests that an active lifestyle through physical activity may reduce the

incidence of osteoporotic hip fractures in the elderly population, suggesting that physical activity

during working hours may be an important determinant of OP diagnosis [33, 36, 37].

Recently, models for determining OP, predicting T-score, or predicting BMD built using deep

learning such as convolutional neural networks using CT and X-ray images have been proposed [38–

40]. Yasaka et al. [38] built a model using 1665 CT images of 183 patients, predicted BMD values

with a correlation coefficient of 0.84-0.85 with actual BMD values, and showed OP prediction

performance of over 96%. Ho et al. [39] built a model with data collected through a total of 3472 pairs

of pelvic X-ray and DXA examinations and predicted BMD values only with femur bone x-ray. The

predicted BMD value showed a correlation r value of 0.85 with the actual value, and OP diagnosis

through the model showed an accuracy of 88% [39]. In addition, Sato et al. [40] constructed a model

to predict T-score with chest x-ray, which is the most common, accessible, and inexpensive

measurement. The correlation coefficient between the BMD value predicted by the model and the

BMD value of the hip was 0.75, and the correlation coefficient of the BMD value of the lumbar spine

was 0.63 [40]. ROC AUC score, a performance indicator of the model that classified normal,

osteopenia, and OP by predicting T-score, was 0.89, 0.70, and 0.84, respectively [40]. The model

constructed in our study did not predict the BMD value, and the predictive performance for OP was

about 75-80%, which was relatively low compared to previous studies. Nevertheless, the advantage of

this model in our study is that it can predict OP with simple physical factors without imaging

techniques such as CT or X-ray. The variables for using this model were easily measurable by

surveying, evaluation tool, digital hand dynamometer, and did not require very high skill.

There are several limitations to this study. First, in this study, since the model was created using

only female data, it is inappropriate to use male data for OP prediction. Second, it is difficult to

quantitatively describe the influence of variables for OP prediction because the hidden layer of the

deep learning model makes it difficult to know the exact contribution of the variables to how they

arrived at the OP prediction. Therefore, it is unclear whether OP can be prevented, or symptoms

improved by controlling for the variables chosen to build this model. Future studies will need to

update the model with additional data or variables to improve predictive performance or to build a

model for the entire population, including males. Along with this, verification of the variables selected

to build this model should be performed by adjusting the influence of other extrinsic variables.

**Conclusion**

In this study, a multi-layer perceptron neural network structure was created to predict the diagnosis

of OP in women aged 35 years or older by using personal characteristics and physical factors as

variables. The variables used to build the model were age, weight, grip strength, quality of life index,

and average working hours per week, which could be easily measured through survey, assessment tool,

and digital hand dynamometer. Therefore, our model will be of great help in performing OP screening

without medical equipment in areas with poor medical facilities or elderly women who have difficulty

accessing hospitals.

**References**

1. Lorentzon M, Cummings SR (2015) Osteoporosis: the evolution of a diagnosis. J Intern Med
277:650–661

2. Clarke BL, Khosla S (2010) Female reproductive system and bone. Arch Biochem Biophys
503:118–128

3. Bruno AG, Broe KE, Zhang X, et al (2014) Vertebral size, bone density, and strength in men
and women matched for age and areal spine BMD. J Bone Miner Res 29:562–569

4. Khosla S, Melton III LJ, Atkinson EJ, et al (1998) Relationship of serum sex steroid levels and
bone turnover markers with bone mineral density in men and women: a key role for
bioavailable estrogen. J Clin Endocrinol Metab 83:2266–2274

5. Felson DT, Zhang Y, Hannan MT, et al (1993) The effect of postmenopausal estrogen therapy
on bone density in elderly women. N Engl J Med 329:1141–1146

6. Popat VB, Calis KA, Vanderhoof VH, et al (2009) Bone mineral density in estrogen-deficient
young women. J Clin Endocrinol Metab 94:2277–2283

7. Lu Y, Genant HK, Shepherd J, et al (2001) Classification of osteoporosis based on bone
mineral densities. J Bone Miner Res 16:901–910

8. Marshall D, Johnell O, Wedel H (1996) Meta-analysis of how well measures of bone mineral
density predict occurrence of osteoporotic fractures. Bmj 312:1254–1259

9. Seeman E (1994) Reduced bone density in women with fractures: contribution of low peak
bone density and rapid bone loss. Osteoporos Int 4:S15–S25

10. Alarcón T, González-Montalvo JI, Gotor P, et al (2011) Activities of daily living after hip
fracture: profile and rate of recovery during 2 years of follow-up. Osteoporos Int 22:1609–
1613

11. Guzon-Illescas O, Perez Fernandez E, Crespí Villarias N, et al (2019) Mortality after

osteoporotic hip fracture: incidence, trends, and associated factors. J Orthop Surg Res 14:1–9

12.   Omsland TK, Holvik K, Meyer HE, et al (2012) Hip fractures in Norway 1999–2008: time

trends in total incidence and second hip fracture rates. A NOREPOS study. Eur J Epidemiol

27:807–814

13.   Solou K, Tyllianakis M, Kouzelis A, et al (2022) Morbidity and mortality after second hip

fracture with and without nursing care program. Cureus 14:

14.   Blake GM, Fogelman I (2007) The role of DXA bone density scans in the diagnosis and

treatment of osteoporosis. Postgrad Med J 83:509–517

15.   Lochmüller E-M, Krefting N, Bürklein D, et al (2001) Effect of fixation, soft-tissues, and scan

projection on bone mineral measurements with dual energy X-ray absorptiometry (DXA).

Calcif Tissue Int 68:

16.   Yamamoto N, Sukegawa S, Kitamura A, et al (2020) Deep learning for osteoporosis

classification using hip radiographs and patient clinical covariates. Biomolecules 10:1534

17.   Zhang B, Yu K, Ning Z, et al (2020) Deep learning of lumbar spine X-ray for osteopenia and

osteoporosis screening: A multicenter retrospective cohort study. Bone 140:115561

18.   Pickhardt PJ, Nguyen T, Perez AA, et al (2022) Improved CT-based Osteoporosis Assessment

with a Fully Automated Deep Learning Tool. Radiol Artif Intell 4:e220042

19.   Manolagas SC (2000) Birth and death of bone cells: basic regulatory mechanisms and

implications for the pathogenesis and treatment of osteoporosis. Endocr Rev 21:115–137

20.   Mueller MJ, Maluf KS (2002) Tissue adaptation to physical stress: a proposed "Physical Stress

Theory" to guide physical therapist practice, education, and research. Phys Ther 82:383–403

21.   Sinaki M, Pfeifer M, Preisinger E, et al (2010) The role of exercise in the treatment of

osteoporosis. Curr Osteoporos Rep 8:138–144

22.   Schmitt NM, Schmitt J, Dören M (2009) The role of physical activity in the prevention of

osteoporosis in postmenopausal women—an update. Maturitas 63:34–38

23.   Herbert AJ, Williams AG, Hennis PJ, et al (2019) The interactions of physical activity,

exercise and genetics and their associations with bone mineral density: implications for injury

risk in elite athletes. Eur J Appl Physiol 119:29–47

24.   Wolman RL (1994) ABC of sports medicine: Osteoporosis and exercise. BMJ 309:400

25.   Voss LA, Fadale PD, Hulstyn MJ (1998) Exercise-induced loss of bone density in athletes. J

Am Acad Orthop Surg Glob Res Rev 6:349–357

26.   Muthukrishnan R, Rohini R (2016) LASSO: A feature selection technique in predictive

modeling for machine learning. In: 2016 IEEE international conference on advances in

computer applications (ICACA). IEEE, pp 18–20

27.   Lee EY, Kim D, Kim KM, et al (2012) Age-related bone mineral density patterns in Koreans

(KNHANES IV). J Clin Endocrinol Metab 97:3310–3318

28.   Morel J, Combe B, Francisco J, Bernard J (2001) Bone mineral density of 704 amateur

sportsmen involved in different physical activities. Osteoporos Int 12:152–157

29.   Cadore EL, Brentano MA, Kruel LFM (2005) Effects of the physical activity on the bone

mineral density and bone remodelation. Rev Bras Med do Esporte 11:373–379

30.   Luo Y, Jiang K, He M (2020) Association between grip strength and bone mineral density in

general US population of NHANES 2013–2014. Arch Osteoporos 15:1–9

31.   Kim SJ, Yang W-G, Cho E, Park E-C (2012) Relationship between weight, body mass index

and bone mineral density of lumbar spine in women. J bone Metab 19:95–102

32.   Devlin NJ, Brooks R (2017) EQ-5D and the EuroQol group: past, present and future. Appl

Health Econ Health Policy 15:127–137

33.   Düppe H, Gärdsell P, Johnell O, et al (1997) Bone mineral density, muscle strength and

physical activity: a population-based study of 332 subjects aged 15-42 years. Acta Orthop

Scand 68:97–103

34.   Bukowska-Damska A, Skowronska-Jozwiak E, Peplonska B (2019) Night shift work and

osteoporosis: evidence and hypothesis. Chronobiol Int 36:171–180

35.   Rutherford OM (1999) Is there a role for exercise in the prevention of osteoporotic fractures?

Br J Sports Med 33:378–386

36.   Jordan KM, Cooper C (2002) Epidemiology of osteoporosis. Best Pract Res Clin Rheumatol

16:795–806

37.   Nguyen T V, Sambrook PN, Eisman JA (1998) Bone loss, physical activity, and weight

change in elderly women: the Dubbo Osteoporosis Epidemiology Study. J bone Miner Res

13:1458–1467

38.   Yasaka K, Akai H, Kunimatsu A, et al (2020) Prediction of bone mineral density from

computed tomography: application of deep learning with a convolutional neural network. Eur

Radiol 30:3549–3557

39.   Ho C-S, Chen Y-P, Fan T-Y, et al (2021) Application of deep learning neural network in

predicting bone mineral density from plain X-ray radiography. Arch Osteoporos 16:1–12

40.   Sato Y, Yamamoto N, Inagaki N, et al (2022) Deep learning for bone mineral density and T-

score prediction from chest X-rays: A multicenter study. Biomedicines 10:2323

Figure 1. Flowchart of the study

Figure 2. Schematic diagram of multi-layer perceptron neural network

Figure 3. Training accuracy and loss curve

Figure 4. Confusion matrix for osteoporosis prediction

Table 1. Description of Variables

| KNHANES | Description | Osteoporosis | Normal |
|---|---|---|---|
| Age | Age | $68.69 \pm 8.39$ | $54.75 \pm 12.17$ |
| HE_wt | Weight | $55.88 \pm 8.43$ | $58.91 \pm 9.25$ |
| HE_ht | Height | $152.77 \pm 5.93$ | $157.12 \pm 6.15$ |
| HE_wc | Waist circumference | $82.79 \pm 9.26$ | $80.73 \pm 9.59$ |
| GS | Grip strength | $18.91 \pm 5.03$ | $22.13 \pm 5.13$ |
| BMI | Body mass index | $23.93 \pm 3.29$ | $23.87 \pm 3.55$ |
| EQ5D | Quality of life | $0.86 \pm 0.16$ | $0.94 \pm 0.11$ |
| EC_wht_23 | Average working hours per week | $12.36 \pm 19.79$ | $21.96 \pm 22.37$ |
| BE3_31 | Walking days per week | $4.41 \pm 2.72$ | $4.75 \pm 2.6$ |
| BE5_1 | Strength training days per week | $1.38 \pm 1.17$ | $1.52 \pm 1.31$ |
| LQ4_00 | Activity restriction | Yes: 410<br>No: 1604 | Yes: 742<br>No: 9173 |
| EC1_1 | Occupational activity | Yes: 673<br>No: 1341 | Yes: 5328<br>No: 4587 |
| BE3_71 | Work-related moderate-intensity physical activity | Yes: 12<br>No: 2002 | Yes: 53<br>No: 9862 |
| BE3_81 | Work-related moderate-intensity physical activity, | Yes: 77<br>No: 1937 | Yes: 512<br>No: 9403 |
| BE3_75 | Leisure-related high-intensity physical activity | Yes: 30<br>No: 1984 | Yes: 633<br>No: 9282 |
| BE3_85 | Leisure-related moderate-intensity physical activity | Yes: 222<br>No: 1792 | Yes: 2049<br>No: 7866 |
| BO1_1 | Weight change | No: 1420<br>Increase: 310<br>Decrease: 284 | No: 6318<br>Increase: 1094<br>Decrease: 2503 |