Diversity of *CFTR* variants across ancestries characterized using 454,727 UK biobank whole exome sequences

Justin E. Ideozu[1], Mengzhen Liu[2], Bridget M. Riley-Gillis[2], Sri R. Paladugu[2], Fedik Rahimov[2], Preethi Krishnan[3], Rakesh Tripathi[3], Patrick Dorr[3], Hara Levy[4], Ashvani Singh[5], Jeffery F. Waring[1,2], and Aparna Vasanthakumar[1].

[1]Genomic Medicine, Genomics Research Center, AbbVie, IL.
[2]Human Genetics, Genomics Research Center, AbbVie, IL.
[3]Precision Medicine, AbbVie, IL.
[4]Department of Pediatrics, Division of Pulmonology and Sleep Medicine, University of Wisconsin School of Medicine and Public Health, WI.
[5]Discovery Research, AbbVie, IL.

Keywords: CFTR, whole exome sequencing, UK biobank, cystic fibrosis

Abstract

Limited understanding of the diversity of *CFTR* variants across ancestries hampers efforts to advance molecular diagnosis of cystic fibrosis (CF). The consequences pose a risk of delayed diagnoses and subsequently worsened health outcomes for patients. Characterizing the distribution of *CFTR* variants across ancestries is, therefore, critical for revolutionizing molecular diagnoses of CF. By interrogating 454,727 UK biobank whole exome Sequences, we detected over 4,000 *CFTR* variants, including novel ancestry-specific variants, across six ancestries. F508del was the most prevalent CF-causing variant found across the ancestries, except in East Asia, where V520F was the most prevalent. Sixteen participants had two CF-causing variants, with two diagnosed as CF. We found 154 participants with a CF-causing and varying clinical consequences (VCC) variant. Overall, participants with multiple clinically relevant variants reported indications significantly associated with CF and its pulmonary phenotypes [Bonferroni adjusted p<0.05]. Potentially, participants with classical CF phenotypes may benefit from current CF therapies.

Introduction

In cystic fibrosis (CF), pathogenic variants in the cystic fibrosis transmembrane conductance regulator (*CFTR*) gene results in system-wide debilitating consequences[1]. Current CFTR modulator therapies targeting the basic CFTR defects have yielded both short- and long-term clinical benefits, but they are approved only for patients with certain *CFTR* variants[2,3]. Even within these patients, the response to current CFTR modulator therapies can be variable. Recent advances in genetic technologies have led to the discovery of many CF-causing variants among populations of European ancestry. Although these findings have led to a better understanding of the disease prevalence, and improved diagnoses[4], there is limited knowledge about the distribution of ancestry-specific *CFTR* variants. Characterizing the broad spectrum of pathogenic and non-pathogenic *CFTR* variants across ancestries holds promise to revolutionize molecular diagnoses of CF and could enable wider access to personalized CFTR modulator therapies.

Most studies that have surveyed the diversity of *CFTR* variants have largely focused on studying persons with CF from European populations[5,6]. This is perhaps driven by the fact that CF predominantly affects individuals from European ancestry, and the disease is less frequent in people from other ancestries. Survey reports from a diverse US population shows CF affects individuals of European ancestry (1:3000 live birth) relative to people from other ethnicities, including Africans (1:15000), Asians (1:35,000), and Native Americans (1:10,900)[7,8]. Globally, CF is presumed to affect about 100,000 people and F508del accounts for most of the cases[6,9]. Recent reports suggest the spectrum of CF-causing variants vary across ancestries and countries[7,10]. CF-causing variants commonly reported in other ethnic groups, include 3120+1G>A (Africans)[10], p.W1282X (Ashkenazim)[6,11], and p.G970D (Chinese)[12]. Although these studies have shed insights on common CF-causing variants in non-European populations, carrier screening panels do not often capture the dominant CF-causing variants in some ethnicities[13]. Thus, the prevalence of the disease in other ethnicities, and worldwide, are possibly underestimated.

Diagnosis of CF can be based on the results of a patients' pilocarpine iontophoresis sweat test (sweat chloride ≥ 60 mmol/L) and/or molecular genetic testing (positive for up to two CF-causing variants)[14]. One of the advantages of molecular genetic testing is that it could detect both symptomatic and pre-symptomatic individuals, including the potential to determine a patients' eligibility for current CFTR modulator therapies. A major drawback of solo reliance on symptomatic diagnoses is that often patients are diagnosed when they are already presenting symptoms, and delayed diagnosis negatively impacts the patients' health outcomes[15]. Early diagnosis and interventions, particularly with triple CFTR therapy, results in improved clinical outcomes and could profoundly affect the trajectory of CF lung disease[16]. Carriers of rare pathogenic variants not included in carrier screening panels are more likely to have delayed diagnosis which poses a risk of increased morbidity. Thus, it has become critical to characterize the broad spectrum of pathogenic *CFTR* variants across ancestral populations.

Determining the clinical and functional relevance of *CFTR* variants is such an inundated task that requires critical examination of the variants' impact on an individual's phenotype. This is compounded by the clinical heterogeneity often seen in patients with the same *CFTR* genotype. Efforts driven by CFTR1/2 have led to the identification of 2,110 *CFTR* variants [http://www.genet.sickkids.on.ca]. Of these, 401 are CF-causing while 49 are variants of varying clinical consequences (VCC), according to the latest CFTR2 annotation [https://cftr2.org]. The CFTR1/2 databases have led to an increased understanding of CF diagnoses and prevalence across populations. However, the variants captured from the CFTR1/2 efforts do not necessarily represent the entirety of potential variants that could possibly exist within the *CFTR* gene across ancestries. Recent annotations from TopMed Whole Genome Sequencing (WGS) efforts that includes diverse populations indicate that the *CFTR* gene is about 250kb with 56,488 variants[17], which is longer than the often referenced 189kb length[18]. Although the vast majority of these newly detected *CFTR* variants are intronic [96%], and their clinical relevance unknown, this underscores the continued need to gain a comprehensive understanding of all pathogenic and non-pathogenic variants that possibly exists within the *CFTR* gene, including those from various ancestries.

The complexities associated with characterizing the clinical relevance of *CFTR* variants can be subdued by leveraging population databases, such as UK Biobank (UKBB), that hosts both genomics and phenotypic information of individuals. In this study, we characterized the diversity of *CFTR* variants across six ancestral populations included in the UKBB. We segregated ancestry-specific *CFTR* variants, including those that are CF-causing from uncharacterized variants. We estimated the age of certain detected CF-causing variants. For participants with two CF-causing variants or a CF-causing variant and a VCC, we performed phenotype analysis to determine the possible impact of pathogenic CF-variants on their health outcomes. Our work provides the foundation for future studies to explore the clinical relevance of the newly detected *CFTR* variants across the described ancestries.

Methods

Study population and whole exome sequences

The datasets used for this study is available in the UKBB public repository and was accessed under application 26041. A total of 454,787 UKBB whole exome sequences were interrogated to characterize the diversity of *CFTR* variants across ancestries. The demographics and characteristics of the UKBB participants has been described elsewhere[19], but relevant information such as gender, age and spirometry measures were recorded for each participant. Details on the calling and quality control can be found in previous publication[19]. To maximize variant discovery, no further filters were applied except the removal of individuals that subsequently withdrew consent (final N = 454,727). Curated disease phenotypes, defined based on participants primary care and hospital in-patient (HESIN) records, were utilized for association tests with pathogenic *CFTR* genotypes.

Ancestry designation

For ancestry designation, we opted to use the classification from the PanUKBB working group (https://pan-dev.ukbb.broadinstitute.org/) which derives the ancestry classification from two large reference datasets, 1000 Genomes Project and Human Genome Diversity Project. Using this classification, the participants were grouped into six major groups: African (AFR), American/American Admixed (AMR), Central South Asia (CSA), East Asian (EAS), European (EUR), and Middle East (MID). Individuals that did not fall neatly into the large continental classifications were classified as MIX. While this classification limits the full extent of genomic diversity in the UKBB, it permitted an estimation of variant frequencies to other populations. Individuals assigned to the MIX group were included in all other counts and analysis.

Variant annotation

Clinically relevant variants (CF-causing and VCCs) detected across the ancestries were classified using CFTR2 annotation. CF-causing variants without an assigned Reference SNP cluster ID (rsID) were excluded since they comprised mostly of large indels and structural variants, which are often challenging to be called by variant calling tools. Functionally relevant variants (high impact variants) were annotated with Variant Effect Predictor[20] and SnpEFF[21] and those high impact variants not known to cause CF were recorded. The canonical transcript of *CFTR*

(ENST00000003084) was used as reference for both tools. CF is an autosomal recessive disease
and molecular diagnosis could be based on the possession of two or more clinically relevant
variants. Thus, participants with two CF-causing variants, a CF-causing variant and a VCC, and
a CF-Causing variant and a non-CF-causing high impact variant, were characterized and
prioritized for further association test. Participants with two CF-causing variants were then
assigned likely disease severity status (pancreatic sufficient or insufficient) based on their *CFTR*
genotype as previously described[22,23].
145
146   **Statistical association analysis**
147
We categorized participants with certain clinical or functionally relevant variant combinations
into three groups: two CF-causing variants, a CF-causing variant and a VCC, a CF-causing
variant and a variant of high impact. We tested each of the groups for association with disease
phenotypes using Fisher-Exact tests. As pulmonary measures such as forced expiratory volume
(FEV) and forced vital capacity (FVC) could reflect pulmonary phenotype, we tested for
differences in these measures between participants with two CF-causing variants and the general
population. Only associations meeting a Bonferroni corrected adjusted p-value < 5% were
considered as statistically significant.
156
157   **Genealogical estimation of CF-causing variants age**
158
To estimate the age of CF-causing variants, we used data from the Atlas of Variant Age (AVA)
(https://human.genome.dating/). Models using data from the 1000 Genomes Project (TGP) and
the Simons Genome Diversity Project (SGDP), implemented in the Genealogical Estimation of
Variant Age (GEVA) tool[24], were deployed. For variant age estimations, joint mutational and
recombinational clocks were considered using an average generation time of 25 years. Estimated
age of variants within the *CFTR* gene locus (Chr7: 117,095,837 – 117,366,025) were then
reported.
166
167   **Result**
168
169   **Over 4000 *CFTR* variants identified across ancestries**.
170
We interrogated 454,787 whole exome sequences available in the UKBB to characterize the
diversity of *CFTR* variants across six ancestries [AFR, CSA, EAS, EUR, AMR, and MID] and
an uncharacterized group (MIX). The median age across all ancestries was greater than 60 years.
Except for CSA and MID where over half of the characterized participants were males, most of
the ancestries were dominated by females (Table 1). Overall, we detected 4,193 variants across
all ancestries in the UKBB, with V470M emerging as the most common variant (Table S1). The
CFTR variants represented a diverse range of variant types, but over 50% were intronic (Fig. 1)
The highest number of *CFTR* variants were detected in Europeans [n=3,192] while the AMR
group had the least number of *CFTR* variants [n=151]. Across the remaining ancestries [CSA,
AFR, EAS, and MID], the number of *CFTR* variants detected [471, 417, 266, and 222
respectively] were also much lower when compared to those detected in Europeans (Table 1).
Many of the variants detected in this study have never been reported as variants in *CFTR*. Next,
we explored if there were *CFTR* variants uniquely represented across the ancestries.

Interestingly, we found several *CFTR* variants that tend to be ancestry specific (Fig. 2, Table S2).
The highest number of unique *CFTR* variants [2212/3192] was found in Europeans, while the
Hispanic/Admixed American group had the least number of unique *CFTR* variants [23/151].
About 8.3% [37850/454,727] of the overall study population did not fall under a specific
ancestry and were categorized as the Mixed group (MIX). The median (interquartile range [IQR]
age for the MIX population was 67 (60, 67), with males accounting for 43% of the population.
Within the MIX group, we detected a total of 1333 *CFTR* variants and 28.6% of these variants
[n=381] were uniquely found in their population (Table 1).
192
193 **CF-causing variants across ancestries**
194
We explored the global distribution of CF-causing variants, variants of varying clinical
consequences (VCC), and high impact variants. As shown in Table 2, we found varying
distribution of CF-causing (Table S3) and VCC (Table S4) variants across the ancestries. The
highest number of CF-causing variants were detected in the EUR population [116/154]. These
variants accounted for 75% of all CF-causing variants with RSIDs captured in the study. Nearly
50% of the CF-causing variants were found in the MIX population, but lower proportions
(<15%) were reported across the other specific ancestries. The AMR group with the least number
of detected *CFTR* variants [151/4193] also had the least number [7/154] of the CF-causing
variants (Table 2). F508del was the most prevalent CF-causing variant observed in all ancestries,
except in EAS where V520F was the most prevalent (Fig. 3). While F508del represented ~90%
of the CF-causing variants found in Europeans, our findings indicate its prevalence was not as
high in other ancestries. We further explored whether there were any CF-causing variants that
were potentially ancestry-specific. As shown in Figure 4, all ancestries had unique CF-causing
variants, but EUR possessed the highest number [n=50] of ancestry-specific CF-causing variants
while four groups (AMR, CSA, EAS, and MID) had only one ancestry-specific variant each
(Table S5). We also detected VCC variants in all ancestries [Table S4]. The number of VCC
variants detected were smaller than the number of CF-causing variants detected across the
ancestries, except for AMR and MID populations. R74W was one of the most common variants
detected across the ancestries. Except for EUR, R74W featured among the top 5 most common
VCC reported in all ancestries. Other VCCs that made the common list included R117H,
D1270N, and L967S (Table S6). Interestingly, we found several high impact variants that are not
known to be CF-causing or clinically relevant in each ancestry. The highest number of these
variants [150/200] were detected in the EUR population. Overall, high impact variants accounted
for only a small fraction of the total number of variants reported in each ancestry [≤ 5%] (Table
S7) and many of these [127/200] are uncharacterized in CF (Table S8).
220
221 **More than one CF-causing variants detected in participants**
222
With CF being an autosomal recessive disease, we explored the prevalence of participants with
two CF-causing variants and those heterozygous for a CF-causing variant and a VCC or high
impact variant. We found 16 UKBB participants possessing at least two CF-causing variants
(Table 3). These participants were found exclusively in the CSA [n=1], EUR (n=9), and MIX
[n=6] populations. The CSA participant was homozygous for V456A, but various CF-causing
variant combinations were found in the EUR and MIX populations. F508del was the most
common CF-causing INDEL detected across the populations. Interestingly, the two individuals

homozygous for F508del were also diagnosed as having CF in UKBB. Many of the participants
with two CF-causing *CFTR* variants had combinations associated with pancreatic sufficiency
(96%). The three variant combinations associated with pancreatic insufficiency were
R851X/F508del and F508del/F508del [n=2] (Table 3). We found 155 participants that harbored
a CF-causing variant and a VCC (Table 4). These participants were found only in the EUR
[n=145] and MIX [n=10] populations. R117H was the most reported VCC [49/154] variant in
this subset of participants, and its combination with F508del was the most common found in the
EUR [38/145] and MIX [5/10] populations. L967S also featured as a common VCC variant. Its
combination with F508del was exclusively reported in the EUR population and dominated as the
second most common combination (Table 4). Since high impact variants are potentially
functionally relevant, we assessed their distribution across the ancestries. Among the 200 high
impact variants detected in the participants, the highest number was found in EUR population
[150/200] while the least number of these variants [4/200] were reported in both the AMR and
EAS populations [Table S7]. We also found a high number of participants heterozygous for a
CF-causing and high impact variant [Table S9]. These findings warranted us to interrogate the
association of these variant with participants health outcomes.
246
247 **Classical CF phenotypes reported in participants with two CF-causing variants and VCCs**
248
Phenotype analysis performed using hospital in-patient records indicated that most of the
participants with at least two CF-causing variants presented classical CF phenotypes. Eight
(International Classification of Diseases 10[th] Revision (ICD-10) codes were significantly
[FDR<0.05] enriched in participants with two CF-causing variants (Fig. 5A). Specifically, ICD
codes directly associated with CF [E840, 848, and 849] and colonization with *Pseudomonas*
aeruginosa [B965] were among the top significantly enriched terms. Interestingly, ICD Code
U837 which corresponds to Resistance to Multiple Antibiotics also featured among the top 10
enriched phenotypes. Although bronchiectasis, a classical consequence of dysfunctional CFTR,
was listed among the top, it did not meet our significance threshold when corrected for multiple
testing (Fig. 4). With VCC variants resulting in variable clinical outcomes when combined with
another CF-causing variant but not often diagnosed as CF, we assessed the phenotypes of
participants that fall within this category. Our analysis indicated that six ICD codes were
significantly [FDR<0.05] associated with CF [E840 and 848]; of these, pulmonary phenotypes
[J47, B441, J998, and J440] were significantly enriched in these participants (Fig. 5B). In an
attempt to characterize the clinical relevance of high impact variants, we then explored the
phenotypes of participants with one CF-causing variant and a high impact variant not known to
cause CF (n=272). Surprisingly, the only significant association [FDR<0.05] returned was
Melanoma in situ of the trunk which is not a classical CF phenotype. Aspergillosis was featured
among the top 10 most enriched phenotypes, but this association failed to meet our significance
threshold (Table S9). Further analysis deployed to compare differences of spirometry measures
(FEV1 and FVC) between participant with two CF-causing variants and the general population
returned no significant results.
271
272 **Chronological age of CF-causing variants**
273
GEVA reported the estimated age of over 4,000 variants within the *CFTR* gene locus (Table
S10). Only four of the dated variants were CF-causing [rs139729994 (3600G>A), rs193922500

(V456A), rs76713772 (1717-1G>A), rs77646904 (V520F)] (Fig. S1). The youngest variant
among these, 1717-1G->A, appears to have originated ~425 generations ago (~10,600 years)
while the earliest variant, V456A, originated about ~215 generations ago (~5,300 years).
Interestingly, V520F and 3600G>A which appeared ~338 and 270 generations ago featured
among the most common CF-causing variants reported in EAS population (Fig. 3). When the
allele frequencies of the CF-causing variants were compared with variants of similar ages within
the respective populations in AVA, there were no significant differences. Thus, an indication that
the variants were not under any selective pressure at the time of origination.
284
285 **Discussion**
286
Molecular diagnosis of CF is confounded by sparse knowledge about the prevalence of
pathogenic and non-pathogenic *CFTR* variants across ancestries, including those that are
potentially ancestry specific. In this study, we interrogated whole-exome sequence datasets
generated from over 450, 000 UKBB participants to characterize the distribution of *CFTR*
variants across six ancestries [AFR, CSA, EAS, EUR, AMR, and MID], including an
uncharacterized group (MIX). We report, for the first time, the detection of over 4000 *CFTR*
variants across the ancestries. Among the detected ancestry-specific variants were several
variants of clinical and functional relevance. Phenotypic characterization of participants
harboring multiple combinations of these variants reported indications associated with CF and its
classical pulmonary phenotypes.
297
Although previous efforts driven by the CFTR1/2 team have led to the identification of just over
2000 *CFTR* variants within the CF population[10], our report of over 4000 *CFTR* variants is a
small fraction of potential variants that could exist within the CFTR gene. *In silico* analyses of
external variant browsers such as BRAVO[17], which is based on ~132k WGS generated by the
TOPMed consortium, and All of Us Research Hub[25], which is based on ~98k WGS and ~165k
genotyping arrays reports a total of 56,448 and 39,797 *CFTR* variants, respectively. The higher
number of *CFTR* variants captured in these databases are perhaps due to WGS having the
capability of capturing the entire length of the *CFTR* gene while WES captures mostly exonic
regions. Overall, the vast majority of the *CFTR* variants found in UKBB [52%], All of Us
Research Hub [89%], and BRAVO [96%], were intronic variants. Overall, c.1408G>A (V470M
[rs213950]) was the most common allele (Table S1). Although in GRCH37/38 the reference
allele is a G, the variation associated with increased risk of chronic pancreatitis indicate at this
position is c.1408A>G (M470V)[26]. Regardless, a combination of this variant with another CF-
causing does not cause CF (https://cftr2.org). CF genetic diagnosis is largely based on
characterized *CFTR* variants, most of which are reported in Europeans[4,10]. Although the UKBB
is dominated by people of European ancestry, which could potentially influence estimates of
genetic variations[27], CF is one of the most common life-threatening genetic disease reported in
this population[28]. In line, with previous reports we also found the highest number of *CFTR*
variants in EUR population. Thus, the mutation rate of the *CFTR* gene is likely higher in the
EUR than other populations. The aspect of precision medicine for CFTR modulators is bound to
make higher impact in this CF population with such variable numbers of mutations.
319
Many clinically (CF-causing and VCC) and functionally (High-Impact) relevant variants were
identified following cross-ancestry annotation of all detected variants. As their global

distribution and frequencies are poorly understood, we characterized the most common and
unique CF-causing variants across populations. Indeed, like previous reports[13,29,30], F508del was
the most common CF-causing variant reported globally. While F508del represented ~90% of the
CF-causing variants found in EUR, our findings indicate its prevalence is much lower in other
populations (Fig. 3). Thus, F508del may be the major cause of CF in EUR population, but not for
people of all ancestries. Lower frequencies of F508del in non-European populations have been
reported in several studies, and the spectrum of variants causing CF in such populations
varies[31,32]. For example, G970D has been reported as the most common cause of CF in the
Chinese population[12]. This variant, along with V520F and 3600G>A, featured among the top
three common variants we found in the EAS group and were more common than F508del (Fig.
3). Although we found V520F to be the most common in EAS, it is conceivable G970D is more
common to China than V520F since the EAS group examined in this study expands beyond the
Chinese population. Like other non-EUR populations, CF is less frequent or likely underreported
in Africans[10]. Although F508del was the most common CF-causing variant found in the AFR
group, 3120+1G>A which is predominantly found in Africans[10], was also found to be common
to the AFR group (Fig. 3). Also, in South Asians, CF is thought to be less frequent. Although
F508del represents about 40-50% of cases, V456A the second most common CF-causing variant
we found in CSA (Fig.3), is a well-characterized cause of CF in South Asians[33,34]. Some other
common variants found in non-EUR populations, such as AFR (3791delC), CSA (1525-1G>A),
and EAS, were ancestry-specific. By comparing our findings with BRAVO reports[17], we found
these variants were also rare and exclusively possessed by similar populations. Unfortunately, we
found most of the common CF-causing variants found in non-European populations are not
included in the American College of Medical Genetics list of 23 variants recommended for CF
carrier screening[35].

Numerous hypotheses have been put forward to explain the persistence of CF-causing variants
across populations, despite their detrimental impact in life outcomes. A leading hypothesis
postulates the heterozygous advantage of CF-causing variants against infectious diseases, such as
cholera, typhoid fever, or tuberculosis[36]. For example, the most common CF-causing variant,
rs113993960 (F508del), has been estimated to have arisen approximately 600 generations
ago[37,38] and its rapid increase in Europeans (MAF = 1.6%) was linked to the tuberculosis
pandemic of the seventeenth century, against which it is thought to increase resistance in
mutation carriers[36]. Among other variants with estimated age data in AVA, we did not detect any
variant that has rapidly increased in allele frequency as the F508del variant in Europeans.
Although, V520F and 3600G>A are more common CF-causing variants in EAS population than
F508del (Fig. 3), their allele frequencies were not significantly different from other variants of
similar ages in EAS population. Thus, F508del is likely a unique variant that confers selective
pressure, which has kept its frequency at high frequency.

Beyond CF-causing variants, we also found several VCCs, and High Impact variants not known
to cause CF but predicted to severely impact protein function. R117H the common VCC, which
results in reduced single-channel activity and open probability[39], was the most prevalent VCC
detected across the ancestries (Table S5). Clinical manifestations of patients with R117H are
heterogenous and largely influenced by its combination with another CF-causing variant or other
common variants of poly-T tract (5T, 7T, and 9T). R117H-T5 is reported to potentially result in
less functional CFTR and pancreatic sufficiency when patients are homozygous for this

combination or when found in compound heterozygosity with a CF-causing variant[40]. Most of
the predicted High Impact variants found in our study were already known to cause CF (deduced
with the CFTR2 annotation). After excluding these variants, we still found some uncharacterized
High Impact variants. Taken together, these findings indicated a handful of UKBB participants
harbored clinically and functionally relevant *CFTR* variants.

Since CF is an autosomal recessive disease, we characterized UKBB participants with more than
one clinical or functional relevant variant. Interestingly, we found sixteen UKBB participants
with at least two CF-causing variants. Collectively, these participants reported ICD codes
associated with classical CF phenotypes [Fig. 5A], despite the fact only two participants
(F508del homozygotes) were diagnosed as CF in UKBB electronic health records. Undiagnosed
cases of CF, especially in patients with milder forms of the disease caused by rare pathogenic
genotypes, is not uncommon[41]. Most of the variant combinations reported in our findings likely
result in pancreatic sufficiency, except for one participant harboring the F508del/R851X
genotype that results in pancreatic insufficiency. Although the F508del/R851X participant was
not diagnosed as CF in UKBB, the retrieved ICD10 code indicated the participant suffered
majorly from Chest Pain [R074]. This variant combination, which was detected in an EUR
participant, is less common in CF, with a frequency of only ~0.0002 recorded in CFTR2
database [https://cftr2.org]. Another rarer genotype with ~0.00002 frequency in CFTR2 which
we found was V456A/V456A. This variant is known to cause CF in South Asians[33] and likewise
was detected in a CSA participant. Some of the detected pathogenic genotypes, such as
L206W/S945L and R1066H/1154insTC, have not yet been recorded in CFTR2. Taken together,
a possible explanation for the undiagnosed cases captured in our study may be due to the
rareness of the variants and milder presentation of the disease.

We recorded about 10-fold more participants harboring a CF-causing variant and a VCC, than
those with two CF-causing variants (154 vs 16, respectively). VCCs contribute majorly to the
complexities associated with genetic diagnosis of CF, because clinical outcomes vary widely
across patient populations[42]. For this sub-group of participants, we also found significant
enrichment of ICD codes associated with classical CF (ICD Codes: E840 and E848) and
pulmonary phenotypes, including bronchiectasis, Aspergillosis, and COPD with acute lower
respiratory infections. Participants with these genotypes were reported only in EUR and MIX
populations, and the most dominant combination was F508del/R117H. This variant combination
on its own does not cause CF but when in *cis* with 5T ploy-Tract variant, then CF diagnosis is
likely[40]. About 1,310 patients have the F508del/R117H variant combination in CFTR2 and 22%
of these are pancreatic insufficient. Meanwhile, 80 patients have the F508del/R117H;5T
combination and 33% of them are pancreatic insufficient [https://cftr2.org]. Although these
participants are not diagnosed as CF, their inclusion as patients in CFTR2 suggest they suffer the
burden of dysfunctional CFTR. The F508del/L967S genotype, which was the second most
common found in this subgroup (CF-causing+VCC), was rare in CFTR2. Only seven patients
had this combo and clinical outcome, though variable, is largely expected to be pancreatic
sufficient. The identification of several UKBB participants harboring such variant combinations
and showing CF-like symptoms, is therefore an indication that a number of undiagnosed or
unreported CF cases may benefit from modulator therapies. Additionally, we found no
differences in pulmonary measures (FEV and FVC) between participants with two CF-causing
variants and the general population. Given that the UKBB recruited individuals that were over 40

years old, the individuals with multiple clinically relevant *CFTR* genotypes likely have a milder form of the disease with a different disease etiology that nevertheless could be potentially either treated or prevented by appropriate CFTR-based interventions.

A major drawback to existing knowledge about pathogenic *CFTR* variants is that most studies have been conducted in European populations. However, CF is also prevalent in people from other ancestries, who may harbor uncharacterized unique variants[4]. Even in individuals of European ancestry, where most of the studies have been conducted, there are several *CFTR* variants with unknown clinical significance. In attempt to decipher the clinical relevance of uncharacterized high impact *CFTR* variants detected across the UKBB population, we performed a phenotype analysis of participants with a CF-causing variant and a non-CF-causing (uncharacterized) high impact variant but found no interesting association. Databases such as the UKBB offer an unprecedented opportunity to correlate uncharacterized genotype and phenotype information at a population scale. Although we utilized the UKBB resource to characterize the ancestral diversity of *CFTR* variants, most of the participants are of European ancestry. Recent genome sequencing efforts that included more diverse populations, such as the NIH All of Us and ToPMed programs, report even more *CFTR* variants in comparison to UKBB. However, limited access to phenotype information at an individual level makes it challenging to unravel the clinical relevance of the detected *CFTR* variants.

Here, we aimed to characterize the natural variation that exists within the *CFTR* gene that is not limited to persons with CF. The large majority (>95%) of variants surveyed have allele frequencies less than 1% and ~9% of all variations are indels. While sequencing technology has improved in accuracy and quality control filters have removed many of the problematic variants, identification, and interpretation of novel rare variants (specifically singletons and doubletons) would warrant more precise methods (such as targeted sequencing) for confirmation. Another limitation of our study is the lack of precise CF-related phenotypes, such as FEV1% predicted and sweat chloride levels in the electronic health records extracted from UKBB. Although we captured some spirometry measures, we lacked information relevant to deduce FEV1% scores and there were no records of sweat chloride levels for the participants.

In summary, we have leveraged the UKBB resource to comprehensively characterize the broad spectrum of *CFTR* variants across ancestries. Until now, most efforts aimed at identifying and characterizing *CFTR* variants have focused more on surveying CF populations in a relatively smaller sample size. For the first time, we report the detection of over 4000 *CFTR* variants, which nearly doubles the number of variants reported by CFTR1/2. However, the higher number of variants reported by the NIH All of Us and ToPMed diverse sequencing programs suggests the 4000 *CFTR* variants represents only a small fraction of variants that exist in the *CFTR* gene. The identification of several ancestry-specific variants, including uncharacterized functionally relevant variants, warrants the need for CF screening panels to consider ethnic specificities. The presentation of classical CF phenotypes seen in some non-CF diagnosed participants, with CF-causing and VCC variants, indicates they may benefit from current *CFTR* therapies. As more diverse *CFTR* sequences are becoming increasingly available in genomic databases, future studies are encouraged to leverage such resources to characterize the clinical relevance of high impact variants not previously known to cause CF.

Supplementary information

Supplementary Tables 1-7: **Table S1.** Detected *CFTR* variants across UKBB populations; **Table S2.** Unique CF-causing variants; **Table S3.** CF-causing *CFTR* variants; **Table S4.** Variable clinical consequence variants detected across UKBB ancestries; **Table S5.** Unique CF-causing variants; **Table S6.** Common varying clinical consequences (VCC) variants; **Table S7.** *CFTR* variants annotated as having High Impact; **Table S8.** Uncharacterized high impact variants; **Table S9.** High impact and CF-Causing variants. **Table S10**. Dated variants within the CFTR gene locus

Supplementary Figures: **Figure S1**. Venn diagram showing genealogical estimation of variant age (GEVA) analysis of CF-causing variants

Data availability

Access to UK biobank resource is available by application (http://www.ukbiobank.ac.uk/). The UK biobank exome sequences analyzed in this study were obtained under application 26401.

Ethics declarations

Competing interests: The authors declare no competing interest. The design, study conduct, and financial support for this research were provided by AbbVie. AbbVie participated in the interpretation of data, review, and approval of the publication. JEI, ML, BRG, SRP, FR, PK, RT, PD, AS, JFW, and AV are employees of AbbVie. HL is an employee of University of Wisconsin School of Medicine and Public Health and provided her expertise on CF genetics and translational research (no funding to disclose).

Acknowledgement

We would like to acknowledge and extend our gratitude to the AbbVie GRC Internal Reviewers; Anneke Den Hollander and Nizar Smaoui.

References

1. Raraigh, K.S. *et al.* Complete CFTR gene sequencing in 5,058 individuals with cystic fibrosis informs variant-specific treatment. *J Cyst Fibros* **21**, 463-470 (2022).
2. Despotes, K.A. & Donaldson, S.H. Current state of CFTR modulators for treatment of Cystic Fibrosis. *Curr Opin Pharmacol* **65**, 102239 (2022).
3. Griese, M. *et al.* Safety and Efficacy of Elexacaftor/Tezacaftor/Ivacaftor for 24 Weeks or Longer in People with Cystic Fibrosis and One or More F508del Alleles: Interim Results of an Open-Label Phase 3 Clinical Trial. *Am J Respir Crit Care Med* **203**, 381-385 (2021).

4. Guo, J., Garratt, A. & Hill, A. Worldwide rates of diagnosis and effective treatment for cystic fibrosis. *J Cyst Fibros* **21**, 456-462 (2022).

5. Corvol, H. *et al.* Genome-wide association meta-analysis identifies five modifier loci of lung disease severity in cystic fibrosis. *Nat Commun* **6**, 8382 (2015).

6. Petrova, N. *et al.* Ethnic Differences in the Frequency of CFTR Gene Mutations in Populations of the European and North Caucasian Part of the Russian Federation. *Front Genet* **12**, 678374 (2021).

7. Schrijver, I. *et al.* The Spectrum of CFTR Variants in Nonwhite Cystic Fibrosis Patients: Implications for Molecular Diagnostic Testing. *J Mol Diagn* **18**, 39-50 (2016).

8. Palomaki, G.E., FitzSimmons, S.C. & Haddow, J.E. Clinical sensitivity of prenatal screening for cystic fibrosis via CFTR carrier testing in a United States panethnic population. *Genet Med* **6**, 405-14 (2004).

9. Shteinberg, M., Haq, I.J., Polineni, D. & Davies, J.C. Cystic fibrosis. *Lancet* **397**, 2195-2211 (2021).

10. Stewart, C. & Pepper, M.S. Cystic Fibrosis in the African Diaspora. *Ann Am Thorac Soc* **14**, 1-7 (2017).

11. Quint, A., Lerer, I., Sagi, M. & Abeliovich, D. Mutation spectrum in Jewish cystic fibrosis patients in Israel: implication to carrier screening. *Am J Med Genet A* **136**, 246-8 (2005).

12. Tian, X. *et al.* p.G970D is the most frequent CFTR mutation in Chinese patients with cystic fibrosis. *Human Genome Variation* **3**, 15063 (2016).

13. Ni, Q. *et al.* Systematic estimation of cystic fibrosis prevalence in Chinese and genetic spectrum comparison to Caucasians. *Orphanet J Rare Dis* **17**, 129 (2022).

14. Levy, H. *et al.* Identification of molecular signatures of cystic fibrosis disease status with plasma-based functional genomics. *Physiol Genomics* **51**, 27-41 (2019).

15. Grosse, S.D. *et al.* Newborn screening for cystic fibrosis: evaluation of benefits and risks and recommendations for state newborn screening programs. *MMWR Recomm Rep* **53**, 1-36 (2004).

16. Coverstone, A.M. & Ferkol, T.W. Early Diagnosis and Intervention in Cystic Fibrosis: Imagining the Unimaginable. *Front Pediatr* **8**, 608821 (2020).

17. Taliun, D. *et al.* Sequencing of 53,831 diverse genomes from the NHLBI TOPMed Program. *Nature* **590**, 290-299 (2021).

18. Moisan, S. *et al.* Analysis of long-range interactions in primary human cells identifies cooperative CFTR regulatory elements. *Nucleic Acids Research* **44**, 2564-2576 (2015).

19. Backman, J.D. *et al.* Exome sequencing and analysis of 454,787 UK Biobank participants. *Nature* **599**, 628-634 (2021).

20. McLaren, W. *et al.* The Ensembl Variant Effect Predictor. *Genome Biol* **17**, 122 (2016).

21. Cingolani, P. *et al.* A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. *Fly (Austin)* **6**, 80-92 (2012).

22. Ideozu, J.E. *et al.* Increased Expression of Plasma-Induced ABCC1 mRNA in Cystic Fibrosis. *Int J Mol Sci* **18**(2017).

23. McKay, I.R. & Ooi, C.Y. The Exocrine Pancreas in Cystic Fibrosis in the Era of CFTR Modulation: A Mini Review. *Front Pediatr* **10**, 914790 (2022).

24. Albers, P.K. & McVean, G. Dating genomic variants and shared ancestry in population-scale sequencing data. *PLoS Biol* **18**, e3000586 (2020).

25.   Denny, J.C. *et al.* The "All of Us" Research Program. *N Engl J Med* **381**, 668-676 (2019).

26.   Zhou, D., Bai, R. & Wang, L. The Cystic Fibrosis Transmembrane Conductance

Regulator 470 Met Allele Is Associated with an Increased Risk of Chronic Pancreatitis in

Both Asian and Caucasian Populations: A Meta-Analysis. *Genetic Testing and Molecular*

Biomarkers **24**, 24-32 (2020).

27.   Sun, Q. *et al.* Analyses of biomarker traits in diverse UK biobank participants identify

associations missed by European-centric analysis strategies. *J Hum Genet* **67**, 87-93

(2022).

28.   Blanchard, A.C. & Waters, V.J. Opportunistic pathogens in cystic fibrosis: epidemiology

and pathogenesis of lung infection. *Journal of the Pediatric Infectious Diseases Society*

11, S3-S12 (2022).

29.   Lima, E.d.S., Pezzin, L.S., Fensterseifer, A.C. & Pinto, L.A. Frequency of CFTR variants

in southern Brazil and indication for modulators therapy in patients with cystic fibrosis.

Genetics and Molecular Biology **45**(2021).

30.   Erdoğan, M. *et al.* The Genetic Analysis of Cystic Fibrosis Patients with Seven Novel

Mutations in the CFTR Gene in the Central Anatolian Region of Turkey. *Balkan Medical*

Journal **38**, 357 (2021).

31.   Ortiz, S.C. *et al.* Spectrum of CFTR gene mutations in Ecuadorian cystic fibrosis

patients: the second report of the p.H609R mutation. *Mol Genet Genomic Med* **5**, 751-757

(2017).

32.   Siryani, I. *et al.* Distribution of Cystic Fibrosis Transmembrane Conductance Regulator

(CFTR) Mutations in a Cohort of Patients Residing in Palestine. *PLoS One* **10**, e0133890

(2015).

33.   Uppaluri, L., England, S.J. & Scanlin, T.F. Clinical evidence that V456A is a Cystic

Fibrosis causing mutation in South Asians. *J Cyst Fibros* **11**, 312-5 (2012).

34.   Indika, N.L.R. *et al.* Phenotypic spectrum and genetic heterogeneity of cystic fibrosis in

Sri Lanka. *BMC Medical Genetics* **20**, 89 (2019).

35.   Deignan, J.L. *et al.* CFTR variant testing: a technical standard of the American College of

Medical Genetics and Genomics (ACMG). *Genetics in Medicine* **22**, 1288-1295 (2020).

36.   Poolman, E.M. & Galvani, A.P. Evaluating candidate agents of selective pressure for

cystic fibrosis. *J R Soc Interface* **4**, 91-8 (2007).

37.   Morris, A.P., Whittaker, J.C. & Balding, D.J. Fine-scale mapping of disease loci via

shattered coalescent modeling of genealogies. *Am J Hum Genet* **70**, 686-707 (2002).

38.   Slatkin, M. & Bertorelle, G. The use of intraallelic variability for testing neutrality and

estimating population growth rate. *Genetics* **158**, 865-74 (2001).

39.   Yu, Y.C., Sohma, Y. & Hwang, T.C. On the mechanism of gating defects caused by the

R117H mutation in cystic fibrosis transmembrane conductance regulator. *J Physiol* **594**,

3227-44 (2016).

40.   Castellani, C. *et al.* Consensus on the use and interpretation of cystic fibrosis mutation

analysis in clinical practice. *J Cyst Fibros* **7**, 179-96 (2008).

41.   Sagesse, G.J., Yadava, S. & Mandava, A. Atypical Cystic Fibrosis: Diagnosis at the Age

of 57 Years. *Cureus* **12**, e10863 (2020).

42.   De Wachter, E., Thomas, M., Wanyama, S.S., Seneca, S. & Malfroot, A. What can the

CF registry tell us about rare CFTR-mutations? A Belgian study. *Orphanet Journal of*

Rare Diseases **12**, 142 (2017).

Approximate global ancestry	Sample Size, n (%)	Gender: Male, n (%)	Age in Years, Median (IQR)	*CFTR* Variants, n (%)	Unique *CFTR* Variants, n (%)
African (AFR)	6278 (1.4%)	2582 (41%)	62 (57,70)	417 (9.9%)	138 (33.1%)
Central South Asia (CSA)	8470 (1.9%)	4572 (54%)	65 (58, 72)	471 (11.2%)	177 (37.6%)
East Asia (EAS)	2608 (0.6%)	886 (34%)	63 (57,70)	266 (6.3%)	104 (39.1%)
European (EUR)	397,073 (87.3%)	182164 (46%)	70 (62,75)	3192 (76.1%)	2212 (69.3%)
American/Admixed American (AMR)	950 (0.2%)	334 (35%)	63 (57,71)	151 (3.6%)	23 (15.2%)
Middle East (MID)	1498 (0.3%)	872 (58%)	63 (57,70)	222 (5.3%)	46 (20.7%)
MIX	37850 (8.3%)	16450 (43%)	67 (60,67)	1333 (31.8%)	381 (28.6%)

Table 1. Diversity of *CFTR* variants across ancestries

Table 2. Number of participants with CF-causing variants across ancestries

Ancestries	Sample Size in UKBB	*CFTR* Variants	CF-Causing	Varying Clinical Consequences (VCC)	High Impact Variants
AFR	6278	417	22	12	16
AMR	950	151	7	10	4
CSA	8470	471	17	6	15
EAS	2608	266	8	6	4
EUR	397,073	3,192	116	29	150
MID	1498	222	8	12	7
MIX	37850	1,333	85	29	70

Table 3. Participants with two CF-causing variants

Ancestry	Variant1	Variant2	Number of Variant	Predicted Phenotype	CFTR2 Participants* [n]
EUR	L206W	R553X	1	Pancreatic Sufficient	3
	L206W	S945L	1	Pancreatic Sufficient	0
	R851X	F508del	1	Pancreatic Insufficient	16
	R1066H	F508del	1	Pancreatic Sufficient	56
	R1066H	1154insTC	1	Pancreatic Sufficient	0
	P67L	G551D	1	Pancreatic Sufficient	12

	P67L	F508del	1	Pancreatic Sufficient	186
	711+3A->G	F508del	1	Pancreatic Sufficient	42
	F508del	F508del	1	Pancreatic Insufficient	33984
MIX	F508del	D110H	2	Pancreatic Sufficient	34
	F508del	L206W	1	Pancreatic Sufficient	235
	F508del	F508del	1	Pancreatic Insufficient	33984
	3600G->A	F508del	1	Pancreatic Sufficient	15
	E474K	F508del	1	Pancreatic Sufficient	3
CSA	V456A	V456A	1	Pancreatic Sufficient	2

610 **Table 4. UKBB participants with a CF-causing and variable clinical consequence (VCC) variant**

Ancestry	Variant1	Variant2	Count
EUR	R117H	F508del	38
	L967S	F508del	30
	D443Y	F508del	13
	P750L	F508del	12
	621+3A->G	F508del	5
	R334Q	F508del	4
	P750L	G551D	3
	R258G	F508del	3
	D1152H	F508del	2
	F1052V	F508del	2
	L967S	1717-1G->A	2
	L967S	G551D	2
	L967S	W1282X	2
	Q1291H	F508del	2
	R1070W	F508del	2
	D1270N	F508del	1
	D443Y	1898+1G->A	1
	D443Y	N1303K	1
	D443Y	Q493X	1
	D443Y	R553X	1
	F575Y	1154insTC	1
	L967S	1138insG	1
	L967S	1898+1G->A	1
	L967S	3659delC	1
	M265R	F508del	1
	P750L	1461ins4	1
	P750L	R560T	1
	P750L	W846X	1
	R1070Q	F508del	1
	R1070W	G542X	1
	R117G	F508del	1
	R117H	1898+1G->A	1
	R117H	2184delA	1
	R117H	3272-26A->G	1
	R117H	I507del	1
	R117H	R347H	1

	R117H	R347P	1
	R352W	N1303K	1
MIX	R117H	F508del	5
	621+3A->G	F508del	1
	D443Y	F508del	1
	D1152H	1898+1G->A	1
	F1052V	F508del	1
	Q1476X	N1303K	1

612
613

Fig. 1: Most *CFTR* variants are intronic

Several variant types with varying consequences were identified in the *CFTR* gene. The labels in the chart represent the number and percentage of the Intronic variants (highlighted in orange) clearly dominated over half of the captured variants despite Whole Exome Sequencing technology capturing mostly exomes.

Fig. 2: Ancestry-specific variants characterized

Intersection of *CFTR* variants across ancestries

Unique *CFTR* variants	
Approximate Global Ancestry	Unique *CFTR* Variants
EUR	2212
Mix	381
CSA	177
AFR	138
EAS	104
MID	46
AMR	23
*Listed in Table S8	

A) Graph of total number of variants (x-axis; Set Size) detected in each ancestry (y-axis). **B)** Intersection of variants detected across the ancestries. Each column corresponds to the number of variants in each intersection. Ancestries present in each intersection are represented by the black dots. Intersections with single dots represents ancestry-specific variants. **C)** A list of ancestry-specific variants. The highest number of unique *CFTR* variants (2212/3192) was detected in Europeans.

Fig. 3: Common CF-causing variants in each ancestry represented in pie charts

Common CF-causing variants across ancestries

For each ancestry, slices vary by color, with blue representing the most abundant CF-causing variant. F508del was the most common CF-causing variant detected in all ancestries, except for East Asia where V520F was the most common.

Fig. 4: Ancestry-specific CF-causing variants characterized

Intersection of CF-causing variants across ancestries

A) Graph of total number of variants (x-axis; Set Size) detected in each ancestry (y-axis). **B)** Intersection of variants detected across the ancestries. Each column corresponds to the number of variants in each intersection. Ancestries present in each intersection are represented by the black dots. Intersections with single dots represents ancestry-specific CF-causing variants. **C)** A list of ancestry-specific CF-causing variants. The highest number of unique CF-causing variants (n=50) was detected in Europeans.

Fig. 5: Classical CF phenotypes enriched in participants with >1 clinically relevant *CFTR* variant

A) Enriched phenotypes observed in patients with two CF-causing variants **B)** Enriched phenotypes observed in patients with one CF-causing and a high impact variant. The top 10 enriched phenotypes are represented in bar graphs. The negative logarithm of the Bonferroni adjusted p-value was used to deduce the enrichment scores shown in the bars. Enrichment score >1.3 corresponds to a significance threshold of *p*<0.05.