1   **TITLE:** The Zero-Corrected, Gravity-Model Multiplier (ZERO-G): A novel method to estimate
2   disease dynamics at the community-scale from passive surveillance data
3
4   Michelle V Evans[1]*, Felana A Ihantamalala[2,3], Mauricianot Randriamihaja[1,2], Andritiana
5   Tsirinomen'ny Aina[2], Matthew H Bonds[2,3], Karen E Finnegan[2,3], Rado JL Rakotonanahary[2,3],
6   Mbolatiana Raza-Fanomezanjanahary[2], Bénédicte Razafinjato[2], Oméga Raobela[4],
7   Sahondraritera Herimamy Raholiarimanana[4], Tiana Harimisa Randrianavalona[4], Andres
8   Garchitorena[1,2]
9
10   1. MIVEGEC, Univ. Montpellier, CNRS, IRD, Montpellier, France
11   2. NGO Pivot, Ranomafana, Ifanadiana, Madagascar
12   3. Department of Global Health and Social Medicine, Blavatnik Institute at Harvard Medical
13   School, Boston, MA, USA
14   4. National Malaria Program, Ministry of Health, Antananarivo, Madagascar
15
16   * Corresponding Author: mv.evans.phd@gmail.com
17
18

ABSTRACT

Data on population health are vital to evidence-based decision making by public health officials,

but are rarely adequately localized, particularly in rural areas where barriers to healthcare can

result in extremely low ascertainment of cases by the health system. Here, we demonstrate a

new method to estimate disease incidence at the community level from passive surveillance

data collected at primary health centers. The zero-corrected, gravity-based multiplier (ZERO-G)

method explicitly models sampling intensity as a function of health facility characteristics and

statistically accounts for extremely low rates of ascertainment, resulting in an unbiased,

standardized estimate of disease incidence at a spatial resolution nearly ten times finer than

typically reported by the facility-based passive surveillance system. We assessed the

robustness of this method by applying it to a case study of malaria incidence in a rural health

district in southeastern Madagascar. ZERO-G decreased geographic and financial bias in the

dataset by over 64% and doubled the agreement rate between spatial patterns in malaria

incidence and prevalence rates. ZERO-G can be applied to other infectious diseases and

settings, increasing the availability of long-term, high quality surveillance datasets at the

community scale.

Key words: health care access; geographic bias; floating catchment area; malaria; Madagascar

INTRODUCTION

Health metrics are vital to public health efforts, allowing decision makers to better understand the state of population health and evaluate the impact of health interventions (Murray et al. 2004, Tatem et al. 2012). Many of these metrics are based on routine passive disease surveillance from facility-based health management information systems (HMIS), which record the number of disease cases received at each facility at a regular frequency. Health records are then aggregated, digitized, and transferred to the district, and eventually national, health offices (Amouzou et al. 2021). While the exact structure differs by country, each scale of spatial aggregation of the data in an HMIS corresponds to a specific level of the health system and corresponding health infrastructure. For example, national-level data are used by international organizations to monitor long-term, multi-country trends and inform policy; regional- and district-level surveillance data may be used by national public health offices to allocate resources within the country; and individual health facility information is used by district health offices for program management.

Missing from most HMIS are routine surveillance data at the scale of individual communities or villages. These data are needed for spatially targeted interventions for disease control in collaboration with community health programs, which primarily serve rural communities and play an integral role in achieving universal health coverage (World Health Organization et al. 2007, Rifkin 2018). While rural primary care facilities typically serve over ten thousand people spread along hundreds of square kilometers, community health workers serve between several hundred to a few thousand and their catchment is generally no bigger than 10 km^2. Due to geographic barriers in particular, systemic lack of access to health facilities for large portions of the population has resulted in community health becoming a central pillar of national health strategies globally (Perry 2020). The lack of long-term surveillance datasets at the community level impedes our ability to monitor changes in disease burdens over time, locally target or evaluate the impact of community-health interventions, create surveillance and forecasting systems at these levels, and generally incorporate health data into decision-making processes. Given the increasing role of community programs in providing primary health care and supporting disease control efforts, the lack of routine surveillance data at this level must be remedied.

There are several barriers to the creation of a routine surveillance system at the community level. First, community health workers often only diagnose and treat common illnesses for children under 5 years old (Young et al. 2012). Therefore, case notifications collected by community health workers are incomplete because they represent a small subset of

the population. Second, community health programs tend to be poorly funded, supported, and
integrated within the larger health system (Kok et al. 2015, Hodgins et al. 2021), with negative
consequences for data completeness and quality. For example, a case study in Malawi found
that over 40% of community health reports contained errors when aggregation was conducted
by CHWs due to lack of training and time available for reporting (Admon et al. 2013). Third, the
existing structure of health system reporting often means that paper reports from the community
level are submitted to district officials and integrated into the electronic HMIS system with
significant delays, which limits their use for disease surveillance. An alternative is the use of
health facility data disaggregated at the community level, which is becoming increasingly
available with the development of new technologies such as eHealth systems. As these
technologies become more common, the availability of data will become less of a barrier, but
even when data remain disaggregated, there are issues of completeness and geographic bias
due to heterogeneous access to care (Afrane et al. 2013, Ohrt et al. 2015, Garchitorena et al.
2021). These problems are exacerbated at fine spatial scales. For example, communities in
rural areas with low access to care may be missed by routine health facility systems (Noor et al.
2003), significantly under-estimating disease burdens in these already vulnerable communities.
Given the current lack of high-quality data at the community level, methods are needed to
account for biases in data while retaining their spatial disaggregation.
At the scale of the government health district and higher, several methods have been
developed to address these issues, particularly under-ascertainment of cases (Table 1).
However, none of these adjustment methods result in estimates of disease incidence that are
available at the spatial scale of individual communities and at a temporal regularity high enough
to allow for rapid responses to epidemics. Existing methods are limited primarily by the
frequency and spatial resolution of external data sources, such as large-scale surveys of
disease prevalence or health-seeking behaviors. For example, information on healthcare
utilization rates, such as that collected via Demographic and Health Surveys, is often collected
nationally at the level of the district or region, and is inappropriate for use within smaller
administrative zones. Prevalence surveys offer only a snapshot of malaria burden in time, and
their inferences, while available at finer spatial scales, often only apply to annual estimates. In
general, both forms of survey data are resource-intensive to obtain and are rarely available at
spatial or temporal scales relevant to community health programs (Cibulskis et al. 2011). There
is a need to leverage the spatial and temporal scale of passive surveillance data while adjusting
for under-ascertainment biases in case reports, resulting in long-term, high-quality data on
disease incidence at resolutions relevant to community health programs.

Table 1. Comparing the ZERO-G method to available methods for adjusting passive
surveillance data.

Method	Temporal Scale	Spatial Scale	Data Required
Standard indirect estimators (e.g. WHO Malaria Report)	Annual	National	Estimates of health-seeking behavior, such as from DHS surveys
Prevalence-based geostatistical models (e.g. Malaria Atlas Project)	Annual	5x5 km	Prevalence survey, environmental variables
ZERO-G multiplier	Monthly or weekly, depending on frequency of data collection	Community or village level	Consultation data, geographic location of health system and population

Here, we introduce the zero-corrected, floating catchment gravity model multiplier

(ZERO-G). This method accounts for under-ascertainment of cases by public health facilities,

resulting in a long-term dataset of disease incidence at the scale of individual communities or

villages. The method builds on work by Hyde et al. (Hyde et al. 2021) and combines an indirect

estimation method with a floating catchment area model of healthcare access based on gravity

models. The method first calculates a sampling intensity multiplier derived from consultation

rates using a floating catchment area model (Delamater et al. 2019), and then scales disease

incidence by this multiplier to adjust for spatio-temporal heterogeneity in access to healthcare.

Importantly, this method relies solely on data available to local stakeholders: all-cause

consultation rates, the focal disease incidence, and health facility characteristics. We

demonstrate the method using a case-study of malaria passive surveillance in a rural health

district in southeastern Madagascar. Then, we validate the method by comparing the estimated

sampling intensity and malaria incidence rates to health-care seeking behavior and malaria

prevalence from a district-representative cohort.

THE ZERO-G MULTIPLIER METHOD

The ZERO-G multiplier is an indirect estimation method that adjusts case data for known biases

in case ascertainment to create high-quality monthly case notification data for multiple

communities. Indirect estimation methods estimate the "true" rate of disease incidence or

prevalence from case data with low or uneven ascertainment rates by including information on

the sampling intensity (e.g. healthcare use) in each administrative region (Hickman and Taylor

2005). ZERO-G specifically uses a benchmark multiplier method, which combines information

on the number of cases recorded by the health system with information on the proportion of

cases that are expected to be observed. In addition, it includes imputation methods for adjusting

for extremely low ascertainment rates that result in zero cases reported. The final result is an

estimation of disease incidence equal to the expected incidence if access to healthcare was

identical across space and time.

136
ZERO-G consists of the following steps (Fig. 1):

1. Estimate sampling intensity using a floating catchment area model

2. Correct for low ascertainment rates that resulted in zero cases via spatio-temporal

imputation methods

3. Identify optimum scaling factor for sampling intensity

4. Apply sampling intensity (multiplier) to zero-adjusted data (benchmark) via benchmark

multiplier method

Figure 1. Workflow for adjusting incidence data using the floating catchment, zero-
corrected multiplier (ZERO-G) method.
148

Step 1. Estimate sampling intensity via a floating catchment area model

Sampling intensity (SI) serves as the multiplier in a benchmark multiplier method, and

can be derived from additional data sources (e.g. proportion in treatment, reported health-

seeking behaviors, mortality rates). Often, these data are collected via surveys of healthcare

seeking behavior and not available at a temporal or spatial scale that matches the case

notification data. ZERO-G overcomes this by estimating a SI for each community for each

month. In this method, SI is proportional to the probability that a disease case in that community

and month is reported to the health system. It is estimated via a temporally-explicit floating

catchment area (FCA) model, an established technique used to estimate spatial access to

healthcare and health facility catchments (Luo and Qi 2009). Based on a gravity model, FCA

models consider both the quantity and spatial accessibility of services at a health center for a

given population by weighting the distance to care by the availability of services provided at

each health center. Access to care in each community is then modeled as an "attractive force"

to each health center and total access to care is the sum of these forces for a given community

(Fig. 1, Step 1).

We used the modified two-step floating catchment area formulation of this metric, which

allows for sub-optimal allocation of health resources via the inclusion of distance-weighted

competition for each healthcare clinic's resources (Delamater 2013). We define the sampling

intensity for each community *i* and month *t (SI$_{i,t}$)* following Equation 1:

$$SI_{i,t} = g(t) \sum_j \frac{S_{j,t} f(d_{ij})^2}{C_j} \qquad \text{Eq. 1}$$

This represents the sum of all healthcare services provided by each health center *j* at time *t* (S$_{j,t}$,

Eq. 2), mediated by distance between each community *i* and the health center *j*, d$_{ij}$, and

competition for services at each health center *j* by the total population across all communities

(C*j*, Eq. 4*)*. Temporal trends in healthcare access are described by g(t), with the exact function

dependent on temporal trends in healthcare access in the region being considered. For

example, seasonal changes in health system use could be represented by a sinusoidal function.

Services provided by each health center *j* represent the "mass" of a gravity model,

defined by Equation 2:

$$S_j = \sum_m \beta_m v_m$$

<div align="right">Eq. 2</div>

180
181
Each health center's mass (S_j) is equal to the sum of characteristics relevant to health services

(v_m), weighted by a unique coefficient β_m. These characteristics will vary by health system, but

should generally describe the quantity and quality of services available at the health center.

They can be continuous variables (e.g. number of staff, number of beds) or binary variables

(e.g. presence of a surgery unit, type of health center (hospital vs. clinic)). These services can

vary over time, resulting in a different value for each month, allowing for changes in healthcare

access following changes to the health system (e.g. opening or closing of clinics, new payment

policies, change in staff numbers).

The effect of distance on spatial access to health center services is described by a

distance-decay function $f(d_{ij})$. While the shape of this function may differ by the local context,

distance-decay is most often described by an exponential function (Khan 1992) and is the

recommended default when using this method (Eq. 3).

$$f(d_{ij}) = e^{-\lambda d_{ij}}$$

<div align="right">Eq. 3</div>

194
195 Competition for services at each health center j is defined as the sum of the population-

196 weighted distance between all administrative zones and the health center, including a scaling

197 factor β_c:

$$C_j = \beta_c \sum_k P_{k,t} f(d_{kj})$$

<div align="right">Eq. 4</div>

198
The calculation of the SI includes several parameters that can be pre-specified (if known) or

estimated by fitting the FCA model to all-cause consultation data with the focal disease removed

(see Table S1 for description of parameters and ranges of values in the case study). Any model

fitting method can be applied, such as maximum likelihood or Bayesian approaches. The

recommended approach is to estimate the number of consultations for each community and

month as a random variable with a binomial distribution with probability equal to the SI and size

n equal to the population size of the community, although other distributions are possible.

Step 2. Correct for low ascertainment rates that resulted in zero cases via spatio-temporal
imputation methods
In many instances, extremely low ascertainment rates combined with small populations in small
geographic areas can result in zero reported cases for a month for a community. Because the
benchmark multiplier is a scaling process, these values would remain unchanged (i.e. report
zero incidence that month) after applying the benchmark multiplier step. The second step of
ZERO-G is therefore replacing a subset of these extreme low incidence months (zero-incidence
months) that are believed to be due to low ascertainment rates with new values using spatio-
temporal imputation that incorporates seasonal and spatial patterns in incidence. Exactly which
values should be replaced is context specific, and should take into account the seasonality of
the focal disease and known spatial patterns in access to health care. For example, in the case
study described below, only those zero-incidence months that occurred during the peak of the
malaria season and in communities outside the catchment of a health-system strengthening
initiative were replaced. Those that occurred outside of the malaria season or in the zone where
health access is known to be higher were assumed to be representative of true spatio-temporal
trends in incidence and were unchanged. Imputation is performed via 100 boosted regression
tree models that estimate monthly incidence as a function of each community's identity,
longitude, and latitude and the specific month of the zero-incidence occurrence, using the
median of 100 imputations as the final imputed value. Imputed values are therefore a function of
both spatial and temporal trends in the incidence data.
227
Step 3. Identify optimal minimum scaling factor for sampling intensity
The benchmark multiplier step of ZERO-G requires a SI multiplier greater than 0 and less than
or equal to 1. A multiplier near zero results in a large change to the estimated incidence and a
multiplier of one results in no change. SI is therefore linearly rescaled to fall between a minimum
value (ω) and 1. For those communities that are considered representative of optimal healthcare
access, the SI can be assigned a value of 1, meaning the incidence in these populations will be
unchanged by the benchmark multiplier. All other communities will be standardized to equal the
sampling intensity in these "high access" communities. The remaining SI values can then be
rescaled between ω and 1. At a minimum, the optimal value of ω is that which most reduces the
effect of geographic bias in the adjusted incidence rates. In addition to geographic biases, other
factors can also be included to identify the optimal ω. In the case study below, we include
financial bias and a limit on upper outliers in incidence rates in our selection of ω.
240

241 *Step 4. Apply sampling intensity (multiplier) to zero-adjusted data (benchmark) via benchmark*

242 *multiplier method*

243 The rescaled sampling intensity (SI) and zero-corrected incidence data (I_{zero}) are then combined

244 via the benchmark multiplier method to result in an adjusted estimate of monthly incidence (I_{adj})

245 for each community (Eq. 5):

$$I_{adj} = I_{zero} * SI^{-1}$$
<div align="right">Eq. 5</div>

In this equation, the zero-corrected incidence data serves as the benchmark while the inverse of

the sampling intensity is the multiplier. Both are calculated to retain the temporal frequency and

spatial resolution of the original dataset, meaning they have been estimated for each community

and each time period. Retaining the frequency of the original dataset allows ZERO-G to account

for temporal changes in ascertainment biases, such as seasonality or clinic closures, that may

act differently across each community.

CASE STUDY: MALARIA INCIDENCE IN IFANADIANA, MADAGASCAR

We applied ZERO-G to malaria incidence in Ifanadiana District, Madagascar to demonstrate its

utility in regions with highly heterogeneous rates of under-ascertainment. Ifanadiana is a district

in the Vatovavy region of southeastern Madagascar. It has an estimated population of 183,000

people spread across 195 fokontany (smallest administrative unit comprising about 1000

people) within 15 communes. Each commune contains one primary health center level 2

(PHC2), and six of the larger communes also contain a primary health center level 1 (PHC1),

which provides more basic care, for a total of 21 PHCs within the district. Beginning in 2014,

the Madagascar Ministry of Public Health (MMoPH) and the non-governmental organization

Pivot began a partnership to strengthen the health system, establishing Ifanadiana as a model

health district. This intervention works across all levels of the health system, from community

health at the household level to tertiary care at the regional hospital. At the level of the PHCs,

the intervention includes renovation of PHC facilities, removal of point-of-care user fees, and

joint recruitment of staff between Pivot and the MMoPH. As of January 2023, PHC-level

interventions have been installed at 15 PHC2 across all 15 communes, and will be expanded to

all levels of PHCs by the end 2023.

The primary barriers to healthcare at PHCs in Ifanadiana are geographical and financial.

The majority of the district is rural and the transportation network is primarily non-motorized;

over 70% of the population lives further than an hour travel time from a PHC (Ihantamalala et al.

2020). As such, geographical access to care at PHCs is highly unequal, and exhibits a strong

distance-decay relationship from PHC locations (Garchitorena et al. 2021). The second barrier

is financial costs at the level of the individual. Nationally, nearly half of the total public health

expenditure in Madagascar is borne by the individual (World Health Organization 2022), and

these user fees are the most cited barrier to healthcare seeking across the district (Garchitorena

et al. 2017). As such, we aimed to reduce the impact of geographic and financial bias in malaria

incidence rates by adjusting the data using ZERO-G.

Data Collection

Monthly consultation data were collected at each PHC for the district of Ifanadiana from January

2016 to December 2021. Photos were taken of handwritten registries at each PHC, and

patients' residences were manually geolocated to the precision of the fokontany. The number of

all-cause consultations were reported by fokontany, as well as the number of malaria cases, as

confirmed by rapid detection test. Because patient ages were provided in these registries, we

were able to divide the number of consultations and malaria cases into three age groups for

analysis: children under 5 years old, juveniles aged 5-14, and adults aged 15 and over.

Ifanadiana suffers from stockouts of diagnostic materials, specifically rapid-detection tests

(RDTs) (Cordier et al. 2020), leading to unconfirmed cases of malaria. We accounted for this

reduced diagnostic capacity by scaling the confirmed malaria cases by the proportion of feverish

patients who were tested via an RDT at each PHC during each month (n = 536). Information on

the characteristics of each clinic by month was provided by Pivot's Monitoring and Evaluation for

Research and Learning team.

Population data came from two sources. For the 80 fokontany that receive community

health program support from Pivot, we used population estimates from a Pivot-led census

conducted in 2021. For the remaining 115 fokontany, population estimates came from a national

census conducted in 2018 by the Madagascar National Institute of Statistics. By interpolating

population values between the 2018 census and the previous 1993 census, we estimated an

average annual population growth rate of 2.0%. We applied this population growth rate to both

datasets to obtain each fokontany's population by year. For both datasets, we assumed 18% of

the population to be under 5 years old, 28.6% of the population to be aged 5 - 14 and the

remainder to be 15 years old or above, based on the average age structure of the 80 fokontany

that were censused in 2021.

Distances between residential areas and PHCs were calculated on a high-resolution

transport network created via crowd-sourced mapping through a collaboration with

Humanitarian OpenStreetMap. Over 20,000 km of footpaths and 100,000 buildings within the

district were mapped through a two-step validation process (Ihantamalala et al. 2020), resulting

in an open-source dataset on OpenStreetMap. Using this dataset, we estimated the distance

between each household and each PHC within the district, and aggregated this to the scale of

the fokontany to result in an average distance to each PHC for each fokontany. Three fokontany

lacked accurate routing information and so were excluded from the analysis.

We evaluated our estimates of the SI and adjusted malaria incidence rates using

external data from a longitudinal cohort survey conducted in the district of Ifanadiana (IHOPE

cohort). The IHOPE cohort has conducted population-representative surveys approximately

every two years from 2014-2021 using a two-stage cluster sampling scheme involving 80 spatial

clusters, each containing 20 households (Miller et al. 2018). We include data from 2016, 2018,

and 2021 in this analysis. The IHOPE cohort is based on the internationally validated

Demographic and Health Surveys and is implemented by the Madagascar National Institute of

Statistics. See Miller et al. (2018) for further details on participant recruitment and study design.

As part of the survey questionnaire, participants were asked if they were ill in the past four

weeks and, if so, if they sought care at a public PHC. This data represented self-reported

health-care seeking behavior, comparable to our estimates of sampling intensity. Malaria

prevalence data was collected via rapid detection tests conducted as part of the IHOPE survey

in 2021. Briefly, children under 15 years old who consented to the study were tested for active

malaria infection using SD One Step Malaria HRP-II(P.f) and pLDH(Pan) Antigen Rapid Tests.

Those who tested positive were provided with a standard treatment of artesunate amodiaquine

and paracetamol, with duration and dosage in accordance with national guidelines. In total, this

resulted in 3774 samples across 80 clusters and 109 fokontany.

Applying the ZERO-G multiplier

Estimating Sampling Intensity

We estimated the sampling intensity for each fokontany and month combination in our dataset

following the methods described above for each age class (children, juveniles, and adults) using

non-malarial consultations at health centers. We included five traits of the health center in our

calculation of S_i:

1. whether the PHC fell within the initial Pivot service catchment,

2. if point-of-care user fees (consultation costs and medications) had been removed,

3. the number of staff at the PHC during each month,

4. level of health clinic (PHC1 or PHC2, with PHC2 providing more services),

5. distance from the PHC to the District office, which provides supplies, medications, and
supervision.
In addition, two new PHC2 were opened in the district during the study period, one in
Ampasinambo in November 2016 and one in Ambiabe in April 2018, which we accounted for in
our calculation of SI. Notably, ZERO-G allows for health center traits that change over time,
which we used to include monthly staffing changes and user fee removal interventions that were
implemented over the study period.
We included linear and seasonal adjustments to the SI via the g(t) function (Eq. 6), to
account for increasing access to healthcare over time and seasonality in major diseases in
Madagascar (i.e. respiratory disease, diarrheal disease). It includes a linear coefficient (β_t) and a
sine-function, to account for seasonality, parameterized by ϕ.
352

$$g(t) = e^{\beta_1 t + \beta_2 \frac{sin(2\pi(t+\phi)/12)+1}{2}}$$ Eq. 6

353
We used an exponential function to describe the distance-decay relationship, allowing
the shape of this relationship to vary between fokontany within the Pivot catchment and those
outside of the Pivot catchment by applying a scaling coefficient (β_s). To reduce computational
time, we set a maximum limit on the distance between a community and the PHC (d_{ij}) at 25 km,
slightly above the maximum distance of a fokontany to the nearest PHC in Ifanadiana (22.1 km).
Finally, we estimated the number of non-malarial consultations y as a random variable
with a binomial distribution with the probability equal to the SI and size n equal to the population
size of the fokontany. A total of 11 parameters were estimated via maximum-likelihood
estimation (MLE) (Table S1). Some fokontany had extremely low consultation rates and
reported zero consultations for over 50% of the study period. We excluded these fokontany (n =
43) from the model fitting exercise, but did estimate their SI from the fit model. We used a two-
step MLE estimation process. First, we performed a grid search via a latin hypercube sample of
1,000 samples of coarse parameter space to identify the ten parameter sets with the lowest
negative log-likelihood. We then performed a second MLE using the BFGS algorithm via the
optim function in the *stats* package in R (R Core Team 2021), using the parameters from the ten
parameter sets with the lowest negative log-likelihood from the first step as the starting
parameters. We assessed each of these ten iterations for convergence and selected the
parameter set with the lowest negative log-likelihood as the optimal fit. From the optimal
parameter set, we estimated an SI for each fokontany-month combination via Eq. 1.
373

Zero-correcting via boosted regression tree imputations

Thirty-seven fokontany reported zero malaria cases during at least half of the study period,

totaling 3468 (28.4%) of fokontany-month samples. We chose to only replace zeros in those

fokontany that were outside of Pivot's initial catchment (n = 140), assuming ascertainment was

high enough in fokontany in Pivot's initial catchment that zeros were due primarily to seasonal

trends in incidence and not under-ascertainment. We also only replaced zeros that occurred

during the malaria season (November through May), assuming zeros outside of this season

were representative of the low incidence characteristic of that time period. This resulted in 8.57

to 9.06% of fokontany-month incidence values being imputed for each age class, an average of

7.65 months per fokontany (range: 0-25), for those fokontany outside of Pivot's initial catchment.

These values were imputed 100 times using boosted regression trees that estimated new

monthly incidence as a function of each foktonany's identity, longitude and latitude and the

specific month of the study period using the *mice* package(Buuren and Groothuis-Oudshoorn

2011). The final imputed value was the median of 100 imputations.

Identifying the optimal scaling factor (ω) and applying the benchmark multiplier

We manually set SI to 1 for those fokontany which had an average annual consultation rate over

1 consultation per capita-year, defined as "high access fokontany" (n = 19). We then explored a

range of minimum values of the scaling factor, from 0.002 to 0.4 with a resolution of 0.002. We

identified the minimum value that met requirements with regards to upper incidence outliers,

geographic bias, and financial bias (Fig. 1):

1. Upper incidence outliers: A very low ω could result in incidence rates that are

unrealistically high. To limit the overcorrection of incidence rates, we limited ω to values

that resulted in a proportion of upper incidence outliers less than 3x the proportion of

upper incidence outliers seen in the high access fokontany in the full dataset. An upper

incidence outlier was defined as an annual incidence rate above the mean annual

incidence rate + 3 *sd* in the high access fokontany.

2. Geographic bias: We calculated geographic bias by estimating the Spearman's ranked

correlation coefficient between the distance to the nearest PHC and the mean adjusted

annual incidence for each fokontany. A correlation coefficient of zero corresponded to

no bias. We calculated this bias for each potential value of ω, and ranked each value by

how well it corrected the bias.

3. Financial bias: We calculated financial bias as the mean ratio of annual incidence rates

in fokontany with user fee reimbursement to incidence in fokontany not benefiting from

user fee reimbursement. A ratio of 1 corresponded to no bias. We calculated this bias for
each potential value of ω, and ranked each value by how well it corrected the bias.
We calculated these three metrics (proportion of upper incidence outliers, geographic bias, and
financial bias) for the full range of minimum values of the scaling factor (ω) and ranked the
minimum scale factors by their performance for each metric, with lower ranks corresponding to
the least bias. To identify the optimum value of ω, we first identified the subset of ω that met our
upper incidence outlier criteria described above. Then, we identified the value of ω that resulted
in the lowest mean rank of correction for geographic and financial bias while minimizing the
difference in performance between the two, measured via the coefficient of variation.
The SI was rescaled between ω and 1 and then used in combination with the zero
adjusted data from step 3 to perform benchmark multiplication (Eq. 4). This resulted in a final
dataset of monthly malaria incidence for each age class from January 2016 - June 2021 for 192
fokontany, adjusted so that the sampling intensity was standardized for all fokontany and
matched that found in the "high access" fokontany.
422
Evaluating Adjusted Datasets
We evaluated our estimates of the SI and adjusted malaria incidence rates using external data
from the IHOPE cohort. Self-reported healthcare seeking behavior was paired spatially to SI
estimates by assigning a value to a fokontany if a village from the cluster was in that fokontany
and was paired temporally by taking the average of the SI during the 6 month period containing
the months when the IHOPE survey was conducted in each year (January through June for
2016 and 2021 and July through December for 2018), to reduce the impact of month outliers in
consultation rates on SI estimates. We assessed the agreement between the two datasets by
calculating the correlation between estimated SI and the proportion of residents reporting illness
who attended PHCs using Clifford's modified t-test, which controls for spatial autocorrelation
(Clifford et al. 1989). We assessed the correlation separately for each year (2016, 2018, 2021),
including 109 fokontany per year.
We evaluated the ability of our adjusted incidence rates to accurately represent malaria
burdens by comparing adjusted incidence rates to malaria prevalence data collected via the
IHOPE cohort in 2021. The two datasets were paired spatially by assigning a value to a
fokontany if a village from the cluster was in that fokontany and was paired overtime by
matching the month of the IHOPE survey to the month of the incidence rates. We compared
adjusted incidence rates for children under 15 years old to prevalence rates of children under 15
years old from the IHOPE cohort for all fokontany with information in both datasets (n= 109) via

Clifford's modified t-test. We also assessed the ability of the adjusted incidence data to correctly

identify hot spots of malaria, defined as the quartile of fokontany with the highest prevalence

values.

Ethics Statement

Use of aggregate monthly consultation counts from PHCs in Ifanadiana District for this study

was authorized by the Medical Inspector of Ifanadiana. The IHOPE longitudinal survey

implemented informed consent procedures approved by the Madagascar National Ethics

Committee and the Madagascar Institute of Statistics. Household-level de-identified data from

the IHOPE survey were provided to the authors for the current study. We recognize that all

research is conducted within the surrounding socio-political context and risks reproducing

existing inequalities within the research team and across research partners. We've chosen to

explicitly reflect on power dynamics and equitable authorship within the context of this research

project in an accompanying reflexivity statement (Supplemental Materials).

RESULTS

Estimating the Sampling Intensity

We estimated the SI by fitting a floating catchment area model to non-malaria consultation data

from January 2016 - December 2021. The resulting model performed well at reproducing the

non-malaria consultation data (under-5: Spearman's ρ = 0.615, p-value<0.001; juvenile:

Spearman's ρ = 0.609, p-value<0.001; adult: Spearman's ρ = 0.704, p-value<0.001). When

averaged over all fokontany per month, it accurately represented the temporal trends in the non-

malaria consultation data (under-5: Spearman's ρ = 0.389, p-value <0.002; juvenile:

Spearman's ρ = 0.608, p-value <0.001; adult: Spearman's ρ = 0.629, p-value <0.001). When

averaged across time to result in one average SI per fokontany, it also was able to capture

spatial and fokontany-specific differences in consultation rates and model predictions closely

matched the non-malaria consultation rates (under-5: Spearman's ρ =0.819, p-value <0.001;

juvenile: Spearman's ρ =0.800, p-value <0.001; adult: Spearman's ρ =0.840, p-value <0.001).

The spatial patterns in the estimated SI mirrored spatial patterns in self-reported

healthcare seeking behavior from the IHOPE longitudinal survey (Fig. 2). The estimated SI and

self-reported healthcare seeking rates were significantly correlated across all years (Clifford's t-

test; 2016: ρ = 0.54 (p=0.002), 2018: ρ = 0.67 (p = 0.008), 2021: ρ = 0.59 (p = 0.01)). Both data

sources estimate higher healthcare access at fokontany nearer the national transportation

network, specifically the paved road that runs east-west through the district, and in close

proximity to PHCs. In addition, the two datasets were in agreement that the majority of the

district has low access to healthcare.

Figure 2. The sampling intensity estimated via the floating catchment area (FCA) method
(bottom row) closely approximates self-reported healthcare seeking rates from the
IHOPE cohort (top row). Shading represents rates grouped into quartiles, with Q1
corresponding to the lowest healthcare utilization rate and Q4 to the highest. Diamond points
show the location of level-2 PHCs. Top row: Cluster-level healthcare seeking rates are
illustrated for each village in a cluster across the three survey years. Bottom row: The scaled
sampling intensity estimated via a floating-catchment area method.

Reduction of Bias in Malaria Incidence due to Geographic and Financial Barriers to Care

We selected a minimum SI scale value (ω) that best corrected geographic bias and bias due to

cost-of-care reimbursement policies (Fig. 1, Fig. 3). The unadjusted dataset showed evidence of

geographic bias; average annual incidence of malaria in children under-5 in a fokontany was

negatively correlated with the distance from that fokontany to the nearest PHC (Spearman's ρ =

-0.617, p-value < 0.001, Fig. 3A), showing an exponential distance decay. The adjusted dataset,

by comparison, demonstrated a much weaker, though still significant, negative relationship

between average annual incidence and distance to the nearest PHC (Spearman's ρ = -0.222, p-

value<0.01, Fig. 3A). Fokontany whose populations attended PHCs where user fees were

reimbursed reported 2.48 times higher incidence than those that did not benefit from the

reimbursement policy in the unadjusted dataset (Fig. 3B). Applying the ZERO-G method

drastically reduced this bias; the average annual incidence in these fokontany was 1.01 times

the incidence in fokontany with cost-of-care-reimbursement (Fig. 3B).

503   Figure 3. The ZERO-G adjustment method greatly reduces geographical and financial
504   bias in malaria incidence rates. Left: Each point represents the average annual malaria
incidence rates for a fokontany over the period of 2016-2020, with the x-axis showing the
distance to the nearest PHC. The smoothed line is the exponential (unadjusted) or linear
(adjusted) fit between average annual incidence and distance to PHC. Right: The median
monthly malaria incidence rates across fokontany whose closest PHC does or does not offer fee
reimbursement. Fee reimbursement began in January 2017.

Comparing Unadjusted and Adjusted Datasets

Comparing the unadjusted and adjusted datasets, we estimated that unadjusted case

notifications are capturing on average 32.4% of symptomatic malaria cases in the district. This

differed by year, with the lowest percentage of 29.2% in 2020 and the highest of 49.6% in 2017.

The level of under-ascertainment also varied across fokontany. On average, the adjusted

annual incidence in a fokontany was 9.82 (range: 1- 1533) times the unadjusted annual

incidence rate. However, when this was calculated omitting fokontany and year combinations

that reported zero malaria cases in a year (26 out of 944), this ratio was reduced to 6.59 (range:

1 - 113).

Validation with Prevalence Data

We validated ZERO-G by comparing incidence rates and prevalence rates in children under 15

years old. Unadjusted incidence rates were negatively correlated with RDT positivity rates

(Spearman's ρ = -18.2, p-value = 0.06) and showed no evidence of a correlation with the

proportion of symptomatic and RDT positive individuals (Spearman's ρ = -0.04, p-value = 0.6).

After adjusting the data, we found a positive correlation between incidence and prevalence rates

(Spearman's ρ = 0.2031, p-value = 0.04). While the estimated correlation coefficient between

incidence rates and the proportion of symptomatic and RDT positive children was positive in the

adjusted dataset, it remained insignificant (Spearman's ρ = 0.127, p-value = 0.2). The adjusted

dataset also doubled the number of correctly-ranked fokontany into quantiles that matched

those from the prevalence data (Fig. 4). The adjusted dataset correctly ranked 38 of 104

fokontany, compared to 19 in the unadjusted dataset.

Figure 4. The adjustment method results in monthly malaria incidence rates in 2021 that
more closely correspond to measures of malaria prevalence in children under 15 years
old. Left: Malaria prevalence as measured by rapid-detection tests (RDT) in children under 15
years old from the IHOPE cohort survey. Colors represent quartiles from Q1 (lowest incidence)
to Q4 (highest incidence). Middle: Monthly malaria incidence in the unadjusted dataset.
Quartiles that match those in the prevalence data are highlighted in black. Right: Monthly
malaria incidence in the adjusted data. Quartiles that match those in the prevalence data are

highlighted in black. Monthly incidence has been chosen to correspond to the month in which
the IHOPE survey was conducted for that fokontany.
544

DISCUSSION

There is a critical need for routine surveillance systems to produce estimates at the spatial scale

of individual communities so that control interventions can be targeted in collaboration with

community health programs. However, HMIS data are rarely kept disaggregated at this scale

and, when they are, they suffer from under-estimation of incidence that varies across space and

time, preventing their use at these scales. We developed an adjustment method called the

ZERO-G multiplier that combines a gravity-model of healthcare access with a benchmark

multiplier method to create long-term routine surveillance data at the community-scale adjusted

for under-ascertainment due to uneven health care access. We demonstrated this method by

applying it to malaria case notification data from 192 communities and over 5 years of

surveillance in a rural District of Madagascar. This method reduced geographical and financial

bias in malaria incidence rates by 64% and 99%, respectively. In addition, we validated this

method with two external, population-representative datasets and found strong agreement with

self-reported healthcare access and malaria prevalence rates. Through this method, we were

able to obtain estimates that approximate long-term active surveillance data at fine-spatial

scales using only passive surveillance data, suggesting that the passive surveillance system

only reported about a third of all symptomatic malaria cases.

ZERO-G greatly reduced bias in malaria incidence rates in passive surveillance data in

our study setting. In Ifanadiana district, per capita health system utilization rates are twice as

high for fokontany within 5km of a health center than those further away (Garchitorena et al.

2021), and we found similar trends in the unadjusted malaria data (Fig. 3). Geographic bias in

the malaria data was therefore primarily reduced by accounting for low sampling intensity at

those fokontany further than 5 km from a PHC (Fig. 2). Financial costs represent a significant

barrier to healthcare seeking, particularly for low-income communities, and differential user fee

reimbursement policies over time (e.g. implementation of universal health coverage) can result

in healthcare access patterns changing as a function of this (James et al. 2005, Yates 2009). In

Ifanadiana, the removal of user fees via reimbursement policies in part of the district led to a

sudden and sustained 65% increase in utilization rates (Garchitorena et al. 2017). ZERO-G

removed this bias, resulting in similar incidence rates regardless of when and where

reimbursement policies were in place. ZERO-G also resulted in data that more accurately

identified malaria prevalence hotspots and coldspots than the unadjusted data, performing twice

as well as the unadjusted data. However, the adjusted dataset only correctly categorized 37% of

fokontany into ranked quantiles, illustrating the difficulty in matching incidence data to
prevalence data. The relationship between malaria prevalence and incidence is non-linear,
particularly for adults (Cameron et al. 2015), which can complicate matched ranking of
prevalence and incidence.

Unlike other methods, which rely on external datasets describing sampling intensity that
are collected at coarse spatial resolutions and infrequently (e.g. DHS, MICS, or other survey
data), ZERO-G uses data that match the spatial and temporal resolution of the case notification
data. This allows it to retain the original spatial and temporal scales at which the data was
collected while relying solely on public health and demographic data that is easily accessible to
public health actors. Population data can be sourced at fine-scale administrative levels via
national census data or via open-source datasets such as PopGrid (Leyk et al. 2019).
Information on PHC locations and services are collected by Ministries of Health or available via
regional, open-source datasets (e.g. Maina et al. 2019). We used a field-verified transport
network created via OpenStreetMap to estimate the distance between a population and a PHC,
which accurately represents patients' distance to PHCs (Ihantamalala et al. 2020); however,
these transportation networks are not globally available. When transportation networks are not
available, open-source databases of populations' distances to PHCs and other services could
serve as suitable substitutes (e.g. Nelson et al. 2019, Weiss et al. 2020). Finally, consultation
rates are commonly tracked by health systems and are increasingly recorded via electronic
health management information systems (Kumar et al. 2018, Siyam et al. 2021), facilitating their
use in these estimates.

ZERO-G differs from existing adjustment methods in several ways. First, it uses monthly
estimates of sampling intensity in the indirect estimate step rather than data from annual or
inter-annual population surveys. Most adjustment methods do not account for changes in
healthcare seeking behavior due to seasonality or temporal shifts to the health system (e.g.
changes in PHC staffing rates), and are therefore limited to inference at an annual frequency
(Gibbons et al. 2014). Second, the resulting dataset is available at the same spatial scale at
which it is collected, rather than spatially interpolated between points or aggregated to coarser
resolutions. We build on work by Hyde et al. (2021), which proposed a similar indirect estimation
adjustment method for malaria data that featured a monthly frequency at the fokontany scale,
but dealt with extreme low incidence values by spatially smoothing estimates between
neighboring fokontany, introducing spatial structure into the adjusted dataset and removing
existing natural variation. In addition, ZERO-G explicitly models the sampling intensity as a
function of geographic and health-system characteristics in all the facilities surrounding a

community via a gravity model instead of using information from the closest facility in a linear

model, as in Hyde et al. (Hyde et al. 2021). Because of this, changes in the health system can

be directly incorporated into calculations of sampling intensity in near real-time. It also allows for

estimation of sampling intensity in unsampled communities or months through these modeled

processes, rather than relying on interpolation.

There are several limitations that should be taken into consideration when evaluating or

implementing ZERO-G. First, active surveillance of malaria in our study area is limited, and we

only had access to one study of malaria prevalence at a spatial-scale finer than 5 x 5 km.

Therefore, we were only able to assess our method's ability to reproduce spatial patterns in

malaria burden, and not temporal patterns. However, our model results agree with national-level

trends in malaria, which witnessed over a 40% increase in confirmed malaria cases in 2020

(World Health Organization 2021), suggesting we are capturing temporal trends as well.

Second, we introduced a further source of uncertainty via the imputation of zero-incidence

samples due to extremely low ascertainment. This replaced months during the malaria season

reporting zero cases with predictions from a spatio-temporal model, mirroring established

methods (Gething et al. 2006), and replaced a small fraction of the overall data (8.77%). Finally,

our method does not include a step to disaggregate consultation rates to a finer spatial scale

than that reported by the PHC, often a major limiting step to accessing disease incidence data

at a fine spatial scale. In Ifanadiana, the standard reporting system aggregates consultations at

the level of the health facility catchment. We manually digitized health registers to obtain

community-level data, a time- and resource-intensive process. However, the increased

availability of electronic systems at the level of primary and community health care represents

an opportunity to apply this method directly and in real time to data at fine spatial scales.

In conclusion, ZERO-G represents a promising new method for adjusting passive

surveillance data for under-ascertainment bias in regions with low and heterogeneous

healthcare seeking rates, developed specifically for use at the community level. This method

can serve as part of a wider toolkit of statistical techniques used to improve targeted health

system responses. In a case study in a rural health district in Madagascar, it was able to reduce

geographic and financial bias in malaria incidence and the resulting dataset more closely

approximated spatial trends in malaria prevalence. ZERO-G can be applied to any case

notification dataset that suffers from geographic bias due to case under-ascertainment,

including both infectious and non-communicable diseases. It is particularly suited to rural areas,

where geographic isolation strongly influences healthcare access (Weiss et al. 2020). As

spatially-explicit health metrics become an increasingly important tool for precision public health

interventions, there is an urgent need to obtain and use quality data sources at the community

scale. Statistical methods such as ZERO-G can be an important tool to support the role of

community health programs in the local targeting of interventions for disease control.

DATA AVAILABILITY STATEMENT

All code and data needed to reproduce this study are available in a figshare repository (doi:

10.6084/m9.figshare.22154492).

ACKNOWLEDGMENTS

We would like to thank the health professionals who collect passive surveillance data in addition

to serving their patients and the Pivot data collection team for their work collecting and digitizing

this data. We would also like to thank the staff at Pivot who supervise much of this work,

particularly Laura Cordier. We would also like to thank Ann Miller and Marius

Randriamanambintsoa for their support of the IHOPE longitudinal survey.

FUNDING STATEMENT

This work was supported by internal funding from Pivot (https://www.pivotworks.org/), which

provided salary for FAI, MR, ATA, MHB, KEF, RJLR, M R-F, and BR. It was also supported by a

grant from the Agence Nationale de la Recherche (Project ANR-19-CE36-0001-01), granted to

AG, which supported AG, MR, and MVE. The funders had no role in study design, data

collection and analysis, decision to publish, or preparation of the manuscript.

REFERENCES

Admon, A. J., J. Bazile, H. Makungwa, M. A. Chingoli, L. R. Hirschhorn, M. Peckarsky, J.

Rigodon, M. Herce, F. Chingoli, P. N. Malani, and B. L. Hedt-Gauthier. 2013. Assessing and

improving data quality from community health workers: a successful intervention in Neno,

Malawi. Public Health Action 3:56–59.

Afrane, Y. A., G. Zhou, A. K. Githeko, and G. Yan. 2013. Utility of Health Facility-based Malaria

Data for Malaria Surveillance. PLOS ONE 8:e54305.

Amouzou, A., C. Faye, K. Wyss, and T. Boerma. 2021. Strengthening routine health information

systems for analysis and data use: a tipping point. BMC Health Services Research 21:618.

Buuren, S. van, and K. Groothuis-Oudshoorn. 2011. mice: Multivariate Imputation by Chained

Equations in R. Journal of Statistical Software 45:1–67.

Cameron, E., K. E. Battle, S. Bhatt, D. J. Weiss, D. Bisanzio, B. Mappin, U. Dalrymple, S. I.

Hay, D. L. Smith, J. T. Griffin, E. A. Wenger, P. A. Eckhoff, T. A. Smith, M. A. Penny, and P.

W. Gething. 2015. Defining the relationship between infection prevalence and clinical

incidence of Plasmodium falciparum malaria. Nature Communications 6:8170.

Cibulskis, R. E., M. Aregawi, R. Williams, M. Otten, and C. Dye. 2011. Worldwide Incidence of

Malaria in 2009: Estimates, Time Trends, and a Critique of Methods. PLOS Medicine

8:e1001142.

Clifford, P., S. Richardson, and D. Hemon. 1989. Assessing the Significance of the Correlation

between Two Spatial Processes. Biometrics 45:123–134.

Cordier, L. F., K. Kalaris, R. J. L. Rakotonanahary, L. Rakotonirina, J. Haruna, A. Mayfield, L.

Marovavy, M. G. McCarty, A. Tsirinomen'ny Aina, B. Ratsimbazafy, B. Razafinjato, T. Loyd,

F. Ihantamalala, A. Garchitorena, M. H. Bonds, and K. E. Finnegan. 2020. Networks of Care

in Rural Madagascar for Achieving Universal Health Coverage in Ifanadiana District. Health

Systems & Reform 6:e1841437.

Delamater, P. L. 2013. Spatial accessibility in suboptimally configured health care systems: A

modified two-step floating catchment area (M2SFCA) metric. Health & Place 24:30–43.

Delamater, P. L., A. M. Shortridge, and R. C. Kilcoyne. 2019. Using floating catchment area (FCA) metrics to predict health care utilization patterns. BMC Health Services Research 19:144.

Garchitorena, A., F. A. Ihantamalala, C. Révillion, L. F. Cordier, M. Randriamihaja, B. Razafinjato, F. H. Rafenoarivamalala, K. E. Finnegan, J. C. Andrianirinarison, J. Rakotonirina, V. Herbreteau, and M. H. Bonds. 2021. Geographic barriers to achieving universal health coverage: evidence from rural Madagascar. Health Policy and Planning 36:1659–1670.

Garchitorena, A., A. C. Miller, L. F. Cordier, R. Ramananjato, V. R. Rabeza, M. Murray, A. Cripps, L. Hall, P. Farmer, M. Rich, A. V. Orlan, A. Rabemampionona, G. Rakotozafy, D. Randriantsimaniry, D. Gikic, and M. H. Bonds. 2017. In Madagascar, Use Of Health Care Services Increased When Fees Were Removed: Lessons For Universal Health Coverage. Health Affairs 36:1443–1451.

Gething, P. W., A. M. Noor, P. W. Gikandi, E. A. A. Ogara, S. I. Hay, M. S. Nixon, R. W. Snow, and P. M. Atkinson. 2006. Improving Imperfect Data from Health Management Information Systems in Africa Using Space–Time Geostatistics. PLOS Medicine 3:e271.

Gibbons, C. L., M.-J. J. Mangen, D. Plass, A. H. Havelaar, R. J. Brooke, P. Kramarz, K. L. Peterson, A. L. Stuurman, A. Cassini, E. M. Fèvre, and M. E. Kretzschmar. 2014. Measuring underreporting and under-ascertainment in infectious disease datasets: a comparison of methods. BMC Public Health 14:147.

Hickman, M., and C. Taylor. 2005. Indirect Methods to Estimate Prevalence. Pages 113–131 *in* Z. Sloboda, editor. Epidemiology of Drug Abuse. Springer-Verlag, New York.

Hodgins, S., M. Kok, D. Musoke, S. Lewin, L. Crigler, K. LeBan, and H. B. Perry. 2021. Community health workers at the dawn of a new era: 1. Introduction: tensions confronting large-scale CHW programmes. Health Research Policy and Systems 19:109.

Hyde, E., M. H. Bonds, F. A. Ihantamalala, A. C. Miller, L. F. Cordier, B. Razafinjato, H.

Andriambolamanana, M. Randriamanambintsoa, M. Barry, J. C. Andrianirinarison, M. N. Andriamananjara, and A. Garchitorena. 2021. Estimating the local spatio-temporal distribution of malaria from routine health information systems in areas of low health care access and reporting. International Journal of Health Geographics 20:8.

Ihantamalala, F. A., V. Herbreteau, C. Révillion, M. Randriamihaja, J. Commins, T. Andréambeloson, F. H. Rafenoarimalala, A. Randrianambinina, L. F. Cordier, M. H. Bonds, and A. Garchitorena. 2020. Improving geographical accessibility modeling for operational use by local health actors. International Journal of Health Geographics 19:27.

James, C., S. S. Morris, R. Keith, and A. Taylor. 2005. Impact on child mortality of removing user fees: simulation model. BMJ 331:747–749.

Khan, A. A. 1992. An integrated approach to measuring potential spatial access to health care services. Socio-Economic Planning Sciences 26:275–287.

Kok, M. C., S. S. Kane, O. Tulloch, H. Ormel, S. Theobald, M. Dieleman, M. Taegtmeyer, J. E. Broerse, and K. A. de Koning. 2015. How does context influence performance of community health workers in low- and middle-income countries? Evidence from the literature. Health Research Policy and Systems 13:13.

Kumar, M., D. Gotz, T. Nutley, and J. B. Smith. 2018. Research gaps in routine health information system design barriers to data quality and use in low- and middle-income countries: A literature review. The International Journal of Health Planning and Management 33:e1–e9.

Leyk, S., A. E. Gaughan, S. B. Adamo, A. de Sherbinin, D. Balk, S. Freire, A. Rose, F. R. Stevens, B. Blankespoor, C. Frye, J. Comenetz, A. Sorichetta, K. MacManus, L. Pistolesi, M. Levy, A. J. Tatem, and M. Pesaresi. 2019. The spatial allocation of population: a review of large-scale gridded population data products and their fitness for use. Earth System Science Data 11:1385–1409.

Luo, W., and Y. Qi. 2009. An enhanced two-step floating catchment area (E2SFCA) method for measuring spatial accessibility to primary care physicians. Health & Place 15:1100–1107.

Maina, J., P. O. Ouma, P. M. Macharia, V. A. Alegana, B. Mitto, I. S. Fall, A. M. Noor, R. W. Snow, and E. A. Okiro. 2019. A spatial database of health facilities managed by the public health sector in sub Saharan Africa. Scientific Data 6:1–8.

Miller, A. C., A. Garchitorena, V. Rabeza, M. Randriamanambintsoa, H.-T. Rahaniraka Razanadrakato, L. Cordier, M. A. Ouenzar, M. B. Murray, D. R. Thomson, and M. H. Bonds. 2018. Cohort Profile: Ifanadiana Health Outcomes and Prosperity longitudinal Evaluation (IHOPE). International Journal of Epidemiology 47:1394–1395e.

Murray, C. J. L., A. D. Lopez, and S. Wibulpolprasert. 2004. Monitoring global health: time for new solutions. BMJ : British Medical Journal 329:1096–1100.

Nelson, A., D. J. Weiss, J. van Etten, A. Cattaneo, T. S. McMenomy, and J. Koo. 2019. A suite of global accessibility indicators. Scientific Data 6:266.

Noor, A. M., D. Zurovac, S. I. Hay, S. A. Ochola, and R. W. Snow. 2003. Defining equity in physical access to clinical services using geographical information systems as part of malaria planning and monitoring in Kenya. Tropical medicine & international health: TM & IH 8:917–926.

Ohrt, C., K. W. Roberts, H. J. W. Sturrock, J. Wegbreit, B. Y. Lee, and R. D. Gosling. 2015. Information Systems to Support Surveillance for Malaria Elimination. The American Journal of Tropical Medicine and Hygiene 93:145–152.

Perry, H. B., editor. 2020. Health for the people: National community health worker programs from Afghanistan to Zimbabwe. USAID MCHIP.

R Core Team. 2021. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Rifkin, S. B. 2018. Alma Ata after 40 years: Primary Health Care and Health for All—from consensus to complexity. BMJ Global Health 3:e001188.

Siyam, A., P. Ir, D. York, J. Antwi, F. Amponsah, O. Rambique, C. Funzamo, A. Azeez, L. Mboera, C. J. Kumalija, S. F. Rumisha, I. Mremi, T. Boerma, and K. O'Neill. 2021. The burden of recording and reporting health data in primary health care facilities in five low- and

lower-middle income countries. BMC Health Services Research 21:691.

Tatem, A. J., S. Adamo, N. Bharti, C. R. Burgert, M. Castro, A. Dorelien, G. Fink, C. Linard, M. John, L. Montana, M. R. Montgomery, A. Nelson, A. M. Noor, D. Pindolia, G. Yetman, and D. Balk. 2012. Mapping populations at risk: improving spatial demographic data for infectious disease modeling and metric derivation. Population Health Metrics 10:8.

Weiss, D. J., A. Nelson, C. A. Vargas-Ruiz, K. Gligorić, S. Bavadekar, E. Gabrilovich, A. Bertozzi-Villa, J. Rozier, H. S. Gibson, T. Shekel, C. Kamath, A. Lieber, K. Schulman, Y. Shao, V. Qarkaxhija, A. K. Nandi, S. H. Keddie, S. Rumisha, P. Amratia, R. Arambepola, E. G. Chestnutt, J. J. Millar, T. L. Symons, E. Cameron, K. E. Battle, S. Bhatt, and P. W. Gething. 2020. Global maps of travel time to healthcare facilities. Nature Medicine 26:1835–1838.

World Health Organization. 2021. World malaria report 2021. World Health Organization, Geneva.

World Health Organization. 2022. Global Health Expenditure Database.

World Health Organization, PEPFAR, and UNAIDS. 2007. Task shifting : rational redistribution of tasks among health workforce teams : global recommendations and guidelines. World Health Organization.

Yates, R. 2009. Universal health care and the removal of user fees. The Lancet 373:2078–2081.

Young, M., C. Wolfheim, D. R. Marsh, and D. Hammamy. 2012. World Health Organization/United Nations Children's Fund Joint Statement on Integrated Community Case Management: An Equity-Focused Strategy to Improve Access to Essential Treatment Services for Children. The American Journal of Tropical Medicine and Hygiene 87:6–10.