

Pre-pandemic inequalities in the burden of disease: a Scottish Burden of Disease study

Ian Grant¹, Neil Chalmers¹, Eilidh Fletcher¹, Fatim Lakha^{1,3}, Gerry McCartney², Diane Stockton¹, Grant Wyper¹

1. *Public Health Scotland, Edinburgh/Glasgow, Scotland*
2. *University of Glasgow, Glasgow, Scotland*
3. *NHS Tayside, Scotland*

Abstract

Background: Health inequalities in Scotland are well documented, including the contribution of different causes to inequalities in mortality. Our aim was to estimate inequalities within a burden of disease framework, accounting for both premature mortality and the effects of morbidity, to understand the contribution of specific diseases to health inequalities prior to the COVID-19 pandemic.

Methods: Disability Adjusted Life Years (DALYs) for 70 individual causes of disease and injury were sourced from the Scottish Burden of Disease Study. Area-level deprivation was measured using the Scottish Index of Multiple Deprivation. Inequalities were measured by the range, Relative Index of Inequality, Slope Index of Inequality, and attributable DALYs were estimated by using the least deprived decile as a reference.

Results: The overall disease burden was double that in the most deprived areas (50,305 vs 20,955 DALYs per 100,000), largely driven by inequalities in premature mortality. The rate in the most deprived decile was around 48% higher than the mean population rate (RII = 0.96), with 35% of DALYs attributed to differences in area-based deprivation. Many leading causes of disease burden in 2019 – heart disease, drug use disorders, lung cancer and COPD – were also the leading drivers of absolute and relative inequalities in the disease burden.

Conclusion: Our study evidences the extent of the stark levels of absolute and relative inequality prior to the COVID-19 pandemic. Given pre-pandemic stalling of mortality trend improvements and widening health inequalities, and the exacerbation of these caused by COVID-19, urgent policy attention is required to address this.

Introduction

It is well evidenced that there is a clear link between levels of deprivation and life expectancy (LE) in Scotland, with those from the most deprived areas and backgrounds having significantly poorer life expectancy and general health outcomes [1, 2, 3]. Recent published data from the National Records of Scotland shows that not only have overall differences in life expectancy between the most and least deprived populations have been worsening, but the gap in the number of years of healthy life expectancy (HLE) is also widening. In terms of overall life expectancy, males in the most deprived areas have 13.5 year's lower LE compared to their least deprived counterparts, and for females, the difference between the most and least deprived is 10.2 years [4]. Furthermore, HLE in the most deprived areas of Scotland was more than 24 years lower than in the least deprived areas for both males and females [5]. COVID-19 has further highlighted and amplified these inequalities: inequalities in COVID-19 mortality rates follow a similar social gradient to that seen for all causes of death i.e., increasing with increasing levels of deprivation, and the causes of inequalities in COVID-19 are similar to the causes of inequalities in health more generally [6,7,8].

Reliable, comparable information about the main causes of disease and injury in populations, and how these are changing and their effect on LE and HLE, is therefore a critical input for decision making, planning and targeting interventions where they are most required. Traditional sources of information about the descriptive epidemiology of diseases, injuries and risk factors are generally incomplete, fragmented and of uncertain reliability and comparability [9]. A lack of a standardized measurement framework to permit comparisons across diseases and injuries and failure to systematically evaluate data quality can impede comparative analyses of the true public health importance of various conditions and risk factors [10]. A burden of disease framework can be used to summarise the debilitating effects of morbidity and premature mortality in a population in a consistent and comparable manner. This is achieved by framing the effects of morbidity and mortality on population health loss as a function of time, in a composite measure called Disability-Adjusted Life Years (DALYs) [11]. By framing health loss in this way, DALYs combine the effects of morbidity and mortality in an equitable way and allow planners and policymakers to have a better understanding of the contribution that different diseases and injuries make to the total burden of disease [12].

Health inequalities in Scotland have been well documented, including the contribution of different causes to inequalities in mortality [1, 2, 13, 14, 15], There is, however, little information on how specific diseases contribute to these health inequalities within a standardised measurement framework, accounting for both premature mortality and the effects of morbidity. Using data from the Scottish Burden of Disease (SBoD) study, we assessed the contribution of individual disease and injuries, across a broad range of causes of morbidity and mortality, to absolute and relative health inequalities in Scotland in 2019.

Methods

Data

Area-level deprivation was measured using the Scottish Index of Multiple Deprivation (SIMD version 2016) [16]. The SIMD quantifies deprivation based on data zones, a geographical unit comparable to a postcode. Using pooled and weighted data from seven domains (employment, income, crime, housing, health, education and geographic access), each data zone is given a composite rank out of 6,505 data zones. The composite rank was then converted to a decile, with 1 assigned to the 10% most deprived data zones and 10 to the 10% least deprived data zones.

Estimates of the number of YLL, YLD, and DALYs for 13 broad disease groupings and 70 individual causes of disease and injury were sourced from the SBoD 2019 study [17]. YLL estimates were derived by multiplying age group-based mortality counts by the aspirational age-conditional life expectancy from the GBD 2019 reference life table [18]. The deprivation decile was derived using the individual's postcode of registered residence. YLD estimates were based on applying five-year age-specific rates, separately for each combination of deprivation decile for males and females, to mid-year population estimates for 2019, sourced from the National Records of Scotland [19]. As the underlying YLD rates were based on 2016 estimates, this approach assumed that any changes in the levels of YLD are entirely due to changes to the underlying demographic structure of the Scottish population.

Analyses

All analyses were carried out at the level of age-group and SIMD decile to facilitate the calculation of age-standardised rates (ASR). ASR using mid-year population estimates were calculated directly to the 2013 European Standard Population [19,20]. ASR removes the contribution of differences in underlying population structure between SIMD deciles to facilitate like-for-like comparisons. All ASR findings are presented per 100,000 population.

We estimated inequalities using several measures. The absolute and relative range differences in ASR were estimated as the difference and ratio respectively, between the most and least deprived deciles. Additionally, inequalities were measured using the Slope Index of Inequality (SII) and Relative Index of Inequality (RII) [21,22]. This involved fitting a linear regression to the ranked area deprivation deciles and the DALY values. An equal rate across deprivation deciles would give a horizontal line with a slope of zero (SII=0). The RII was estimated by dividing the SII rate of DALYs by the overall Scottish rate of DALYs. An equal rate across deprivation deciles would give a horizontal line with a slope of zero (RII=0). Disease DALYs attributable to area deprivation were estimated by using the least deprived decile as the reference group and summing all DALYs in excess of this reference across the other nine deprivation deciles. All DALYs in excess of this reference group, were summed and expressed as a percentage of the overall disease burden [23].

Results for all inequality measurements are presented for 13 disease groups and the 20 leading individual causes of disease and injury in Scotland. All analyses were performed using R and RStudio [24,25].

Results

Inequalities in the disease burden

There were approximately 1,732,800 DALYs lost in Scotland in 2019 (ASR 32,093 per 100,000 population), with YLL accounting for 63.5% of total DALYs. We found evidence of marked inequalities in DALYs across areas experiencing different levels of deprivation (Figure 1). DALYs in the most deprived areas were double that of the least deprived areas (ASR: 50,305 vs. 20,955). Additionally, there were inequalities in the composition of DALYs across areas experiencing different levels of deprivation with YLL accounting for 67.8% of total DALYs in the most deprived areas, compared to 57.1% in the least deprived areas (Figure 2). The impact of this meant that the contribution of YLD to total DALYs increased with decreasing levels of deprivation, from 32.1% in the most deprived decile, rising to 42.9% in the least deprived decile. Furthermore, YLL in the most deprived decile was substantially higher (34,922 per 100,000 population) than total DALYs (20,955 per 100,000 population) in the least deprived decile (*Table 1, Supplementary file*).

Fig 1. All-cause DALYs (per 100, 100) by SIMD decile, Scotland, 2019





Fig 2. Proportion of YLL and YLD by SIMD decile, Scotland 2019



Absolute and relative inequalities in all cause DALYs

The absolute difference in all-cause DALYs between the most and least deprived areas in Scotland was approximately 29,350 per 100,000 population (Table 1). The relative difference in DALYs between the most and least deprived areas was 2.4, indicating that DALYs were 2.4 times higher in our most, compared to least, deprived areas. The RII was 0.95, which when multiplied by 0.5 and expressed as a percentage means that the rate in the most deprived areas was 47.5% higher than the mean rate of DALYs in the Scottish population. The difference between the most and least deprived areas, measured by SII was 30,502 DALYs per 100,000 population. DALYs attributable to differences in area deprivation accounted for 35% of total DALYs.

Absolute and relative inequalities in disease groupings

In terms of absolute rate differences of DALYs, between the most and deprived areas, substance use disorders had the largest difference (6,837 DALYs per 100,000 population), followed by cancers (4,783 DALYs per 100,000 population) and cardiovascular diseases (4,583 DALYs per 100,000 population) (Table 1). For all others disease groups, except sensory conditions, there were positive absolute rate differences, indicating that the rate of DALYs were higher in the most, compared to least deprived areas.

Relative rate differences between the most and least deprived areas were also evident. In over half (9 out of 17) of the disease groups, the relative rate difference was ≥ 2.0 . Substance use disorders had the highest relative rate difference (20.5), followed by chronic respiratory diseases (4.0) and digestive diseases (3.5). These three disease groups also had the highest RII values indicating that the rate in most deprived communities was higher than the mean

rate of population: substance use disorders (RII = 2.82); chronic respiratory diseases (RII=1.49), digestive diseases (RII=1.41).

The disease groups with the largest estimates of DALYs attributable to inequalities in area deprivation were: substance use disorders (84.5%); chronic respiratory diseases (54.0%) and digestive diseases (49.7%). Conversely, we found a negative association for DALYs inequality attributable DALYs for sensory conditions (-13.8%). The estimates of inequality attributable DALYs for all other disease groups ranged from 13.7% for musculoskeletal disorders to 43.5% for injuries.

Table 1. Inequalities in the broad causes of DALYs, per 100, 00 population, Scotland, 2019

Disease group	ASR of DALYs per 100,000	Absolute rate difference between most and least deciles	Relative rate difference between most and least deciles	Slope Index of Inequality	Relative Index of Inequality	Attributable DALYs (%)
Cancers	6,460	4,780	2.0	4,831	0.75	27.6
Cardiovascular diseases	5,416	4,583	2.3	5,206	0.96	36.3
Neurological disorders	3,534	1,009	1.4	1,069	0.30	22.3
Mental health disorders	2,494	2,464	2.6	2,622	1.05	38.3
Musculoskeletal disorders	2,263	399	1.2	530	0.23	13.7
Substance use disorders	2,250	6,837	20.5	6,334	2.82	84.5
Chronic respiratory diseases	1,871	2,627	4.0	2,788	1.49	54.0
Injuries	1,646	1,980	3.1	2,061	1.25	43.5
Other non-communicable diseases	1,506	826	1.7	922	0.61	24.2
Digestive diseases	1,503	1953	3.5	2,118	1.41	49.7
Diabetes and chronic kidney diseases	1,133	1,028	2.5	1,108	0.98	38.3
Communicable, maternal, neonatal, and nutritional diseases	1,119	1,069	2.5	1,092	0.98	38.3
Sensory conditions	815	-204	-0.8	-169	-0.21	-13.8
All causes	32,093	29,350	2.4	30,512	0.95	35.3

Absolute and relative inequalities in individual causes

Of the 20 leading causes of diseases burden, drug use disorders had the largest absolute rate difference (5,452 DALYs per 100,000 population) between the most and least deprived

areas, followed by ischaemic heart disease (3,027 DALYs per 100,000 population), lung cancer (2,475 DALYs per 100,000 population) and chronic obstructive pulmonary disorder (2,358 DALYs per 100,000 population) (Table 2). For most of the remaining causes, there were positive absolute rate differences in DALYs, indicating DALYs were higher in the most compared to least deprived areas. The only exception to this was headache disorders, which had a negligible negative estimate of -15 DALYs per 100,000 population.

Relative rate differences between the most and least deprived areas were also evident across most of the leading individual causes of disease burden. Drug use disorders had the highest relative rate difference (28.3) then followed by alcohol use disorders (10.2), chronic obstructive pulmonary disorder (6.4), cirrhosis and other chronic liver diseases (5.4) and self-harm and suicide (4.9).

The diseases with the highest relative rate differences also had some of the highest RII values, indicating that the rate in most deprived communities was substantially higher than the mean rate of population (Scotland-level RII=0.96) for these diseases: substance use disorders (RII=2.99), alcohol use disorders (RII=2.31), chronic obstructive pulmonary disorder (RII=1.89), cirrhosis and other chronic liver diseases (RII=1.86), self-harm and suicide (RII=1.6) and lung cancer (RII=1.6). Notably, several conditions out with the leading causes of disease burden in Scotland had high RII values in the most deprived decile compared to mean rate of population, including schizophrenia (RII=5.5) and epilepsy (RII=4.7) (Table 2, Supplementary file).

The estimate of all-cause inequality attributable DALYs was 35.3%, however this varied considerably by individual cause (Table 2). Of the twenty leading causes of disease burden, DALYs attributable to inequalities in area deprivation ranged from 88.0% for drug use disorders, 74.4% for alcohol use disorders DALYs and 67.6 % for chronic obstructive pulmonary disorder, to 17.5% for low back and neck pain and 10% of headache disorder DALYs.

Table 2. Inequalities in the 20 leading causes of DALYs, Scotland, 2019

Leading causes	ASR of DALYs per 100,000	Absolute rate difference between most and least deciles	Relative rate difference between most and least deciles	Slope Index of Inequality	Relative Index of Inequality	Attributable DALYs (%)
Ischaemic heart disease	2,572	3,027	3.2	3,165	1.23	46.2
Alzheimer's disease and other dementias	1,764	492	1.4	576	0.33	25.0
Drug use disorders	1,673	5,452	28.3	5,003	2.99	88.0
Tracheal, bronchus, and lung cancer	1,596	2,475	4.3	2,518	1.58	54.9
Cerebrovascular disease	1,450	790	1.8	1,022	0.70	30.7
Chronic obstructive pulmonary disease	1,293	2,358	6.4	2,442	1.89	67.6
Low back and neck pain	1,269	355	1.3	430	0.34	17.5
Depression	1,243	1,212	2.6	1,282	1.03	38.2
Other cancers	1183	955	2.1	865	0.73	24.8
Headache disorders	982	-19	1.0	20	0.02	15.0
Other cardiovascular and circulatory diseases	908	574	1.8	738	0.81	30.2
Anxiety disorders	852	809	2.5	861	1.01	37.9
Self-harm and suicide	789	1,315	4.9	1,279	1.62	56.8
Colon and rectum cancer	752	226	1.3	219	0.29	21.4
Diabetes mellitus	744	688	2.5	760	1.02	39.7
Lower respiratory infections	699	670	2.5	717	1.03	37.9
Cirrhosis and other chronic liver diseases	626	1,107	5.4	1,164	1.86	62.6
Alcohol use disorders	577	1,385	10.2	1,332	2.31	74.4
Breast cancer	555	84	1.2	77	0.14	8.0
Other digestive diseases	431	465	3.0	502	1.17	47.9
All cause DALY	32,093	29,350	2.4	30,512	0.96	35.3

Discussion

Our study evidences the extent of stark levels of absolute and relative inequality, using range and distributional measures, in Scotland in 2019, prior to the COVID-19 pandemic.

The overall disease burden was double that in the most deprived areas in Scotland, largely driven by inequalities in mortality. We found that the gap in the burden of disease is so wide that the fatal burden of disease (YLL) in the most deprived areas was substantially higher than the total burden of disease (DALYs) in the least deprived areas. We already know from life expectancy figures that people in the most deprived communities die at younger age and spend more than a third of their lives in poor health compared to the least deprived communities [4,5]. Our findings further strengthen our understanding of the cause specific diseases driving the interaction between stalling mortality improvements, HLE and health inequalities [26,27].

Many of the leading causes of disease burden in 2019 – heart disease, drug use disorders, lung cancer and chronic obstructive pulmonary disease – were also the leading drivers of absolute and relative inequalities in the disease burden. Causes such as: alcohol, and drug use disorders; liver cirrhosis; chronic obstructive pulmonary disease; lung cancer; and, self-harm and suicide, exhibited the largest relative inequalities in disease burden which is consistent with results published elsewhere in the literature [2, 13, 15]. On the other hand, there were health conditions where there were smaller, to no, inequalities, for example: sensory conditions; headaches; and some musculoskeletal conditions. Triangulating these results with estimates of inequality-attributable DALY estimates for COVID-19, indicate that although COVID-19 infection has likely increased all-cause inequalities, there are several other causes of disease which have a much larger impact in driving levels of all-cause inequality in the burden of disease [8].

The COVID-19 pandemic has caused almost unprecedented change across health, education and the economy in Scotland and across the world. COVID-19 has been shown to have a substantial impact on the population health in Scotland, however this impact was not shared equally across areas experiencing different levels of socio-economic deprivation; in Scotland, the marked inequalities in COVID-19 YLL reported in 2020 were further exacerbated in 2021, to the extent that approximately half of COVID-19 YLL was attributable to inequalities in area deprivation. This widening of inequalities in 2021 was confirmed across all measures of absolute and relative inequality [7, 8]. Furthermore, health problems that existed before the COVID pandemic, have not gone away, and in many cases have been exacerbated, and unless future harm is prevented, this trend will continue. Integrating this impact, quantitatively, is difficult, and trends may not be comparable due to the major disruptions, and changes in access to, and delivery of, services, and the competition between causes of mortality [28]. Prior to the pandemic, improvements in life expectancy and healthy life expectancy in Scotland had stalled since 2012, with a slowdown in the overall progress of reducing mortality and widening of socioeconomic inequalities in mortality [5, 29,30, 35]. The results presented in this paper are therefore likely to be representative of wider pre pandemic trends given that YLD does not change quickly over time and that YLL estimates ultimately drive changes in the overall DALYs. Our estimates help us to understand the state of pre-pandemic inequalities, providing important baseline positions that can be triangulated with robust, and locally relevant, emerging evidence, to assess the likely impact of the pandemic, and other public health crisis such as the current cost-of-living crisis, on non-COVID and COVID-related health inequalities.

Strengths and Limitations

This study has several important strengths. Firstly, this study assesses the impact of inequalities using DALYs for over 70 individual diseases, conditions and injuries. DALYs provide a composite, internally consistent measure of population health loss which can be used to evaluate the proportionate burden of different diseases and injuries and compare population health by geographic region and over time. By combining information on fatal and non-fatal burden, BOD studies allow planners and policymakers to have a better understanding of the contribution that different diseases and injuries make to the total burden of disease and how this burden varies by levels of deprivation. This in turn supports decisions about where prevention and service activity should be focused to address health inequalities in Scotland.

YLD estimates are derived from surveys, and routine administrative data and as such may not be fully representative of the underlying burden by deprivation decile, for example administrative data are reflective of demand rather than need. This is also relevant for severity distributions, which are internationally derived with cancers being the main exception. In terms of impact of study results by deprivation decile, on the absolute scale this is most relevant for the causes with the largest YLD estimates, or those where YLD is the main contributor to disease-specific DALYs. For those causes which have a larger YLL component, it's plausible that any biases in YLD from unmet need or survey-specific bases will have a smaller impact on the DALY estimate if this unmet need causes disease-specific excess mortality.

Our measure of deprivation is area-based rather than individual measure and is therefore likely to misclassify many individuals into categories that do not reflect their individual experiences [32]. Although the SIMD includes health indicators in its range of domains (thereby raising the theoretical possibility of reverse causality whereby people are ordered by the health outcome rather than socioeconomic deprivation), the employment-income deprivation index (i.e., excluding the health measures) is very highly correlated with the overall SIMD index and so this is unlikely to have changed the results [33,34]. Furthermore, The SII and RII have the advantage that they are based on data about the whole population, rather than just the extremes, and so take into account inequalities across the entire distribution of inequality. They do, however, require a reasonably linear relationship between outcomes and exposures [31].

Conclusion

This study demonstrates that even before the COVID-19 pandemic there was a steep stepwise gradient in the burden of disease experienced across the Scottish population. This was evident for YLL and YLD, and almost every specific cause of death. By exploring cause-specific inequalities within a standardized measurement framework, to permit equitable comparisons across diseases and injuries, we confirm the stark gap between the most and least deprived areas in Scotland in 2019 prior to the COVID-19 pandemic and show that many of the leading causes of disease burden in 2019 were also the leading drivers of absolute and relative health inequalities in the disease burden.

Given the pre-pandemic stalling of mortality trend improvements and widening health inequalities, and the exacerbation of these caused by COVID-19, urgent policy attention is required to reduce the burden of fatal and non-fatal conditions in Scotland.

References

- 1 Audit Scotland. Health Inequalities in Scotland. Audit Scotland 2012 URL: http://www.audit-scotland.gov.uk/docs/health/2012/nr_121213_health_inequalities.pdf (Accessed on 30 January 2023).
- 2 Leyland AH, Dundas R, McLoone P, Boddy FA. Cause-specific inequalities in mortality in Scotland: two decades of change. A population-based study. BMC Public Health. 2007;7:172. <https://doi.org/10.1186/1471-2458-7-172>.
- 3 Mesalles-Naranjo O, Grant I, Wyper GMA, *et al*. Trends and inequalities in the burden of mortality in Scotland 2000–2015. PLoS One. 2018;13(8):e0196906. <https://doi.org/10.1371/journal.pone.0196906>.
- 4 National Records of Scotland. Life expectancy in Scotland 2018-2020 NRS 2021. URL:<https://www.nrscotland.gov.uk/files//statistics/life-expectancy-in-scotland/18-20/life-expectancy-18-20-report.pdf> (Accessed on 30 January 2023).
- 5 National Records of Scotland. Healthy Life Expectancy in Scotland, 2018-2020. NRS 2022 <https://www.nrscotland.gov.uk/statistics-and-data/statistics/statistics-by-theme/life-expectancy/healthy-life-expectancy-in-scotland/2018-2020> (Accessed on 30 January 2023).
- 6 Marmot M, Allen J, Goldblatt P *et al* (2020) Build Back Fairer: The COVID-19 Marmot Review The Pandemic, Socioeconomic and Health Inequalities in England. URL: <https://www.instituteofhealthequity.org/resources-reports/build-back-fairer-the-covid-19-marmot-review> (Accessed on 30 January 2023).
- 7 Wyper GMA, Fletcher E, Grant I, Harding O, de Haro Moro MT, Stockton DL, McCartney G. Correction: Inequalities in population health loss by multiple deprivation: COVID-19 and pre-pandemic all-cause disability-adjusted life years (DALYs) in Scotland. Int J Equity Health. 2022 Aug 25;21(1):117. <https://doi.org/10.1186/s12939-022-01714-4>.
- 8 Wyper GMA, Fletcher E, Grant I, McCartney G, Harding O, de Haro Moro MT, Stockton DL. Socioeconomic inequalities in COVID-19 DALYs in Scotland, 2020. Eur J Public Health. 2021 Oct 20;31(Suppl 3):ckab164.210. <https://doi.org/10.1093/eurpub/ckab164.210>.
- 9 Lopez, A.D. The evolution of the Global Burden of Disease framework for disease, injury and risk factor quantification: developing the evidence base for national, regional and global public health action. Global Health 1, 5 (2005). <https://doi.org/10.1186/1744-8603-1-5>.
- 10_Frank JW and Matsunaga E (2022) National monitoring systems for health inequalities by socioeconomic status – an OECD snapshot, Critical Public Health, 32:4, 580-587, DOI:[10.1080/09581596.2020.1862761](https://doi.org/10.1080/09581596.2020.1862761).

- 11 Murray CJL, Lopez AD. The Global Burden of Disease: A comprehensive assessment of mortality and disability from diseases, injuries and risk factors in 1990 and projected to 2020. World Health Organization and the World Bank; 1996. Cambridge, MA: Harvard University Press.
- 12 Lopez AD, Mathers CD, Ezzati M, *et al.* Global Burden of Disease and Risk Factors. Washington (DC): The International Bank for Reconstruction and Development / The World Bank; 2006. URL: <https://www.ncbi.nlm.nih.gov/books/NBK11812/>.
- 13 National Records of Scotland. Scotland's Population 2021 - The Registrar General's Annual Review of Demographic Trends. NRS 2022. URL: <https://www.nrscotland.gov.uk/statistics-and-data/statistics/stats-at-a-glance/registrar-generals-annual-review/2021> (Accessed on 30 January 2023).
- 14 Brown D, Allik M, Dundas R, Leyland AH, Inequalities in mortality in Scotland using a small area measure of deprivation. European Journal of Public Health, Volume 24, Issue suppl_2, October 2014, cku166–022, <https://doi.org/10.1093/eurpub/cku166.022>.
- 15 Allik M, Brown D, Dundas R, Leyland AH. Deaths of despair: cause-specific mortality and socioeconomic inequalities in cause-specific mortality among young men in Scotland. Int J Equity Health. 2020 Dec 4;19(1):215. doi: 10.1186/s12939-020-01329-7.
- 16 The Scottish Government Scottish Index of Multiple Deprivation 2016. URL: <https://www.data.gov.uk/dataset/a448dd2a-9197-4ea0-8357-c2c9b3c29591/scottish-index-of-multiple-deprivation-simd-2016> (Accessed on 30 January 2023).
- 17 Public Health Scotland. The Scottish Burden of Disease Study URL: <https://www.scotpho.org.uk/comparative-health/burden-of-disease/overview>. (Accessed on 30 January 2023).
- 18 Global Burden of Disease Collaborative Network. Global Burden of Disease Study 2019 (GBD 2019) Reference Life Table. Seattle, United States: Institute for Health Metrics and Evaluation (IHME), 2021.
- 19 National Records Scotland URL:(<https://www.nrscotland.gov.uk/statistics-and-data/statistics/statistics-by-theme/population>) (Accessed on 30 January 2023).
- 20 Eurostat. Revision of the European Standard Population Report of Eurostat's task force. Luxembourg: Publications Office of the European Union; 2013.
- 21 Moreno-Betancur M, Latouche A, Menvielle G, *et al.* Relative index of inequality and slope index of inequality: a structured regression framework for estimation. Epidemiology. 2015 Jul;26(4):518-27. doi: 10.1097/EDE.0000000000000311.
- 22 Pamuk ER. Social class inequality in mortality from 1921 to 1972 in England and Wales. Popul Stud (Camb). 1985 Mar;39(1):17-31. doi: 10.1080/0032472031000141256.
- 23 Carr-Hill R, Chalmers-Dixon P. The Public Health Observatory Handbook of Health Inequalities Measurement. South East Public Health Observatory 2005. URL: http://snspsms.ro/UserFiles/File/doc/the_public_health_observatory_handbook.pdf. (Accessed on 30 January 2023).
- 24 Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing 2021, Vienna, Austria. URL: <https://www.R-project.org/>.(Accessed 28 February 2023)

25 RStudio Team (). RStudio: Integrated Development for R. RStudio 2016, Inc., Boston, MA URL <https://posit.co/>. (Accessed 28 February 2023).

26. Ramsay J, Minton J, Fischbacher C, *et al.* How have changes in death by cause and age group contributed to the recent stalling of life expectancy gains in Scotland? Comparative decomposition analysis of mortality data, 2000-2002 to 2015-2017. *BMJ Open*. 2020 Oct 7;10(10):e036529. doi: 10.1136/bmjopen-2019-036529.

27 Walsh D, Wyper GMA, McCartney G. Trends in healthy life expectancy in the age of austerity. *J Epidemiol Community Health*. 2022 Jun 6;76(8):743–5. doi: 10.1136/jech-2022-219011.

28 Shah SA, Brophy S, Kennedy J *et al.* Impact of first UK COVID-19 lockdown on hospital admissions: Interrupted time series study of 32 million people. *EClinicalMedicine*. 2022 Jul;49:101462. <https://doi.org/10.1016/j.eclinm.2022>.

29 Fenton L, Minton J, Ramsay J, *et al.* Recent adverse mortality trends in Scotland: comparison with other high-income countries. *BMJ Open*. 2019;9:e029936. <https://doi.org/10.1136/bmjopen-2019-029936>.

30 Fenton L, Wyper GMA, McCartney G, Minton J. Socioeconomic inequality in recent adverse all-cause mortality trends in Scotland. *J Epidemiol Community Health*. 2019; 73:971–4. <https://doi.org/10.1136/jech-2019-212300>.

31 Scottish Government. Long-term monitoring of health inequalities: Scottish Government, 2022 URL: <https://www.gov.scot/collections/long-term-monitoring-of-health-inequalities/> (Accessed on 30 January 2023).

32 McLoone P. Targeting deprived areas within small areas in Scotland: population study. *BMJ*. 2001 Aug 18;323(7309):374-5. <https://doi.org/10.1136/bmj.323.7309.374>.

33 Fischbacher C. Identifying “deprived individuals”: are there better alternatives to the Scottish Index of Multiple Deprivation (SIMD) for socioeconomic targeting in individually based programmes addressing health inequalities in Scotland? Scottish Public Health Observatory/Information Services Division 2014, URL: <https://www.scotpho.org.uk/media/1166/scotpho140109-simd-identifyingdeprivedindividuals.pdf> Jan 2014 (Accessed on 30 January 2023).

34 Bradford DRP, Allik M, Brown D, McMahon AD. Assessing the risk of endogeneity bias in health and mortality inequalities research using composite measures of multiple deprivation which include health-related indicators: A case study using the Scottish Index of Multiple Deprivation and population health and mortality data medRxiv 2022.12.01.22282961; doi: <https://doi.org/10.1101/2022.12.01.22282961>.

35 Walsh D and McCartney G. Changing mortality rates in Scotland and the UK: an updated summary. Glasgow Centre for Population Health February 2023, URL: https://www.gcph.co.uk/assets/0000/9349/Changing_mortality_rates_in_Scotland_and_the_UK_-_an_updated_summary.pdf. (Accessed 28 February 2023).