How have mathematical models contributed to understanding the transmission and control of SARS-CoV-2 in healthcare settings? A systematic search and review

Authors: David R M Smith*[1,2,3], Sophie Chervet*[1,2,4], Théo Pinettes[3,5], George Shirreff[1,2,3], Sofía Jijón[3,5], Ajmal Oodally[1,2,3], Kévin Jean[3,5], Lulla Opatowski**[1,2], Solen Kernéis**[4,6], Laura Temime**[3,5]

*Contributed equally

**Contributed equally

Institutional affiliations :

[1] Institut Pasteur, Epidemiology and Modelling of Antibiotic Evasion (EMAE), Paris, France

[2] INSERM, Centre de recherche en épidémiologie et santé des populations (CESP), UMR 1018, "Anti-infective Evasion and Pharmacoepidemiology" team, UVSQ, Université Paris-Saclay, Montigny-Le-Bretonneux, France

[3] Conservatoire national des Arts et Métiers, Laboratoire Modélisation, épidémiologie et surveillance des risques sanitaires (MESuRS), Paris, France

[4] INSERM, Infection, Antimicrobials, Modelling, Evolution (IAME), UMR 1137, Université Paris-Cité, Paris, France

[5] Conservatoire national des Arts et Métiers, Institut Pasteur, Unité Pasteur-Cnam sur les Risques Infectieux et émergents (PACRI), Paris, France

[6] AP-HP, Hôpital Bichat, Equipe de Prévention du Risque Infectieux (EPRI), Paris, France

Abstract

Background: Since the onset of the COVID-19 pandemic, mathematical models have been

widely used to inform public health recommendations regarding COVID-19 control in

healthcare settings.

Objectives: To systematically review SARS-CoV-2 transmission models in healthcare

settings, and summarise their contributions to understanding nosocomial COVID-19.

Methods: Systematic search and review.

Data sources: Published articles indexed in PubMed.

Study eligibility criteria: Modelling studies describing dynamic inter-individual transmission

of SARS-CoV-2 in healthcare settings, published by mid-February 2022.

Participants and interventions: Any population and intervention described by included

models.

Assessment of risk of bias: Not appropriate for modelling studies.

Methods of data synthesis: Structured narrative review.

Results: Models have mostly focused on acute care and long-term care facilities in high-

income countries. Models have quantified outbreak risk across different types of individuals

and facilities, showing great variation across settings and pandemic periods. Regarding

surveillance, routine testing – rather than symptom-based testing – was highlighted as

essential for COVID-19 prevention due to high rates of silent transmission. Surveillance

impacts were found to depend critically on testing frequency, diagnostic sensitivity, and turn-

around time. Healthcare re-organization was also found to have large epidemiological

impacts: beyond obvious benefits of isolating cases and limiting inter-individual contact,

more complex strategies such as staggered staff scheduling and immune-based cohorting

reduced infection risk. Finally, vaccination impact, while highly effective for limiting COVID-

19 burden, varied substantially depending on assumed mechanistic impacts on infection

acquisition, symptom onset and transmission. Studies were inconsistent regarding which

individuals to prioritize for interventions, probably due to the high diversity of settings and

populations investigated.

Conclusions: Modelling results form an extensive evidence base that may inform control

strategies for future waves of SARS-CoV-2 and other viral respiratory pathogens. We

propose new avenues for future models of healthcare-associated outbreaks, with the aim of

enhancing their efficiency and contributions to decision-making.

Introduction

SARS-CoV-2 transmission in healthcare settings has contributed significantly to the global health-economic burden of COVID-19. Healthcare settings are particularly vulnerable to COVID-19, due to dense concentrations of frail patients, high frequencies of at-risk medical procedures, and high rates of inter-individual contact. Both patients and healthcare workers (HCWs) have been at high risk of SARS-CoV-2 infection throughout the pandemic, resulting in major nosocomial outbreaks worldwide [1,2]. In England, for instance, an estimated 20% of patients hospitalized with COVID-19 before August 2020 acquired their infection in hospital [3], while 95,000-167,000 patients became infected during their hospital stay between June 2020 and March 2021 [4]. Further, HCWs have experienced an estimated 1.6- to 3.4-fold higher risk of infection relative to the general population [5,6]. Long-term care facilities (LTCFs) have been most severely impacted, with one- to two-thirds of COVID-related deaths in Europe by May 2020 estimated to have occurred among LTCF residents [7].

Healthcare facilities have undergone extensive organizational changes to combat SARS-CoV-2 transmission, particularly during local surges in COVID-19 cases. Many facilities have adopted non-pharmaceutical infection prevention and control (IPC) measures, including social distancing, reinforced contact precautions and hand hygiene procedures, use of personal protective equipment (PPE), banning of visitors, and systematic test-trace-isolate protocols. HCWs and residents of LTCFs have also been among the first populations targeted for vaccination. However, despite these interventions, nosocomial COVID-19 risk has not been eliminated, as evidenced by ongoing outbreaks in healthcare facilities worldwide. A key barrier to effective COVID-19 prevention in healthcare settings is imperfect understanding of transmission routes among patients and HCWs, and of the relative impacts of different control strategies, which depend on setting-specific organizational and demographic characteristics, immunological histories of the specific population concerned, and virological properties of locally circulating variants.

Throughout the pandemic, mathematical models (Box 1) have proven useful both to better

understand the epidemiological processes underlying SARS-CoV-2 outbreaks and to

support public health decision-making. Modelling studies focusing on nosocomial risk in

particular, although less publicized than those focusing on community risk, have addressed

a broad range of epidemiological questions [8] and aided public health decision-making [9].

However, epidemiological insights and public health recommendations from nosocomial

SARS-CoV-2 models have not previously been reviewed or synthesized systematically.

Here, we systematically search and review mathematical models developed to investigate

SARS-CoV-2 transmission dynamics and control strategies in healthcare settings over the

critical phase of the pandemic, present their main contributions, synthesize their conclusions,

and discuss their limits.

Methods

We conducted a systematic search and review of mathematical/mechanistic models of inter-

individual SARS-CoV-2 transmission within healthcare settings published up to February 14,

2022. Details on the search, inclusion and exclusion criteria, screening process and data

extraction are provided in the Supplementary File, and results are reported according to the

PRISMA guidelines [10] (see Supplementary File).

Results

Model characteristics

Overall, our search identified 1,431 studies, of which 59 were included after title, abstract

and full-text screening (Suppl. Fig. S1 and Suppl. Table). Most (43/59) were posted in a

publicly accessible pre-print server, with a median delay of 8 months (range: 1 to 24 months)

between initial preprint posting and publication.

The majority of models were stochastic (85%; 50/59), and about half were agent-based or

network models (53%; 31/59). At early stages of the pandemic, when testing resources were

highly limited, most studies focused on surveillance and healthcare organization (Fig. 1A).

By comparison, impacts of PPE have been assessed less frequently, and vaccination
strategies only began to be explored towards the end of 2020, as the first vaccines became
available.

While models were mostly developed by teams from the USA, the UK, and other European
countries, these models explored locations representing a wider range of countries
worldwide (Fig. 1B). Acute care facilities and long-term care facilities were the main types of
healthcare facilities considered, although this varied depending on the country of study (Fig.
1C) and the subject addressed (Fig. 1D).

Insights on SARS-CoV-2 acquisition routes and transmission risk

Few studies have attempted to estimate the reproduction number of SARS-CoV-2 in
healthcare settings, despite evidence that nosocomial and community risk may differ widely
[11]. Estimates of nosocomial R_0 range from 0.45 (0.36-0.56) in English acute care hospitals
using a relatively simple approach [12] to 8.72 (5.14-16.32) in a French LTCF using a
stochastic dynamic model accounting for imperfect surveillance [13]. Interestingly, in the
latter study, R_0 was estimated to drop to 1.33 (0.68-2.04) after introduction of control
measures.

Several models have quantified the relative burden of SARS-CoV-2 infection and
transmission among patients and HCWs over the course of the pandemic. HCWs were
identified as being at high risk of occupational exposure to infection, either from colleagues
or patients [14–16]. During the first wave in early 2020, they have been estimated to be the
most important source of onward nosocomial transmission, both to patients and other HCWs
[17,18]. However, other studies found that patient infection could result primarily from
transmission from other patients [4,16,18].

Insights on SARS-CoV-2 infection control

Evaluating and optimizing surveillance strategies

Models have been widely used to assess and compare the epidemiological impacts of

SARS-CoV-2 testing strategies. Because SARS-CoV-2 spreads extensively through pre-

symptomatic and asymptomatic transmission [19,20], the identification of non-symptomatic

infections is a key component of nosocomial IPC. Several studies have highlighted difficulty

controlling outbreaks when targeting only symptomatic individuals for testing [21–28].

Limited impact of only testing patients upon their admission has also been identified,

suggesting that more thorough screening methods are required to limit SARS-CoV-2

introductions from the community, visitors, HCWs or ancillary staff [24,27,29,30].

Many studies have evaluated the impact of routine testing of non-symptomatic individuals.

The most universal finding is that more frequent testing leads to greater reductions in

nosocomial transmission [12,14,21,23,24,30–40]. Similarly, increasing daily testing capacity

has been found to limit nosocomial transmission [27,41]. In the context of limited test

availability early in the pandemic, effective strategies identified for optimizing nosocomial

outbreak detection include pooling samples via group testing [27] and distributing tests over

several batches instead of using them all at once [42].

Modelling results are less consistent concerning which subpopulations to target for routine

non-symptomatic testing. Many conclude that targeting HCWs is most effective

[12,25,32,43], while others suggest targeting facility patients or residents [22,27,39].

Divergence owes to underlying modelling assumptions on how patients and HCWs differ,

regarding their infectiousness, susceptibility to infection, contact behaviour, and degree of

interaction with visitors and other individuals in the community. For instance, testing staff

proved more effective in a model of English care homes where the main source of SARS-

CoV-2 introductions was staff members infected in the community [25]. Conversely, testing

patients was more effective in models of a French rehabilitation hospital in which high rates

of contact among ambulatory patients translated to high rates of patient-to-patient

transmission [27,39]. In nursing homes, patient testing likely becomes increasingly important

when visitors or other interactions with the community are permitted [33].

Lastly, in addition to testing frequency, studies have quantified the critical impact of the

sensitivity and turnaround time of the test being used [24,28,30,32,33,37,38,44–47]. Several

studies have identified that gains in turnaround time tend to outweigh gains in test sensitivity,

explaining why rapid diagnostic tests (e.g. Ag-RDT) may be more effective than laboratory-

based tests (e.g. RT-PCR) for routine non-symptomatic testing [23,25,32,39]. Conversely, if

same-day test results are achievable, or if individuals effectively isolate while awaiting test

results, more sensitive laboratory tests likely outperform rapid tests [33,34,44].

Evaluating impacts of personal protective equipment (PPE)

Several studies have found that, when available, PPE use is highly effective for reducing

infection risk among both HCWs and patients. Although predicted reductions in infection risk

naturally depend on assumptions underlying PPE's impact on viral transmissibility, which

vary considerably across studies and for which data are sorely lacking, several studies

suggest that widespread PPE use remains a key SARS-CoV-2 prevention strategy, even

when conferring comparatively low levels of protection [15,29,37,48–50]. By preventing

infection, PPE use has also been shown to reduce HCW workplace absence [37] and

workday loss [22]. Very few studies have tackled the question of who should be given priority

to PPE access when in limited supply, although PPE has been shown to be particularly

effective when accessible to all HCWs [48].

Regarding different types of PPE that may be used, Hüttel et al. [15] found hand sanitizer to

be an effective means of reducing risk as a supplement to other strategies. Regarding

timing, earlier introduction of PPE was found to allow a much more efficient response [22]

and to enable prevention of large outbreaks [25]. However, further analyses suggest that the

level of protection PPE confers can be occasionally overwhelmed in the context of large

numbers of infected people in a room [51]. Finally, waning PPE adherence due to pandemic

fatigue could significantly impact the efficacy of PPE-based interventions [52].

Evaluating and optimizing healthcare organization

Many modelling studies have assessed the epidemiological impacts of healthcare facilities

adapting their structures of care and workplace organization. Larger facilities have been

found to be at greater risk of nosocomial SARS-CoV-2 outbreaks [43], and splitting facilities

into smaller independent units has been shown to reduce the total number of infected

individuals [36,48], except when transmissibility is high [53]. Forbidding visitors was

identified as not having a significant effect on outbreak probability [25], except when infection

prevalence among visitors' contacts in the community is higher than that of HCW community

contacts [54].

Models have highlighted that rapid isolation of positive cases is an effective strategy for

infection prevention [36,55,56]. Interestingly, models suggest that intermixing recovered

individuals with potentially susceptible individuals could reduce outbreak size [32], and that

sufficient spacing between patient beds is needed to limit transmission risk [36]. Results are

less consistent regarding isolation upon admission. Models have highlighted the efficacy of

isolating all newly admitted patients for a given duration [29] or while awaiting test results

[12]. Conversely, others report no additional benefit of front-door screening or quarantine

upon admission when other strategies are already in place [43,55].

Regarding staff organization, models have demonstrated benefits of cancelling HCW

gatherings [57,58] and of forcing shorter and fewer patient-HCW interactions [51,58],

although surprisingly this latter result was not confirmed by others [22]. Reducing between-

ward staff sharing also seems to reduce both the number of wards with infected individuals

[55] and the global reproduction number [37]. More complex staffing strategies have also

shown potential benefits, like splitting staff into two teams that do not interact [59,60],

synchronizing rather than staggering staff rotations [61], or immunity-based staffing [32,35],

e.g. assigning recovered staff to infected patients [35]. Finally, admitting all COVID-19

patients to specialized quarantine hospitals in which HCWs continuously resided for 1-to-2

week-long shifts did not necessarily increase occupational HCW risk [62].

Evaluating vaccination strategies

All models exploring vaccination found that it could help reduce COVID-19 morbidity and
mortality, especially in LTCFs [23,28,31,33,52,63–67]. However, some studies also noted
that vaccination benefits could be hindered by high levels of community SARS-CoV-2
circulation [31,65] or by reduced adherence to contact precautions within facilities
concomitant with vaccine rollout, for instance due to pandemic fatigue or risk compensation
[52].

A major focus of these models has been evaluation of which groups of individuals to target
or prioritize for vaccination in a context of limited vaccine availability, yielding sometimes
inconsistent results. Some found that LTCF residents should be prioritized over staff,
especially in LTCFs with low adherence to IPC measures [33,52]. Conversely, staff
vaccination was shown to be particularly effective for decreasing the overall attack rate,
especially in the absence of a testing protocol [64]. Targeting staff for vaccination may also
be preferable when the risk of virus importation from the community is high [31]. Finally, it
has been shown that vaccine rollout should first target staff members most exposed to
potential COVID-19 patients (e.g. staff from emergency departments) [63].

It should be noted that the conclusions reached by these models depend strongly on
modelling assumptions underlying vaccine action. Across models, assumed mechanisms
related to vaccination effectiveness included one or several of the following: a reduction in
acquisition risk, a reduction in symptom risk, and a reduction in the infectiousness of infected
vaccinated individuals. For instance, it was shown that if a vaccine only reduces symptom
risk, then increasing vaccination among nursing home staff leads to an increase in the
proportion of asymptomatic infections among staff, which subsequently leads to increased
infection risk for residents [23]. Additionally, no model considered vaccine impact over the
long-term, which is particularly relevant in the context of waning immunity and the
emergence of novel variants with vaccine-escape properties.

Discussion

Mathematical models have become ubiquitous tools to help understand the dynamics of

infectious disease outbreaks and to support public health decision-making. Here, we have

reviewed how models have helped to inform COVID-19 risk management in healthcare

settings, in particular by providing a better understanding of nosocomial transmission

dynamics and control strategy effectiveness.

The generation of *in silico* evidence to inform infection control strategies has been the

leading motivation for nosocomial SARS-CoV-2 transmission modelling. Although real-world

evidence collected during randomized controlled trials is the gold-standard, such data are

extremely difficult to generate in the context of a rapidly evolving epidemic. Beyond the

obvious costs and time involved, great heterogeneity in population characteristics and

exposure risk across different healthcare settings means that a large number of centres

must be enrolled to achieve sufficient cluster randomisation. Several high-impact

randomized controlled trials have nonetheless been successfully conducted in healthcare

settings despite these challenges, in particular to evaluate COVID-19 vaccines, therapies

and pre- or post-exposure prophylactic agents [68–71]. However, trials evaluating impacts of

common IPC interventions, including surveillance testing, PPE and healthcare

reorganization, are scarce [72,73].

In this context, mathematical modelling approaches have been particularly helpful to

investigate critical IPC questions in (near) real-time, since they allow for the simulation of

diverse scenarios at relatively high speed and low cost, while accounting for all available

knowledge and uncertainty at a given place and time. Model-based evaluations also allow

for levels of granularity in intervention arms that may be infeasible in real trial designs. Our

review highlights the range of modelling studies published before the end of 2020, at a time

when the scientific and medical communities were in particularly great need of evidence to

inform optimal allocation of limited infection prevention resources, including RT-PCR tests,

face masks and, later, vaccines.

However, two common means of SARS-CoV-2 transmission prevention with important

implications for the field have been notably under-addressed. First, modelling studies on the

relative impact of different types of face masks (e.g. surgical masks, N95 respirators) are

scarce [74], tied to a lack of precise data on how they impact viral transmissibility, as well as

their potential indirect roles as transmission vectors. Second, although ubiquitous in practice

at various stages of the pandemic, explicit social distancing interventions have rarely been

assessed [22,39]. This is probably because accurate modelling of social distancing requires

fine-scaled simulation of inter-individual contact networks, typically using an agent-based

approach, which is beyond the scope of most models. When faced with both epistemic

uncertainty and a need for relative computational simplicity, modelers may be reluctant to

include and formalize specific interventions that require arbitrary, highly sophisticated and/or

potentially wrongheaded assumptions about their mechanistic impacts on SARS-CoV-2

transmission. Instead, a common modelling strategy has been to include generic non-

pharmaceutical interventions that simply reduce the viral transmission rate, and which are

assumed to represent any combination of basic infection prevention interventions, including

face masks, gloves, gowns, face shields, hand hygiene or social distancing.

Relative to the evaluation of infection control strategies, modelling studies have more rarely

focused on the estimation of epidemiological parameters using statistical inference. In

particular, in the event of the sudden emergence of a novel infectious disease, R_0 estimation

is essential for epidemic forecasting and emergency response planning, and relies largely on

mathematical modelling approaches. Although estimates of R_0 quickly became available for

SARS-CoV-2 across various community settings in early 2020, [75], estimates specific to

healthcare settings remain scarce. Yet there is a great need for robust estimates across

diverse settings, as underlying levels of epidemic risk vary substantially across facilities due

to their intrinsic heterogeneity (e.g. types of care provided, population sizes, contact

behaviour of these populations). For instance, assuming R_0=3.5 in the community, Temime

et al. [11] estimated that nosocomial R_0 could range from 0.7 to 11.7, depending on the type

of ward and the density of contacts among and between patients and HCWs. This

heterogeneity in nosocomial R_0 is consistent with the range of estimates derived from

models described in this review [12,13], and has critical implications, informing which

facilities and populations are most at risk for explosive outbreaks and thus most in need of

urgent infection control measures.

This lack of evidence likely stems from both data limitations and remaining uncertainty about

the relative importance of precise paths of transmission (e.g. through direct person-to-person

contact; transient viral carriage on hands, garments or shared medical devices; stagnant air

in poorly ventilated spaces). Particularly early in the pandemic, nosocomial COVID-19 data

came primarily from contexts of emergency outbreak management rather than routine data

collection through stable surveillance systems. For future waves of SARS-CoV-2 and other

infectious diseases, the estimation of epidemiological parameters may be made easier by

harnessing large databases that systematically gather patient and HCW tests results,

administrative data and healthcare exposures across diverse healthcare facilities over time.

Researchers have faced significant challenges when developing SARS-CoV-2 transmission

models. First, data limitations, particularly early in the pandemic, forced many modelers to

make assumptions that oversimplify healthcare facility structure, population behaviour and

SARS-CoV-2 transmission dynamics, limiting the applicability of some results to real-world

settings. Second, the shifting epidemiological landscape – characterized not only by the

rapid spread of SARS-CoV-2, but also rapid change in population behaviours, sudden

resource shortages, consecutive changes in public health recommendations, rapid approval

of novel diagnostics, therapies and vaccines, and the successive emergence of distinct

variants of concern – required researchers to continually adapt their models in order to

remain useful, with relevant data required to parameterize these updates often lagging

behind.

Greater interdisciplinarity will be required to maximize the utility of mathematical modelling in

the future. More direct lines of contact between modelers, hospital infection control teams,

clinicians and decision-makers should guide modelers in their research. First, this may help

to steer studies towards the questions that are most clinically relevant, as informed by the

needs and issues faced in real clinical settings. Second, this may help modelers to evaluate

strategies that are feasible in practice, considering logistical constraints such as workforce or

equipment availability and hospital structure. Third, these discussions may inspire modelers

to account for outcomes beyond transmission risk and infection burden, such as cost-

effectiveness or mental health. Indeed, interventions such as visitor restrictions or staff re-

organization can have a great impact on the social isolation of patients or workload of

HCWs, which is difficult to take into account explicitly in mathematical models. Cost-

effectiveness is increasingly considered in modelling studies; for instance, several studies

have quantified the health-economic efficiency of frequent non-symptomatic testing

[12,24,38,39,45]. However, more frequent estimation of health-economic outcomes may

increase their usefulness for decision-makers, who must balance the competing priorities of

maximizing population health and minimizing monetary cost. Finally, increased

communication across disciplines may facilitate more timely sharing of modelling results to

those who may benefit from them most, including infection control teams and hospital

administrators. The use of social media and the surge in posting of articles on pre-print

repositories during the COVID-19 pandemic have helped to facilitate the timely sharing of

results, but there remains an onus on academic publishers to ensure a timely peer review

process so that modelling results are shared quickly enough to maximize their impact.

This review has several limitations. First, we chose to exclude all statistical, mathematical, or

computational models not including inter-individual SARS-CoV-2 transmission.

Consequently, other types of models such as physical or biomechanical models of airborne

transmission were excluded [76–78]. Second, we excluded articles posted on public archives

such as arXiv, medrXiv or biorXiv [79], which are not subject to peer review and can be

difficult to track. Although we did include some preprints in our review, we were unable to

systematically explore these archives. Third, we may have missed articles published in

journals not referenced in PubMed (e.g. computer science or mathematical journals).
However, since our focus was on epidemiological insights and public health
recommendations, we do not believe that this significantly impacted our main findings.

Conclusion

Often developed in the face of great epidemiological uncertainty, mathematical models have
come to form a rich evidence base describing how SARS-CoV-2 spreads in healthcare
settings and informing which nosocomial COVID-19 control strategies are optimal, in
particular with regards to healthcare reorganization and the allocation of limited supplies of
PPE, diagnostic testing and vaccines. Into the future, epidemiological models may continue
to inform control strategies for outbreaks of SARS-CoV-2 and other viral respiratory
pathogens, but increased collaboration should be sought between modelers, hospital
infection control teams, clinicians, and public health decision-makers to help maximize their
utility.

Conflict of interest

Authors report no conflict of interest.

Funding

This study received funding from the MODCOV project from the Fondation de France (Grant
106059) as part of the alliance framework "Tous unis contre le virus", the Université Paris-
Saclay (AAP Covid-19 2020) and the French government through its National Research
Agency project SPHINX-17-CE36-0008-01 and the "Investissement d'Avenir" program,
Laboratoire d'Excellence "Integrative Biology of Emerging Infectious Diseases" (Grant ANR-
10-LABX-62- IBEID). The work was also supported directly by internal resources from the
French National Institute for Health and Medical Research (Inserm), Institut Pasteur, le
Conservatoire National des Arts et Métiers, and l'Université Versailles Saint-Quentin-en-
Yvelines/Université Paris-Saclay.

Author contributions

LO, LT, SK conceived the study, LO and LT acquired the funding. Screening of the articles

was performed by DRMS, SC, GS, SJ, KJ, LO and LT. Data extraction and analysis were

performed by DRMS, SC, GS, SJ, LO and LT. DRMS, SC and TP rendered figures and

statistics. All authors contributed to writing.

References

[1]   Zhou Q, Gao Y, Wang X, Liu R, Du P, Wang X, et al. Nosocomial infections among
patients with COVID-19, SARS and MERS: a rapid review and meta-analysis. Ann
Transl Med 2020;8:629. https://doi.org/10.21037/atm-20-3324.

[2]   Leclerc QJ, Fuller NM, Knight LE, CMMID COVID-19 Working Group, Funk S,
Knight GM. What settings have been linked to SARS-CoV-2 transmission clusters?
Wellcome Open Res 2020;5:83. https://doi.org/10.12688/wellcomeopenres.15889.2.

[3]   Knight GM, Pham TM, Stimson J, Funk S, Jafari Y, Pople D, et al. The contribution
of hospital-acquired infections to the COVID-19 epidemic in England in the first half
of 2020. BMC Infect Dis 2022;22:556. https://doi.org/10.1186/s12879-022-07490-4.

[4]   Cooper B, Evans S, Jafari Y, Pham TM, Lim C, Pritchard M, et al. The burden and
dynamics of hospital-acquired SARS-CoV-2 in England. Res Sq 2021.
https://doi.org/10.21203/rs.3.rs-1098214/v1.

[5]   Iversen K, Bundgaard H, Hasselbalch RB, Kristensen JH, Nielsen PB, Pries-Heje M,
et al. Risk of COVID-19 in health-care workers in Denmark: an observational cohort
study. Lancet Infect Dis 2020;20:1401–8. https://doi.org/10.1016/S1473-
3099(20)30589-2.

[6]   Nguyen LH, Drew DA, Graham MS, Joshi AD, Guo C-G, Ma W, et al. Risk of
COVID-19 among front-line health-care workers and the general community: a
prospective cohort study. Lancet Public Health 2020;5:e475–83.
https://doi.org/10.1016/S2468-2667(20)30164-X.

[7]   ECDC Public Health Emergency Team, Danis K, Fonteneau L, Georges S, Daniau C,
Bernard-Stoecklin S, et al. High impact of COVID-19 in long-term care facilities,
suggestion for monitoring in the EU/EEA, May 2020. Euro Surveill 2020;25.
https://doi.org/10.2807/1560-7917.ES.2020.25.22.2000956.

[8]   Zhang W, Liu S, Osgood N, Zhu H, Qian Y, Jia P. Using simulation modelling and
systems science to help contain COVID-19: A systematic review. Syst Res Behav Sci
2022. https://doi.org/10.1002/sres.2897.

[9]   Opatowski L, Temime L. Contributions of modelling for the control of COVID-19
405   nosocomial transmission. Anaesth Crit Care Pain Med 2022;41:101054.
406   https://doi.org/10.1016/j.accpm.2022.101054.

407   [10]   Page, McKenzie, Bossuyt, Boutron, Hoffmann, Mulrow CD, et al. The PRISMA
408   2020 statement: an updated guideline for reporting systematic reviews. BMJ
409   2021;372:n71. https://doi.org/10.1136/bmj.n71.

410   [11]   Temime L, Gustin M-P, Duval A, Buetti N, Crépey P, Guillemot D, et al. A
411   conceptual discussion about the basic reproduction number of severe acute respiratory
412   syndrome coronavirus 2 in healthcare settings. Clin Infect Dis 2021;72:141–3.
413   https://doi.org/10.1093/cid/ciaa682.

414   [12]   Evans S, Agnew E, Vynnycky E, Stimson J, Bhattacharya A, Rooney C, et al. The
415   impact of testing and infection prevention and control strategies on within-hospital
416   transmission dynamics of COVID-19 in English hospitals. Philos Trans R Soc Lond B
417   Biol Sci 2021;376:20200268. https://doi.org/10.1098/rstb.2020.0268.

418   [13]   Shirreff G, Zahar J-R, Cauchemez S, Temime L, Opatowski L, EMEA-MESuRS
419   Working Group on the Nosocomial Modelling of SARS-CoV-22. Measuring Basic
420   Reproduction Number to Assess Effects of Nonpharmaceutical Interventions on
421   Nosocomial SARS-CoV-2 Transmission. Emerging Infect Dis 2022;28:1345–54.
422   https://doi.org/10.3201/eid2807.212339.

423   [14]   Sanchez-Taltavull D, Castelo-Szekely V, Murugan S, Hamley JID, Rollenske T,
424   Ganal-Vonarburg SC, et al. Regular testing of asymptomatic healthcare workers
425   identifies cost-efficient SARS-CoV-2 preventive measures. PLoS ONE
426   2021;16:e0258700. https://doi.org/10.1371/journal.pone.0258700.

427   [15]   Hüttel FB, Iversen A-M, Bo Hansen M, Kjær Ersbøll B, Ellermann-Eriksen S,
428   Lundtorp Olsen N. Analysis of social interactions and risk factors relevant to the
429   spread of infectious diseases at hospitals and nursing homes. PLoS ONE
430   2021;16:e0257684. https://doi.org/10.1371/journal.pone.0257684.

431   [16]   Evans S, Stimson J, Pople D, Bhattacharya A, Hope R, White PJ, et al. Quantifying
432   the contribution of pathways of nosocomial acquisition of COVID-19 in English
433   hospitals. Int J Epidemiol 2022;51:393–403. https://doi.org/10.1093/ije/dyab241.

434   [17]   Abbas M, Robalo Nunes T, Cori A, Cordey S, Laubscher F, Baggio S, et al.
435   Explosive nosocomial outbreak of SARS-CoV-2 in a rehabilitation clinic: the limits of
436   genomics for outbreak reconstruction. J Hosp Infect 2021;117:124–34.
437   https://doi.org/10.1016/j.jhin.2021.07.013.

438   [18]   Lindsey BB, Villabona-Arenas ChJ, Campbell F, Keeley AJ, Parker MD, Shah DR, et
439   al. Characterising within-hospital SARS-CoV-2 transmission events: a retrospective
440   analysis integrating epidemiological and viral genomic data from a UK tertiary care
441   setting across two pandemic waves. MedRxiv 2021.
442   https://doi.org/10.1101/2021.07.15.21260537.

[19]   Buitrago-Garcia D, Egli-Gany D, Counotte MJ, Hossmann S, Imeri H, Ipekci AM, et
al. Occurrence and transmission potential of asymptomatic and presymptomatic
SARS-CoV-2 infections: A living systematic review and meta-analysis. PLoS Med
2020;17:e1003346. https://doi.org/10.1371/journal.pmed.1003346.

[20]   Yanes-Lane M, Winters N, Fregonese F, Bastos M, Perlman-Arrow S, Campbell JR,
et al. Proportion of asymptomatic infection among COVID-19 positive persons and
their transmission potential: A systematic review and meta-analysis. PLoS ONE
2020;15:e0241536. https://doi.org/10.1371/journal.pone.0241536.

[21]   Chin ET, Huynh BQ, Chapman LAC, Murrill M, Basu S, Lo NC. Frequency of
Routine Testing for Coronavirus Disease 2019 (COVID-19) in High-risk Healthcare
Environments to Reduce Outbreaks. Clin Infect Dis 2021;73:e3127–9.
https://doi.org/10.1093/cid/ciaa1383.

[22]   Huang Q, Mondal A, Jiang X, Horn MA, Fan F, Fu P, et al. SARS-CoV-2
transmission and control in a hospital setting: an individual-based modelling study. R
Soc Open Sci 2021;8:201895. https://doi.org/10.1098/rsos.201895.

[23]   Kahn R, Holmdahl I, Reddy S, Jernigan J, Mina MJ, Slayton RB. Mathematical
Modeling to Inform Vaccination Strategies and Testing Approaches for Coronavirus
Disease 2019 (COVID-19) in Nursing Homes. Clin Infect Dis 2022;74:597–603.
https://doi.org/10.1093/cid/ciab517.

[24]   Litwin T, Timmer J, Berger M, Wahl-Kordon A, Müller MJ, Kreutz C. Preventing
COVID-19 outbreaks through surveillance testing in healthcare facilities: a modelling
study. BMC Infect Dis 2022;22:105. https://doi.org/10.1186/s12879-022-07075-1.

[25]   Rosello A, Barnard RC, Smith DRM, Evans S, Grimm F, Davies NG, et al. Impact of
non-pharmaceutical interventions on SARS-CoV-2 outbreaks in English care homes: a
modelling study. BMC Infect Dis 2022;22:324. https://doi.org/10.1186/s12879-022-
07268-8.

[26]   Schmidt AJ, García Y, Pinheiro D, Reichert TA, Nuño M. Using Non-Pharmaceutical
Interventions and High Isolation of Asymptomatic Carriers to Contain the Spread of
SARS-CoV-2 in Nursing Homes. Life (Basel) 2022;12.
https://doi.org/10.3390/life12020180.

[27]   Smith DRM, Duval A, Pouwels KB, Guillemot D, Fernandes J, Huynh B-T, et al.
Optimizing COVID-19 surveillance in long-term care facilities: a modelling study.
BMC Med 2020;18:386. https://doi.org/10.1186/s12916-020-01866-6.

[28]   Vilches TN, Nourbakhsh S, Zhang K, Juden-Kelly L, Cipriano LE, Langley JM, et al.
Multifaceted strategies for the control of COVID-19 outbreaks in long-term care
facilities in Ontario, Canada. Prev Med 2021;148:106564.
https://doi.org/10.1016/j.ypmed.2021.106564.

[29]   Baek YJ, Lee T, Cho Y, Hyun JH, Kim MH, Sohn Y, et al. A mathematical model of
COVID-19 transmission in a tertiary hospital and assessment of the effects of

482    different intervention strategies. PLoS ONE 2020;15:e0241169.
483    https://doi.org/10.1371/journal.pone.0241169.

484 [30] Tsoungui Obama HCJ, Adil Mahmoud Yousif N, Alawam Nemer L, Ngougoue
485    Ngougoue PM, Ngwa GA, Teboh-Ewungkem M, et al. Preventing COVID-19 spread
486    in closed facilities by regular testing of employees-An efficient intervention in long-
487    term care facilities and prisons? PLoS ONE 2021;16:e0249588.
488    https://doi.org/10.1371/journal.pone.0249588.

489 [31] Gómez Vázquez JP, García YE, Schmidt AJ, Martínez-López B, Nuño M. Testing
490    and vaccination to reduce the impact of COVID-19 in nursing homes: an agent-based
491    approach. BMC Infect Dis 2022;22:477. https://doi.org/10.1186/s12879-022-07385-4.

492 [32] Holmdahl I, Kahn R, Hay JA, Buckee CO, Mina MJ. Estimation of Transmission of
493    COVID-19 in Simulated Nursing Homes With Frequent Testing and Immunity-Based
494    Staffing. JAMA Netw Open 2021;4:e2110071.
495    https://doi.org/10.1001/jamanetworkopen.2021.10071.

496 [33] Lasser J, Zuber J, Sorger J, Dervic E, Ledebur K, Lindner SD, et al. Agent-based
497    simulations for protecting nursing homes with prevention and vaccination strategies. J
498    R Soc Interface 2021;18:20210608. https://doi.org/10.1098/rsif.2021.0608.

499 [34] Love J, Wimmer MT, Toth DJA, Chandran A, Makhija D, Cooper CK, et al.
500    Comparison of antigen- and RT-PCR-based testing strategies for detection of SARS-
501    CoV-2 in two high-exposure settings. PLoS ONE 2021;16:e0253407.
502    https://doi.org/10.1371/journal.pone.0253407.

503 [35] Lucia-Sanz A, Magalie A, Rodriguez-Gonzalez R, Leung C-Y, Weitz JS. Modeling
504    shield immunity to reduce COVID-19 transmission in long-term care facilities. Ann
505    Epidemiol 2023;77:44–52. https://doi.org/10.1016/j.annepidem.2022.10.013.

506 [36] Martos DM, Parcell BJ, Eftimie R. Modelling the transmission of infectious diseases
507    inside hospital bays: implications for COVID-19. Math Biosci Eng 2020;17:8084–
508    104. https://doi.org/10.3934/mbe.2020410.

509 [37] Pham TM, Tahir H, van de Wijgert JHHM, Van der Roest BR, Ellerbroek P, Bonten
510    MJM, et al. Interventions to control nosocomial transmission of SARS-CoV-2: a
511    modelling study. BMC Med 2021;19:211. https://doi.org/10.1186/s12916-021-02060-
512    y.

513 [38] See I, Paul P, Slayton RB, Steele MK, Stuckey MJ, Duca L, et al. Modeling
514    Effectiveness of Testing Strategies to Prevent Coronavirus Disease 2019 (COVID-19)
515    in Nursing Homes-United States, 2020. Clin Infect Dis 2021;73:e792–8.
516    https://doi.org/10.1093/cid/ciab110.

517 [39] Smith DRM, Duval A, Zahar JR, EMAE-MESuRS Working Group on Nosocomial
518    SARS-CoV-2 Modelling, Opatowski L, Temime L. Rapid antigen testing as a reactive
519    response to surges in nosocomial SARS-CoV-2 outbreak risk. Nat Commun
520    2022;13:236. https://doi.org/10.1038/s41467-021-27845-w.

521   [40]   Zhang Y, Cheng S-R. Evaluating the Need for Routine COVID-19 Testing of
522   Emergency Department Staff: Quantitative Analysis. JMIR Public Health Surveill
523   2020;6:e20260. https://doi.org/10.2196/20260.

524   [41]   Ding Y, Agrawal SK, Cao J, Meyers L, Hasenbein JJ. Surveillance Testing for Rapid
525   Detection of Outbreaks in Facilities. ArXiv 2021.
526   https://doi.org/10.48550/arxiv.2110.00170.

527   [42]   Yuval J, Nitzan M, Tannenbaum NR, Barak B. Optimizing testing policies for
528   detecting COVID-19 outbreaks. ArXiv 2020.

529   [43]   Nguyen LKN, Howick S, McLafferty D, Anderson GH, Pravinkumar SJ, Van Der
530   Meer R, et al. Evaluating intervention strategies in controlling coronavirus disease
531   2019 (COVID-19) spread in care homes: An agent-based model. Infect Control Hosp
532   Epidemiol 2021;42:1060–70. https://doi.org/10.1017/ice.2020.1369.

533   [44]   Kendall EA, Arinaminpathy N, Sacks JA, Manabe YC, Dittrich S, Schumacher SG, et
534   al. Antigen-based Rapid Diagnostic Testing or Alternatives for Diagnosis of
535   Symptomatic COVID-19: A Simulation-based Net Benefit Analysis. Epidemiology
536   2021;32:811–9. https://doi.org/10.1097/EDE.0000000000001400.

537   [45]   Stevenson M, Metry A, Messenger M. Modelling of hypothetical SARS-CoV-2
538   point-of-care tests on admission to hospital from A&E: rapid cost-effectiveness
539   analysis. Health Technol Assess 2021;25:1–68. https://doi.org/10.3310/hta25210.

540   [46]   Stevenson M, Metry A, Messenger M. Modelling of hypothetical SARS-CoV-2 point
541   of care tests for routine testing in residential care homes: rapid cost-effectiveness
542   analysis. Health Technol Assess 2021;25:1–74. https://doi.org/10.3310/hta25390.

543   [47]   Wilmink G, Summer I, Marsyla D, Sukhu S, Grote J, Zobel G, et al. Real-Time
544   Digital Contact Tracing: Development of a System to Control COVID-19 Outbreaks
545   in Nursing Homes and Long-Term Care Facilities. JMIR Public Health Surveill
546   2020;6:e20828. https://doi.org/10.2196/20828.

547   [48]   Qiu X, Miller JC, MacFadden DR, Hanage WP. Evaluating the contributions of
548   strategies to prevent SARS-CoV-2 transmission in the healthcare setting: a modelling
549   study. BMJ Open 2021;11:e044644. https://doi.org/10.1136/bmjopen-2020-044644.

550   [49]   Jang H, Polgreen PM, Segre AM, Pemmaraju SV. COVID-19 modeling and non-
551   pharmaceutical interventions in an outpatient dialysis unit. PLoS Comput Biol
552   2021;17:e1009177. https://doi.org/10.1371/journal.pcbi.1009177.

553   [50]   Procter SR, Abbas K, Flasche S, Griffiths U, Hagedorn B, O'Reilly KM, et al. SARS-
554   CoV-2 infection risk during delivery of childhood vaccination campaigns: a modelling
555   study. BMC Med 2021;19:198. https://doi.org/10.1186/s12916-021-02072-8.

556   [51]   Dy LF, Rabajante JF. A COVID-19 infection risk model for frontline health care
557   workers. Netw Model Anal Health Inform Bioinform 2020;9:57.
558   https://doi.org/10.1007/s13721-020-00258-3.

[52] Love J, Keegan LT, Angulo FJ, McLaughlin JM, Shea KM, Swerdlow DL, et al.
Continued need for non-pharmaceutical interventions after COVID-19 vaccination in
long-term-care facilities. Sci Rep 2021;11:18093. https://doi.org/10.1038/s41598-021-
97612-w.

[53] Nguyen LKN, Megiddo I, Howick S. Hybrid simulation modelling of networks of
heterogeneous care homes and the inter-facility spread of Covid-19 by sharing staff.
PLoS Comput Biol 2022;18:e1009780. https://doi.org/10.1371/journal.pcbi.1009780.

[54] Nguyen LKN, Howick S, McLafferty D, Anderson GH, Pravinkumar SJ, Van Der
Meer R, et al. Impact of visitation and cohorting policies to shield residents from
covid-19 spread in care homes: an agent-based model. Am J Infect Control
2021;49:1105–12. https://doi.org/10.1016/j.ajic.2021.07.001.

[55] Oodally A, Hammami P, Reilhac A, Guérineau de Lamérie G, Opatowski L, Temime
L. Hospital-level work organization drives the spread of SARS-CoV-2 within
hospitals: insights from a multi-ward model. MedRxiv 2021.
https://doi.org/10.1101/2021.09.09.21262609.

[56] Saurabh S, Kumar R, Kumar N, Bhardwaj P, Nag VL, Garg MK, et al. Dynamics of
SARS-CoV-2 Transmission Among Indian Nationals Evacuated From Iran. Disaster
Med Public Health Prep 2022;16:627–33. https://doi.org/10.1017/dmp.2020.393.

[57] Tofighi M, Asgary A, Merchant AA, Shafiee MA, Najafabadi MM, Nadri N, et al.
Modelling COVID-19 transmission in a hemodialysis centre using simulation
generated contacts matrices. PLoS ONE 2021;16:e0259970.
https://doi.org/10.1371/journal.pone.0259970.

[58] Cardinal RN, Meiser-Stedman CE, Christmas DM, Price AC, Denman C, Underwood
BR, et al. Simulating a Community Mental Health Service During the COVID-19
Pandemic: Effects of Clinician-Clinician Encounters, Clinician-Patient-Family
Encounters, Symptom-Triggered Protective Behaviour, and Household Clustering.
Front Psychiatry 2021;12:620842. https://doi.org/10.3389/fpsyt.2021.620842.

[59] Sánchez-Taltavull D, Castelo-Szekely V, Candinas D, Roldán E, Beldi G. Modelling
strategies to organize healthcare workforce during pandemics: Application to COVID-
19. J Theor Biol 2021;523:110718. https://doi.org/10.1016/j.jtbi.2021.110718.

[60] Cheng T, Liu J, Liu Y, Zhang X, Gao X. Measures to prevent nosocomial
transmissions of COVID-19 based on interpersonal contact data. Prim Health Care
Res Dev 2022;23:e4. https://doi.org/10.1017/S1463423621000852.

[61] Kluger DM, Aizenbud Y, Jaffe A, Parisi F, Aizenbud L, Minsky-Fenick E, et al.
Impact of healthcare worker shift scheduling on workforce preservation during the
COVID-19 pandemic. Infect Control Hosp Epidemiol 2020;41:1443–5.
https://doi.org/10.1017/ice.2020.337.

[62] Jijón S, Al Shafie A, Hassan E, EMAE-MESuRS working group on nosocomial
SARS-CoV-2 modeling, Temime L, Jean K, et al. Estimating the risk of incident
SARS-CoV-2 infection among healthcare workers in quarantine hospitals: the

Egyptian example. Sci Rep 2022;12:19773. https://doi.org/10.1038/s41598-022-
23428-x.

[63] Bosbach WA, Heinrich M, Kolisch R, Heiss C. Maximization of Open Hospital
Capacity under Shortage of SARS-CoV-2 Vaccines-An Open Access, Stochastic
Simulation Tool. Vaccines (Basel) 2021;9. https://doi.org/10.3390/vaccines9060546.

[64] Fosdick BK, Bayham J, Dilliott J, Ebel GD, Ehrhart N. Model-based evaluation of
policy impacts and the continued COVID-19 risk at long term care facilities. Infect
Dis Model 2022;7:463–72. https://doi.org/10.1016/j.idm.2022.07.003.

[65] Holmdahl I, Kahn R, Slifka KJ, Dooling K, Slayton RB. Modeling the impact of
vaccination strategies for nursing homes in the context of increased severe acute
respiratory syndrome coronavirus 2 community transmission and variants. Clin Infect
Dis 2022;75:e880–3. https://doi.org/10.1093/cid/ciac062.

[66] Blumberg S, Lu P, Kwan AT, Hoover CM, Lloyd-Smith JO, Sears D, et al. Modeling
scenarios for mitigating outbreaks in congregate settings. PLoS Comput Biol
2022;18:e1010308. https://doi.org/10.1371/journal.pcbi.1010308.

[67] Liu T, Bai Y, Du M, Gao Y, Liu Y. Susceptible-Infected-Removed Mathematical
Model under Deep Learning in Hospital Infection Control of Novel Coronavirus
Pneumonia. J Healthc Eng 2021;2021:1535046.
https://doi.org/10.1155/2021/1535046.

[68] Boulware DR, Pullen MF, Bangdiwala AS, Pastick KA, Lofgren SM, Okafor EC, et
al. A Randomized Trial of Hydroxychloroquine as Postexposure Prophylaxis for
Covid-19. N Engl J Med 2020;383:517–25. https://doi.org/10.1056/NEJMoa2016638.

[69] Simonovich VA, Burgos Pratx LD, Scibona P, Beruto MV, Vallone MG, Vázquez C,
et al. A Randomized Trial of Convalescent Plasma in Covid-19 Severe Pneumonia. N
Engl J Med 2021;384:619–29. https://doi.org/10.1056/NEJMoa2031304.

[70] Abella BS, Jolkovsky EL, Biney BT, Uspal JE, Hyman MC, Frank I, et al. Efficacy
and Safety of Hydroxychloroquine vs Placebo for Pre-exposure SARS-CoV-2
Prophylaxis Among Health Care Workers: A Randomized Clinical Trial. JAMA
Intern Med 2021;181:195–202. https://doi.org/10.1001/jamainternmed.2020.6319.

[71] Abarca K, Iturriaga C, Urzúa M, Le Corre N, Pineda A, Fernández C, et al. Safety
and Non-Inferiority Evaluation of Two Immunization Schedules with an Inactivated
SARS-CoV-2 Vaccine in Adults: A Randomized Clinical Trial. Vaccines (Basel)
2022;10. https://doi.org/10.3390/vaccines10071082.

[72] Jafari Y, Yin M, Lim C, Pople D, Evans S, Stimson J, et al. Effectiveness of infection
prevention and control interventions, excluding personal protective equipment, to
prevent nosocomial transmission of SARS-CoV-2: a systematic review and call for
action. Infection Prevention in Practice 2022;4:100192.
https://doi.org/10.1016/j.infpip.2021.100192.

[73]   Chou R, Dana T, Jungbauer R, Weeks C, McDonagh MS. Masks for Prevention of
638   Respiratory Virus Infections, Including SARS-CoV-2, in Health Care and Community
639   Settings: A Living Rapid Review. Ann Intern Med 2020;173:542–55.
640   https://doi.org/10.7326/M20-3213.

641   [74]   Ferris M, Ferris R, Workman C, O'Connor E, Enoch DA, Goldesgeyme E, et al.
642   Efficacy of FFP3 respirators for prevention of SARS-CoV-2 infection in healthcare
643   workers. ELife 2021;10. https://doi.org/10.7554/eLife.71131.

644   [75]   Du Z, Liu C, Wang C, Xu L, Xu M, Wang L, et al. Reproduction Numbers of Severe
645   Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Variants: A Systematic
646   Review and Meta-analysis. Clin Infect Dis 2022;75:e293–5.
647   https://doi.org/10.1093/cid/ciac137.

648   [76]   Zhou Y, Ji S. Experimental and numerical study on the transport of droplet aerosols
649   generated by occupants in a fever clinic. Build Environ 2021;187:107402.
650   https://doi.org/10.1016/j.buildenv.2020.107402.

651   [77]   Prajapati S, Mehta N, Chharia A, Upadhyay Y. Computational fluid dynamics-based
652   disease transmission modeling of SARS-CoV-2 Intensive Care Unit. Mater Today
653   Proc 2022;56:2967–72. https://doi.org/10.1016/j.matpr.2021.11.013.

654   [78]   Saw LH, Leo BF, Nor NSM, Yip CW, Ibrahim N, Hamid HHA, et al. Modeling
655   aerosol transmission of SARS-CoV-2 from human-exhaled particles in a hospital
656   ward. Environ Sci Pollut Res Int 2021;28:53478–92. https://doi.org/10.1007/s11356-
657   021-14519-9.

658   [79]   Fraser N, Brierley L, Dey G, Polka JK, Pálfy M, Nanni F, et al. The evolving role of
659   preprints in the dissemination of COVID-19 research and their impact on the science
660   communication landscape. PLoS Biol 2021;19:e3000959.
661   https://doi.org/10.1371/journal.pbio.3000959.

Mathematical model of SARS-CoV-2 transmission

Observed data on COVID-19 cases in patients and HCWs

Estimate the unobserved burden of COVID-19

Identify the most effective control measures

Legend

↔ *SARS-CoV-2 transmission*

🔴 *Interventions that limit transmission*

🔵 *transmission*

Control measures

- **Vaccination**
- **Enhanced ventilation**
- **Surveillance**
  - **A** : Testing HCWs and/or patients
  - **B** : Testing HCWs
  - **C** : Testing patients

- **Contact restrictions**
  - **1** : Patients/HCW cohorting – PPE
  - **2** : HCW cohorting – PPE
  - **3** : Hospital re-organization
  - **4** : Visitation restrictions

Mathematical models are theoretical constructs that mechanistically formalize the dynamic processes underlying SARS-CoV-2 transmission. A typical model splits the individuals present in a healthcare facility into sub-populations, including one or more categories of patients (or residents) and HCWs. Mathematical or statistical tools are used to describe the natural history of infection through the definition of different infection states or "compartments". For instance, in the specific case of SARS-CoV-2, which is characterized by an incubation period subsequent to exposure and the acquisition of (partial) immunity after infection, the main compartments considered are: Susceptible to infection (S), Exposed to infection or incubating (E), Infectious (I), and Recovered or immunized (R). As a large share of infectious individuals may be asymptomatic, the I compartment is often subdivided into asymptomatic (IA) and symptomatic (IS) compartments. Various other sources of heterogeneity may also be considered, including different levels of viral shedding among infectious individuals, or different trajectories of care among symptomatic individuals (e.g., isolation, mechanical ventilation, admission to intensive care, death). Specific contact patterns between individuals of different sub-populations and infection statuses can further be accounted for through the definition of contact matrices.

Compartmental models are most frequent, but agent-based models (also known as individual-based models) are another common formalism, in which each unique individual in the population is explicitly modelled. This enables more detailed integration of heterogeneity in contact patterns, disease progression, transmission risk and other epidemiological processes. Models can further be categorized as either deterministic or stochastic. In deterministic models, there is no randomness in epidemiological processes, and a particular set of initial conditions always results in identical outbreaks. By contrast, in stochastic models, it is possible to account for randomness in the parameters or processes included, resulting in different outbreak trajectories each time the model is run. Stochasticity is particularly relevant in models of healthcare settings, where population sizes are small and randomness can have a strong impact on outbreak dynamics.

Models are used for a variety of purposes. They are widely used to simulate virus transmission in specific settings and populations, allowing for the quantification of virus burden in particular epidemiological scenarios (e.g. after the introduction of a novel SARS-CoV-2 variant into a hospital via newly admitted patients, short-stay visitors, or members of staff infected in the community). Models are also used to enable the *in silico* assessment of public health interventions through the mechanistic implementation of interventions (e.g. testing, isolation, PPE provisioning, contact restrictions, vaccine deployment). Intervention impact can then be evaluated by simulating counterfactual outbreaks with and without the intervention in place. Finally, models are key to analysing reported data from real outbreaks (e.g. time series data, individual line lists of SARS-CoV-2 cases), accounting not only for unobserved processes (e.g., virus transmission) but also incomplete infection data, whether due to the presence of asymptomatic infections, imperfect reporting, or limited surveillance capacity. This allows for the retrospective assessment of true disease burden from a given outbreak (e.g., cumulative infection incidence) as well as estimation of important epidemiological parameters such as R_0, the basic reproduction number, which describes the average number of secondary cases caused by an index case in an immunologically naïve population. R_0 is particularly helpful to understand the epidemic potential of an emerging pathogen, though its value may vary across distinct sub-populations and settings, such as particular groups of patients and HCWs in particular healthcare facilities.