Nasopharyngeal Angiotensin Converting Enzyme 2 (ACE2) Expression as a Risk-Factor for SARS-CoV-2 Transmission in Concurrent Hospital Associated Outbreaks

Aidan M. Nikiforuk[1,4], Kevin S. Kuchinski[1,2], Katy Short[3], Susan Roman[3], Mike A. Irvine[1], Natalie Prystajecky[1,2], Agatha N. Jassem[1,2], David M. Patrick[1,4] and Inna Sekirov[1,2]

[1] British Columbia Centre for Disease Control, Vancouver, BC V5Z 4R4, Canada

[2] Department of Pathology and Laboratory Medicine, University of British Columbia, Vancouver, BC V6T 1Z4, Canada

[3] Fraser Health Authority, New Westminster, BC V3L 3C2, Canada

[4] School of Population and Public Health, University of British Columbia, Vancouver, BC V6T 1Z4, Canada

[5] Faculty of Health Sciences, Simon Fraser University, Burnaby, BC V5A 1S6, Canada

Abstract:

Background: Widespread human-to-human transmission of the severe acute respiratory syndrome coronavirus two (SARS-CoV-2) stems from a strong affinity for the cellular receptor angiotensin converting enzyme two (ACE2). We investigate the relationship between a patient's nasopharyngeal *ACE2* transcription and secondary transmission within a series of concurrent hospital associated SARS-CoV-2 outbreaks in British Columbia, Canada.

Methods: Epidemiological case data from the outbreak investigations was merged with public health laboratory records and viral lineage calls, from whole genome sequencing, to reconstruct the concurrent outbreaks using infection tracing transmission network analysis. *ACE2* transcription and RNA viral load were measured by quantitative real-time polymerase chain reaction. The transmission network was resolved to calculate the number of potential secondary cases. Bivariate and multivariable analyses using Poisson and Negative Binomial regression models was performed to estimate the association between *ACE2* transcription the number of

SARS-CoV-2 secondary cases.

Results: The infection tracing transmission network provided n = 76 potential transmission

events across n = 103 cases. Bivariate comparisons found that on average *ACE2* transcription did

not differ between patients and healthcare workers (P = 0.86). High *ACE2* transcription was

observed in 98.6% of transmission events, either the primary or secondary case had above

average *ACE2*. Multivariable analysis found that the association between *ACE2* transcription and

the number of secondary transmission events differs between patients and healthcare workers. In

health care workers Negative Binomial regression estimated that a one unit change in *ACE2*

transcription decreases the number of secondary cases (B = -0.132 (95%CI: -0.255 to -0.0181)

adjusting for RNA viral load. Conversely, in patients a one unit change in *ACE2* transcription

increases the number of secondary cases (B = 0.187 (95% CI: 0.0101 to 0.370) adjusting for

RNA viral load. Sensitivity analysis found no significant relationship between *ACE2* and

secondary transmission in health care workers and confirmed the positive association among

patients.

Conclusion: Our study suggests that *ACE2* transcription has a positive association with SARS-

CoV-2 secondary transmission in admitted inpatients, but not health care workers in concurrent

hospital associated outbreaks, and it should be further investigated as a risk-factor for viral

transmission.

Keywords: ACE2, SARS-CoV-2, Infection Tracing, Transmission Network, Outbreak
Investigation, Multivariable Analysis, Poisson Regression Model, Negative Binomial Regression

49
Introduction:
The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) causes potentially

life threatening lower respiratory and systemic inflammatory disease defined as COVID-19[1–3].

SARS-CoV-2 has spread widely since late 2019 causing a global pandemic. Two previous public

health emergencies have provided the opportunity to study human-to-human transmission of

pathogenic coronaviruses. In 2003, the severe acute respiratory syndrome emerged infecting

eight-thousand four-hundred and twenty-two people [4]. In 2012 and 2015 the Middle East

respiratory syndrome coronavirus (MERS-CoV) caused epidemics in the Kingdom of Saudi

Arabia and South Korea with two-thousand five hundred and sixty-two laboratory confirmed

cases reported to the World Health Organization[5]. SARS- and MERS-CoV spread

predominately in health care settings and community spread was controlled by public health

interventions or self-limited[4,5]. The transmission pattern of SARS-CoV-2 contrasts that of the

other pathogenic coronaviruses (SARS-CoV and MERS-CoV). SARS-CoV-2 has predominately

spread within the community, the virus has evolved to become more infectious over time and

vaccination does not protect against re-infection or transmission of newer strains [6].

Transmission of respiratory viruses like SARS-CoV-2 involves a complex interplay of social,

environmental, and biological variables[7]. Numerous observational studies have described risk

factors of SARS-CoV-2 transmission across a variety of settings including hospitals[8],

households[9], and schools[10,11]. Cumulative evidence suggests that the increased human-to-

human transmissibility of SARS-CoV-2 in comparison to SARS-CoV stems from stronger

affinity for the primary host receptor angiotensin converting enzyme 2 (ACE2)[12–14]. SARS-

CoV-2 binds to ACE2 through its spike glycoprotein, the spike protein undergoes proteolytic

cleavage before binding with ACE2 prompting endocytosis and membrane fusion[13]. The

Omicron variant of SARS-CoV-2 has a stronger affinity for ACE2 and less reliance on host

proteases biasing cell entry towards endocytosis, consequently less cell fusion and syncytia

formation occurs[15]. ACE2 is highly expressed in the upper and lower respiratory tract,

upregulation may occur in response to stimulation with interferon[16]. SARS-CoV-2 exploits

upregulation of *ACE2* in an interferon rich environment (like the early stages of an innate

immune response) to spread from cell-to-cell producing a high viral titre[17]. In the later stages

of replication, SARS-CoV-2 downregulates *ACE2,* preventing any benefits to the host mediated

by interferon stimulated upregulation[18,19]. SARS-CoV-2 has a shifting relationship with

ACE2 and like viral load, ACE2 expression fluctuates through the course of viral infection[20–

22]. Regardless, SARS-CoV-2 replicates most efficiently in tissues rich with ACE2, the strong

positive association between viral load and ACE2 expression at the time of diagnostic testing

increase the degree of viral shedding and thereby, the risk of ongoing transmission [23].

To understand the role of *ACE2* expression in SARS-CoV-2 transmission, we conducted

an infection tracing network analysis of positive testers in concurrent, single-site, hospital

associated COVID-19 outbreaks in British Columbia during late 2020 and early 2021. The study

aims to: (i) describe the outbreaks in context of person, place, and time, (ii) visualize an infection

tracing transmission network to infer the number of secondary cases per primary case and (iii)

quantify the relationship between the number of potential secondary cases (outcome) and

nasopharyngeal transcription of transmembrane ACE2 (exposure) while adjusting for viral RNA

load, and interaction by case designation (patient or healthcare worker). We build from previous

work which demonstrated no relationship between transmembrane nasopharyngeal *ACE2*

transcription and age, biological sex or *TMPRSS2* transcription in a sample of COVID-19

negative persons[23]. This study contributes novel evidence of how nasopharyngeal *ACE2*

transcription may drive SARS-CoV-2 transmission and highlights the potential role of

respiratory masks and infection prevention and control measures in limiting the viruses'

nosocomial spread.

Methods:

Study Design and Participants:

We conducted surveillance for COVID-19 cases in a British Columbian hospital over a

series of outbreaks, from declaration of the first facility associated outbreak on 07-11-2020 to the

date the last outbreak was declared over on 04-01-2021 (dd-mm-yyyy). Participants included

patients and health care workers who tested positive for COVID-19 upon self-reporting or

showing indicative symptoms (e.g. fever, fatigue, cough and loss of taste or smell) or through

asymptomatic point prevalence testing during the outbreak investigations. Inclusion criteria were

applied to select study participants whose diagnostic specimens were tested centrally at the

British Columbia Center for Disease Control Public Health Laboratory, underwent SARS-CoV-2

whole genome sequencing, and had adequate remaining volume of nucleic acid extract to assay

ACE2 transcription. In the case that a participant was tested for COVID-19 more than once, their

first diagnostic specimen from the study period was sampled. Participants who met the inclusion

criteria (n = 202) were excluded from the study if their specimen collection container was not

identifiable (n = 22), whole genome sequencing failed and was not able to classify the viral

lineage (n = 63) or *ACE2* transcription was unmeasurable (n = 14) (Figure S1). An analytic

dataset of n= 103 participants was used for analysis. Demographic variables of age, biological

sex, case description (patient or health care worker) and site of infection/ site of transmission

(hospital unit) were drawn from public health laboratory data or the outbreak report of the

investigating epidemiologist. Outbreaks were defined by the presence of one or more confirmed

cases of COVID-19 which were epidemiologically linked within hospital units (Table S3). The

laboratory procedures were performed in a laboratory accredited by the College of American

Pathologists and BC's Diagnostic Accreditation Program using validated RT-qPCR and whole

genome sequencing protocols[24–26].

Ethics Statement:

Ethical approval for the study was sought from the University of British Columbia human

ethics board (#H20-01110) which was harmonized with the Fraser Health Authority. Written

informed consent was not required. All data was de-identified prior to analysis and the results

were not linked back to any identifying records. This study was deemed as minimal risk to the

participants involved. To ensure privacy the site of the outbreak series will not be disclosed.

Procedures:

Participant's diagnostic specimens were collected by nasopharyngeal swab and stored in

Universal Transport Medium^{TM} (UTM®) (COPAN) and stored at 4$\square$C before RNA extraction.

RNA extraction was performed on the MagMAX-96^{TM} platform with the Viral RNA isolation kit

(ThermoFisher). Host and viral gene transcription was assayed by quantitative real-time

polymerase chain reaction (RT-qPCR) on the Applied Biosystems 7500 Real-Time PCR

platform using TaqMan FastVirus 1-step polymerase (ThermoFisher). Reaction volumes were 20

μl, with 5 μl of RNA template, 1μl of 20X primer/probe, 5 μl FastVirus and 9 μl of nuclease free

water. Cycling conditions were set to: 50$\square$C for 5min, 95$\square$C for 20s followed by 40 cycles of

95$\square$C for 15s and 60$\square$C for 1 min. As previously described, multiplex RT-qPCR reactions were

used to detect host (*ACE2*, *GAPDH*, *RNaseP*) and viral (*E* gene) transcription[27]. Participants

were diagnosed COVID-19 positive with an *E* gene cycle threshold value < 38 and a *RNaseP*

gene cycle threshold value < 40. The *E* gene Ct values were transformed to genome equivalents

per millilitre using a standard curve of SARS-CoV-2 synthetic RNA (MN908947.3) (Twist

Bioscience)[23]. Relative gene transcription for *ACE2* was calculated in proportion to the

control gene *GAPDH* using the $2^{-\Delta\Delta Ct}$ method[28]. SARS-CoV-2 whole genome sequencing was

performed on all diagnostic specimens which tested positive for SARS-CoV-2 by RT-qPCR.

Viral genomes were amplified using a 1200bp amplicon scheme and sequenced on an Illumina

NextSeq instrument[24]. Genome assembly was performed via a modified ARTIC Nextflow

pipeline[29]. Quality control did not pass any genomes with < 85% completeness or < 10X depth

of coverage. Viral lineages were assigned using the PANGOLIN tool (Version 1.15.1)[30]. All

molecular and genomic testing was performed at the BCCDC-PHL.

Variable Definition:

Demographic variables of interest were drawn from public health laboratory data, or the

outbreak reports provided by the investigating epidemiologist. The measures included in the

study are age, biological sex, case description, diagnostic specimen collection date (dd-mm-

yyyy), SARS-CoV-2 *E* gene cycle-threshold (Ct) value, *ACE2* gene transcription, SARS-CoV-2

lineage (PANGO lineage), site of transmission (hospital unit) and site of infection (hospital unit).

Age, *E* gene Ct value (transformed to log10 GE/mL[23]) and *ACE2* gene transcription are

continuous numeric variables. Biological sex, case description, viral lineage, collection date and

site of transmission/infections are categorical variables. We assumed that collection date equals

date of symptom onset for this study because once the hospital declared an outbreak, patients and

health care workers were required to be tested at the onset of symptoms. *ACE2* gene

transcription was transformed to a categorical value using the mean transcription value ($\bar{X}$ =

0.00, SD = 1.08) of COVID-19 negative testers, from a previous study which collected a random

sample of n = 212 nasopharyngeal specimens in British Columbia during 2020[23]. The number

of potential secondary cases per primary case was determined by infection tracing transmission

network analysis. *ACE2* gene transcription and number of potential secondary cases are the

exposure and outcome of interest. Viral load and case description were included in multivariable

analysis because, they meet the definition of a confounder and are a common cause of the

exposure (*ACE2* gene transcription) and outcome (number of secondary cases)[31]. Viral load

and *ACE2* gene transcription share a strong positive association, people with high *ACE2* gene

transcription were found to have high viral loads[32]. Viral load relates to transmission in that

people with high viral loads may shed more infectious virus[33,34]. Case description (patient or

healthcare worker) relates to *ACE2* gene transcription because patients may have comorbidities

or receive medications which affect *ACE2* transcription[35]. Health care workers have additional

social connections relative to in-patients, as they can leave work and be exposed to SARS-CoV-2

outside of clinical areas or in the community[8].

Statistical Analysis:

<u>Descriptive Statistics:</u>

The analytic data (n = 103) was used for bivariate analysis. The variables of interest were

stratified by case description and categorical *ACE2* transcription. An epidemiological curve was

visualized for the series of concurrent hospital outbreaks from 07-11-2020 to 04-01-2021 (dd-

mm-yyyy). Parametric statistical tests were used given the large sample size of our study[36].

     <u>Infection Tracing Network Analysis:</u>

A matrix of all possible transmission pairs was permuted using SARS-CoV-2 lineage

classifications, for two cases to form a pair they had to share the same lineage. SARS-CoV-2

lineages circulating in the hospital were diverse enough that the minimal difference of single

nucleotide polymorphisms between them was unlikely to have occurred by mutation during a

single transmission event, suggestion multiple independent introductions (Table S4). Four

assumptions (i-iv) were used to select potential transmission pairs from all possible

permutations. The first assumption (i) stipulated that specimen collection date (symptom onset

date) of the primary case was before that of the secondary case. The second assumption (ii)

affirms that the primary case's unit of transmission equals the secondary case's unit of infection.

The third assumption (ii) holds that the collection date of the secondary case is at least one-serial

interval (5 days) from that of the primary case[37]. The fourth assumption (iv) dictates that the

collection date of the secondary case is not more than three-serial intervals (15 days) from the

primary case[38]. Potential transmission pairs, which met the four-assumptions, were stratified

by categorical *ACE2* transcription (High/Low) or viral lineage and plotted overtime.

<u>Primary Analysis:</u>

Multivariable analysis was performed to estimate the relationship between *ACE2*

transcription and number of secondary cases using a Poisson generalized linear regression

model. Variable importance was assessed conceptually using the common cause criterion and

statistically by the partial F-test. Collinearity was measured using the variable inflation factor,

variables with a value greater than five were excluded[39]. Effect modification terms were

included if they were statistically significant and supported conceptually. The two assumptions

of Poisson models; overdispersion and zero inflation were checked using a ratio of residual

deviance to degrees of freedom (theta)[40] and the score test[41], respectively. If the specified

Poisson model failed to meet either of these assumptions than an alternative quasi-Poisson or

Negative Binomial (NB) model was tested, and the assumptions re-examined. Model fit was

measured by the Akaike information criterion[42].

Sensitivity Analysis:

Sensitivity analysis was performed to measure the impact of assumptions (ii, iii and iv)

used to construct the transmission network on the relationship between *ACE2* transcription and

number of secondary cases. The assumptions were excluded independently, and the transmission

network was iteratively reconstructed with (n-1) assumptions. Additionally, the fourth

assumption (iv) was challenged by changing the infectious period from 15 days (three serial

intervals) to 10 days (two serial intervals). The number of secondary cases was computed for

each transmission network and used in multivariable analysis.

Statistical analysis was performed in R version 4.04 using the packages: readxl,

tidyverse, dataexplorer, ggpubr, car, ggsci, stringr, tableone, rio, remotes, lubridate, dplyr,

epicontacts, AER, devtools, rlang, DHARMa, MASS, pscl, epiR, EpiCurve[43].

Results:

Descriptive Statistics:

The analytic dataset contains (n = 103) cases of COVID-19 associated with a single-site

concurrent series of hospital outbreaks from 07-11-2020 to 04-01-2021 (dd-mm-yyyy) in British

Columbia, Canada. Participant characteristics are displayed in Table 1, where the variables of

interest are stratified by case description (Patients or Health Care Workers) (Table 1). The

bivariate relationship between variables and categorical *ACE2* transcription was also investigated

(Table S1). Most cases occurred in patients n = 57 (55%), while n = 48 infections were observed

in health care workers (45%). The mean age of patients associated with the hospital outbreaks

was 78.61 years, health care workers were significantly younger with a mean age of 41.11 years

(P < 0.001). Most of the healthcare workers (n = 41, 89%) and patients were biologic females (n

= 33, 58%). *ACE2* transcription and viral load did not differ between case descriptions (P = 0.86,

P = 0.30). Six SARS-CoV-2 lineages were characterized by whole genome sequencing, two of

which did not have more than one case (B.1.128 and B.1.36.38). The highest proportion of

observed SARS-CoV-2 infections were either SARS-CoV-2 lineage AL.1 (n = 56) or B.1.2 (n =

40) (Table 1), these two lineages have at least a five SNP difference between them (Table S4).

An epidemiological curve was made to show the incidence of positive testers per week

over the course of the hospital associate outbreaks. The outbreaks spanned ten epidemiological

weeks (Week-#45-2020 to Week-#1-2021) and possesses a non-normal distribution (Figure 1A

& B). We are cautious to directly interpret the incidence curve as infection prevention and

control practices implemented during the outbreaks (like point prevalence testing and stoppage

of admissions) likely biases the observed incidence of SARS-CoV-2 cases overtime. A first peak

in cases occurred on Week #46-2020 and a second on Week #52-2020. In the first phase of the

outbreaks, many cases occurred on the second floor of the hospital in units 2A, 2B and 2C. Later

transmission was predominately observed on floors four or five in units 4B, 4C, 4D and 5A

(Figure 1A). Multiple introductions of SARS-CoV-2 into the hospital environment occurred over

the surveillance period; however, timely and effective infection prevention and control measures

limited transmission of several variants like B.1.36.38, B.1.128 and B.1.279 (Table S3, Figure

S2).

<u>Transmission Network:</u>

Permuting all possible transmission pairs in which the primary and secondary case share

the same SARS-CoV-2 lineage yielded n = 9389 combinations. Applying the first assumption (i)

filtered the possible transmission pairs to n = 2167. The second assumption (ii) eliminated

another n = 1480 possible pairs for a total of n = 687. The third and fourth assumptions restricted

the possible pairs to n = 382 and n = 76, respectively. A directed transmission network was

constructed from n = 103 cases with n = 76 contacts. The network captures n= 55 transmission

events for viral lineage AL.1 and n = 39 for B.1.2; transmission of the other viral lineages was

either too low (only one case reported) or did not meet the assumption criteria (Figure 2B). The

nodes of the transmission network are shown stratified by categorical ACE2 transcription level

(Low or High) (Figure 2A). The distribution of *ACE2* transcription among hospital associated

cases was tested with a single sample proportion test, 82% of cases had above average *ACE2*

transcription (84/103, P<0.001).   Transmission pairs were further stratified into primary (n = 34)

and secondary cases (n = 28). Primary cases possessed predominately high *ACE2* transcription

(29/34, P< 0.001). *ACE2* transcription was similarly enriched in secondary cases (23/28,

P<0.01). Overall, 98.6% of transmission events involved at least one case with high *ACE2*

transcription (Table S2).

 Primary Analysis:

        Multivariable NB regression estimated that a one-unit change in *ACE2* transcription

 decreases the number of secondary cases in health care workers by -0.13 (95%CI: -0.255 to -

 0.0181) adjusting for RNA viral load and case description (Table 2). Effect modification was

 observed between *ACE2* transcription and case description; in patients, a one-unit change in

 ACE2 increased the number of secondary cases by 0.187 (95% CI: 0.0101 to 0.370). Therefore, a

 5 unit increase in *ACE2* transcription may lead to approximately one more SARS-CoV-2

 secondary case per primary patient case. Poisson and quasi-Poisson regression models were

 constructed using the same variables, neither provided a better fit to the NB option (Table 2).

 Sensitivity Analysis:

        A sensitivity analysis was performed to measure the impact of assumptions ii, iii and iv

 on construction of the transmission network and number of secondary cases. The first

 assumption- that the symptom onset date of the primary case occurs before the secondary case-

 was not included in the sensitivity analysis, as breaking this assumption would make the

 secondary case the primary case. Omission of the second assumption (ii), so that the primary

 case did not share a hospital unit with the secondary case, increased the number of possible

 transmission pairs to n = 538 (Figure S3). The network was not clearly resolved and no

 significant relationship between *ACE2* transcription and the number of secondary cases was

 estimated using a Poisson (B= -0.17 [95%CI: -0.903 to 0.558], P = 0.64) or NB regression model

 (B = -0.01 [95%CI: -0.082 to 0.0613], P = 0.8) (Figure 3). Both models had a worse fit then

 when used in the primary analysis ($Poisson_{AIC}$ = 705 and NB_{AIC} = 691). Leaving out the third

 assumption (iii), to appreciate that the secondary case could have been infected in less time than

the average serial interval, increased the number of possible transmission pairs to n = 246 (Figure

S4). A significant relationship between *ACE2* transcription and the number of secondary cases

was estimated using a Poisson (B = -0.60 [95%CI: -1.08 to -0.125], P = 0.015) and NB

regression model (B = -0.11 [95%CI: -0.215 to -0.0189], P = 0.03) (Figure 3). However, neither

model fit better then when used in the primary analysis (Poisson$_{AIC}$ = 618 and NB$_{AIC}$ = 551). Re-

analysis without the fourth assumption that the infectious period of SARS-CoV-2 does not

eclipse 15 days increased the possible transmission pairs to n = 158 (Figure S5), no significant

relationship between *ACE2* transcription and the number of secondary cases was estimated using

a Poisson (B = 0.0143 [95%CI: -0.284 to 0.313], P = 0.64) or NB regression model (B = 3.90 x

10^{-3} [95%CI: -0.0849 to 0.0911], P = 0.8) (Figure 3). Additionally, assumption four (iv) was

challenged by decreasing the infectious period to 9.5 days or ~ 2 serial intervals (Figure S6,

transmission pairs n = 48). Applying a shorter infectious period provided the following estimates

using a Poisson (B = -0.270 [95%CI: -0.454 to -0.0852], P = 0.00) and NB regression model (B

= -0.160 [95%CI: -0.324 to -0.0118], P = 0.09) (Figure 3). Importantly, the effect modification

term was significant in the NB model, and it provided a better fit than when used in the primary

analysis (AIC = 269). In patients a one-unit change in ACE2 transcription increases the number

of secondary cases by 0.268 (95% CI: 0.0315 to 0.518, P = 0.04), when adjusting for RNA viral

load (Figure 3).

Discussion:

This study describes case-to-case transmission within a series of hospital associated

SARS-CoV-2 outbreaks in British Columbia during late 2020 and early 2021. We used a

combination of epidemiological data from outbreak reports, laboratory measurements of ACE2,

and SARS-CoV-2 lineages derived from whole genome sequencing to recreate the outbreaks

using infection tracing transmission network analysis[44]. The transmission network was

resolved to provide the possible number of transmission events (secondary cases) per primary

case. Bivariate and multivariable analysis was employed to estimate the relationship between

nasopharyngeal *ACE2* transcription and the number of potential secondary cases.

Bivariate analysis found that nasopharyngeal *ACE2* transcription did not differ between

patients and health care workers. However, *ACE2* transcription was enriched in cases, more than

fifty percent of the cases had 'High' ACE2. The transmission network constructed with all four

assumptions provided n = 76 transmission events for analysis. The simplest recreation of the

outbreaks was provided by a sensitivity analysis where the fourth assumption (iv) was changed

to stipulate an infectious period of 9.5 days (~2 serial intervals), which yielded n = 48

transmission events. Multivariable analysis with Poisson regression models was inferior to NB

regression as the number of secondary cases was over dispersed. In the primary analysis, NB

regression estimated that a one-unit change in nasopharyngeal *ACE2* transcription decreases the

number of secondary cases in health care workers by B = -0.132 (95%CI: -0.255 to -0.0181, P =

0.04) and increases transmission in patients by B = 0.187 (95% CI: 0.0101 to 0.370, P = 0.04).

Sensitivity analysis using a shorter infectious period of 9.5 days, provided comparable results, no

association was found between nasopharyngeal *ACE2* transcription and SARS-CoV-2

transmission from health care workers (P = 0.09). In patients, nasopharyngeal ACE2 expression

was associated with secondary transmission B = 0.27 (95% CI: 0.0315 to 0.518, P = 0.04)

adjusting for RNA viral load (Figure 3A & B).

Viral transmission exemplifies a complex system where understanding the mechanism of

transmission requires deeper analysis than observing the sum of its components. This

observational study describes nasopharyngeal *ACE2* transcription as a component of SARS-

CoV-2 transmission. In a series of hospital associated outbreaks in British Columbia

nasopharyngeal *ACE2* transcription was positively associated with the number of secondary

cases (ongoing transmission) in patients but not health care workers. The reason for this

difference may stem from health care workers wearing personal protective equipment while

providing care. The Public Health Agency of Canada recommends that health care workers

providing care to COVID-19 positive patients, as a minimum, adhere to droplet and contact

precautions [47][45]. If care requires aerosol generating procedures, then the health care worker

dons a N95 respirator with a gown, gloves, and eye protection[45]. Surgical masks and N95

respirators provide a physical barrier between the mucosal membrane of the nasopharyngeal

passage and virus carrying droplets or particles suspended in the air. The design and intended use

of respirators and surgical masks differ. Respirators are designed to prevent inhalation of

airborne particles and must fit tightly to the user's face. Surgical masks are designed to protect

others from aerosolized droplet production from the wearer's upper respiratory system[46]. In

either case, physical barrier provided by a mask can prevent SARS-CoV-2 viral particles from

binding to ACE2 proteins on the surface of host cells and initiating the viral replication cycle.

Masking may not completely prevent exposure to SARS-CoV-2 yet still decreases the infectious

dose below the level necessary for infection[47]. Various studies have demonstrated the utility of

surgical masks and N95 respirators at preventing COVID-19 or infection with other respiratory

pathogens[47–49]. Use of surgical masks and social distancing was associated with a reduction

of 44.9 COVID-19 cases per 1000 students and staff in Boston area school districts. This

reduction represents a 4.49% decrease in the incidence of SARS-CoV-2[50]. A clinical trial

investigating the use of surgical masks or N95 respirators to prevent SARS-CoV-2 infection in

health care workers found no difference in the masks' efficacy to protect from infection[51].

This suggests that when worn by health care workers, surgical masks and N95 respirators have

comparable efficacy to reduce respiratory infections transmitted by small, aerosolized particles

<5 μm. Taken together, this evidence supports the hypothesis that health care workers in our

study were sufficiently protected by their personal protective equipment to negate the association

between nasopharyngeal *ACE2* transcription and secondary transmission. In contrast,

hospitalized patients did not wear PPE while receiving care and nasopharyngeal *ACE2*

transcription possessed a positive association with secondary transmission. The principle of a

mask protecting the wearer from causing transmission instead of protecting contacts from

acquiring infection parallels how high nasopharyngeal *ACE2* transcription in an infected person

may produce more secondary cases. High nasopharyngeal *ACE2* transcription may promote

viral shedding in the upper airway and thus the production of finer aerosols from talking,

singing, or sneezing. Fine aerosols can spread further than the large ones, produced lower in the

respiratory tract, and pose an increased risk of infection to contacts[52]. As PPE use was not

measured in our study competing hypotheses could explain the apparent difference in the

relationship between *ACE2* transcription and transmission among health care workers and

patients. An unmeasured confounding variable present for health care workers but not patients

could bias the relationship between *ACE2* transcription and secondary transmission. For

example, health care workers have variable shift lengths and leave the hospital daily after work;

therefore, their contact time with an infected patient differs from that of an inpatient who shares

the same hospital unit. This example also serves to demonstrate the complexity of measuring

SARS-CoV-2 and other respiratory pathogen's transmission. We acknowledge that SARS-CoV-

2 transmission occurs due to an intricate balance of social, biological, and environmental risk-

factors of which nasopharyngeal *ACE2* transcription may contribute a singular role[7].

Regulators of *ACE2* transcription in the lower respiratory tract are well documented a

lesser evidence base exists for the upper respiratory tract. Importantly, upper, and lower airway

ACE2 transcription are not strongly corelated indicating that they may occur independently of

each other[53]. In previous work we found that nasopharyngeal transcription of *ACE2* did not

differ by age (within adults over the age of eighteen) or biological sex[27], a finding which was

further supported by this study. High nasopharyngeal ACE2 expression has been associated with

long term inhaled corticosteroid use and exposure to fine particulate matter $PM_{2.5}$. Below

average, low nasopharyngeal *ACE2* transcription is associated with type 2 inflammation

mediated by the cytokines IL-4, IL-5, and IL-13[54,55]. Therefore, individuals with asthma or

allergic rhinitis may have lower upper airway *ACE2* expression than those that do not. Further

studies are required to understand additional predictors of upper airway ACE2 expression and if

it relates to that of the lower airway.

Strengths and limitations:

The described study has several limitations in design, data collection and analysis.

Sampling cases of SARS-CoV-2 but not their contacts prevented us from constructing a more

robust transmission network[44]. Contact tracing data would benefit our analysis by increasing

power of the study and allow estimation of the relationship between nasopharyngeal *ACE2*

transcription and lack of transmission. Estimating the relationship between nasopharyngeal

ACE2 transcription and secondary transmission in only confirmed cases may have resulted in

selection bias, where we selected for *ACE2* measurements in persons that had been exposed and

infected by SARS-CoV-2[56]. Sampling from a population of hospitalized patients may also

have introduced Berkson's bias (admission rate bias) into the study and restricts our ability to

generalize the results to other sub-groups or the community[56]. The hospitalized patients could

share an unmeasured exposure, comorbidity or drug treatment which unknowingly affected their

nasopharyngeal *ACE2* transcription. Performing a sensitivity analysis helped us understand the

importance of our assumptions in building the infection tracing transmission network. The

assumption that the primary and secondary case shared the same hospital unit, proved the most

important. Without knowing the place of exposure, the transmission network would not have

resolved transmission pairs. Whole genome sequencing provided limited specificity of

transmission events when the SARS-CoV-2 outbreak was short and genomes were classified by

lineage and not at the resolution of single nucleotide polymorphisms (SNPs). Incorporating SNPs

into the analysis may increase resolution of the transmission network by more accurately

determining transmission pairs. Using RT-qPCR to measure *ACE2* transcription and RNA viral

load overapproximates available ACE2 protein and viral particles[23]. Future work should aim

to re-estimate the association between nasopharyngeal *ACE2* transcription and secondary cases

using data from a SARS-CoV-2 household transmission study. The design of a household

transmission study has the benefits of selecting participants from a non-hospitalized population,

observing secondary infections and contacts within family clusters in a context with limited use

of personal protective equipment.

Conclusion:

We estimate the association between nasopharyngeal *ACE2* transcription and secondary

SARS-CoV-2 transmission in a series of hospital associated outbreaks in British Columbia from

late 2020 to early 2021. Analysis shows that 98.6% of transmission pairs and 85% of primary

cases had high nasopharyngeal *ACE2* transcription. Multivariable analysis adjusting for RNA

viral load and interaction by case description found that nasopharyngeal *ACE2* transcription was

positively associated with SARS-CoV-2 transmission in hospital patients but not health care

workers. We postulate that use of masks among health care workers explains this difference,

transmission from health care workers with high ACE2 was interrupted by barrier protection.

SARS-CoV-2 transmission remains a complex and unresolved phenomenon driven by biological,

environmental, and social risk factors. Differential nasopharyngeal *ACE2* transcription has been

observed in COVID-19 negative persons and having high- above average- expression of ACE2

may serve as a risk factor for transmission of SARS-CoV-2.

References:

1. Long, B.; Brady, W.J.; Koyfman, A.; Gottlieb, M. Cardiovascular Complications in COVID-19. *Am J Emerg Med* **2020**, *38*, 1504–1507, doi:10.1016/j.ajem.2020.04.048.

2. Dhar, D.; Dey, T.; Samim, M.M.; Padmanabha, H.; Chatterjee, A.; Naznin, P.; Chandra, S.R.; Mallesh, K.; Shah, R.; Siddiqui, S.; et al. Systemic Inflammatory Syndrome in COVID-19–SISCoV Study: Systematic Review and Meta-Analysis. *Pediatr Res* **2022**, *91*, 1334–1349, doi:10.1038/s41390-021-01545-z.

3. Guo, T.; Fan, Y.; Chen, M.; Wu, X.; Zhang, L.; He, T.; Wang, H.; Wan, J.; Wang, X.; Lu, Z. Cardiovascular Implications of Fatal Outcomes of Patients With Coronavirus Disease 2019 (COVID-19). *JAMA Cardiol* **2020**, *5*, 811, doi:10.1001/jamacardio.2020.1017.

4. CHAN-YEUNG, M.; XU, R.-H. SARS: Epidemiology. *Respirology* **2003**, *8*, S9–S14, doi:10.1046/j.1440-1843.2003.00518.x.

5. Zhang, A.-R.; Shi, W.-Q.; Liu, K.; Li, X.-L.; Liu, M.-J.; Zhang, W.-H.; Zhao, G.-P.; Chen, J.-J.; Zhang, X.-A.; Miao, D.; et al. Epidemiology and Evolution of Middle East Respiratory Syndrome Coronavirus, 2012–2020. *Infect Dis Poverty* **2021**, *10*, 66, doi:10.1186/s40249-021-00853-0.

6. Singanayagam, A.; Hakki, S.; Dunning, J.; Madon, K.J.; Crone, M.A.; Koycheva, A.; Derqui-Fernandez, N.; Barnett, J.L.; Whitfield, M.G.; Varro, R.; et al. Community Transmission and Viral Load Kinetics of the SARS-CoV-2 Delta (B.1.617.2) Variant in Vaccinated and Unvaccinated Individuals in the UK: A Prospective, Longitudinal, Cohort Study. *Lancet Infect Dis* **2022**, *22*, 183–195, doi:10.1016/S1473-3099(21)00648-4.

7. Leung, N.H.L. Transmissibility and Transmission of Respiratory Viruses. *Nat Rev Microbiol* **2021**, *19*, 528–545, doi:10.1038/s41579-021-00535-6.

8. Lindsey, B.B.; Villabona-Arenas, Ch.J.; Campbell, F.; Keeley, A.J.; Parker, M.D.; Shah, D.R.; Parsons, H.; Zhang, P.; Kakkar, N.; Gallis, M.; et al. Characterising Within-Hospital SARS-CoV-2 Transmission Events Using Epidemiological and Viral Genomic Data across Two Pandemic Waves. *Nat Commun* **2022**, *13*, 671, doi:10.1038/s41467-022-28291-y.

9. Madewell, Z.J.; Yang, Y.; Longini, I.M.; Halloran, M.E.; Dean, N.E. Household Transmission of SARS-CoV-2. *JAMA Netw Open* **2020**, *3*, e2031756, doi:10.1001/jamanetworkopen.2020.31756.

10. Cowger, T.L.; Murray, E.J.; Clarke, J.; Bassett, M.T.; Ojikutu, B.O.; Sánchez, S.M.; Linos, N.; Hall, K.T. Lifting Universal Masking in Schools — Covid-19 Incidence among Students and Staff. *New England Journal of Medicine* **2022**, doi:10.1056/NEJMoa2211029.

11. Cordery, R.; Reeves, L.; Zhou, J.; Rowan, A.; Watber, P.; Rosadas, C.; Crone, M.; Storch, M.; Freemont, P.; Mosscrop, L.; et al. Transmission of SARS-CoV-2 by Children to Contacts in Schools and Households: A Prospective Cohort and Environmental Sampling Study in London. *Lancet Microbe* **2022**, *3*, e814–e823, doi:10.1016/S2666-5247(22)00124-0.

12. Li, F.; Li, W.; Farzan, M.; Harrison, S.C. Structure of SARS Coronavirus Spike Receptor-Binding Domain Complexed with Receptor. *Science (1979)* **2005**, *309*, 1864–1868, doi:10.1126/science.1116480.

13. Mannar, D.; Saville, J.W.; Zhu, X.; Srivastava, S.S.; Berezuk, A.M.; Tuttle, K.S.; Marquez, A.C.; Sekirov, I.; Subramaniam, S. SARS-CoV-2 Omicron Variant: Antibody

Evasion and Cryo-EM Structure of Spike Protein–ACE2 Complex. *Science (1979)* **2022**, *375*, 760–764, doi:10.1126/science.abn7760.

14. Shang, J.; Ye, G.; Shi, K.; Wan, Y.; Luo, C.; Aihara, H.; Geng, Q.; Auerbach, A.; Li, F. Structural Basis of Receptor Recognition by SARS-CoV-2. *Nature* **2020**, *581*, 221–224, doi:10.1038/s41586-020-2179-y.

15. Meng, B.; Abdullahi, A.; Ferreira, I.A.T.M.; Goonawardane, N.; Saito, A.; Kimura, I.; Yamasoba, D.; Gerber, P.P.; Fatihi, S.; Rathore, S.; et al. Altered TMPRSS2 Usage by SARS-CoV-2 Omicron Impacts Infectivity and Fusogenicity. *Nature* **2022**, *603*, 706–714, doi:10.1038/s41586-022-04474-x.

16. Ziegler, C.G.K.; Allon, S.J.; Nyquist, S.K.; Mbano, I.M.; Miao, V.N.; Tzouanas, C.N.; Cao, Y.; Yousif, A.S.; Bals, J.; Hauser, B.M.; et al. SARS-CoV-2 Receptor ACE2 Is an Interferon-Stimulated Gene in Human Airway Epithelial Cells and Is Detected in Specific Cell Subsets across Tissues. *Cell* **2020**, *181*, 1016-1035.e19, doi:10.1016/j.cell.2020.04.035.

17. Sui, Y.; Li, J.; Venzon, D.J.; Berzofsky, J.A. SARS-CoV-2 Spike Protein Suppresses ACE2 and Type I Interferon Expression in Primary Cells From Macaque Lung Bronchoalveolar Lavage. *Front Immunol* **2021**, *12*, doi:10.3389/fimmu.2021.658428.

18. Verdecchia, P.; Cavallini, C.; Spanevello, A.; Angeli, F. The Pivotal Link between ACE2 Deficiency and SARS-CoV-2 Infection. *Eur J Intern Med* **2020**, *76*, 14–20, doi:10.1016/j.ejim.2020.04.037.

19. Groß, S.; Jahn, C.; Cushman, S.; Bär, C.; Thum, T. SARS-CoV-2 Receptor ACE2-Dependent Implications on the Cardiovascular System: From Basic Science to Clinical Implications. *J Mol Cell Cardiol* **2020**, *144*, 47–53, doi:10.1016/j.yjmcc.2020.04.031.

20. Jia, H.P.; Look, D.C.; Shi, L.; Hickey, M.; Pewe, L.; Netland, J.; Farzan, M.; Wohlford-Lenane, C.; Perlman, S.; McCray, P.B. ACE2 Receptor Expression and Severe Acute Respiratory Syndrome Coronavirus Infection Depend on Differentiation of Human Airway Epithelia. *J Virol* **2005**, *79*, 14614–14621, doi:10.1128/JVI.79.23.14614-14621.2005.

21. Ni, W.; Yang, X.; Yang, D.; Bao, J.; Li, R.; Xiao, Y.; Hou, C.; Wang, H.; Liu, J.; Yang, D.; et al. Role of Angiotensin-Converting Enzyme 2 (ACE2) in COVID-19. *Crit Care* **2020**, *24*, 422, doi:10.1186/s13054-020-03120-0.

22. Ziegler, C.G.K.; Allon, S.J.; Nyquist, S.K.; Mbano, I.M.; Miao, V.N.; Tzouanas, C.N.; Cao, Y.; Yousif, A.S.; Bals, J.; Hauser, B.M.; et al. SARS-CoV-2 Receptor ACE2 Is an Interferon-Stimulated Gene in Human Airway Epithelial Cells and Is Detected in Specific Cell Subsets across Tissues. *Cell* **2020**, *181*, 1016-1035.e19, doi:10.1016/j.cell.2020.04.035.

23. Nikiforuk, A.M.; Kuchinski, K.S.; Twa, D.D.W.; Lukac, C.D.; Sbihi, H.; Basham, C.A.; Steidl, C.; Prystajecky, N.A.; Jassem, A.N.; Krajden, M.; et al. The Contrasting Role of Nasopharyngeal Angiotensin Converting Enzyme 2 (ACE2) Transcription in SARS-CoV-2 Infection: A Cross-Sectional Study of People Tested for COVID-19 in British Columbia, Canada. *EBioMedicine* **2021**, *66*, 103316, doi:10.1016/j.ebiom.2021.103316.

24. Freed, N.E.; Vlková, M.; Faisal, M.B.; Silander, O.K. Rapid and Inexpensive Whole-Genome Sequencing of SARS-CoV-2 Using 1200 Bp Tiled Amplicons and Oxford Nanopore Rapid Barcoding. *Biol Methods Protoc* **2020**, *5*, doi:10.1093/biomethods/bpaa014.

25.   Messmann, R.; Dietl, A.; Wagner, S.; Domenig, O.; Jungbauer, C.; Luchner, A.; Maier,
L.S.; Schopka, S.; Hirt, S.; Schmid, C.; et al. Alterations of the Renin Angiotensin System
in Human End-Stage Heart Failure before and after Mechanical Cardiac Unloading by
LVAD Support. *Mol Cell Biochem* **2020**, *472*, 79–94, doi:10.1007/s11010-020-03787-7.
26.   LeBlanc, J.J.; Gubbay, J.B.; Li, Y.; Needle, R.; Arneson, S.R.; Marcino, D.; Charest, H.;
Desnoyers, G.; Dust, K.; Fattouh, R.; et al. Real-Time PCR-Based SARS-CoV-2
Detection in Canadian Laboratories. *Journal of Clinical Virology* **2020**, *128*, 104433,
doi:10.1016/j.jcv.2020.104433.
27.   Nikiforuk, A.M.; Kuchinski, K.S.; Twa, D.D.W.; Lukac, C.D.; Sbihi, H.; Basham, C.A.;
Steidl, C.; Prystajecky, N.A.; Jassem, A.N.; Krajden, M.; et al. The Contrasting Role of
Nasopharyngeal Angiotensin Converting Enzyme 2 (ACE2) Transcription in SARS-CoV-
2 Infection: A Cross-Sectional Study of People Tested for COVID-19 in British
Columbia, Canada. *EBioMedicine* **2021**, *66*, 103316, doi:10.1016/j.ebiom.2021.103316.
28.   Livak, K.J.; Schmittgen, T.D. Analysis of Relative Gene Expression Data Using Real-
Time Quantitative PCR and the 2−ΔΔCT Method. *Methods* **2001**, *25*, 402–408,
doi:10.1006/meth.2001.1262.
29.   British Columbia Centre for Disease Control Ncov2019-Artic-Nf (accessed on 22
November 2022).
30.   O'Toole, Á.; Scher, E.; Underwood, A.; Jackson, B.; Hill, V.; McCrone, J.T.; Colquhoun,
R.; Ruis, C.; Abu-Dahab, K.; Taylor, B.; et al. Assignment of Epidemiological Lineages in
an Emerging Pandemic Using the Pangolin Tool. *Virus Evol* **2021**,
doi:10.1093/ve/veab064.
31.   VanderWeele, T.J. Principles of Confounder Selection. *Eur J Epidemiol* **2019**, *34*, 211–
219, doi:10.1007/s10654-019-00494-6.
32.   Gutiérrez-Chamorro, L.; Riveira-Muñoz, E.; Barrios, C.; Palau, V.; Nevot, M.; Pedreño-
López, S.; Senserrich, J.; Massanella, M.; Clotet, B.; Cabrera, C.; et al. SARS-CoV-2
Infection Modulates ACE2 Function and Subsequent Inflammatory Responses in Swabs
and Plasma of COVID-19 Patients. *Viruses* **2021**, *13*, 1715, doi:10.3390/v13091715.
33.   Marc, A.; Kerioui, M.; Blanquart, F.; Bertrand, J.; Mitjà, O.; Corbacho-Monné, M.;
Marks, M.; Guedj, J. Quantifying the Relationship between SARS-CoV-2 Viral Load and
Infectiousness. *Elife* **2021**, *10*, doi:10.7554/eLife.69302.
34.   Carraturo, F.; del Giudice, C.; Morelli, M.; Cerullo, V.; Libralato, G.; Galdiero, E.; Guida,
M. Persistence of SARS-CoV-2 in the Environment and COVID-19 Transmission Risk
from Environmental Matrices and Surfaces. *Environmental Pollution* **2020**, *265*, 115010,
doi:10.1016/j.envpol.2020.115010.
35.   Onweni, C.L.; Zhang, Y.S.; Caulfield, T.; Hopkins, C.E.; Fairweather, D.L.; Freeman,
W.D. ACEI/ARB Therapy in COVID-19: The Double-Edged Sword of ACE2 and SARS-
CoV-2 Viral Docking. *Crit Care* **2020**, *24*, 475, doi:10.1186/s13054-020-03195-9.
36.   Kwak, S.G.; Kim, J.H. Central Limit Theorem: The Cornerstone of Modern Statistics.
Korean J Anesthesiol **2017**, *70*, 144, doi:10.4097/kjae.2017.70.2.144.
37.   Alene, M.; Yismaw, L.; Assemie, M.A.; Ketema, D.B.; Gietaneh, W.; Birhan, T.Y. Serial
Interval and Incubation Period of COVID-19: A Systematic Review and Meta-Analysis.
BMC Infect Dis **2021**, *21*, 257, doi:10.1186/s12879-021-05950-x.
38.   Byrne, A.W.; McEvoy, D.; Collins, A.B.; Hunt, K.; Casey, M.; Barber, A.; Butler, F.;
Griffin, J.; Lane, E.A.; McAloon, C.; et al. Inferred Duration of Infectious Period of
SARS-CoV-2: Rapid Scoping Review and Analysis of Available Evidence for

614   Asymptomatic and Symptomatic COVID-19 Cases. *BMJ Open* **2020**, *10*, e039856,
615   doi:10.1136/bmjopen-2020-039856.

616 39. Craney, T.A.; Surles, J.G. Model-Dependent Variance Inflation Factor Cutoff Values.
617   *Qual Eng* **2002**, *14*, 391–403, doi:10.1081/QEN-120001878.

618 40. Cameron, A.C.; Trivedi, P.K. Regression-Based Tests for Overdispersion in the Poisson
619   Model. *J Econom* **1990**, *46*, 347–364, doi:10.1016/0304-4076(90)90014-K.

620 41. van den Broek, J. A Score Test for Zero Inflation in a Poisson Distribution. *Biometrics*
621   **1995**, *51*, 738–743.

622 42. Portet, S. A Primer on Model Selection Using the Akaike Information Criterion. *Infect Dis*
623   *Model* **2020**, *5*, 111–128, doi:10.1016/j.idm.2019.12.010.

624 43. R Core Team R: A Language and Environment for Statistical Computing. 2013.

625 44. Keeling, M.J.; Eames, K.T.D. Networks and Epidemic Models. *J R Soc Interface* **2005**, *2*,
626   295–307, doi:10.1098/rsif.2005.0051.

627 45. Public Health Agency of Canada Infection Prevention and Control for COVID-19: Interim
628   Guidance for Acute Healthcare Settings.

629 46. Vincent, M.; Edwards, P. Disposable Surgical Face Masks for Preventing Surgical Wound
630   Infection in Clean Surgery. *Cochrane Database of Systematic Reviews* **2016**, *2016*,
631   doi:10.1002/14651858.CD002929.pub3.

632 47. Leung, N.H.L.; Chu, D.K.W.; Shiu, E.Y.C.; Chan, K.-H.; McDevitt, J.J.; Hau, B.J.P.;
633   Yen, H.-L.; Li, Y.; Ip, D.K.M.; Peiris, J.S.M.; et al. Respiratory Virus Shedding in
634   Exhaled Breath and Efficacy of Face Masks. *Nat Med* **2020**, *26*, 676–680,
635   doi:10.1038/s41591-020-0843-2.

636 48. Andrejko, K.L.; Pry, J.; Myers, J.F.; Openshaw, J.; Watt, J.; Birkett, N.; DeGuzman, J.L.;
637   Barbaduomo, C.M.; Dong, Z.N.; Fang, A.T.; et al. Predictors of Severe Acute Respiratory
638   Syndrome Coronavirus 2 Infection Following High-Risk Exposure. *Clinical Infectious*
639   *Diseases* **2022**, *75*, e276–e288, doi:10.1093/cid/ciab1040.

640 49. Fennelly, K.P. Particle Sizes of Infectious Aerosols: Implications for Infection Control.
641   *Lancet Respir Med* **2020**, *8*, 914–924, doi:10.1016/S2213-2600(20)30323-4.

642 50. Cowger, T.L.; Murray, E.J.; Clarke, J.; Bassett, M.T.; Ojikutu, B.O.; Sánchez, S.M.;
643   Linos, N.; Hall, K.T. Lifting Universal Masking in Schools — Covid-19 Incidence among
644   Students and Staff. *New England Journal of Medicine* **2022**, *387*, 1935–1946,
645   doi:10.1056/NEJMoa2211029.

646 51. Loeb, M.; Bartholomew, A.; Hashmi, M.; Tarhuni, W.; Hassany, M.; Youngster, I.;
647   Somayaji, R.; Larios, O.; Kim, J.; Missaghi, B.; et al. Medical Masks Versus N95
648   Respirators for Preventing COVID-19 Among Health Care Workers. *Ann Intern Med*
649   **2022**, *175*, 1629–1638, doi:10.7326/M22-1966.

650 52. Wang, C.C.; Prather, K.A.; Sznitman, J.; Jimenez, J.L.; Lakdawala, S.S.; Tufekci, Z.;
651   Marr, L.C. Airborne Transmission of Respiratory Viruses. *Science (1979)* **2021**, *373*,
652   doi:10.1126/science.abd9149.

653 53. Hou, Y.J.; Okuda, K.; Edwards, C.E.; Martinez, D.R.; Asakura, T.; Dinnon, K.H.; Kato,
654   T.; Lee, R.E.; Yount, B.L.; Mascenik, T.M.; et al. SARS-CoV-2 Reverse Genetics Reveals
655   a Variable Infection Gradient in the Respiratory Tract. *Cell* **2020**, *182*, 429-446.e14,
656   doi:10.1016/j.cell.2020.05.042.

657 54. Kimura, H.; Francisco, D.; Conway, M.; Martinez, F.D.; Vercelli, D.; Polverino, F.;
658   Billheimer, D.; Kraft, M. Type 2 Inflammation Modulates ACE2 and TMPRSS2 in

659   Airway Epithelial Cells. *Journal of Allergy and Clinical Immunology* **2020**, *146*, 80-88.e8, doi:10.1016/j.jaci.2020.05.004.

661 55.   Takabayashi, T.; Yoshida, K.; Imoto, Y.; Schleimer, R.P.; Fujieda, S. Regulation of the Expression of SARS-CoV-2 Receptor Angiotensin-Converting Enzyme 2 in Nasal Mucosa. *Am J Rhinol Allergy* **2022**, *36*, 115–122, doi:10.1177/19458924211027798.

664 56.   Westreich, D. Berkson's Bias, Selection Bias, and Missing Data. *Epidemiology* **2012**, *23*, 159–164, doi:10.1097/EDE.0b013e31823b6296.

Ethical Approval:

Ethical approval for the study was sought from the University of British Columbia human ethics board (#H20-01110) which was harmonized with the Fraser Health Authority. Written informed consent was not required. All data was de-identified prior to analysis and the results were not linked back to any identifying records. This study was deemed as minimal risk to the participants involved. To ensure privacy the site of the outbreak series will not be disclosed.

Funding:

This work was funded by the Canadian Institutes of Health Research (#434951) and Genome British Columbia (COV-55).

Competing Interests:

The authors have no conflicts of interest to declare.

Author Contributions:

A.M.N., K.S.K., K.S. and I.S. conceived and designed the study. Epidemiological reporting, data recording and stewardship was performed by K.S. and S.R. Laboratory data reporting and stewardship was performed by N.P. and A.N.J. Laboratory measurements were taken by A.M.N and K.S.K. Data analysis was performed by A.M.N and K.S.K., D.M.P and

688   M.A.I assisted in interpretation of the results. The manuscript was written by A.M.N, all authors

689   helped with editing and provided their approval for publication.

730 **Table 1:** Characteristics of analytic data stratified by case designation (n = 103).

Variable	Level	Health Care Worker n = 46	Patient n = 57	P-Value
Age (mean (SD))		41.1 (10.3)	78.6 (12.1)	<0.001
Biological Sex (%)	Female	41.0 (89.1)	33.0 (57.9)	0.001
	Male	5.00 (10.9)	24.0 (42.1)	
ACE2 Transcription (mean (SD))		2.11 (2.88)	2.00 (3.06)	0.861
RNA Viral Load (mean (SD))		6.94 (1.79)	7.33 (1.98)	0.300
Viral Lineage (%)				0.385
	AL.1	28.0 (60.9)	27.0 (47.4)	
	B.1.128	0.00 (0.00)	1.00 (1.80)	
	B.1.2	14.0 (30.4)	25 (43.9)	
	B.1.279	0.00 (0.00)	2.00 (3.50)	
	B.1.36	2.00 (4.30)	1.00 (1.80)	
	B.1.36.36	1.00 (2.20)	0.00 (0.00)	
	B.1.36.38	1.00 (2.20)	0.00 (0.00)	

731 Variables included in the study are stratified by case designation health care worker or patient. P-values for
732 parametric statistical test performed on continuous (t-test) or categorical data (X^2 test) are reported. Viral lineage
733 calls are from whole genome sequencing data classified using the PANGOLIN tool (Version 1.15.1). Data is also
734 available stratified by 'High' or 'Low' nasopharyngeal ACE2 transcription (Table S1).

744   **Table 2:** Multivariable analysis of the relationship between ACE2 transcription and the number
745   of SARS-CoV-2 secondary cases.
746

Multivariable Regression Models
*(Secondary Cases ~ ACE2 Transcription + RNA Viral Load + Case
Description + ACE2 Transcription : Case Description)*

Beta Coefficients *(95% CI)*	Poisson	Quasi-Poisson	Negative Binomial (NB)
ACE2 Transcription in HCW	-0.164 *(-0.236- [-0.0919])*	-0.164 *(-0.273-[-0.0528])*	-0.132 *(-0.255- [-0.0181])*
RNA Viral Load	$2.73*10^{-3}$ *($5.34*10^{-3}$ – 0.0976)*	$2.73*10^{-3}$ *(-0.141-0.150)*	-0.0197 *(-0.164- [0.123])*
Case (Patient)	-0.808 *(-1.20- [-0.421])*	-0.808 *(-1.43- [-0.235])*	-0.782 *(-1.42- [-0.162])*
ACE2 Transcription in Patients	0.208 *(0.0958-0.320)*	0.208 *(0.0382-0.382)*	0.187 *(0.01011-0.370)*
AIC	394	--	346
Theta	2.44	2.44	0.922
Score Test (P Value)	$P < 0.001$	--	$P = 0.976$

748
749
750
Figure 1: Incidence curve of several hospital associated SARS-CoV-2 outbreaks in British Columbia by
epidemiological week from 07-11-2020 to 04-01-2021 (n = 103 laboratory confirmed cases). SARS-CoV-2
incidence follows a non-normal distribution where an initial peak caseload was observed in W46-2020 and a second
lesser case surge occurred in W52-2020. **A**) SARS-CoV-2 cases are stratified by hospital floor, early in the
surveillance period transmission occurred mostly on floor number two and later moved to floors four and five. If the
floor where transmission occurred was not determined by the investigating epidemiologist, then it was coded as
'unknown' for our analysis. **B**) SARS-CoV-2 cases are stratified by viral variant, n = 7 unique SARS-CoV-2
variants were identified by whole genome sequencing participants diagnostic specimens. SARS-CoV-2 variants
AL.1 and B.1.2 caused the highest percentage of cases: AL.1, 55/103, 53% and B.1.2, 39/103, 38%.
760
761

762
Figure 2: Infection tracing transmission network analysis of a series of hospital associated SARS-CoV-2
outbreaks in British Columbia by epidemiological week from 07-11-2020 to 04-01-2021 (n = 103 laboratory
confirmed cases). Nodes represent confirmed cases while the edge show the direction of outgoing transmission, link
a primary case with at least one secondary case. **A)** Nodes in the transmission network are stratified by ACE2
transcription (high or low), 82% of cases (84/103) had above average, high ACE2 transcription (P < 0.001). **B)**
Nodes in the transmission network are classified by SARS-CoV-2 viral variant. The viral variants AL.1 and B.1.2
caused multiple secondary infections thorough the outbreaks, in the first phase (before Dec.15[th], 2020) of the study
period most cases were AL.1 and later transitioned to B.1.2 in phase two (after Dec.15[th], 2020).
771

Figure 3: Forest plot of point estimates from the association between nasopharyngeal *ACE2* transcription of the primary case and the SARS-CoV-2 secondary cases. Poisson and Negative Binomial (NB) regression models were used for the primary analysis and sensitivity analysis of the assumptions used to construct the infection tracing networks. **A)** Beta-coefficients and ninety-five percent confidence intervals for the association between nasopharyngeal *ACE2* transcription and secondary cases within health care workers, adjusting for viral load. The best fit NB model shows no association in the primary analysis or sensitivity analysis (ii, iv or v). A significant association between nasopharyngeal *ACE2* transcription and the number of SARS-CoV-2 secondary cases was found for healthcare workers in sensitivity analysis iii, where the serial interval of 5.6 days was not used to build the transmission network. **B)** Beta-coefficients and ninety-five percent confidence intervals for the association between nasopharyngeal *ACE2* transcription and secondary cases within patients, adjusting for viral load. The best fit NB model finds a positive association between in the primary analysis, and sensitivity analysis iii, and v. No association between nasopharyngeal *ACE2* transcription and the number of SARS-CoV-2 secondary cases was found for patients in sensitivity analysis ii and iv, where assumptions of transmission occurring within hospital units and a 15.6-day infectious period were excluded from constructing the transmission network, respectively.