

## ***Research Article***

### **Validation of the Brazilian version of the Processes of Change Questionnaire in Weight Management in adults with overweight and obesity in Brazil**

Quênia de Carvalho<sup>a,b</sup>; Paola Rampelotto Ziani<sup>a,b</sup>; Bruno Braga Montezano<sup>b,c</sup>; Jeferson Ferraz Goularte<sup>b,c</sup>; Adriane R Rosa<sup>a,b,c</sup>

<sup>a</sup> Graduation Program in Pharmacology and Therapeutics: Institute of Basic Health Sciences. University Federal of Rio Grande do Sul (UFRGS), Porto Alegre, Brazil.

<sup>b</sup> Laboratory of Molecular Psychiatry, Experimental Research Center, Clinical Hospital of Porto Alegre (HCPA), University Federal of Rio Grande do Sul (UFRGS), Porto Alegre, Brazil.

<sup>c</sup> Graduation Program in Psychiatry and Behavioral Sciences: Medical School. University Federal of Rio Grande do Sul (UFRGS), Porto Alegre, Brazil.

Short Title: **Validation of the Brazilian version of the Processes of Change**

Corresponding Author: Dra. Adriane Ribeiro Rosa

Full name - Dra. Adriane Ribeiro Rosa

Department - Laboratory of Molecular Psychiatry, Hospital das Clínicas de Porto Alegre (HCPA), University Federal of Rio Grande do Sul.

Institute/University/Hospital - Hospital das Clínicas de Porto Alegre (HCPA), University Federal of Rio Grande do Sul.

Street Name & Number - Avenida Ramiro Barcelos, 2350

City, State, Postal code, Country – Porto Alegre, RS, Postal Code 90035-903, Brazil.

Tel: +55 (51) 3359.6328

E-mail: [adrianerrosa@gmail.com](mailto:adrianerrosa@gmail.com)

Number of Tables: 7

Number of Figures: 3

Word count: 5761

Keywords: P-Weight, S-Weight, Transtheoretical model, psychometry, weight loss

## 1 **Abstract**

2 **Introduction:** This study aimed to analyze the psychometric properties of the Brazilian version of  
3 the Processes of Change Questionnaire in Weight Management (P-Weight) questionnaire.

4 **Methods:** A total of 656 adults participated in the study, including people in weight loss treatment  
5 and people from the general community. All participants responded to the Process of Change  
6 Questionnaire (P-Weight), the Stages of Change Questionnaire (S-Weight), and the Eating Attitudes  
7 Test (EAT-26), which assesses risk of eating disorder used as a measure of external validity. Socio-  
8 demographic variables were also investigated.

9 **Results:** The 33-item P-weight questionnaire showed satisfactory psychometric properties with high  
10 internal consistency (Cronbach's  $\alpha=0.959$ ). Exploratory and confirmatory factor analyses  
11 revealed a 4-factor model similar to the original Spanish version of P-Weight with a slightly  
12 rearrangement of the items ( $KMO=0.92$ ,  $df(528, n=328) = 8,401.015$ ;  $p<0.0001$ ). We found significant  
13 associations between processes and stages of change ( $p<0.001$ ) and a moderate correlation between  
14 the four subscales of P-weight and EAT-26 ( $p<0.001$ ). Finally, the mean score of P-Weight were  
15 higher in clinical sample compared to the general community, suggesting the sensitivity to  
16 discriminate cases and controls ( $p<0.001$ ).

17 **Conclusion:** This study showed the validity and reliability of the Brazilian version of the P-Weight  
18 scale. Therefore, the P-Weight is readily available to help professionals to employ precision  
19 interventions to weight loss considering the patient's motivational stage in combination with their  
20 individual use of the cognitive processes of change.

## 21 **Introduction**

22 In Brazil, the percentage of adult people who living with overweight is 57.5% for men and 62.6% for  
23 women [1], which suggests that obesity is a public health problem. The long-term consequences of  
24 obesity have revealed increased risk for other primary lifestyle diseases, including coronary heart  
25 disease, hypertension, stroke, type 2 diabetes mellitus, sleep apnea, osteoarthritis, prostate disease,  
26 autoimmune conditions, neuropsychiatric disorders, and others [2]. Current approaches emphasize  
27 that lifestyle changes with the adoption of low-energy dietary patterns are effective in reducing body  
28 weight and may prevent obesity-associated chronic diseases [3]. However, most people living with  
29 obesity do not stay on weight loss programs long enough and only a few of them lose and maintain  
30 the 'new' weight [4]. Thus, a study ( $n=50,081$ ) showed that 95% of people who try to lose weight,  
31 regain it in a period of five years after a dietary, pharmacological, or behavioural intervention; the  
32 faster the weight loss, the faster the weight recovery; with this behaviour more evident in cases than  
33 control groups [5]. Therefore, the long-term success of weight loss programs depends on the ability  
34 of individuals living with obesity to change their behaviours, which is extremely difficult for most  
35 people, especially regarding to maintain a new diet pattern and to exercise regularly [6].

36 Considering behaviour as the key to weight loss, The Transtheoretical Model (TTM) proposed by  
37 James Prochaska in 1979, has been used in many fields of Medicine, including obesity [7]. TTM is  
38 based on behaviour change considering the individual's readiness to change and the process of  
39 change [8]. According to the TTM, stages of change represent the temporal, motivational, and  
40 constancy aspects of change that focus on 'when' people change [4], indicating that the individual  
41 moves through these stages when modifying their behaviour. The stages of change can be modelled  
42 in five stages: 1) Pre-contemplative (PC), individual does not want to make any change; 2)  
43 Contemplative (C), individual is ambivalent about change; 3) Preparation (P), individual begins to plan  
44 and commit to change; 4) Action (A), individual practices actions aimed at change; and 5)  
45 Maintenance (M), individual works to maintain and sustain change for a long time [9]. On the other  
46 hand, the processes of change refer to "how" people change and consist of activities in which  
47 individuals engage when trying to modify complex behaviours [4]. Although the TTM has been  
48 initially applied to smoking cessation attempts [10], it has been applied against obesity with relative  
49 success [11]. Furthermore, processes of change have been identified as predictors of behaviour  
50 change in interventions aimed at health promotion, such as quitting smoking, healthy eating  
51 behaviour, physical exercises, and weight control [4].

52 The identification of stages and processes of change is critical to individualize interventions and may  
53 help to increase the treatment effectiveness. For that, specific questionnaires to assess stages and  
54 processes of change in weight management have been proposed by the scientific community [12].  
55 The Stages of Change Questionnaire (S-Weight) and Process of change questionnaire in weight  
56 management (P-Weight) seem to be effective to assess the readiness for change and the processes  
57 faced during change in patients who are controlling their weight [13]. Both instruments were  
58 validated in populations of Spain [4] and United Kingdom (UK) [14] with acceptable psychometric  
59 properties, which allow their use to implement behaviour change interventions for weight loss. When  
60 applied in individuals living with obesity on the waiting list for bariatric surgery, P-weight and S-  
61 weight showed they were in the Action and Maintenance stages for change. There was also  
62 relationship between stages for change and the body mass index (BMI), since the BMI is lower in  
63 individuals who are in the Action and Maintenance stages, compared to those who reported being in  
64 the Pre-Contemplation stage for change [6]. Therefore, the identification of these stages and  
65 processes of change may contribute to weight loss and increased postoperative success rates.

66 Using the P-Weight and S-Weight scales, A Thai randomized clinical trial that aimed at determining  
67 the effects of individualized nutrition counselling, combined with a TTM model on weight loss in  
68 people with overweight and obese found that individuals who received the combined nutritional

69 counselling with TTM experienced greater weight loss than the control group, which received only an  
70 educational handbook [15]. Another recent study evaluated the stages and processes of change for  
71 weight loss in patients living with obesity who had a recent stroke. It showed that patients who are  
72 emotionally invested in weight loss and take action regarding their weight are more likely to be in an  
73 active phase of behaviour change for weight loss. Although studies using the P-Weight and S-Weight  
74 scales have shown promising results, a Cochrane review reported that the effectiveness of TTM in  
75 promoting weight loss is still unclear due to the small sample and heterogeneity in the  
76 methodologies used, making it difficult to draw effective conclusions [10].

77 Thus, considering the high prevalence of overweight and obesity in Brazil and the absence of  
78 validated scales capable to assess stages and processes of changes in individuals with overweight and  
79 obesity, the aim of this case-control study was: 1) to analyze the psychometric properties of the  
80 transculturally adapted Brazilian version of the P-Weight; 2) to assess the relationship between  
81 stages and process of change; and 3) to assess the relationship between P-Weight and external  
82 measures regarding eating disorders such as EAT.

## 83 **Materials and Methods**

### 84 *Translation, transcultural adaptation, and content validity*

85 Initially, the P-Weight scale was independently translated from Spanish to Portuguese by two  
86 bilingual translators (QC and ARR) and then the back translation was done by a native speaker of  
87 Brazilian Portuguese (DB) that was unaware of the study objectives. Then, the adaptation of  
88 semantic and idiomatic equivalences was carried out by the responsible researchers (QC, JG and  
89 ARR). In the next step, the instruments were administered to the ten specialists (clinical nutritionists)  
90 to detect possible divergences in the understanding of the item by Brazilians. In addition, each item  
91 was evaluated regarding cultural equivalence. All considerations made by specialists were reviewed  
92 by our research team. Then, a final Brazilian Portuguese version was obtained and applied to a  
93 sample of ten patients who were undergoing weight loss treatment to verify its applicability.

94 The P-Weight is composed of 34 items to measure the processes of behaviour change. For each  
95 question, by a 5-point Likert scale, the person scores according to their level of agreement for the  
96 question. The P-Weight suggests that four processes of change are involved in this setting: i)  
97 emotional re-evaluation (EmR; emotional reactions to being overweight and what will happen if they  
98 engage in weight management actions); ii) weight consequences evaluation (WCE; the individual's  
99 awareness of consequences that overweight has on their life, becoming aware that actually they

100 have a weight problem); iii) weight management actions (WMA; those specific actions that people  
101 engage when trying to manage their weight); iv) environmental restructuring (EnR; actions aimed to  
102 modify the individual's environment to promote weight management) [16]. The scores for each of  
103 the four change processes are calculated by adding the scores obtained on the items belonging to  
104 the same subscale. None of the items are reverse scored. To make the scores of the different  
105 subscales comparable, these scores are transformed into a scale from 0 to 100, with 0 being the non-  
106 use of a given process of change, and 100 being the full use of such process [16]. The highest use of a  
107 process is represented by scores above 50, whereas the lowest use of a process, by scores below 50  
108 [15].

109 *Sample, inclusions criteria, and recruitment:*

110 A cross-sectional observational study was carried out using an anonymous online questionnaire. The  
111 sample was non-probabilistic by convenience (n=656), composed of people from the general  
112 community (community sample) and participants in weight loss programs at the Hospital das Clínicas  
113 de Porto Alegre (HCPA) and private clinics in Porto Alegre (clinical sample). The inclusion criteria for  
114 the community sample were individuals aged over 18 years and BMI above 18.5 kg/m<sup>2</sup>; and the  
115 inclusion criteria for the clinical sample were individuals aged over 18 years, BMI above 18.5 kg/m<sup>2</sup>,  
116 and participation in nutritional or pharmacological treatment for weight loss or waiting for a bariatric  
117 surgery. In the clinical sample, pregnant and lactating women were excluded.

118 The community sample was recruited via an online questionnaire available on social networks, via an  
119 access link on the REDCAP platform; and the clinical sample was collected at the HCPA Nutrition and  
120 Bariatric Surgery outpatient clinics and private weight loss treatment clinics. In the clinical sample,  
121 the research team contacted patients, inviting them to participate in the research, and an individual  
122 and exclusive link was generated on the platform for each invitee, which was made available only  
123 when the person agreed to participate. To minimize selection bias and to ensure that each  
124 participant answered the questionnaire only once—to avoid duplicates—a question was inserted at  
125 the end of the consent form: “- Have you answered this questionnaire before?” When answering yes,  
126 the questionnaire was automatically excluded. At the end of the survey, all participants who wished  
127 to, received, as a form of thanks, an online material on Mindful Eating. The study followed the  
128 conditions established in Resolution 466/12, of the Brazilian National Health Council (CNS) and was  
129 approved by the HCPA Research Ethics Committee (CAAE: 54352521.8.0000.5327 N. 2021.0447).  
130 Before entering, the survey participants were asked to confirm their eligibility and to sign an  
131 informed consent form.

132 To calculate the sample size, a P-Weight questionnaire, with 34 items was considered, with a 0.05  
133 margin of error, a 95% confidence level, and a 0.85 expected Cronbach's alpha [16]. Thus, the total  
134 sample was initially estimated at 640 individuals, 320 for the clinical sample and 320 for the  
135 community sample.

#### 136 *Other Instruments*

137 *S-Weight*: The S-Weight consists of five mutually exclusive items, among which participants had to  
138 choose in order to be allocated to one of the five stages of change (Pre-contemplative (PC)  
139 Contemplative (C) Preparation (P) Action (A) Maintenance (M), as proposed by the TTM [17].

140 *Eating attitudes test-26 - (EAT-26)*: This scale is used to screen for Anorexia (AN) and Bulimia (BN)  
141 (Freitas et al., 2002). In Brazil, the EAT-26 was validated in female adolescents with satisfactory  
142 psychometric properties [18]. The EAT was used in the UK and Spanish study in adults with obesity  
143 [16].

#### 144 *Data Analysis*

145 Descriptive analyses were performed using the PAWS Statistics 18 software, whereas the R software  
146 version 4.1.3 was used for exploratory (EFA) and confirmatory (CFA) factor analyses. For the EFA and  
147 CFA analyses, the total sample (n=656) and participants from the community and clinic samples were  
148 all grouped into the same group and then randomized to obtain two comparable subgroups  
149 regarding age (p=0.6) and gender (p=0.81). Following the cross-validation method used in the original  
150 articles [4,16], the Subgroup 1 (n=328) was used to perform EFA and the Subgroup 2 (n=328) for CFA.  
151 A cross-validation is desirable for EFA and CFA when the sample is large enough to randomly  
152 distribute participants into at least two groups, and it is potentially useful to perform EFA with half  
153 the sample and then use CFA to validate the factor structure with the other half [19]. The Oblimin  
154 rotation with eigenvalues >1 with extraction of four factors, and degree of item extraction >0.3 [4]  
155 was applied to the EFA for the Subgroup 1. Then, the CFA was used to the other half of the sample  
156 (Subgroup 2) to confirm the internal structure of the questionnaire. Third, internal consistency was  
157 performed for both subgroups using Cronbach's alpha. Analysis of variance (ANOVA) was applied to  
158 the total sample (n=656) to assess the relationship between processes (dependent variables) and  
159 stages of change (independent variables) to analyze the trend in the use of processes of change.  
160 Spearman's correlations were used to assess the convergent external validity between the P-Weight  
161 and the EAT-26.

## 162 **Results**

### 163 *Participants*

164 A total of 1,018 individuals were initially invited, of which only 656 correctly completed the  
165 questionnaire; so, the participation rate was 64.44%. Most participants (70%) were from the city of  
166 Porto Alegre. The mean age was 41.53 (SD=11.28) with BMI of 30.8 kg/m<sup>2</sup> (SD=7.10). As shown in  
167 Table 1 other sample characteristics.

### 168 *Internal Structure*

169 EFA was applied to Subgroup 1 (n=328). The Kaiser–Meyer–Olkin test indicated the sample adequacy  
170 for the analysis (KMO=0.92, df (528, n=328) =8,401.015; p<0.0001) and the Bartlett’s test showed a  
171 significant value with X<sup>2</sup> (528) =11,809.46 and p<0.001, indicating that the correlations among the  
172 items are sufficient to perform the analysis (polycor () function of the package psych of the R  
173 programming language version 4.1.3 was used). The communalities ranged from 0.412 to 0.914,  
174 indicating a good proportion of shared variance among the analyzed items. The percentage of  
175 variance explained with four factors was 73%. The factor loadings of these items were acceptable as  
176 they reached values above 0.30, ranging from 0.374 to 1.007, as shown in Table 2.

177 The content analysis revealed that the items belonging to the same construct presented a slight  
178 difference in relation to the Spanish version regarding the items arrangement. The four factors  
179 revealed at EFA were labelled according to their specific content, following the same idea of the  
180 original nomenclature given by Andrés 2011: 1) Emotional Re-evaluation (RE) with 11 items; 2)  
181 Actions for weight control (ACP) with 12 items; 3) Consequences of excess weight (CEP) with five  
182 items; and 4) Environmental Restructuring (RA) with five items. One item (“*I think I should eat food*  
183 *with less fat*”) was excluded since the degree of extraction was less than 0.3 and the communality  
184 was less than 0.2. Therefore, the final Brazilian version of P-weight consists of 33 items. Figure 1  
185 shows the diagram representing the four process factors of change in weight management

186 CFA was performed with Subgroup 2 (n=328), excluding data from Subgroup 1 for this analysis. The  
187 measurement model consisted of four factors loaded according to the results found in the EFA,  
188 showing acceptable fit of the 33 items. As shown in Table 3 the CFA structure with unstandardized  
189 parameters. The complete table is in the Supplementary Material (library function, lavaan package, R  
190 program version 4.3.1).

191 The Comparative Fit Index and Tucker-Lewis Index indices calculate the relative fit of the observed  
192 model when comparing it with a base model, i.e., it measures a relative improvement of the study

193 model in relation to the standard model. The values equal to or greater than 0.90 indicate adequate  
194 adjustment, even in small samples [20]. The root mean square error of approximation is the  
195 difference index between the covariance matrix observed by degrees of freedom and the  
196 hypothetical matrix, indicating a satisfactory fit with values from 0.05 to 0.08. The standardized root  
197 mean square residual is the average of the standardized residuals between the hypothetical  
198 covariance matrices and the observed covariance; values lower than 0.10 indicates that the model is  
199 accepted [21]. As shown in Figure 1 the diagram representing the four process factors of change in  
200 weight management.

#### 201 *Internal Consistency:*

202 Internal consistency was assessed for the four subdomains of P-weight derived from internal  
203 structure analysis and for the total scale in both Subgroups 1 (sample 1) and 2 (sample 2). As shown  
204 in Table 4, Cronbach's alpha was satisfactory. Corrected item-total correlations were adequate for all  
205 items since all of them reached values over 0,30.

#### 206 *Relationship between Processes and Stages of Change*

207 As shown in Figure 2 and Table 5, ANOVA revealed significant differences in processes of change  
208 scores across the five stages of change. These differences are illustrated in the very strong linear  
209 association, with P-weight subscales scores increasing from PC to A stages of changes, except for the  
210 "consequences of excess weight (CEP)" that reached the peak in P stage. As expected, in the  
211 maintenance stage (M), the P-weight scores were lower than other stages, suggesting that the use of  
212 processes of change stabilizes or even decreases during the maintenance stage of weight  
213 management. Furthermore, in the Pre-Contemplation phase, all processes are used less than the  
214 Action stage.

215 Regarding the S-Weight, most participants (223) were in Action stage, which suggests that most of  
216 them were making efforts to manage their weight. From these, 155 participants (23.6%) were from  
217 the clinical group, whereas 68 (10.4%) participants were from the community group.



218 *Relationship of processes of change with other variables*

219 As shown in Table 6, the subdomains RE, RA, ACP, and CEP of the P-Weight were significantly  
220 correlated with EAT-26, suggesting the convergent validity of the P-weight ( $p < 0,0001$ ).

221 *Sensitivity*

222 Mean differences between clinical and community samples were calculated to assess whether the P-  
223 Weight and S-Weight were sensitive enough to detect differences between these groups. Results  
224 showed that statistically significant differences were obtained in all subscales, with participants in  
225 the clinical sample obtaining higher scores. Finally, participants from the clinical sample were  
226 allocated at more advanced stages of change for weight management. Table 7 and Figure 3 shows it.

227 **Discussion**

228 The Brazilian version of the P-Weight showed satisfactory psychometric properties with very high  
229 internal consistency and a positive relationship between processes and stages of change (S-Weight)  
230 and convergent validity, as indicated by a significant correlation with the EAT-26. As expected, the P-  
231 Weight mean score was higher in the clinical sample compared to the community sample, suggesting  
232 the discriminative validity of the instrument. Furthermore, the P-Weight resulted in a four-  
233 dimensional model, similar to the original Spanish version [4,16], that is, this instrument is fit to  
234 measure attitudes and behaviors involved in weight control. Therefore, the Brazilian version of the P-  
235 Weight is readily available for its use in both research and clinical practice.

236 Regarding internal structure, the subscales and the total scale showed adequate results, with a  
237 Cronbach's alpha of 0.959 for the total sample, characterizing the reliability of our instrument. These  
238 results corroborate the findings in the Spanish and the UK validation studies, which showed  
239 Cronbach's alpha of 0.960 [4] and values ranging from 0.849 to 0.955, respectively [16]. The Brazilian  
240 version of the P-Weight has 33 items, which has one item less than the original version. The item "I  
241 think I should eat food with less fat" was excluded since it has a 0.081 factor loading (less than 0.3),  
242 as recommended by Floyd and Widaman [22]. We believe that this happened due to different eating  
243 patterns between Europe and Latin America. In Brazil, people present a high consumption of fast-  
244 assimilated carbohydrates and low consumption of fatty foods, although carbohydrates- and sugar-  
245 rich foods are often also fatty foods. However, it is possible that respondents have not been aware  
246 about this technical information, which leads to a misunderstanding of this question.

247 Factorial analysis identified and confirmed that a model of four correlated processes of change was  
248 the most appropriate to explain behavior change in weight management. The four following  
249 subscales of the P-Weight obtained were: *Emotional re-evaluation (RE)*, *Action for weight control*  
250 *(ACP)*, *Consequences of excess weight (CEP)*, and *Environmental restructuring (RA)*. Compared with  
251 the original Spanish version, the number of items in each domain was similar with some exchange of  
252 items between domains, which was also observed in the UK validation study [16]. For example, the  
253 question “I try to put food away to avoid nibbling” in the original Spanish version is in the  
254 Environmental restructuring (RA) domain and in the UK and Brazilian versions was rearranged to the  
255 Actions for weight control (ACP) domain. This is justified because it is a specific Action item, which is  
256 taken by the individual to lose weight and, at the same time, it is related to the environment in which  
257 the individual is inserted. Likewise, the question “I now realize I have a weight problem,” which in the  
258 Spanish and the UK versions is included in the Consequences of excess weight (CEP) domain, in the  
259 Brazilian version, was rearranged to the Emotional Reevaluation (RE) domain; this can be explained  
260 because besides its representation as a consequence, this item also represents the individual’s  
261 feeling when they understand that they have a weight problem. Therefore, an item can either be in  
262 the RE or RA domain or in the CEP or ACP domain since they are closely related and focus on the  
263 actions that individuals take when trying to lose weight, how they feel in this context, how the  
264 environment influences in this process, and what they understand about the effects generated by  
265 excess weight. Thus, 12 items were rearranged in different domains in comparison to previous  
266 works; all of them were assessed by the research group that considered this version satisfactory  
267 since the subscales are strongly related to each other.

268 The relationship between the processes and the stages of change was also a noteworthy finding. We  
269 observed a gradual increase in the use of processes of change as individuals moved from pre-  
270 contemplation (PC) to action (A) stages. People who were in PC stage used fewer processes of  
271 change compared to people in later stages. The highest use of processes of change was observed in  
272 the action (A) stage, before declining in the maintenance (M) stage. These results are in line with  
273 Andrés’s study that demonstrated that the use of processes of change was lower in the PC stage  
274 compared to subsequent stages [4]. Our results also corroborate the findings of a 12-week  
275 randomized controlled trial that concluded that the RA, CEP and ACP processes are higher in the  
276 Action stage compared to the previous stages [15]. Another recent study, using the S-Weight and P-  
277 Weight scales, revealed that patients with higher scores in the RE and ACP subscales are more likely  
278 to be in the Action stage for weight loss [23].

279 These findings suggest that when individuals shift their focus from *trying to lose weight* to *trying to*

280 *maintain weight*, they invest less effort in processes of change. It is also possible that individuals who  
281 are closer to their ideal body weight have improved self-esteem and, therefore, remain in the Pre-  
282 Contemplation stages [13]. However, this finding may partly explain the high rates of weight regain  
283 observed in long-term studies for obesity.

284 Furthermore, we found a significant relationship between the four subscales of the P-Weight and  
285 measures of concern about thinness through the EAT-26, which supports the convergent validity of  
286 the scale. As expected, participants who achieved higher scores on using processes of change also  
287 achieved higher scores on external measures. These results are in line with previous findings that  
288 showed a correlation between P-Weight and EAT-40 among university students and general  
289 population [4] that participated in a program for weight loss.

290 The Transtheoretical Model has been used to improve changes in health-related behaviors since the  
291 1980s, including smoking cessation and, more recently, weight management [6,24]. Thus,  
292 interventions designed to encourage behavior change are more effective when tailored to the  
293 individual's stage of change [8]. Measuring an individual's readiness for change in the weight  
294 management setting before undertaking an intervention can increase the effectiveness of weight loss  
295 interventions, especially in the long term [25]. Therefore, it is crucial to have reliable and validated  
296 measurement tools to assess the processes and stages of behavior change in weight management.  
297 The Brazilian version of the P-Weight has shown adequate psychometric properties and can be a  
298 useful tool to identify the individual's behavior change processes in relation to their weight.  
299 Personalized interventions that promote specific change processes and encourage progress through  
300 the motivational stages can aid in weight management.

301 This study has some limitations. Firstly, the non-random sampling procedure may not accurately  
302 represent the Brazilian population, so the results should not be generalized to populations with  
303 different characteristics. Secondly, participants' self-reported weight and height may be biased and  
304 could affect questionnaire responses. Finally, we highlight that the correlation between P-Weight  
305 scores and EAT-26 should be viewed as evidence of the scale convergent validity, and not as a  
306 measure of mental health issues.

### 307 **Conclusion**

308 The results of this study show that the Brazilian version of the P-Weight has satisfactory  
309 psychometric properties, making it a valuable application for weight loss interventions based on the  
310 Transtheoretical Model (TTM). The use of motivational strategies is crucial to promote adherence to

311 treatments for overweight and obesity. The availability of validated instruments that allow the  
312 evaluation of change processes, based on the TTM, is a great advance and should contribute  
313 enormously to this important public health problem. We highlight that individualizing weight  
314 management interventions according to each patient's motivational stage and processes of change  
315 allows for health professionals to employ more effective treatments according to each individual's  
316 motivational reality.

## 317 **Statements**

### 318 **Acknowledgments**

319 The authors thank to the Higher Education Personnel Improvement Coordination - Brazil (CAPES),  
320 National Council for Scientific and Technological Development and Research Incentive Fund of the  
321 Clinical Hospital of Porto Alegre for their funding support (FIPE/HCPA 2021.0447). Adriane R. Rosa  
322 would like to thank CNPq, PQ number 305707-2015/09.

### 323 **Statement of Ethics**

324 The study followed the conditions established in Resolution 466/12, of the Brazilian National Health  
325 Council (CNS) in accordance with the World Medical Association Declaration of Helsinki.

326 Study approval statement: This study protocol was reviewed and approved by the HCPA Research  
327 Ethics Committee, approval number (CAAE: 54352521.8.0000.5327 N. 2021.0447).

328 Consent to participate statement: Before entering, the survey participants were asked to confirm  
329 their eligibility and to sign an informed consent form. The study followed the bioethical principles of  
330 autonomy, beneficence, non-maleficence, veracity and confidentiality. All participants were informed  
331 about the risks related to psychological discomfort when answering questions about eating habits,  
332 their relationship with food and their body weight. To minimize these discomforts, subjects were  
333 informed that they could discontinue their participation in the study at any time, since participation  
334 is voluntary. To minimize the chance of breach of confidentiality, the questionnaires were  
335 anonymous. In addition, we use a specific tool for this type of study, which is the REDCAP. The  
336 completed online questionnaires were kept electronically and confidentially for a period of 5 years.  
337 All participants completed the free and informed consent form (TCLE) online and only after that the  
338 research started and they were free to discontinue the study at any time without this implying any  
339 harm to the participant. The research group declares to know and comply with the requirements of  
340 the General Data Protection Law (Law No. 13,709, of August 14, 2018) regarding the processing of  
341 personal data and sensitive personal data that will be used for the execution of this research project.

342 **Conflict of Interest Statement**

343 The authors have no conflicts of interest to declare.

344 **Funding Sources**

345 This study did not receive any specific grant from funding agencies in the public, commercial, or not-  
346 for-profit sectors.

347 **Author Contributions**

348 ARR and QC conceived the manuscript idea. QC and BBM performed the statistical analysis. PRZ,  
349 BBM, JFG, and QC wrote the manuscript. ARR supervised the findings of this work. All authors  
350 discussed the results and contributed to the final manuscript.

351 **Data Availability Statement**

352 All data generated or analyzed during this study are included in this article. Further enquiries can be  
353 directed to the corresponding author.

## References

- 1 Obesidade AB para E da. Diretrizes brasileiras de obesidade 2016. VI Diretrizes Brasileiras de Obesidade. 2016;7–186.
- 2 Safaei M, Sundararajan EA, Driss M, Boulila W, Shapi'i A. A systematic literature review on obesity: Understanding the causes & consequences of obesity and reviewing various machine learning approaches used to predict obesity. *Comput Biol Med*. 2021 Sep;136. DOI: 10.1016/j.combiomed.2021.104754
- 3 Jospe MR, Roy M, Brown RC, Haszard JJ, Meredith-Jones K, Fangupo LJ, et al. Intermittent fasting, Paleolithic, or Mediterranean diets in the real world: Exploratory secondary analyses of a weight-loss trial that included choice of diet and exercise. *American Journal of Clinical Nutrition*. 2020 Mar;111(3):503–14.
- 4 Andrés A, Saldaña C, Gómez-Benito J. The transtheoretical model in weight management: Validation of the processes of change questionnaire. *Obes Facts*. 2011 Dec;4(6):433–42.
- 5 Frie K, Hartmann-Boyce J, Jebb SA, Aveyard P. Effectiveness of a self-regulation intervention for weight loss: A randomized controlled trial. *Br J Health Psychol*. 2020;25(3):652–76.
- 6 Lecube A, Sánchez E, Andrés A, Saldaña C, Morales MJ, Calañas A, et al. Assessing Motivational Stages and Processes of Change for Weight Management Around Bariatric Surgery: a Multicenter Study. *Obes Surg*. 2019 Oct;29(10):3348–56.
- 7 Prochaska JO. Strong and Weak Principles for Progressing From Precontemplation to Action on the Basis of Twelve Problem Behaviors. 1994.
- 8 Sarkin JA, Johnson SS, Prochaska JO, Prochaska JM. Applying the transtheoretical model to regular moderate exercise in an overweight population: Validation of a stages of change measure. *Prev Med (Baltim)*. 2001;33(5):462–9.
- 9 di Noia J, Prochaska JO. Dietary stages of change and decisional balance: A meta-analytic review. *Am J Health Behav*. 2010;34(5):618–32.
- 10 Mastellos N, Gunn LH, Felix LM, Car J, Majeed A. Transtheoretical model stages of change for dietary and physical exercise modification in weight loss management for overweight and obese adults. *Cochrane Database of Systematic Reviews*. 2014 Feb;2014(2). DOI: 10.1002/14651858.CD008066.pub3
- 11 Andrés A, Gómez J, Saldaña C. The transtheoretical model and obesity: A bibliometric study. *Scientometrics*. 2007;73(3):289–301.
- 12 Bennett GG, Steinberg D, Askew S, Levine E, Foley P, Batch BC, et al. Effectiveness of an App and Provider Counseling for Obesity Treatment in Primary Care. *Am J Prev Med*. 2018 Dec;55(6):777–86.

- 13 Ceccarini M, Borrello M, Pietrabissa G, Manzoni GM, Castelnuovo G. Assessing motivation and readiness to change for weight management and control: An in-depth evaluation of three sets of instruments. *Front Psychol*. 2015;6(MAY). DOI: 10.3389/fpsyg.2015.00511
- 14 Andrés A, Saldaña C, Beeken RJ. Assessment of processes of change for weight management in a UK sample. *Obes Facts*. 2015 Mar;8(1):43–53.
- 15 Karintrakul S, Angkatavanich J. A randomized controlled trial of an individualized nutrition counseling program matched with a transtheoretical model for overweight and obese females in Thailand. *Nutr Res Pract*. 2017 Aug;11(4):319–26.
- 16 Andrés A, Saldaña C, Beeken RJ. Assessment of processes of change for weight management in a UK sample. *Obes Facts*. 2015 Mar;8(1):43–53.
- 17 Andrés A, Saldaña C, Gómez-Benito J. Establishing the stages and processes of change for weight loss by consensus of experts. *Obesity*. 2009 Sep;17(9):1717–23.
- 18 Bighetti F. “Tradução e validação do Eating Attitudes Test (EAT-26) em adolescentes do sexo feminino na cidade de Ribeirão Preto - SP.” Tese (Mestrado em Enfermagem em Saúde Pública). 2003;123.
- 19 Floyd FJ, Widaman KF. *Factor Analysis in the Development and Refinement of Clinical Assessment Instruments*. 1995.
- 20 Bentler PM. *QUANTITATIVE METHODS IN PSYCHOLOGY Comparative Fit Indexes in Structural Models*. 1990.
- 21 Cangur S, Ercan I. Comparison of model fit indices used in structural equation modeling under multivariate normality. *Journal of Modern Applied Statistical Methods*. 2015;14(1):152–67.
- 22 Damásio BF. *Uso da análise fatorial exploratória em psicologia*. 2012.
- 23 Wilson M, Orgass H, Dearborn-Tomazos J. Stages and Processes of Change for Weight Loss in Acute Stroke or TIA Patients Living with Obesity. *Cerebrovasc Dis Extra*. 2022 Dec;12(1):1–6.
- 24 Prochaska JO, Diclemente CC, Norcross JC. *In Search of How People Change Applications to Addictive Behaviors*. 1992.
- 25 Johnson SS, Paiva AL, Cummins CO, Johnson JL, Dyment SJ, Wright JA, et al. Transtheoretical Model-based multiple behavior intervention for weight management: Effectiveness on a population basis. *Prev Med (Baltim)*. 2008;46(3):238–46.

## Figure Legends

Fig. 1. Legend text. Diagram representing the four first-order factors model for processes of change in weight management. Emotional=RE=emotional re-evaluation. Action=ACP= actions for weight control. Environment=RA=environmental restructuring. Consequences=CEP=consequences of excess weight.

Fig. 2. Legend text. Analysis of processes of change scores across stages of change. PC=pre-contemplation; C=contemplation; P=preparation; A=action; M=maintenance.

Fig. 3. Legend text. Analysis of stages of change in clinical and community samples. 1=pre-contemplation (PC) 2=contemplation (C) 3=preparation (P) 4=action (A) 5=maintenance (M). Y axis=S-Weight mean scores.



**Table 1.** Sociodemographic characteristics of the sample. (n=656).

Characteristic	Community (n=330)	Clinical (n=326)	Total (n=656)
Gender, %Female	81.5	74.5	78
Age, years (a)	41.21 (11.46)	41.84 (11.11)	41.53 (11.28)
BMI (a,b), Kg/m <sup>2</sup>	28.41 (5.45)	33.24 (7.71)	30.81 (7.10)
<b>%Educational level</b>			
Higher education	63.63	70.85	67.22
High School	36.36	29.14	32.77
<b>%Skin-color</b>			
White	85	83.12	84.29
Non-white	15	16.87	15.70
<b>%Marital status</b>			
Married	65.45	59.50	62.5
Not married	34.54	40.50	37.5
<b>%Family income</b>			
<R\$ 7,272.00	66.06	85.27	75.61
>R\$ 7,272.00	33.93	14.72	24.39
	a=Standard deviations	b=Self-reported BMI	

**Table 2. Factor loadings of P-Weight items to factors in the exploratory factor analysis (n=328).**

item	Question	Communalities	Factor loadings*			
			RE	ACP	RA	CEP
4	I now realize I have a weight problem	0.629	0.525			
9	I am worried about gaining more weight	0.699	0.772			
13	Losing weight would help me improve my relationships with others	0.777	0.493			
17	Being overweight makes me feel bad	0.830	0.618			
21	I feel guilt when I overeat	0.563	0.644			
23	If I lost weight, I would feel better about myself	0.914	1.007			
25	I'm not happy with my current weight	0.756	0.828			
27	If I lost weight, I would be happier	0.839	0.833			
29	I feel good when I am able to control my eating habits	0.654	0.620			
31	When I lose weight, I feel proud of myself	0.767	0.813			
34	I am committed to losing weight	0.627	0.468			
2	I look for information about the types of food that could help me lose weight	0.412		0.497		
3	I try to put food away to avoid nibbling	0.506		0.726		
6	I tell myself positive things to avoid overeating	0.595		0.728		
7	I try not to have food in sight	0.637		0.685		
10	When I really want to eat, I do activities to avoid it	0.651		0.702		
11	There are no snack foods in my fridge or cupboards	0.651		0.749		
14	I have learned to control my appetite	0.620		0.790		
15	I avoid places where people eat a lot	0.724		0.816		
18	I have learned skills that reduce my desire to eat (e.g., distracting myself)	0.685		0.705		
19	When I am on a diet <sup>1</sup> I avoid eating with people who I overeat with	0.662		0.693		
22	I avoid buying high-calorie food	0.585		0.728		
26	To avoid overeating I prefer eating at home or cooking my own food	0.754		0.645		
24	I am aware that there are more and more people who encourage me to lose weight	0.821			0.682	
28	My family and friends praise me for not overeating	0.811			0.566	

30	My family and friends congratulate me when I manage to lose weight	0.770			0.529	
32	People around me support me in trying to lose weight	0.852			0.723	
33	I have someone who listens to me when I need to talk about my being overweight	0.510			0.374	
5	Society's view of obese people affects me emotionally	0.527			0.463	
8	My weight restricts my relationships	0.818			0.609	
12	My current weight makes my daily life difficult	0.846			0.554	
16	My family and friends are worried about my weight	0.879			0.614	
20	Most of my health problems are due to my being overweight	0.762			0.424	
	Eigenvalues		17.55	3.83	1.63	1.13
	Explained variance proportion (%)		53	65	70	73

RE=emotional re-evaluation ACP=actions for weight control RA=environmental restructuring CEP=Consequences of excess weight

\*Extraction of factor loadings using function `fa()` psych package maximum likelihood method (ML), R program version 4.1.3

**Table 3. Indices obtained in the confirmatory factor analysis.**

	CFI	NFI/TLI	SRMR	RMSEA
	comparative fit index	normed fit index /Tucker-Lewis Index	standardized root means square residual	Root means square error of approximation
Model 1	<b>0.863</b>	<b>0.852</b>	<b>0.077</b>	<b>0.083</b>

**Table 4.** Internal consistency analysis of the 4 factors P-Weight questionnaire in both subgroups and total sample.

	SAMPLE 1 (N=328)			SAMPLE 2 (N=328)			Total sample (N=656)	
	Number of Items	Cronbach's alpha	Item-total correlations (range)	Cronbach's alpha	Item-total correlations (range)	Cronbach's alpha	Item-total correlations (range)	
RE	11	0.933	0.558-0.827	0.925	0.541-0.803			
ACP	12	0.922	0.480-0.746	0.921	0.477-0.763			
RA	5	0.884	0.515-0.788	0.871	0.436-0.763			
CEP	5	0.904	0.576-0.834	0.900	0.547-0.811			
						0.959	0.377 a 0.776	

RE=emotional re-evaluation. ACP=Actions for weight control. RA=environmental restructuring. CEP=consequences of excess weight.

**Table 5. Analysis of variance of processes of change scores across stages of change (ANOVA).**

	PC mean (SD)	C mean (SD)	P mean (SD)	A mean (SD)	M mean (SD)	F*(Sig)
RE	54.92 (17.02)	77.76 (12.85)	86.24 (9.86)	86.24 (12.43)	80.67 (14.93)	60.55 (0.000)
ACP	50.83 (14.15)	55.03 (14.91)	59.48 (14.73)	74.40 (19.47)	70.87 (17.14)	40.24 (0.000)
CEP	44.80 (18.23)	58.00 (19.76)	75.66 (22.32)	75.04 (23.92)	69.15 (24.94)	25.05 (0.000)
RA	47.12 (18.00)	63.25 (16.78)	79.98 (19.15)	74.43 (21.26)	72.44 (21.35)	36.20 (0.000)
Clinical sample (n) (%)	<b>1 (0.15)</b>	<b>7 (1.06)</b>	<b>42 (6.40)</b>	<b>155 (23.6)</b>	<b>121 (18.4)</b>	<b>326 (49.7)</b>
Community sample (n) (%)	<b>49 (7.47)</b>	<b>95 (14.5)</b>	<b>42 (6.40)</b>	<b>68 (10.4)</b>	<b>76 (11.6)</b>	<b>330 (50.3)</b>
Total (n)	<b>50</b>	<b>102</b>	<b>84</b>	<b>223</b>	<b>197</b>	<b>656</b>

PC=pre-contemplation C=contemplation P=preparation A=action M=maintenance RE=emotional re-evaluation ACP=actions for weight control RA=environmental restructuring CEP=consequences of excess weight

**Table 6. Spearman's correlations between P-Weight and EAT-26**

	RE	ACP	RE	CEP
<b>EAT 26* – Correlation Coefficient</b>	0.579	0.593	0.584	0.638
<b>p (Sig)</b>	<0.001	<0.001	<0.001	<0.001

RE=emotional re-evaluation. ACP=actions for weight control. RA= environmental restructuring. CEP=consequences of excess weight. \*EAT-26 =Eating attitude test-26; validated for the Brazilian population (FORTES et al., 2016)

**Table 7. Difference means scores of the P-Weight subscales between the community and clinical sample**

P-Weight domains	Community sample (n=330)	Clinical sample (n=326)	Overall sample (n=656)
<b>RE</b>			
Mean (SD)	73.6 (16.4)	88.2 (10.7)	80.9 (15.7)
Median [Min, Max]	75.4 [23.1, 100]	89.2 [43.1, 100]	83.1 [23.1, 100]
<b>ACP</b>			
Mean (SD)	56.4 (14.7)	76.9 (17.5)	66.6 (19.1)
Median [Min, Max]	56.7 [20.0, 93.3]	76.7 [28.3, 100]	65.0 [20.0, 100]
<b>RA</b>			
Mean (SD)	58.4 (17.6)	86.7 (14.3)	72.4 (21.4)
Median [Min, Max]	60.0 [20.0, 100]	88.0 [36.0, 100]	76.0 [20.0, 100]
<b>CEP</b>			
Mean (SD)	53.6 (19.9)	83.4 (19.5)	68.4 (24.7)
Median [Min, Max]	52.0 [20.0, 100]	88.0 [20.0, 100]	72.0 [20.0, 100]

RE=emotional re-evaluation. ACP=actions for weight control. RA=environmental restructuring CEP=consequences of excess weight









