Investigation of acute encephalitis syndrome with implementation of metagenomic next generation sequencing in a low-middle-income setting

Shrestha R*[Ψ,1,2], Katuwal N*[1], Tamrakar D[1,3], Tato CM[4], Vanaerschot M[5], Ahyong V[4], Gil J[4], Madhup SK[6], Gupta BP[7]

Center for Infectious Diseases, Dhulikhel Hospital Kathmandu University Hospital
Department of Pharmacology, Kathmandu University School of Medical Sciences
Department of Community Medicine, Kathmandu University School of Medical Sciences
Rapid Response Team, Chan Zuckerberg Biohub
Roche Diagnostics
Department of Microbiology, Kathmandu University School of Medical Sciences
World Health Organisation-Nepal

* Contributed equally
Ψ Corresponding Author

Word Count (Intro, Methods, Results, Discussion and Conclusion): 3264
Abstract Word Count: 371
Figures: 4
Tables: 1

Correspondence:
Dr. Rajeev Shrestha
Chief, Center for Infectious Diseases, Dhulikhel Hospital Kathmandu University Hospital
Professor, Department of Pharmacology, Kathmandu University School of Medical Sciences
Dhulikhel, Kavre, Nepal
Email: rajeev.shrestha@kusms.edu.np
Ph: +977-9801002489

Abstract

Background

Acute Encephalitis Syndrome is characterised by acute onset of fever and altered sensorium followed by a rapidly worsening clinical condition and even death. The causative agents vary with season and geographic location, while the etiologies remain unknown in 68-75% of the cases. In Nepal, the cases of acute encephalitis syndrome are tested only for Japanese encephalitis, which constitutes about 15% of the cases globally. However, there could be several causative organisms, including vaccine-preventable etiologies that cause acute encephalitis, which upon identification could direct public health efforts for prevention, including expanded use of vaccines or address gaps in vaccine coverage.

Objective

This study employs metagenomic next generation sequencing in the exploration of infectious etiologies contributing to acute encephalitis syndrome in a low-and-middle-income setting.

Methods

In this study, we have investigated 90, Japanese encephalitis-negative, banked retrospective cerebrospinal fluid samples that were collected as part of a national surveillance network in 2016 and 2017. The randomisation was done to include three age groups of <5 years, 5-14 years, and >15 years. Only some metadata related to the subjects (age and gender) were known. The analysis was performed in two batches which included total nucleic acid extraction, followed by individual library preparation for DNA and RNA and sequencing on Illumina iSeq100. The generated genomic data were interpreted using CZID and confirmed with polymerase chain reaction.

Results

Human alphaherpesvirus 2 and Enterovirus B were seen in two of the ninety samples. These hits were confirmed by qPCR and seminested PCR respectively. Most of the other samples were marred by low sample quality, lack of clinical metadata, storage issues, and possible ambiguous collection procedures.

Conclusion

From this study, two documented, causative agents were revealed through metagenomic next generation sequencing. Insufficiency of clinical metadata, process controls, and absence of standard operating procedure to collect and store samples in nucleic acid protectants impeded the study and incorporated ambiguity while correlating the identified hits to infections. Therefore, there is dire need of implementing stringent collection procedures for sample collection, including proper process controls. Despite the challenging conditions, this study highlights the usefulness of mNGS to investigate diseases with unknown etiologies, such as AES, and guide the development of adequate clinical management algorithms and outbreak investigations in low-middle income settings.

Keywords: acute encephalitis syndrome, enterovirus, Human alphaherpes virus, low-middle-income country, metagenomic next generation sequencing

INTRODUCTION

Acute Encephalitis Syndrome (AES) is defined by acute onset of fever and a change in mental status (including symptoms such as confusion, disorientation, coma, or inability to talk) and/or new onset of seizures (excluding simple febrile seizures) in a person of any age at any time of year.[1] This term was coined by World Health Organization (WHO) in 2008.[1] Globally, based on various studies, the incidenceof AES has ranged from 3.5 to 7.4 per 100,000 patients-years, with a higher incidence among children.[2]

The patients suffering from AES usually present acute onset of fever and altered sensorium. This is followed by rapidly worsening clinical conditions and death.[3] The survivors can suffer from long term health issues, including neurological sequelae.[3,4] The etiologies behind AES can be grouped under infective and non-infective categories, with the infective category comprising of a broad range of organisms (bacteria, virus, parasites).[2,5] The causative agents of AES also vary with season and geographic location.[6] Research has shown that the etiologies of AES remain unknown in 68-75% of the cases, while Japanese encephalitis (JE) constitutes about 15% of the cases.[7,8,9,10] The landscape of AES, in terms of etiology, has changed in India as well, where outbreak investigations and surveillance studies have increasingly reported non-JEV etiologies.[11]

In Nepal, JE is majorly associated with mortality and morbidity among children.[12] Therefore, since 2004, the Ministry of Health and Population of Nepal, supported by the Office of Infection Prevention Division, World Health Organisation (WHO), has integrated JE surveillance with Acute Flaccid Paralysis, Neonatal Tetanus, and Measles in its National Surveillance Network.[13] Until 2011, over 23,000 AES cases were reported by the surveillance network.[14] Due to a lack of knowledge in etiology, AES cases are only tested for JE and clinical management is performed based on this result. The incidence of undiagnosed AES etiology contributes to a high rate of death and morbidity.[14] There could be several etiologies, including vaccine preventable etiologies, that cause acute encephalitis, which upon identification could direct public health efforts for prevention, including expanded use of vaccines or addressing gaps in vaccine coverage. Herpes Simplex Virus (HSV), Varicella-Zoster Virus (VSV), Enterovirus, Adenovirus, and Rubella, as well as emerging pathogens such as Nipah, Chandipura and Chikungunya have all been reported as causative viral agents of AES, while Neisseria meningitidis, Streptococcus pneumoniae, Listeria sp, and Brucella have been reported as causative bacterial agents.[2,15,16,17]

While molecular methods such as PCR require prior genetic information on causative agents, genomic

methods such as metagenomic Next Generation Sequencing (mNGS) can simultaneously identify

minute amounts of infections and co-infections of varying origin in a single investigation and assist in

the investigation of transmission of such infections.[18, 19] With the recent dramatic decrease in

sequencing costs, this technology provides access to genomic information in a scale that can be

implemented to fill gaps in routine clinical practice and address epidemiological questions. In addition

to identification (identifying genotypes, virulence or pathogenesis), NGS provide information

epidemiological investigation (comparative genomics, phylogenetic analysis).[20, 21, 22, 23]

This study, employing mNGS to explore the infective etiologies behind AES, complements a growing

number of studies that have used a similar approach to investigate encephalitis, including in a LMIC

context.[16,24,25,26] The identification of such etiologies is an important step in developing effective

prevention and treatment measures which in turn will reduce disability and morbidity in LMICs and

beyond.

METHODOLOGY

Sample Collection and Selection

The investigation included a random selection of 90 retrospective cerebrospinal fluid (CSF) samples

that were collected by WHO-IPD (World Health Organisation-Immunization Preventable Diseases),

throughout Nepal, as a part of the National AES Surveillance Network in collaboration with FWD

(Family Welfare Department) in 2016 and 2017. These samples had been tested for JE at NPHL

(National Public Health Laboratory) and stored at low but undocumented temperature. For this study,

only those samples that tested negative for JE were selected. A randomisation was done to include

three age groups of <5 years, 5-14 years, and >15 years. Only some metadata related to the subjects

(age and gender) were known. Each sample was provided with unique study codes to maintain privacy.

Nucleic Acid Extraction and mNGS

Total Nucleic Acid was extracted from the CSF samples using Zymo Quick-DNA/RNA™ Pathogen

MiniPrep (R1042).

The total nucleic acid samples were aliquoted into two sub samples for RNA and DNA library

preparation, respectively. The library preparations were done using NEB Library Prep Kit Ultra II RNA

for RNA (New England Biolabs, E7770S) library preparation, and NEB Library Prep Kit Ultra II FS

DNA for DNA library preparation (New England Biolabs, E7805S). The library preparation for the
first 30 samples was done in a single batch (at Chan Zuckerberg Biohub, USA) while the remaining 60
samples were done in three batches of 20 samples each (at Dhulikhel Hospital Kathmandu University
Hospital, Nepal). Negative extraction and library preparation controls were included in each batch. The
library preparation included 10ng of input nucleic acid, followed by fragmentation, adapter ligation,
cleanup (Solid Phase Reversible Immobilization beads), barcoding and amplification of library for 12-
16 cycles. With subsequent library preparations, quality control was done using agarose gel
electrophoresis and Tapestation 4200 platform from Agilent Technologies and later by qPCR using
Kapa Illumina Library Amplification (KK2702) and Quantitation Complete Kit (KK4923). It was made
sure that the length of DNA in the libraries was around 350-400bp and had concentration >1nM. In
RNA library preparation, ERCC (External Control Controls Consortium, 4456740) RNA Spike-in
controls were used.

The libraries that passed quality control filters were pooled and run on an Illumina iSeq100 sequencer.
The sequencing was performed for 2x146bp length using custom unique dual indices of 12 bp length.
5% PhiX was added as an internal control for sequencing. The loading concentration of pooled libraries
was maintained at 100-120pM.

Data Analysis
The analysis was performed on the CZ ID (formally known as IDSeq) platform developed by Chan
Zuckerberg Initiative and CZ Biohub. CZID accepts raw sequencing data, perform host and quality
filteration, followed by execution of assembly based alignment pipeline.[27]The samples are analysed
based on number of reads per million (Number of reads aligning to the taxon in the NCBI NR/NT
database, per million reads sequenced), reads (Number of reads aligning to the taxon in the NCBI
NT/NR database), contig number (Number of assembled contigs aligning to the taxon in the NCBI
NT/NR database), id% and z-score. The samples were also visualized using a rpm heatmap where
samples and controls are cross-matched against each other. Respective background models were
created, from negative extraction and library preparation control, for RNA and DNA Libraries.

PCR Confirmation
Human alphaherpes virus confirmation was done by qPCR (KAPA HiFi HotStart Ready Mix) using
two primer sets: established primers (FP: 5'TGCAGTTTACGTATAACCACATACAGC 3' and RP: 5'
AGCTGCGGGCCTCGTT 3') and self-designed primers (FP: 5' GACTCAAACACGTGCACCAC

and RP: 5' CCATCGCGTACAGCCTACAT 3').[28] The primer sets were designed using NCBI primer

blast and Gene Script, then checked with Beacon Designer Free and Snap Gene Viewer.

Similarly, for confirmation of Enterovirus, modified protocol with established primers from

Enterovirus Surveillance Guidelines were used to perform semi-nested Polymerase Chain Reaction

(snPCR).[29] The protocol followed visualisation of the bands, for confimration, in agarose (1.5%) gel

electrophoresis.

Ethics

This study was ethically cleared from Nepal Health Research Council (NHRC) under id: 903 – 2019.

RESULTS

Subject Metadata

The samples selected for this study were banked, retrospective CSF samples collected in 2016 and

2017 with limited metadata such as age and gender. Out of the 90 subjects, 31 (34.4%) were female

while the age distribution has been presented below:

Table 1: Age Distribution between subjects with AES

SN	Age Group	Number
1	<5 years	23
2	5-14 years	14
3	>15 years	53

The median age of the subjects infected with AES was 20 years (IQR: 4-79 years)

Sample collection

The samples were collected in glass bottles without any preservative and transported to NPHL where

they, first, had been tested for JE and subsequently stored at undocumented but low temperature. As

these samples were collected in 2016 and 2017 and banked, negative controls were not available during

collection and transportation.

Nucleic Acid Extraction and mNGS

The extracted nucleic acid had a concentration ranging from too low to detect to 222 ng/ul. As the

analysis was done in two sets. Each set was processed for DNA and RNA Library Preparation and has

been presented accordingly. Out of 90 samples, only two samples showed confirmed hits from
Enterovirus B and Human alphaherpesvirus 2, respectively.

mNGS of RNA Libraries

The results from RNA libraries showed some distinct organisms hit in CZID, and also provided a broad
picture of the landscape of taxa across the samples. The following are heatmaps generated from
through RNA library preparation.

Figure 1: Heatmap depicting top hits from sequencing of RNA library for S01 to S90. The organisms (at the genus level)
that were seen in the samples are shown on the left. The names of the samples are on the top of the heat map. The samples
marked by green are water controls during extraction (NEC) and library preparation (NLC). The heatmap was generated
using the threshold of NT rPM (nucleotide reads per million) >=10 and NT L (alignment length in basepairs: length of the
aligned sequence)>=50

Figure 2: Heatmap depicting hits for Enterovirus (unnamed genus taxon 12059 and 138949) from sequencing of
AES_S47_RNA. The names of the samples are on the top of the heat map. The samples marked by green are water controls
during extraction (NEC) and library preparation (NLC). The heatmap was generated using the threshold of NT rPM
(nucleotide reads per million) >=10 and NT L (alignment length in basepairs: length of the aligned sequence)>=50

In figures 1 and 2, we can see top hits of organisms in the heat map that shows various organisms
which are seen at similar levels in the water controls as well. Nevertheless, Pseudomonas genus is seen

in all of the samples including few negative controls. There was similar trend, in both sets, with other

organism such as Sphingomonas, Acinetobacter, Escherichia and others.

Interestingly, only AES_S47_RNA showed a hit to Enterovirus B. This hit was particular to sample 47

and not seen in any negative controls. The metrics such as rPM of 359,409.1 (provides information of

the abundance of a specific microbe within the sample), NT L (depicts the length of aligned sequence

in base pair), Z score of 99 (shows the significance of any hit compared to the background), coverage

visualization (assess breadth and depths of reads) and id of 85.8% signify that the organism hit is

highly similar to the reference organism.[30] The figure below shows the abundance of Enterovirus B in

Sample 47 (NT rPM >=10 and NT L >=50). The coverage breadth of this hit was 98.7% with depth of

700.4x as seen in Figure 3.

Figure 3: Coverage visualisation of Enterovirus in AES_S47_RNA from CZID, depicting the coverage metrics of the contigs

generated for the particular hit along with coverage depth and width. The hit was visualised with the threshold of NT rPM

>=10 and NT L >=50.

mNGS of DNA Libraries

In mNGS of DNA libraries, hits were observed for Human alphaherpesvirus 2 [AES_S28_DNA] from

the first set. The same sample showed hit for Human alphaherpesvirus 1, but in a very low abundance.

Additionally, background contaminations (laboratory and hospital) were seen in the water controls in

this DNA sequencing result as well. Similar to RNA Libraries, most of the samples showed hits for

Sphingomonas spp., Pseudomonas spp, or Acinetobacter spp. Nevertheless, the following figure depicts

the result of the hit where there were 2,598.6 rPM for Human alphaherpesvirus 2.

Figure 4: Result from CZID depicting the details of top hits in AES_S28_DNA along with various metrics related to the hit. The hits were visualised with the threshold of NT rPM >=10 and NT L >=50.

However, due to lower coverage, contig visualization was not available for this hit.

PCR Confirmation

Confirmation of Human alphaherpesvirus 2

Out of the two primer sets used, the established primers fared better providing Ct value of 23.11 for Human alphaherpesvirus 2.

Confirmation of Enterovirus B

After completion of snPCR for Enterovirus, the band was seen between 700-800bp after first amplification and between 300-400bp after final amplification. This confirmed the presence of Enterovirus as per the Enterovirus Surveillance Guidelines.[29]

DISCUSSION

Demography of Acute Encephalitis Syndrome

Most of the subjects suffering from acute encephalitis syndrome (AES) were young male population of median age 20 years. This gender distribution was concurrent to previous studies done, in Nepal, on epidemiology of AES.[31,32] It has been observed that AES affects individuals from both gender and all ages, however, most of the studies have been done in younger population, as they pose high risk due to lack of developed antibodies.[33,34,35,36] Another study done in Nepal also observed the young median age (19 years) for AES, while others observed older population.[31,32,37,38]

Metagenomic Next Generation Sequencing

In this study, out of the 90 samples tested, most (n=88) of them could not be specified as specific hits. This was due to high level of background contaminants resulting in low confidence in calling organism

hits within the experimental samples. Nevertheless, two samples showed confirmed hits for Enterovirus

B and Human alphaherpesvirus 2, respectively. The absence of causative agent in remaining samples

indicate that either the samples did not have intact nucleic to start with or were degraded.[16,39] Extended

storage in uncontrolled conditions could also be a likely contributing factor for the degradation.

As per the result of mNGS, the high Z score (99) for Enterovirus B shows that hit for the organism is

present significantly in our sample, when compared to the background. The average length of

alignment (as shown by L metrics) is long (L=7258.7), which confirms for a good local alignment to

reference.[40] The id% is also higher (85.8%) meaning that the organism is highly similar to the reference

organism in the database. Additionally, when the genome coverage is seen in detail, we can see that our

sequenced genome depicts good coverage breadth and depth (depth of 700x and breadth of 98.7%),

which is the range and uniformity of sequencing coverage for the particular hit.[39] The presence of

ENVB was also confirmed through snPCR followed by visualization of product size specific for all

enteroviruses.[29]

The hit for alphaherpesvirus 2 was considered significant because it was not present in the control

samples at the thresholds used to analyse the sample (high Z score of 100%, L value of 128.9, id% of

99.9%) considered reliable.[24,27,41] The low contig value, for this hit, could be because of the organism

being present at such a low abundance that the sequencer did not sequence enough reads to generate a

contig. The contig value is dependent upon the total number of reads and the size of organism's

genome.[42] Additionally, the decreased sensitivity of mNGS due to low abundance of pathogen has been

studied for CSF.[43] Several methods have been reported that can be used to increase the abundance of

pathogen sequences or remove the unwanted <u>host</u> sequences.[44,45] Nevertheless, as this genus is

associated with encephalitis, the sample was taken further for analysis.[46,47] During confirmation, the

lower Ct value of 23.11 indicates presence of alphaherpesvirus 2, a known causative agent, in the

sample.

Enterovirus B and Human Alphaherpesvirus 2

Enterovirus B is a known causative agent of encephalitis.[16,48,49,50] Enteroviruses are named by their

transmission-route through the intestine.[51] Studies have shown that enterovirus can cause various

diseases in the nervous system, including aseptic meningitis, acute paralysis, encephalitis, meningo-

encephalomyelitis among others, in children.[52,53,54] Interestingly, studies in India have linked

Enterovirus, among other pathogens, to AES, by various studies.[55,56,57] For instance, Enterovirus

outbreak was first reported from Uttar Pradesh, India in 2006 with seasonal outbreaks with high fatality
occurring for several years.[55,58,59] Southern Nepal borders with UP, India and due to open borders and
similar climate, it is no surprise that Enterovirus was found in AES infected CSF sample from Nepal.
Additionally, some studies in Nepal have reported Enterovirus as possible etiology of AES for
Nepal.[60,61]

Similarly, Human alphaherpesvirus 2 is known to cause encephalitis in neonates and
immunocompromised patients. Herpes simplex encephalitis (HSE) has significant morbidity and
mortality, even with early diagnosis and treatment.[62,63] HSV is found to be one of the predominant
causes of AES in the western world.[64,65,66] Among herpes simplex encephalitis, the vast majority of the
encephalitis is caused by HSV-1, with HSV-2 being the etiology in less than 10% of the cases.[63] Studies
in India and Nepal have reported the presence of HSV-2 as causative agent of encephalitis, with
varying range of incidence. [61,67,68,69,70, 71]

Clinical Data and Process Control
However, due to lack of proper clinical metadata, the presence of Enterovirus B and Human
alphaherpes 2 virus could not be clinically correlated. Clinical metadata such as onset of fever, date of
infection, fatality, WBC counts, adjoining infection, etc are vital to correspond with the presence of
infections.[16,72,73] Additionally, usual environmental contaminants such as Sphingomonas spp.,
Pseudomonas spp, or Acinetobacter spp were seen. For instance, Sphingomonas are widely distributed
in nature, having been isolated from many different land and water habitats, as well as from plant root
systems, clinical specimens, and other sources. This is essentially due to their ability to survive in low
concentrations of nutrients.[74,75] Background contaminations of laboratory and hospital origin were also
seen in the water controls. With appropriate use of background or negative controls, a background
model can be created and subsequently subtracted from the results.[16,24]

Collection Procedures
The lack of identification of causative agent in other 88 samples could be because the majority of
samples were degraded prior to extraction. The samples that were analysed were as old as 2016 and
2017, and were collected without stringent or consistent guidelines. They could have gone through
numerous freeze and thaw cycles. Moreover, the sample could have gone through degradation prior to
storage depending upon how they were transported from the site of collection. The causative agents

could also have left the cerebrospinal fluid prior to collection. Collection of samples in nucleic acid protectant such as Zymo RNA/DNA Shield would have protected the nucleic acid from degradation after sampling.[76,77]

We postulate that the results of this study should not be corroborated as limitations as the CSF samples analysed were not collected specifically for mNGS. The lack of stringent collection procedures and sample quality are likely limiting to the analysis, resulting in ambiguous interpretation of some samples. Additionally, the sequencing was done in Illumina iSeq100 which has a maximum of approximately 4 million reads per run and can only accommodate a certain number of organisms with adequate coverage breadth and depth.[78]Additionally, host depletion and pathogen enrichment methods can be applied for samples with low pathogen abundance.[44,45]

CONCLUSION

From this study, two documented, causative agents were revealed through metagenomic next generation sequencing and subsequently confirmed by PCR. Insufficiency of clinical metadata, process controls, and absence of standard operating procedure to collect in nucleic acid protectant incorporates ambiguity when correlating identified pathogens to infections. Therefore, there is a dire need of implementing stringent collection procedures for sample collection, including proper process controls. Additionally, delicate samples such as CSF should be collected in a protectant and transported in a controlled and sterile environment. Collection of clinical metadata such as WBC count, primary diagnosis, discharge type and presence of other known organisms are vital for the analysis.

Identification and investigation of etiologies behind AES is essential for developing clinical management algorithms, improving surveillance with region-specific treatment and prevention policy as well as outbreak investigation. We do not expect the administration of mNGS as a regular diagnostic tool but rather an investigational and exploration tool to identify causative etiologies and develop molecular methods (such as qPCR) for diagnosis.

Declaration of competing interest

We declare that we have no potential conflicts of interest.

Acknowledgements

We thank Bill and Melinda Gates Foundation, Grand Challenges Explorations Grant for the support. We appreciate the guidance from Chan Zuckerberg Biohub, San Francisco. We express our gratitude to

412 Family Welfare Department and National Public Health Laboratory (NPHL) for supporting the study
413 by providing the samples. We also thank WHO-IPD for providing data for the study.
414

REFERENCES:

1. Guidelines for Surveillance of Acute Encephalitis Syndrome (With Special Reference to Japanese
Encephalitis) NVBDCP. 2006. Accessed: 18 July 2022. Available at:
http://www.nvbdcp.gov.in/Doc/AES%20guidelines.pdf

2. Granerod J, Crowcroft NS. The epidemiology of acute encephalitis. Neuropsychol Rehabil. 2007;
17:406–28.

3. Narain JP, Dhariwal AC, MacIntyre CR. Acute encephalitis in India: an unfolding tragedy. Indian J
Med Res. 2017. 145:584–7.

4. Srivastava N, Deval H, Mittal M, Kant R, Bondre VP. The Outbreaks of Acute Encephalitis
Syndrome in Uttar Pradesh, India (1978-2020) and Its Effective Management: A Remarkable Public
Health Success Story. Front Public Health. 2022 Feb 9;9:793268.

5. Tripathy SK, Mishra P, Dwibedi B, Priyadarshini L, Das RR. Clinico-epidemiological Study of Viral
Acute Encephalitis Syndrome Cases and Comparison to Nonviral Cases in Children from Eastern India.
J Glob Infect Dis. 2019 Jan-Mar;11(1):7-12

6. National Health Mission. Routine Immunization, Government of India. Available at:
http://www.nrhm.gov.in/nrhmcomponents/rmncha/immunization/background.html

7. AES/JE Cases and Deaths in the Country. National Vector Borne Disease Control Programme.
Directorate General of Health Services. Ministry of Health and Family Welfare, Government of India.
2012.

8. Director, Child Health Division. Teku, Kathmandu, Nepal: Department of Health Services, Ministry
of Health and Population. 2012. Acute Encephalitis Syndrome/Japanese Encephalitis Data of Nepal.

9. Cizman M, Jazbec J. Etiology of acute encephalitis in childhood in Slovenia. Pediatr Infect Dis J.
1993. 12:903–8.

10. Potharaju NR. Incidence Rate of Acute Encephalitis Syndrome without Specific Treatment in India
and Nepal. Indian J Community Med. 2012 Oct. 37(4):240-51.

11. Joshi R, Kalantari SP, Reingold A et al. Changing landscape of acute encephalitis syndrome in
India:A systematic review. Natl Med J India 2012. 25:212-20.

12. Rayamajhi A, Singh R, Prasad R, Khanal B, Singhi S: Study of Japanese encephalitis and other
viral encephalitis in Nepali children. Pediatr Int. 2007, 49 (6): 978-984.

13. Pant GR: A serological survey of pigs, horses, and ducks in Nepal for evidence of infection with

Japanese encephalitis virus. Ann N Y Acad Sci. 2006, 1081: 124-129.

14. Rayamajhi A, Ansari I, Ledger E, Bista KP, Impoinvil DE, Nightingale S, Kumar R, Mahaseth C,

Solomon T, Griffiths MJ. Clinical and prognostic features among children with acute encephalitis

syndrome in Nepal; a retrospective study. BMC Infect Dis. 2011 Oct 28;11:294

15. Hsu, V. P., Hossein, M. J., Parashar, U. D., Ali, M. M., Ksiazek, T. G., Kuzmin, I., Niezgoda, N.,

Rupprecht, C., Bresee, J., & Breiman, R. F. Nipah virus encephalitis reemergence, Bangladesh.

Emerging Infectious Diseases. 2004. 10(12), 2082 – 2087.

16. Saha S, Ramesh A, Kalantar K, Malaker R, Hasanuzzaman M, Khan LM, Mayday MY, Sajib MSI,

Li LM, Langelier C, Rahman H, Crawford ED, Tato CM, Islam M, Juan YF, de Bourcy C, Dimitrov B,

Wang J, Tang J, Sheu J, Egger R, De Carvalho TR, Wilson MR, Saha SK, DeRisi JL. Unbiased

Metagenomic Sequencing for Pediatric Meningitis in Bangladesh Reveals Neuroinvasive Chikungunya

Virus Outbreak and Other Unrealized Pathogens. mBio. 2019 Dec 17;10(6):e02877-19.

17. Booss, J. & Esiri, M. M. Viral encephalitis in humans. 2003. Washington, DC: American Society

for Microbiology Press.

18. Patricia J Simner, Steven Miller, Karen C Carroll, Understanding the Promises and Hurdles of

Metagenomic Next-Generation Sequencing as a Diagnostic Tool for Infectious Diseases, Clinical

Infectious Diseases, Volume 66, Issue 5, 1 March 2018, Pages 778–788

19. Ryan C. Shean & Alexander L. Greninger. One future of clinical metagenomic sequencing for

infectious diseases, Expert Review of Molecular Diagnostics. 2019. 19:10, 849-851.

20. Phillips KA, Douglas MP, Marshall DA. Expanding Use of Clinical Genome Sequencing and the

Need for More Data on Implementation. JAMA. 2020. Nov 24;324(20):2029-2030.

21. Gilbert GL. Molecular diagnostics in infectious diseases and public health microbiology: cottage

industry to postgenomics. Trends Mol Med. 2002. 8(6):280-7.

22. Dwivedi, S., Purohit, P., Misra, R. et al. Diseases and Molecular Diagnostics: A Step Closer to

Precision Medicine. Ind J Clin Biochem. 2007. 32, 374–398

23. Krishna NK, Cunnion KM. Role of molecular diagnostics in the management of infectious disease

emergencies. Med Clin North Am. 2012. 96(6):1067-78.

24. Wilson MR, O'Donovan BD, Gelfand JM, Sample HA, Chow FC, Betjemann JP, Shah MP, Richie

MB, Gorman MP, Hajj-Ali RA, Calabrese LH, Zorn KC, Chow ED, Greenlee JE, Blum JH, Green G,

Khan LM, Banerji D, Langelier C, Bryson-Cahn C, Harrington W, Lingappa JR, Shanbhag NM, Green

AJ, Brew BJ, Soldatos A, Strnad L, Doernberg SB, Jay CA, Douglas V, Josephson SA, DeRisi JL.

Chronic meningitis investigated via metagenomic next-generation sequencing. JAMA Neurol. 2018.
75:947–955.

25. Langelier C, Kalantar KL, Moazed F, Wilson MR, Crawford E, Deiss T, Belzer A, Bolourchi S,
Caldera S, Fung M, Jauregui A, Malcolm K, Lyden A, Khan L, Vessel K, Quan J, Zinter M, Chiu CY,
Chow ED, Wilson J, Miller S, Matthay MA, Pollard KS, Christenson S, Calfee CS, DeRisi JL.
Integrating host response and unbiased microbe detection for lower respiratory tract infection diagnosis
in critically ill adults. Proc Natl Acad Sci USA. 2018. 115:E12353–E12362.

26. Wilson MR, Sample HA, Zorn KC, Arevalo S, Yu G, Neuhaus J, Federman S, Stryke D, Briggs B,
Langelier C, Berger A, Douglas V, Josephson SA, Chow FC, Fulton BD, DeRisi JL, Gelfand JM,
Naccache SN, Bender J, Dien Bard J, Murkey J, Carlson M, Vespa PM, Vijayan T, Allyn PR, Campeau
S, Humphries RM, Klausner JD, Ganzon CD, Memar F, Ocampo NA, Zimmermann LL, Cohen SH,
Polage CR, DeBiasi RL, Haller B, Dallas R, Maron G, Hayden R, Messacar K, Dominguez SR, Miller
S, Chiu CY. Clinical Metagenomic Sequencing for Diagnosis of Meningitis and Encephalitis. N Engl J
Med. 2019 Jun 13;380(24):2327-2340.

27. Kalantar KL, Carvalho T, de Bourcy CFA, Dimitrov B, Dingle G, Egger R, Han J, Holmes OB,
Juan YF, King R, Kislyuk A, Lin MF, Mariano M, Morse T, Reynoso LV, Cruz DR, Sheu J, Tang J,
Wang J, Zhang MA, Zhong E, Ahyong V, Lay S, Chea S, Bohl JA, Manning JE, Tato CM, DeRisi JL.
IDseq-An open source cloud-based pipeline and analysis service for metagenomic pathogen detection
and monitoring. Gigascience. 2020 Oct 15;9(10):giaa111.

28. Namvar L, Olofsson S, Bergström T, Lindh M. Detection and typing of Herpes Simplex virus
(HSV) in mucocutaneous samples by TaqMan PCR targeting a gB segment homologous for HSV types
1 and 2. J Clin Microbiol. 2005 May;43(5):2058-64.

29. WHO. Enterovirus Surveillance Guidelines. WHO Regional Office for Europe. 2015. Available at:
https://www.euro.who.int/__data/assets/pdf_file/0020/272810/EnterovirusSurveillanceGuidelines.pdf

30. CZID Portal. https://czid.org. Accessed July 2022

31. Rayamajhi, et al. Evaluating cognitive outcomes in adult patients with acute encephalitis syndrome:
a prospective study from a tertiary care center in Nepal. Encephalitis. 2022; 2(2): 36-44

32. Thapa et al. Clinical profile and outcome of acute encephalitis syndrome (AES) patients treated in
College of Medical Sciences-Teaching Hospital. Journal of College of Medical Sciences. 2013. Vol 9,
No.2, 31-37.)

33. Rayamajhi, A., Ansari, I., Ledger, E. et al. Clinical and prognostic features among children with
acute encephalitis syndrome in Nepal; a retrospective study. BMC Infect Dis 11, 294 (2011).
https://doi.org/10.1186/1471-2334-11-294 AND

34. Griffiths MJ, Lemon JV, Rayamajhi A, Poudel P, Shrestha P, Srivastav V, et al. (2013) The
Functional, Social and Economic Impact of Acute Encephalitis Syndrome in Nepal – a Longitudinal
Follow-Up Study. PLoS Negl Trop Dis 7(9): e2383. AND

35. Roy DB, Khatri HV. Study of Demographic Profile, Etiology, and Clinical Outcome in Patients
Admitted With Acute Encephalitis Syndrome From the Western Part of India. Cureus. 2022 Mar
11;14(3):e23085. doi: 10.7759/cureus.23085. PMID: 35464588; PMCID: PMC9001832. And

36. Jmor, F., Emsley, H.C., Fischer, M. et al. The incidence of acute encephalitis syndrome in Western
industrialised and tropical countries. Virol J 5, 134 (2008). https://doi.org/10.1186/1743-422X-5-134)

37. Clinical presentation, etiology, and survival in adult acute encephalitis syndrome in rural Central
India. Joshi R, Mishra PK, Joshi D, et al. https://doi.org/10.1016/j.clineuro.2013.04.008. Clin Neurol
Neurosurg. 2013;115:1753–1761.

38. Roy DB, Khatri HV. Study of Demographic Profile, Etiology, and Clinical Outcome in Patients
Admitted With Acute Encephalitis Syndrome From the Western Part of India. Cureus. 2022 Mar
11;14(3):e23085. doi: 10.7759/cureus.23085. PMID: 35464588; PMCID: PMC9001832.

39. Thomas T, Gilbert J, Meyer F. Metagenomics - a guide from sampling to data analysis. Microb
Inform Exp. 2012 Feb 9;2(1):3.

40. CZID Portal. h https://chanzuckerberg.zendesk.com/hc/en-us. Accessed July 2022.

41. Zinter MS, Dvorak CC, Mayday MY, Iwanaga K, Ly NP, McGarry ME, Church GD, Faricy LE,
Rowan CM, Hume JR, Steiner ME, Crawford ED, Langelier C, Kalantar K, Chow ED, Miller S,
Shimano K, Melton A, Yanik GA, Sapru A, DeRisi JL. Pulmonary Metagenomic Sequencing Suggests
Missed Infections in Immunocompromised Children. Clin Infect Dis. 2019 May 17;68(11):1847-1855

42. Ayling M, Clark MD, Leggett RM. New approaches for metagenome assembly with short reads.
Brief Bioinform. 2020 Mar 23;21(2):584-594.

43. Miller S, Naccache SN, Samayoa E, Messacar K, Arevalo S, Federman S, Stryke D, Pham E, Fung
B, Bolosky WJ, Ingebrigtsen D, Lorizio W, Paff SM, Leake JA, Pesano R, DeBiasi R, Dominguez S,
Chiu CY. Laboratory validation of a clinical metagenomic sequencing assay for pathogen detection in
cerebrospinal fluid. Genome Res. 2019. 29:831–842.

44. Gu W, Crawford ED, O'Donovan BD, Wilson MR, Chow ED, Retallack H, DeRisi JL. Depletion
of abundant sequences by hybridization (DASH): using Cas9 to remove unwanted high-abundance
species in sequencing libraries and molecular counting applications. Genome Biol. 2016. 17:41.

45. Quan J, Langelier C, Kuchta A, Batson J, Teyssier N, Lyden A, Caldera S, McGeever A, Dimitrov
B, King R, Wilheim J, Murphy M, Ares LP, Travisano KA, Sit R, Amato R, Mumbengegwi DR, Smith
JL, Bennett A, Gosling R, Mourani PM, Calfee CS, Neff NF, Chow ED, Kim PS, Greenhouse B,
DeRisi JL, Crawford ED. FLASH: a next-generation CRISPR diagnostic for multiplexed detection of
antimicrobial resistance sequences. Nucleic Acids Res. 2019. 47:e83.

46. Carneiro, V.C.d., Alves-Leon, S.V., Sarmento, D.J.d. et al. Herpesvirus and neurological
manifestations in patients with severe coronavirus disease. Virol J 2022. 19, 10.

47. Patil S, Beck P, Nelson TB, Bran A, Roland W. Herpes Simplex Virus-2 Meningoencephalitis With
Abducens Nerve Palsy With Literature Review. Cureus. 2021 Jun 8;13(6):e15523

48. Kumar A, Shukla D, Kumar R, Idris MZ, Misra UK, Dhole TN. Molecular epidemiological study
of enteroviruses associated with encephalitis in children from India. J Clin Microbiol. 2012. 50:3509–
3512.

49. Jain S, Patel B, Bhatt GC. Enteroviral encephalitis in children: clinical features, pathophysiology,
and treatment advances. Pathog Glob Health. 2014 Jul;108(5):216-22

50. Calvo C, Gallardo P, Torija P, Bellón S, Méndez-Echeverría A, Del Rosal T, Baquero-Artigao F,
Sainz T, Romero M, Cabrerizo M. Enterovirus neurological disease and bacterial coinfection in very
young infants with fever. J Clin Virol. 2016 Dec;85:37-39

51. "Genus: Enterovirus". International Committee on Taxonomy of Viruses (ICTV). Accessed: 18
July 2022. Derivation of names Entero: from Greek enteron, 'intestine'

52. Chaudhary MC, Bronze MS. Enterovirus Infection. Medscape: Infectious Diseases. 2019.
Accessed: 18 July 2022.

53. Glaser CA, Gilliam S, Schnurr D et al. In search of encephalitis etiologies: diagnostic challenges in
the California Encephalitis Project, 1998–2000. Clin. Infect. Dis. 2003. 36(6), 731–742.

54. Gofshteyn J, Cárdenas AM, Bearden D. Treatment of chronic enterovirus encephalitis with
fluoxetine in a patient with X-linked agammaglobulinemia. Pediatr. Neurol. 2016. 64, 94–98.

55. Sapkal GN Sapkal GN Bondre VP Fulmali PV. Enteroviruses in patients with acute encephalitis,
Uttar pradesh, India. Emerg Infect Dis. 2009. 15: 295-298.

56. Mittal M, Kushwaha KP, Pandey AK, Fore MM. A clinico-epidemiological study of acute

encephalitis syndrome with multi organ dysfunction. International Journal of Contemporary Pediatrics.

2017. 4(3).

57. Ravi V, Hameed SKS, Desai A, Mani RS, Reddy V, Velayudhan A, Yadav R, Jain A, Saikia L,

Borthakur AK, Sharma A, Mohan DG, Bhandopadhyay B, Bhattacharya N, Inamdar L, Hossain S,

Daves S, Sejvar J, Dhariwal AC, Sen PK, Venkatesh S, Prasad J, Laserson K, Srikantiah P. An

algorithmic approach to identifying the aetiology of acute encephalitis syndrome in India: results of a

4-year enhanced surveillance study. Lancet Glob Health. 2022 May;10(5):e685-e693.

58. Mittal M, Bondre V, Murhekar M, Deval H, Rose W, Verghese VP, Mittal M, Patil G, Sabarinathan

R, Vivian Thangaraj JW, Kanagasabai K, Prakash JAJ, Gupta N, Gupte MM, Gupte MD. Acute

Encephalitis Syndrome in Gorakhpur, Uttar Pradesh, 2016: Clinical and Laboratory Findings. Pediatr

Infect Dis J. 2018 Nov;37(11):1101-1106

59. Kumar A, Shukla D, Kumar R, Idris MZ, Misra UK, Dhole TN. An epidemic of encephalitis

associated with human enterovirus B in Uttar Pradesh, India, 2008. J Clin Virol. 2011 Jun;51(2):142-5.

60. Giri A, Arjyal A, Koirala S, Karkey A, Dongol S, Thapa SD, Shilpakar O, Shrestha R, van Tan L,

Thi Thuy Chinh BN, Krishna K C R, Pathak KR, Shakya M, Farrar J, Van Doorn HR, Basnyat B.

Aetiologies of central nervous system infections in adults in Kathmandu, Nepal: a prospective hospital-

based study. Sci Rep. 2013;3:2382

61. Säll O, Thulin Hedberg S, Neander M, Tiwari S, Dornon L, Bom R, Lagerqvist N, Sundqvist M,

Mölling P. Etiology of Central Nervous System Infections in a Rural Area of Nepal Using Molecular

Approaches. Am J Trop Med Hyg. 2019 Jul;101(1):253-259.

62. Bookstaver PB, Mohorn PL, Shah A, Tesh LD, Quidley AM, Kothari R, Bland CM, Weissman S.

Management of viral central nervous system infections: a primer for clinicians. J Cent Nerv Syst Dis.

2017. 9.

63. AK AK, Mendez MD. Herpes Simplex Encephalitis. 2022 Mar 15. In: StatPearls [Internet].

Treasure Island (FL): StatPearls Publishing; 2022 Jan–. PMID: 32491575.

64. Khetsuriani, N., Holman, R. C. & Anderson, L. J. Burden of encephalitis-associated

hospitalizations in the United States, 1988–1997. Clinical infectious diseases : an official publication of

the Infectious Diseases Society of America. 2002. 35. 175–182.

65. Davison, K. L., Crowcroft, N. S., Ramsay, M. E., Brown, D. W. & Andrews, N. J. Viral

encephalitis in England, 1989–1998: what did we miss? Emerging infectious diseases. 2003. 9. 234–

240.

66. Huppatz, C. et al. Etiology of encephalitis in Australia, 1990–2007. Emerging infectious diseases.

2009. 15. 1359–1365.

67. Panagariya A, Jain RS, Gupta S, Garg A, Sureka RK, Mathur V. Herpes simplex encephalitis in

North West India. Neurol India. 2001 Dec;49(4):360-5

68. Bradshaw MJ, Venkatesan A. Herpes Simplex Virus-1 Encephalitis in Adults: Pathophysiology,

Diagnosis, and Management. Neurotherapeutics. 2016 Jul;13(3):493-508.

69. Kubo T, Rai SK, Nakanishi M, Yamano T. Seroepidemiological study of herpes viruses in Nepal.

Southeast Asian J Trop Med Public Health. 1991 Sep;22(3):323-5

70. Adhikari A. A Case Report on Herpes Simplex Encephalitis. Medical Journal of Shree Birendra

Hospital. 2012. 11(1).

71. Rayamajhi P, Nepal G, Ojha R, Rajbhandari R, Gajurel BP, Karn R. Evaluating cognitive outcomes

in adult patients with acute encephalitis syndrome: a prospective study from a tertiary care center in

Nepal. 2022. 2(2). 36-44.

72. Ramesh A, Nakielny S, Hsu J, Kyohere M, Byaruhanga O, de Bourcy C, Egger R, Dimitrov B,

Juan YF, Sheu J, Wang J, Kalantar K, Langelier C, Ruel T, Mpimbaza A, Wilson MR, Rosenthal PJ,

DeRisi JL. Metagenomic next-generation sequencing of samples from pediatric febrile illness in

Tororo, Uganda. PLoS One. 2019 Jun 20;14(6):e0218318.

73. Institute of Medicine (US) Roundtable on Value & Science-Driven Health Care. Clinical Data as

the Basic Staple of Health Learning: Creating and Protecting a Public Good: Workshop Summary.

Washington (DC): National Academies Press (US); 2010. Available from:

https://www.ncbi.nlm.nih.gov/books/NBK54302/

74. Leys NM, Ryngaert A, Bastiaens L, Verstraete W, Top EM, Springael D. Occurrence and

phylogenetic diversity of Sphingomonas strains in soils contaminated with polycyclic aromatic

hydrocarbons. Appl Environ Microbiol. 2004 Apr;70(4):1944-55

75. Cavicchioli R, Fegatella F, Ostrowski M, Eguchi M, Gottschal J. Sphingomonads from marine

environments. J Ind Microbiol Biotechnol. 1999 Oct;23(4-5):268-272.

76. Thraenhart O, Jursch C. Virucidal activity if the nucleic acid preservation product "DNA/RNA

Shield" against the murine parvovirus (MVM) at 20°C. 2018. Eurovir.

77. Phommanivong, V., Kanda, S., Shimono, T. et al. Co-circulation of the dengue with chikungunya

virus during the 2013 outbreak in the southern part of Lao PDR. Trop Med Health. 2016. 44. 24

78.   Iseq100 Specifications. Accessed: 18 July 2022. Available at

https://support.illumina.com/bulletins/2020/04/maximum-read-length-for-illumina-sequencing-

platforms.html