# Long-term Exposure to Ambient PM<sub>2.5</sub> and Hospitalizations for Myocardial Infarction among U.S. Residents: A Difference-in-Differences Analysis

Yichen Wang<sup>a\*</sup>, BS; Xinye Qiu<sup>a</sup>, PhD; Yaguang Wei<sup>a</sup>, PhD; Joel D. Schwartz<sup>a,b</sup>, PhD

<sup>a</sup> Department of Environmental Health, Harvard T.H. Chan School of Public Health, Boston,

### MA, USA

<sup>b</sup> Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA, USA

\*Corresponding author: Yichen Wang, Department of Environmental Health, Harvard T.H. Chan School of Public Health, 401 Park Drive, Boston, MA 02215 USA. Email:

yichenwang@hsph.harvard.edu

Total word count: 6710

1

### 1 Abstract

Background: Air pollution has been recognized as an untraditional risk factor for myocardial
infarction (MI). However, the MI risk attributable to long-term exposure to fine particulate matter
(PM<sub>2.5</sub>) is unclear, especially in younger populations, and few studies represented the general
population.

6 **Methods:** We applied the difference-in-differences approach to estimate the relationship between 7 annual PM<sub>2.5</sub> exposure and hospitalizations for MI among U.S. residents and further identified 8 potential susceptible subpopulations. All hospital admissions for MI in ten U.S. states over the 9 period 2002-2016 were obtained from the Healthcare Cost and Utilization Project State Inpatient 10 Database.

11 **Results:** In total, 1,914,684 MI hospital admissions from 8,106 ZIP codes in ten states from 2002 12 to 2016 were included in this study. We observed a 1.35% (95% CI: 1.11-1.59%) increase in MI 13 hospitalization rate for 1  $\mu$ g/m<sup>3</sup> increase in annual PM<sub>2.5</sub> exposure. The estimate was robust to 14 adjustment for surface pressure, relative humidity and co-pollutants. In the population with 15 exposure at or below 12  $\mu$ g/m<sup>3</sup>, there was a larger increment of 2.17% (95% CI: 1.79-2.56%) in hospitalization rate associated with 1  $\mu$ g/m<sup>3</sup> increase in PM<sub>2.5</sub>. Young people (0-34 years) and 16 17 elderly people ( $\geq$ 75 years) were the two most susceptible age groups. Residents living in more 18 densely populated or poorer areas and individuals with comorbidities were observed to be at a 19 greater risk.

Conclusions: This study indicates long-term residential exposure to PM<sub>2.5</sub> could lead to increased
 risk of MI among U.S. general population. The association persists below current standards.

Keywords: Long-term PM<sub>2.5</sub> exposure; Difference-in-differences; Myocardial infarction;
 Hospitalizations

# 2

# 24 Clinical Perspective:

# 25 What is new?

26	• Long-term exposure to PM <sub>2.5</sub> increased the risk of myocardial infarction in the general U.	S.
27	population.	
28	• Young individuals aged 0-34 years had the highest relative risk from long-term exposure	re
29	to PM <sub>2.5</sub> , and elderly people aged $\geq$ 75 years were the second most susceptible to the effect	ïs.
30	• Individuals with iron deficiency anemia, psychosis, and renal failure were more susceptib	le
31	to the long-term effects of $PM_{2.5}$ on MI.	
32	What are the clinical implications?	
33	• Long-term PM <sub>2.5</sub> exposure is one of the important modifiable environmental risk facto	rs
34	for myocardial infarction, therefore, air pollution control and behavioral intervention	ns
35	should be taken to prevent the occurrence of myocardial infarction.	
36		
37		
38		
39		
40	1. Introduction	
41	Myocardial infarction (MI) is responsible for years of life lost and reduced life expectance	су
42	across the globe (Ojha and Dhamoon, 2022). Over the last few decades, the morbidity ar	ıd
43	mortality burden of MI has declined in developed countries due to improved healthcare system	ns
44	and preventive actions (Camacho et al., 2022). However, an increasing incidence of MI over tim	ne
45	has been reported in younger populations with its underlying reasons unclarified (Arora et al., 201	9;

46 Yang et al., 2020).

3

47 Over 90% of the risk for acute MI is accounted for by modifiable risk factors (Chadwick Jayaraj et al., 2019). Air pollution is an important driver of MI with a population attributable 48 49 fraction of 5-7%, (Nawrot et al., 2011). This is similar to well-recognized behavioral factors (e.g., 50 physical exertion, alcohol, and coffee). To date, the epidemiological evidence on the relationship 51 between long-term exposure to ambient fine particulate matter  $\leq 2.5 \,\mu$ m in aerodynamic diameter 52  $(PM_{2.5})$  and MI remains inconclusive. Some studies suggested a positive association (Hartiala et 53 al., 2016; Madrigano et al., 2013), whereas other research did not observe a significant risk 54 associated with PM<sub>2.5</sub>. According to a recent review article, the risk of incident MI could be increased by 8% (95% CI: -1-18%) per 10-ug/m<sup>3</sup> increase in long-term PM<sub>2.5</sub> exposure, but the 55 56 association is evidently weaker as compared to other cardiovascular end points (Alexeeff et al., 57 2021). In addition, most of the studies used data from cohorts that were not representative of the 58 general population and mostly involving the participants over middle age, such as the 59 EuroTARGET cohort (Gandini et al., 2018) and the Nurses' Health Study cohort (Puett et al., 60 2009). Seldom has the effect modification by age been explored in the general population, which, 61 however, is important for understanding the role of air pollution in MI occurrence at earlier ages 62 and potentially different risk profiles among age groups.

Furthermore, concerns are raised in terms of the lack of causal interpretability in the existing literature. In contrast to randomized controlled trials (RCTs) that balance covariates across treatment classes via randomization, observational studies are the most powerful method available to study the health effect of air pollution, since randomization to exposure is not possible. Standard observational studies control for potential confounders in the outcome regression, but are often criticized for the potential for omitted confounders. Several alternative causal modeling approaches have been applied in air pollution epidemiology to attempt to mimic randomized trials

4

70 (Reich et al., 2021). The generalized propensity score is one example for investigating the health 71 effects of continuous pollution exposure. It can remove the confounding bias by measured 72 covariates and give population marginal effect estimates by balancing the covariates among 73 different exposure levels (Schwartz et al., 2021; Wei et al., 2020). Nevertheless, the propensity 74 score method is not able to control for the unmeasured confounders. The difference-in-differences 75 (DID) is a classical quasi-experimental estimator in econometrics designed to control for 76 unmeasured variables (Ashenfelter and Card, 1985; McEwan, 2010). To date, several studies have 77 applied the DID approach to evaluate the effects of time-varying PM<sub>2.5</sub> exposure and industrial air 78 pollution on mortality (Leogrande et al., 2019; J. Schwartz et al., 2021; Wang et al., 2016; Yitshak-79 Sade et al., 2019). This modeling design only examines variations in exposure and outcome within 80 geographic locations (in our case ZIP codes), thereby removing any confounding by measured or 81 unmeasured confounders that differ between ZIP codes. It controls for unmeasured confounders 82 that vary over time similarly in all ZIP codes using indicator variables for each calendar year. 83 Measured covariates are controlled for as usual. Therefore, the model framework provides a robust 84 tool to add control for many unmeasured confounders to traditional observational epidemiology.

Using the DID modeling approach, we investigated the effects of annual ambient PM<sub>2.5</sub> on the hospital admission rate for MI in residents of all ages across ten states of the United States during 2002-2016. Hospitalization data were accessed from the Health Cost and Utilization Project State Inpatient Databases. Furthermore, we examined whether age could modify the relationships. We also explored the effect modification by community socioeconomic status (SES) indicators and individual comorbidities.

91 2. Methods

### 92 2.1. Assessment of air pollution

5

93 Daily predictions of ambient fine particulate matter ( $PM_{2.5}$ ), nitrogen dioxide ( $NO_2$ ), and 94 ozone (O<sub>3</sub>) were derived from an ensemble of machine learning algorithms using geographically 95 weighted regression at a resolution of  $1 \text{ km} \times 1 \text{ km}$  in the contiguous United States. The prediction 96 model involved critical environmental measures from air monitoring data, meteorological 97 conditions, chemical transport model simulations, land-use features, and satellite remote sensing 98 data of aerosol optical depth, and the model was validated with a 10-fold cross-validation (Di et 99 al., 2020, 2019). We computed the exposure for each ZIP code and each year by averaging the grid 100 cell predictions whose centroids were inside the ZIP code polygons or assigning the nearest grid 101 cell predictions for the ZIP codes that do not have polygon representations, and then linked the 102 exposure data to participants based on their residential ZIP code and admission year. To explore 103 the potential lag effects of air pollution on MI hospitalizations, we used moving averages of air 104 pollution data in the admission year and one year before.

### 105 2.2. Outcome measurement

The inpatient care records used in this study were obtained from the Healthcare Cost and Utilization Program (HCUP) State Inpatient Database (SID) (HCUP Databases, 2021; Qiu et al., 2023, 2022; Steiner et al., 2002). The database has a 97% completeness of all U.S. community hospital discharges. More details about the inpatient databases and data availability are provided in the Supplement.

We analyzed the hospital admission data in the following listed states and years: Arizona (AZ), Michigan (MI), North Carolina (NC), New York (NY), Rhode Island (RI), Washington (WA), New Jersey (NJ) from 2002 to 2016, Maryland (MD, 2009-2016), Georgia (GA, 2010-2016), and Wisconsin (WI, 2012-2016). The inconsistent inclusion of the study years across these states was based on data availability from the States. The hospitalizations for principal diagnosis

6

of MI were identified with reference to the International Classification of Diseases (ICD) codes(ICD-9: 410; ICD-10: I21).

118 *2.3. Covariates* 

119 Demographic information for each participant comprising age, sex, race, admission year, 120 state, and ZIP code of residence for that year were obtained from the SIDs. Other covariates 121 included in the models were selected based on our prior knowledge, such as ambient seasonal 122 temperature, and area-level SES and behavioral factors. The SES data at the ZIP code tabulation-123 area level (ZCTA) were downloaded from the U.S. Census for 2002 Summary File 3 and 2011 124 Summary File 1 (Bureau, U. S. C., 2011, 2002). They were interpolated annually between census 125 years, and further extrapolated using the American Community Surveys after 2010 (Bureau, U. S. 126 C., 2020). They included the percent of female residents, percent of black residents, median 127 household income, percent of residents having high school education or less, and population 128 density. The percent of residents over the age of 65 years living in poverty was additionally used 129 as a proxy for SES. We defined the ZIP codes where the percent of persons  $\geq 65$  years old living 130 in poverty was within the lowest 25<sup>th</sup> percentile among all areas (i.e., <7%) as low-level poverty 131 areas and other ZIP codes as poorer areas. We also linked in the county-level yearly percentage of 132 residents who ever smoked and mean body mass index (BMI) from the Centers for Disease Control 133 and Prevention (CDC) Behavioral Risk Factor Surveillance System (BRFSS) (Bureau, U. S. C., 134 2020), which were converted to ZCTA-level based on the county and updated annually. Ambient 135 daily surface temperature and daily average surface pressure at a 12 km  $\times$  12 km resolution were 136 obtained from the National Aeronautics and Space Administration's Land Data Assimilation 137 System (NLDAS-2). In light of the seasonal effect of temperature on cardiovascular health (Shi et 138 al., 2015; Stewart et al., 2017), we generated two metrics of temperature, i.e., the average

7

temperature in the warm months (April-September) and the average temperature in the cold months (October-March). The ambient annual average level of daily maximum relative humidity at a 4 km × 4 km resolution was obtained from the gridMET dataset (Abatzoglou, 2013). Temperature, surface pressure, and relative humidity data were later aggregated to ZIP codes and years.

### 144 2.4. Comorbidity data

145 In this study, we considered several coexisting medical conditions that are likely to impact 146 the vulnerability of MI patients exposed to  $PM_{2.5}$  in the secondary analysis, which included: 147 hypertension (Yang et al., 2022), diabetes mellitus (DM), chronic obstructive pulmonary disease 148 (COPD) (Croft et al., 2022; Rich et al., 2010; Wynands et al., 2022), renal failure (Bo et al., 2021), 149 iron deficiency anemias (IDA) (Pereira and Sarnak, 2003), obesity (Weichenthal et al., 2014), 150 peripheral vascular disorders (PVD) (Bauersachs et al., 2019; Gwon et al., 2021), depression 151 (Gładka et al., 2018), other neurological disorders (Shi et al., 2020), and psychoses (Larsen and 152 Christenfeld, 2009; Qiu et al., 2023, 2022). The comorbidity data was assigned to the MI inpatients 153 using the AHRQ comorbidity software and stored in the Disease Severity Measures file as a part 154 of SIDs. The comorbidity data are available from 2005 to 2015. The identification of comorbidities 155 is based on International Classification of Diseases, 9th revision, Clinical Modification (ICD-9-156 CM) diagnoses and the Diagnosis-Related Group (DRG) in effect on the discharge date.

#### 157 2.5. Statistical analysis

We used the DID approach to estimate the relationship between long-term exposure to ambient  $PM_{2.5}$  and the incidence of hospital admissions for MI. The analysis was limited to the ZIP code-year combinations with a population of more than 100 to reduce the noise from lowpopulation areas and increase the analytical power (Qiu et al., 2022). First, we calculated the

8

annual aggregated counts of hospital admissions for MI for each ZIP code, year, and age group (034, 35-54, 55-64, 65-74, ≥75 years). We fitted a quasi-Poisson regression to account for the
overdispersion of MI hospitalization counts. The equation is given below:

165 
$$\ln(E(Y_{z,t,k})) = \beta_0 + \beta_1 P M_{2.5\,z,t} + \beta_2 Z_z + \beta_3 U_t + \beta_4 I_k + \beta_5 W_{z,t} + \ln(P_{z,t,k})$$

166 where  $Y_{z,t,k}$  represents the aggregated count of hospitalizations for MI in ZIP code z, year 167 t and age group k;  $PM_{2.5 z,t}$  represents the mean ambient  $PM_{2.5}$  concentration for the same stratum of ZIP code z and year t;  $Z_z$  is a dummy variable for ZIP code which captures all time-invariant or 168 slowly changing variables that vary across ZIP code areas, measured or unmeasured;  $U_t$  is a 169 170 dummy variable for year t which represents the time-varying variables whose temporal variation 171 is similar across ZIP code areas;  $I_k$  is a dummy variable for age groups;  $W_{z,t}$  represents the variables that may vary differently over time across ZIP code areas;  $\ln (P_{z,t,k})$  is an offset term 172 173 that represents the natural log of the population in ZIP code z, year t and age group k.

The essence of the DID method is to account for all differences across ZIP codes using indicator variables, and only examine the within ZIP code variation in exposure and outcome. This controls for all slowly changing covariates, measured or unmeasured, that vary across ZIP codes. Omitted confounders that vary similarly over time across ZIP codes are controlled using indicator variables for year. Time-varying confounders that vary over time differently across ZIP codes must be measured and controlled as usual. More statistical details can be found elsewhere (Wang et al., 2016).

In our DID model, the spatial confounding was controlled for by fitting individual intercepts for each ZIP code. The time trends were controlled by using an indicator for each year., We controlled for time-varying estimates of average warm-season and cold-season temperatures, percentage of female residents, percentage of black residents, median household income, percent

9

of education level under high school, population density, percentage of residents who ever smoked, percentage aged  $\geq 65$  years old living in poverty, and mean BMI in this paper. We assumed no other covariates that are both correlated with exposure and outcome can change differentially over time across ZIP codes were left out other than these adjusted variables. In the ecological design, the individual-level risk factors that displayed temporal variabilities were unrelated to the exposure and therefore would not confound the association. Therefore, the estimates given by the model can reflect the changes in MI morbidity attributable to the changes in ambient PM<sub>2.5</sub>.

To test the robustness of the effect estimate, we further controlled for surface pressure and relative humidity, respectively. We also constructed two- and multi-pollutant models to control for potential confounding by NO<sub>2</sub>, and O<sub>3</sub>. Moreover, we explored the variation in the effect estimates across age subgroups by including an interaction term for an indicator of age subgroups and the PM<sub>2.5</sub> term. We additionally examined the potential effect modification by some typical area-level SES indicators that could illustrate the degree of urbanity, i.e., poverty level, medium household income, population density, and education level.

199 We performed two secondary analyses. First, we restricted the analysis to the residents who 200 lived in the areas with annual exposure consistently at or below the annual standard of  $12 \,\mu g/m^3$ 201 set by National Ambient Air Quality Standards (NAAQS) over the study period. For the above 202 analysis, the percent change in hospitalization rate for MI and their 95% confidence intervals (CIs) 203 per  $\mu g/m^3$  increase in long-term average PM<sub>2.5</sub> level were reported. Second, we employed the case-204 only analysis to examine whether the following individual-level comorbidities could be effect 205 modifiers: hypertension, DM, COPD, renal failure, IDA, obesity, PVD, depression, other 206 neurological disorders, and psychoses. The case-only method was suggested by Armstrong (2003) 207 to study modification by effectively time-invariant individual factors in research of time-varying

208 environmental factors to identify the susceptible groups. For each comorbidity type, we performed 209 a logistic regression model of the ambient  $PM_{2.5}$  on the comorbidity status adjusted for the time 210 trend among the MI inpatients. The exposure-modifier interaction odds ratio (OR) and 95% CIs 211 were reported.

- All analyses were performed using R software version 4.1.2. A two-sided p-value <0.05</li>
  was considered statistically significant.
- 214 **3. Results**

# 215 3.1. Characteristics of community conditions and MI hospitalizations

216 In total, we included 1,914,684 hospital admissions for MI from 8,106 ZIP codes (with a population over 100) in 10 U.S. states from 2002 to 2016. The air pollution, temperature, 217 218 demographic and SES characteristics averaged over the study period in these ZIP codes are 219 presented in Table 1. Over the study period, the average long-term annual  $PM_{2.5}$  concentration 220 was  $8.9\pm2.6 \,\mu\text{g/m}^3$  among all the ZIP code areas. The average temperatures in the warm season 221 and in the cold season were 19.0±3.7 and 4.7±4.8 °C, respectively. On average, female residents, 222 ever smokers, or residents who received high school education or less accounted for about half of 223 the ZIP code-level population. In addition, there was evidence of spatial heterogeneity in several 224 community-level demographic and SES covariates. For example, the population density differed 225 substantially across ZIP codes with its 10<sup>th</sup> and 90<sup>th</sup> percentiles of 9 and 2,062 persons/mi<sup>2</sup>, respectively. The 10<sup>th</sup> and 90<sup>th</sup> percentiles in the percentage of the low-education population (*section*) 226 227 school) ranged between 25.2% and 63.6%.

Table 2 shows the total and age subgroup-specific number of hospitalizations for MI and the corresponding average hospital admission rate among the included ZIP codes during the study period. Among all 1,914,684 MI cases, the highest hospital admission rate was occurring in

11

patients aged at least 75 years old, which accounted for 35.98% of total MI hospitalizations. The proportions of MI hospital admissions in 45-54, 55-64, and 65-74 age groups were lower than that in the oldest group, being 14.20, 21.92, and 22.51%, respectively. In contrast, the age group aged 0-34 years old only included 0.79% of total hospitalizations for MI. The annual admission rate in the general population across all ZIP code areas shows an increasing pattern with age group: The highest ZIP code-level annual hospitalization rate for MI of 10.62% was observed in the oldest age group ( $\geq 75$  years), while lower hospitalization rates were seen in younger age groups.

### 238 3.2. Main effects of long-term PM<sub>2.5</sub> exposures on MI risk

239 Table 3 shows the estimated effects of PM<sub>2.5</sub> on MI risk in the general population using 240 different models. If the model assumptions are met, these are causal estimates. Overall, we 241 observed a 1.35% (95% CI: 1.11-1.59%) increase in the hospitalization rate for every  $1-\mu g/m^3$ 242 increase in PM<sub>2.5</sub> exposure. The estimate was stable after further adjusting for surface pressure or 243 relative humidity. In the model further adjusted for  $NO_2$  or  $O_3$ , the adverse effects of  $PM_{2.5}$  on the 244 risk of MI were slightly attenuated, which corresponded to 1.23% (95% CI: 0.99-1.47%) and 1.18% 245 (95% CI: 0.94-1.42%) increase in hospital admission rate associated with 1 µg/m<sup>3</sup> increase in PM<sub>2.5</sub>, 246 respectively. In the model adjusted for  $NO_2$  and  $O_3$  simultaneously, the effect estimate was similar 247 and remained significant. Moreover, we also restricted the main analysis to the ZIP codes where 248 the annual PM<sub>2.5</sub> exposures were always below 12  $\mu$ g/m<sup>3</sup> over the study period. A higher increment 249 of 2.17% (95% CI: 1.79-2.56%) in hospitalization rate for MI was observed per 1-µg/m<sup>3</sup> increase 250 in the low-level  $PM_{2.5}$  exposures.

### 251 3.3. Heterogeneity of susceptibilities among age subgroups

Figure 1 and Table S1 show the variation in estimates for annual  $PM_{2.5}$  and hospitalization for MI across six age subgroups. In detail, the subgroups aged 0-34 and  $\geq$ 75 years appeared to have the highest relative risks for chronic exposures to  $PM_{2.5}$ . For every 1-µg/m<sup>3</sup> increase in annual PM<sub>2.5</sub>, there was a 2.40% (95% CI: 1.62-3.20%) and a 1.94% (95% CI: 1.69-2.20%) increase in the hospitalization rate in 0-34 years and  $\geq$ 75 age subgroups, respectively. Every 1-µg/m<sup>3</sup> increase in PM<sub>2.5</sub> was associated with an increment of 1.32% (95% CI: 0.93-1.70%) in the hospitalization rate for MI among the subpopulation aged 35-44 years old. In contrast, the lowest relative risks for PM<sub>2.5</sub> were observed in the subpopulations with ages 45-54, 55-64, and 65-74 years.

# 260 3.4 Effect modifications by area-level SES covariates

261 Figure 2 and Table S2 illustrate the potential effect modification by several area-level SES 262 covariates on the relationship between PM<sub>2.5</sub> and the risk of MI. We found significant effect 263 modification by area-level poverty ( $p_{\text{interaction}}=0.0497$ ). Specifically, the population living in the 264 poorer areas was more susceptible to long-term  $PM_{2.5}$  exposures as compared to those living in 265 lower-poverty areas. In addition, population density significantly modified the association (p 266 interaction < 0.0001). The result suggests that people living in more populated geographical areas 267 tended to have a higher risk of MI attributed to long-term PM<sub>2.5</sub> exposures. However, the 268 association was not modified by either median household income (*p* interaction=0.5190) or education 269 level ( $p_{\text{interaction}}=0.6640$ ), despite a tendency of the higher risk occurring in the population with 270 lower median household income and lower education level.

### 271 3.5. Effect modifications by individual comorbidities

We also investigated whether the association between long-term  $PM_{2.5}$  and hospitalizations for MI could differ by individual-level comorbidities using the data from 2005 to 2015. The results of the case-only analysis are shown in **Figure 3** and **Table S3**. The risk of MI due to chronic exposures to  $PM_{2.5}$  was significantly higher in individuals with most of the common comorbidities than in those without it: hypertension, DM, COPD, renal failure, IDA, obesity, PVD, other

13

277 neurological disorders, and psychoses. The strongest modifying effect was observed in individuals 278 with comorbid IDA ( $p_{\text{interaction}} < 0.0001$ ). The odds of hospitalizations for MI was increased by 5.6% 279 (95% CI: 5.3-5.8%) for each 1-µg/m<sup>3</sup> increase in PM<sub>2.5</sub> associated with comorbid psychoses. 280 However, we found that having depression may decrease the susceptibility to PM<sub>2.5</sub> exposures 281 (OR=0.995; 95% CI: 0.991-0.998).

### 282 **4. Discussion**

283 In this study, we used hospital inpatient records from HCUP SIDs covering all admissions 284 in ten states in the U.S. and performed a DID analysis to estimate the relationship between long-285 term exposure to ambient PM<sub>2.5</sub> and the risk of hospitalizations for MI. In the entire population, 286 we found that the incidence of MI would increase by 1.35% (95% CI: 1.11-1.59%) for every 1-287  $\mu g/m^3$  increase in long-term residential PM<sub>2.5</sub> exposure. The effect was robust to adjustment for 288 surface pressure, relative humidity and co-pollutants (i.e., NO<sub>2</sub> and/or O<sub>3</sub>). Effects remained and 289 were stronger when limited to locations that never exceeded U.S. EPA's air quality standards, were 290 larger in poorer neighborhoods, and were increased in persons with chronic conditions most of 291 which are more common in minorities. A key aspect of this analysis is that the population studied 292 is the total population of those states, and not the selected sample of most cohort studies. This 293 makes the findings more generally applicable and the coefficients more useful for calculating 294 attributable risks. The findings that people with iron deficiency anemia, psychosis, and renal 295 failure were more susceptible to the effects of PM<sub>2.5</sub> on MI are novel, and the other effect 296 modification (hypertension, diabetes, COPD) findings add to a thin literature.

Less attention has been paid to the long-term effects of  $PM_{2.5}$  as compared to their acute effects (Cai et al., 2016; Zhu et al., 2021); however, an additional 10% risk is estimated for longterm exposure than for short-term exposures to  $PM_{2.5}$ , partially due to deteriorated cardiometabolic

14

300 conditions over time (Rajagopalan et al., 2018). Therefore, our research supplements the limited 301 evidence in this regard. The risk of hospitalization for MI was slightly attenuated when NO<sub>2</sub> and/or 302 O<sub>3</sub> were accounted for as confounding factors. It is important to note that NO<sub>2</sub> and O<sub>3</sub> are 303 associated with certain components of PM2.5. In particular, NO2 is associated with secondary 304 organic particles and O<sub>3</sub> with secondary inorganic (sulfate and nitrate) particles. Hence controlling 305 for them may remove the effects of important PM components. Indeed a personal exposure study 306 reported that ambient NO<sub>2</sub> and O<sub>3</sub> were not associated with personal exposure to NO<sub>2</sub> and O<sub>3</sub> but 307 were associated with personal exposure to PM<sub>2.5</sub> (Valli, 2001). This suggests that three pollutant 308 models risk biasing downward the PM<sub>2.5</sub> effect as the gases may not be confounders. This change 309 may also be due to the potential positive association for  $NO_2$  with MI (Roswall et al., 2017). The 310 possible confounding by  $O_3$  might be explained by its potential to amplify cardiovascular effects 311 (Malik et al., 2019; Weichenthal et al., 2017), which, however, remains unclear because of its 312 complex dynamics and interaction with other typical air pollutants that may vary with seasons (Ito 313 et al., 2007; Mustafić et al., 2012).

314 Moreover, we observed that the increased risk was stronger at or below annual  $PM_{2.5}$ concentrations of 12  $\mu$ g/m<sup>3</sup> which is an indication of a curvilinear concentration-response function 315 316 (CRF). Such a CRF has been noted previously. For example, Vodonos et al. (2018) in a meta-317 analysis of 54 cohort studies reported a similar shape in the association of both all-cause mortality 318 and cardiovascular mortality. The Global Exposure Mortality Model study reported a similar 319 finding, with a steeper slope for ischemic heart disease mortality below 12  $\mu$ g/m<sup>3</sup> (Burnett et al., 2018). Possible reasons for such a pattern include adaptation, different chemical compositions of 320 321 particles in locations with lower exposure, a different ratio of surface area to mass in higher 322 concentration locations where particles tend to be larger due to agglomeration, etc. Overall, our

15

results highlight that there is a significant health threat even under low-level long-term exposuresand that pollution control measures should be prioritized to protect public health.

325 Our main finding is in accordance with some recent evidence. For example, Danesh Yazdi 326 et al. (2019) observed a 2.6% (95% CI: 2.4-2.8%) increase in the hazard of first hospital admission 327 for MI associated with each 1 µg/m<sup>3</sup> increase in average annual PM<sub>2.5</sub> concentration among 328 Medicare participants over 65 years or older in seven southeastern states in the U.S. A study 329 targeted the entire adult population in Ontario, Canada reported an  $\sim 3\%$  increase in the risk of MI 330 incidence per  $\mu g/m^3$  increase in long-term exposure to PM<sub>2.5</sub> (Chen et al., 2020). Among 118,229 331 individuals in a large-scale and population-based Chinese cohort, an increment of 10 µg/m<sup>3</sup> 332 increase in PM<sub>2.5</sub> was associated with a higher risk of acute MI (HR=1.28; 95% CI, 1.19-1.39) (Li 333 et al., 2020). Hystad et al. (2020) found that a hazard ratio of 1.03 (95% CI: 1.00-1.05) per 10 334  $\mu g/m^3$  increase in long-term ambient PM<sub>2.5</sub> and a population-attributable fraction of 8.4% (95%) 335 CI: 0.0-15.4%) for MI in the Prospective Urban and Rural Epidemiology cohort comprising 336 157,436 adults from 21 countries. Nevertheless, some researchers reported null associations of 337 long-term PM<sub>2.5</sub> with MI (Cesaroni et al., 2014; Lipsett et al., 2011; Miller et al., 2007). The 338 nationwide Danish Nurse Cohort Study reported a higher risk of fatal MI induced by PM<sub>2.5</sub>, but 339 did not suggest any association with overall incident MI (Cramer et al., 2020). Aside from the 340 statistical approaches and study period, the differential magnitudes of the effect among the studies 341 may be due to differences in demographic structure, prevailing concentrations, since the 342 association appears curvilinear but linear models were fit, physiological characteristics (e.g., 343 medical history, comorbidities), or SES factors (e.g., poverty, healthcare accessibility) of the study 344 population. Diverse sources and constituents of ambient PM<sub>2.5</sub> across different regions can also 345 explain the difference in the toxicity and the triggering of MI (Rich et al., 2013).

16

346 Multiple pathways by which PM<sub>2.5</sub> can affect the myocardium have been heavily 347 elaborated, which are linked to oxidative stress, endothelial dysfunction, systemic immunity 348 response, systemic inflammatory response, and autonomic nervous system injuries (Langrish et al., 349 2012; Rajagopalan et al., 2018; Wang et al., 2013; Wang et al., 2016; Zhao et al., 2013); these 350 physiological responses could further contribute to the development of atherosclerosis, the widely 351 believed main trigger and underlying disease process of MI onset (Palasubramaniam et al., 2019). 352 MI progression could be accelerated by disturbed cardiac mitochondrial function and dynamics 353 followed by impaired redox metabolism and inflammation in the lung due to chronic exposure to 354 PM<sub>2.5</sub> inhalation (Marchini et al., 2022). More pathological evidence of specific molecular 355 signaling pathways is also emerging. For example, toxicological experiments revealed that the 356 down-regulation of microRNA-205 induced by PM<sub>2.5</sub> could stimulate myocardial inflammation 357 and cardiac dysfunction (Feng et al., 2020). Hu et al. (2021) demonstrated that NLRP3 358 inflammasome activation is responsible for PM2.5-induced endothelial cell dysfunction. These 359 uncovered biological mechanisms well support our findings.

360 We provide novel evidence into the health disparities across age groups in terms of the 361 relationship under investigation. Our evidence of high sensitivity in the oldest age group is in line 362 with prior studies (Bai et al., 2019; Li et al., 2020). It could result from reduced physiological, 363 metabolic and compensatory processes, as well as the heavier health burden of cardiopulmonary 364 disorders in the elderly (Shumake et al., 2013). It might be also attributed to declined antioxidant 365 defense ability of the respiratory tract lining fluid in the elderly population (Kelly et al., 2003). We 366 also found a surprisingly high susceptibility in young individuals, which has been seldom explored 367 or discussed before. A similar finding was suggested in a Canadian study where the researchers 368 observed a strong association of MI incidence in relation to PM<sub>2.5</sub> between 35-44 years-old adults

17

369 and 75-85 years-old adults (Bai et al., 2019). In comparison, we extended the scope to a large 370 sample size of multiple age groups and found that the susceptibility in individuals aged 0-34 years 371 was higher than that in the middle-aged groups and slightly higher than that in those aged >75. It 372 is important to note that the baseline risk in the younger individuals was two orders of magnitude 373 lower than in the oldest category, so a high relative risk is still a small attributable risk. These 374 findings might be partially due to the longer time spent outdoors by younger people than by elder 375 people. It's also possible that a potentially immature immune system in adolescents may increase 376 their vulnerability (Frcpc et al., 2006). More importantly, while the traditional risk factors for MI 377 such as underlying atherosclerosis are less common in younger populations, their unique risk 378 profile is linked to behavioral and psychosocial factors (e.g., recreational drug use) which may add 379 to the risk of cardiovascular events (Gulati et al., 2020). Abundant evidence concludes cocaine use 380 is a strong risk factor for MI (DeFilippis et al., 2018; Phillips et al., 2009; Schwartz et al., 2010). 381 These risk factors may convey susceptibility to these individuals to the inflammatory response 382 induced by  $PM_{2.5}$ . On the other hand, if cocaine usage is correlated with  $PM_{2.5}$  exposure, it may be 383 a confounder.

384 In addition, we observed that population living in poorer areas or densely populated areas 385 may be more susceptible to the risk of MI attributed to chronic exposures to PM<sub>2.5</sub>. The evidence 386 of differential susceptibility to exposure in low-income areas, combined with the already 387 established higher exposure levels and poorer healthcare accessibility in socioeconomically 388 disadvantaged areas (Colmer et al., 2020; Wang et al., 2017) identifies this as a key environmental 389 justice issue. Although increased susceptibility to PM<sub>2.5</sub>-related health impacts accompanied by 390 SES disadvantage has been well recognized (Bravo et al., 2016; Chi et al., 2016; Xu et al., 2020), 391 the effect modification by neighboring SES factors in the association of MI risk with long-term

18

392  $PM_{2.5}$  exposures is little discussed, and present conclusions on this topic are mixed. Several studies 393 did not identify significant differences across levels of neighborhood-level sociodemographic 394 characteristics (Hystad et al., 2020; Madrigano et al., 2013; Weaver et al., 2019). In contrast, the 395 evidence of effect modification by annual household income was suggested in long-term adult 396 residents of Ontario (Bai et al., 2019), which differed from our study. Our present finding can 397 provide a novel understanding of potential susceptible subpopulations and facilitate clarification 398 of social drivers for substantial geographic disparities in MI across the U.S. (Camacho et al., 2022; 399 Yu et al., 2021).

400 In line with prior epidemiological studies, we identified a statistically elevated risk of MI 401 associated with long-term PM<sub>2.5</sub> exposure among patients with diabetes than those without 402 (Cramer et al., 2020; Hart et al., 2015; McGuinn et al., 2016). We also found that hypertension, 403 COPD, renal failure, IDA, obesity, PVD, and unfavorable neurological conditions may increase 404 the vulnerability to the risk of MI in relation to PM<sub>2.5</sub>, which has not been widely appreciated or 405 documented in the past. Higher insulin resistance and oxidative stress may enhance the alteration 406 of ventricular repolarization and systemic inflammation triggered by PM<sub>2.5</sub> and thus exacerbate 407 myocardial vulnerability to arrhythmias (Schneider et al., 2010), which might provide a possible 408 explanation for our conclusion. The strongest effect of IDA may be due to stimulated oxidative 409 stress, decreased blood supply, myocardial hypoxia, and the close interplay of the outcomes 410 (Inserte et al., 2021; Padda et al., 2021). Additionally, we assume the intake of psychotropic 411 medications may exacerbate the susceptibility since they are known to cause undesirable 412 cardiometabolic effects (Abosi et al., 2018; Mohamed et al., 2019). A much lower chance of 413 adherence to cardiovascular disease medication and healthcare-seeking behaviors to prevent 414 physical illness among many psychiatric patients (especially those with severe mental illness)

19

415 might also contribute to an increased risk of MI events (Druss et al., 2001; Newcomer and 416 Hennekens, 2007). However, comorbid depression, which showed a protective effect, remains 417 unclear. Further investigations on the specific impacts and mechanisms of comorbidities are 418 warranted.

419 There are multiple strengths in the present study. To begin with, we used high coverage 420 (97%) medical records across multiple U.S. states over a long period, which enabled the statistical 421 power to be sufficient and expanded the generalizability of our findings. Furthermore, in the DID 422 modeling framework, we eliminated confounding by variables that change slowly over time and 423 minimized the confounding by the variables that showed parallel variations in time across area 424 units, thus overcoming the limitations of data availability by design. We also fully adjusted for 425 multiple area-level demographic, SES, and behavioral covariates to relax the assumptions. 426 Additionally, in light of underappreciated cardiovascular risk among younger individuals, we 427 performed the analysis with fine divisions of age and revealed the strong susceptibility in young 428 people. Additionally, we determined multiple common individual comorbid conditions as potential 429 susceptibility factors. With this investigation, we aimed to raise the public health concern towards 430 sensitive groups and implement more targeted preventative measures both in pollution control and 431 healthcare services.

However, some limitations should also be acknowledged. First, despite high resolution of the air pollution predictions, the PM<sub>2.5</sub> estimates only represent the ambient exposure level within a specific ZIP code area and year but were not based on personal exposures which are influenced by activity patterns, air exchange rates, second-hand tobacco smoke, and other factors that lead to them differing, often substantially, from neighborhood exposure. However, Weisskopf and Webster (2017) have pointed out that personal exposure is correlated with many potential

20

438 confounders (e.g. stress while driving) that are not associated with outdoor concentration, and 439 hence outdoor exposures can act as instrumental variables for the personal exposure. Further, the 440 National Human Activity Pattern Survey in the U.S. reported that U.S. adults spent 69% of their 441 time at home and 8% of the time immediately outside their home (Klepeis et al., 2001), making it 442 reasonable to use neighborhood ambient pollution to capture their exposure to ambient pollution. 443 Nevertheless, our model clearly does not capture personal exposure. While there is likely a 444 considerable difference between the personal exposure of individuals living in a neighborhood and 445 the ambient concentration in that neighborhood, the difference between the personal and ambient 446 is likely to be mostly Berksonian exposure error since it represents individual variation around a 447 group mean. As such, it should not produce much bias in the effect estimates in an epidemiology 448 study. A recent simulation study showed that even in the presence of substantial Berksonian 449 exposure error, little bias would be found in an epidemiology study (Wei et al., 2022). Second, the 450 DID approach should be better understood as a more robust approach to confounding control that 451 addresses some omitted confounders rather than as an approach that addresses all of them. 452 However, since we used indicator variables for every ZIP code, our analysis is only examining 453 year-to-year exposure within ZIP code, and hence ZIP code differences between e.g. high and low 454 SES ZIP codes are controlled by design. Within ZIP code, we have controlled for time-varying 455 SES variables, but we acknowledge we were not able to control for all. For example, we did not 456 have data on occupation. However, since the DID approach eliminates any contrasts between ZIP 457 codes, and the exposure assigned to all individuals within a ZIP code is the same, it is unlikely that 458 such factors could be associated with exposure. Third, differential exposure histories and other 459 risk factors such as diets across areas are overlooked because of the unknown human migration 460 patterns in this study, which may have resulted in a wrong recognition of the true cause of MI

21

hospitalizations. Finally, we aggregated the hospital admissions by strata defined by ZIP code,
year, and age, therefore, potentially low numbers in some age subpopulation in less populated
geographical areas could add noise to the analysis.

464 **5.** Conclusions

In this study, we found that long-term exposures to ambient PM<sub>2.5</sub> increased the risk of MI 465 466 hospitalization. The relationship remained significant after controlling for surface pressure, 467 relative humidity and concurrent exposures to  $NO_2$  and  $O_3$ . A more pronounced adverse 468 association was observed at or below the annual  $PM_{2.5}$  limit enacted by NAQQS. In addition, we 469 observed higher susceptibility to the PM<sub>2.5</sub>-related MI risk among young people and elderly people, 470 residents living in poorer or highly populated areas, and individuals with a comorbid burden. Our 471 study highlights the roles of air pollution being one important environmental risk factor for MI, 472 gives insights into the underlying susceptible subpopulations, and informs the future strategies to 473 control air pollution and mitigate related health disparities.

474

### 475 Acknowledgments

We want to offer special thanks to U.S. Agency for Healthcare Research and Quality Health
Cost and Utilization Project (HCUP) for giving us access to the State Inpatient Databases (SIDs)
to conduct this study.

479 Sources of Funding

This work was made possible by U.S. Environmental Protection Agency (EPA) grant RD-835872. Its contents are solely the responsibility of the grantee and do not necessarily represent the official views of the U.S. EPA. Further, U.S. EPA does not endorse the purchase of any commercial products or services mentioned in the publication. This work was also supported by

484	National	Institutes	of	Health	(NIH)	grant	R01	ES032418-01	and	National	Institute	of
485	Environm	nental Heal	th S	ciences	(NIEHS	) grant	(P30 )	ES000002).				
486	Disclosu	res										
487	Ν	one.										
488												
489												

- 1)]

# **Table 1.** Distribution of ZIP code-level annual PM<sub>2.5</sub> and covariates from 2002 through 2016.

ZIP code-level covariates*	Mean (SD)	Percentile						
		10th	25th	50th	75th	90th		
Annual PM <sub>2.5</sub> (µg/m <sup>3</sup> )	8.9 (2.6)	5.4	7.2	8.9	10.5	12.4		
Annual NO <sub>2</sub> (ppb)	17.6 (10.0)	7.4	10.1	14.8	23.3	31.9		
Annual O <sub>3</sub> (ppb)	39.0 (3.9)	34.7	37.3	38.9	40.5	44.0		
Warm season temperature	10.0 (2.7)	15.0	16.2	10 1	21.6	22.0		
(°C)	19.0 (3.7)	15.0	10.5	18.1	21.0	23.9		
Cold season temperature (°C)	4.7 (4.8)	-0.8	0.9	4.3	7.8	11.1		
Surface pressure (Pa)	95,668 (2,980)	94,187	94,267	94,330	98,118	100,097		
Relative humidity (%)	84.8 (10.6)	74.9	83.8	87.6	90.4	92.9		
Percent female (%)	50.3 (3.6)	47.7	49.4	50.6	51.8	53.0		

						2
Percent black (%)	9.9 (17.1)	0.2	0.5	2.2	10.5	31.6
Median household income (\$)	54,044 (23,198)	32,364	39,049	48,130	63,069	83,717
Percent ≥65 below poverty	13.2 (8.5)	4.33	7.06	11.53	17.06	24.17
Population density (persons/mi <sup>2</sup> )	1,109 (3,866)	9	23	96	753	2,062
Percent education ≤high school	46.0 (14.6)	25.2	35.9	47.7	56.9	63.6
Ever smokers (%)	48.0 (7.5)	39.0	43.3	47.6	52.5	57.4
Mean BMI (kg/m <sup>2</sup> )	27.6 (1.1)	26.3	26.9	27.5	28.2	29.0

496

\* Air pollutant concentration data were the moving averages of the year and the previous year.

497

498

499 **Table 2.** Summary of total and age-specific hospitalizations for MI from 2002 through 2016.

	Number of bespitalizations (9/)*	Annual rate of
	Number of hospitalizations (%)	hospitalization (‰) <sup>**</sup>
Total	1,914,684	1.40
Age (years)		
0-34	15,215 (0.79)	0.03
35-44	87,893 (4.59)	0.68
45-54	271,937 (14.20)	1.91
55-64	419,733 (21.92)	3.55
65-74	431,071 (22.51)	5.70

~	
2	4
-	•

- 501 \*\*The annual rate of hospitalization was averaged across ZIP code areas.
- 502
- 503 **Table 3.** Percent change in hospitalization rate for MI per 1-µg/m<sup>3</sup> increase in long-term
- 504 exposure to PM<sub>2.5</sub> in the main model, in models adjusted for surface pressure (SP) or relative

505 humidity (RH), in models adjusted for NO<sub>2</sub> and/or O<sub>3</sub>, and in the low-level exposure analysis.

	Percent change in MI
Models	hospitalization rate (95%
	CI)
Main model	1.35 (1.11, 1.59)
Adjusted for SP	1.34 (1.10, 1.58)
Adjusted for RH	1.35 (1.10, 1.59)
Adjusted for NO <sub>2</sub>	1.23 (0.99, 1.47)
Adjusted for O <sub>3</sub>	1.18 (0.94, 1.42)
Adjusted for NO <sub>2</sub> and O <sub>3</sub>	1.18 (0.94, 1.43)
Low-level exposures to	2 17 (1 70 2 56)
PM <sub>2.5</sub> *	2.17 (1.79, 2.30)

<sup>\*</sup>The analysis for low-level exposures to  $PM_{2.5}$  was restricted to the areas where the population

507 was continuously exposed to  $PM_{2.5}$  concentrations at or below 12  $\mu$ g/m<sup>3</sup> over 2002-2016.

508





511 **Figure 1.** Percent change in hospitalization rate for MI per 1-µg/m<sup>3</sup> increase in long-term



513

514



516 Figure 2. Percent change in hospitalization rate for MI per  $1-\mu g/m^3$  increase in long-term exposure

517 to PM<sub>2.5</sub> at the different percentiles of potential SES modifiers.

- \* indicates the statistical significance of the modifier ( $p_{\text{interaction}} < 0.05$ ). For the percent of residents  $\geq 65$  years old living in poverty, the data below the 25<sup>th</sup> percentile (i.e., <7% percent; defined as low-level poverty areas) compared to the data at or above the 25<sup>th</sup> percentile (i.e.,  $\geq 7\%$  percent; defined as poorer areas). For median household income, population density, and percent education level  $\leq$ high school, the effects of PM<sub>2.5</sub> were compared between the 10<sup>th</sup> and 90<sup>th</sup> of the modifier level.
- 524
- 525







- 528 exposure to PM<sub>2.5</sub> comparing individuals with and without comorbidities.
- 529 DM, diabetes mellitus; COPD, chronic obstructive pulmonary disease; IDA, iron deficiency
- anemia; PVD, peripheral vascular disorders.

27

### 532 **References:**

- 533 Abatzoglou, J.T., 2013. Development of gridded surface meteorological data for ecological
- applications and modelling. Int. J. Climatol. 33, 121–131.
- 535 https://doi.org/10.1002/joc.3413
- 536 Abosi, O., Lopes, S., Schmitz, S., Fiedorowicz, J.G., 2018. Cardiometabolic effects of
- 537 psychotropic medications. Hormone Molecular Biology and Clinical Investigation 36.
  538 https://doi.org/10.1515/hmbci-2017-0065
- 539 Alexeeff, S.E., Liao, N.S., Liu, X., Van Den Eeden, S.K., Sidney, S., 2021. Long-Term PM 2.5
- 540 Exposure and Risks of Ischemic Heart Disease and Stroke Events: Review and Meta-

541 Analysis. JAHA 10, e016890. https://doi.org/10.1161/JAHA.120.016890

- 542 Armstrong, B.G., 2003. Fixed Factors that Modify the Effects of Time-Varying Factors:
- 543 Applying the Case-Only Approach: Epidemiology 14, 467–472.
- 544 https://doi.org/10.1097/01.ede.0000071408.39011.99
- 545 Arora, S., Stouffer, G.A., Kucharska-Newton, A.M., Qamar, A., Vaduganathan, M., Pandey, A.,
- 546 Porterfield, D., Blankstein, R., Rosamond, W.D., Bhatt, D.L., Caughey, M.C., 2019.
- 547 Twenty Year Trends and Sex Differences in Young Adults Hospitalized With Acute
- 548 Myocardial Infarction: The ARIC Community Surveillance Study. Circulation 139,

549 1047–1056. https://doi.org/10.1161/CIRCULATIONAHA.118.037137

- Ashenfelter, O., Card, D., 1985. Using the Longitudinal Structure of Earnings to Estimate the
  Effect of Training Programs. MIT Press 67, 648–660.
- 552 Bai, L., Shin, S., Burnett, R.T., Kwong, J.C., Hystad, P., van Donkelaar, A., Goldberg, M.S.,
- 553 Lavigne, E., Copes, R., Martin, R.V., Kopp, A., Chen, H., 2019. Exposure to ambient air

554	pollution and the incidence of congestive heart failure and acute myocardial infarction: A
555	population-based study of 5.1 million Canadian adults living in Ontario. Environment
556	International 132, 105004. https://doi.org/10.1016/j.envint.2019.105004
557	Bauersachs, R., Zeymer, U., Brière, JB., Marre, C., Bowrin, K., Huelsebeck, M., 2019. Burden
558	of Coronary Artery Disease and Peripheral Artery Disease: A Literature Review.
559	Cardiovascular Therapeutics 2019, e8295054. https://doi.org/10.1155/2019/8295054
560	Bo, Y., Brook, J.R., Lin, C., Chang, L., Guo, C., Zeng, Y., Yu, Z., Tam, T., Lau, A.K.H., Lao,
561	X.Q., 2021. Reduced Ambient PM2.5 Was Associated with a Decreased Risk of Chronic
562	Kidney Disease: A Longitudinal Cohort Study. Environ. Sci. Technol. 55, 6876-6883.
563	https://doi.org/10.1021/acs.est.1c00552
564	Bravo, M.A., Son, J., de Freitas, C.U., Gouveia, N., Bell, M.L., 2016. Air pollution and mortality
565	in São Paulo, Brazil: Effects of multiple pollutants and analysis of susceptible
566	populations. J Expo Sci Environ Epidemiol 26, 150–161.
567	https://doi.org/10.1038/jes.2014.90
568	Bureau, U. S. C., 2020. Behavioral Risk Factor Surveillance System (2005–2019).
569	Bureau, U. S. C., 2011. Summary File 1 Dataset.
570	Bureau, U. S. C., 2002. Summary File 3 Dataset.
571	Burnett, R., Chen, H., Szyszkowicz, M., Fann, N., Hubbell, B., Pope, C.A., Apte, J.S., Brauer,
572	M., Cohen, A., Weichenthal, S., Coggins, J., Di, Q., Brunekreef, B., Frostad, J., Lim,
573	S.S., Kan, H., Walker, K.D., Thurston, G.D., Hayes, R.B., Lim, C.C., Turner, M.C.,
574	Jerrett, M., Krewski, D., Gapstur, S.M., Diver, W.R., Ostro, B., Goldberg, D., Crouse,
575	D.L., Martin, R.V., Peters, P., Pinault, L., Tjepkema, M., van Donkelaar, A., Villeneuve,
576	P.J., Miller, A.B., Yin, P., Zhou, M., Wang, L., Janssen, N.A.H., Marra, M., Atkinson,

577	R.W.,	Tsang.	H., 0	Duoc	Thach.	Т.,	Cannon.	J.B.,	Allen.	<b>R.T.</b>	Hart.	J.E.,	Laden.	F
• • • •				X		,					,,	· · _ · ,		

- 578 Cesaroni, G., Forastiere, F., Weinmayr, G., Jaensch, A., Nagel, G., Concin, H., Spadaro,
- 579 J.V., 2018. Global estimates of mortality associated with long-term exposure to outdoor
- 580 fine particulate matter. Proceedings of the National Academy of Sciences 115, 9592–
- 581 9597. https://doi.org/10.1073/pnas.1803222115
- 582 Cai, X., Li, Z., Scott, E.M., Li, X., Tang, M., 2016. Short-term effects of atmospheric particulate
- 583 matter on myocardial infarction: a cumulative meta-analysis. Environ Sci Pollut Res 23,
- 584 6139–6148. https://doi.org/10.1007/s11356-016-6186-3
- 585 Camacho, X., Nedkoff, L., Wright, F.L., Nghiem, N., Buajitti, E., Goldacre, R., Rosella, L.C.,
- 586 Seminog, O., Tan, E.J., Hayes, A., Hayen, A., Wilson, N., Blakely, T., Clarke, P., 2022.
- 587 Relative contribution of trends in myocardial infarction event rates and case fatality to
- 588 declines in mortality: an international comparative study of 1.95 million events in 80.4
- 589 million people in four countries. The Lancet Public Health 7, e229–e239.
- 590 https://doi.org/10.1016/S2468-2667(22)00006-8
- 591 Cesaroni, G., Forastiere, F., Stafoggia, M., Andersen, Z.J., Badaloni, C., Beelen, R., Caracciolo,
- 592 B., Faire, U. de, Erbel, R., Eriksen, K.T., Fratiglioni, L., Galassi, C., Hampel, R., Heier,
- 593 M., Hennig, F., Hilding, A., Hoffmann, B., Houthuijs, D., Jöckel, K.-H., Korek, M.,
- 594 Lanki, T., Leander, K., Magnusson, P.K.E., Migliore, E., Ostenson, C.-G., Overvad, K.,
- 595 Pedersen, N.L., J. J.P., Penell, J., Pershagen, G., Pyko, A., Raaschou-Nielsen, O., Ranzi,
- 596 A., Ricceri, F., Sacerdote, C., Salomaa, V., Swart, W., Turunen, A.W., Vineis, P.,
- 597 Weinmayr, G., Wolf, K., Hoogh, K. de, Hoek, G., Brunekreef, B., Peters, A., 2014. Long
- 598 term exposure to ambient air pollution and incidence of acute coronary events:

30

599	prospective cohort study and meta-analysis in 11 European cohorts from the ESCAPE
600	Project. BMJ 348, f7412. https://doi.org/10.1136/bmj.f7412

- 601 Chadwick Jayaraj, J., Davatyan, K., Subramanian, S.S., Priya, J., 2019. Epidemiology of
- 602 Myocardial Infarction, in: Pamukçu, B. (Ed.), Myocardial Infarction. IntechOpen.
- 603 https://doi.org/10.5772/intechopen.74768
- 604 Chen, H., Zhang, Z., van Donkelaar, A., Bai, L., Martin, R.V., Lavigne, E., Kwong, J.C.,
- Burnett, R.T., 2020. Understanding the Joint Impacts of Fine Particulate Matter
- 606 Concentration and Composition on the Incidence and Mortality of Cardiovascular
- 607 Disease: A Component-Adjusted Approach. Environ. Sci. Technol. 54, 4388–4399.
- 608 https://doi.org/10.1021/acs.est.9b06861
- 609 Chi, G., Hajat, A., Bird, C., Cullen, M., Griffin, B.A., Miller, K., Shih, R., Stefanick, M., Vedal,
- 610 S., Whitsel, E., Kaufman, J., 2016. Individual and Neighborhood Socioeconomic Status
- 611 and the Association between Air Pollution and Cardiovascular Disease. Environmental

612 Health Perspectives 124. https://doi.org/10.1289/EHP199

- 613 Colmer, J., Hardman, I., Shimshack, J., Voorheis, J., 2020. Disparities in PM <sub>2.5</sub> air pollution in
- the United States. Science 369, 575–578. https://doi.org/10.1126/science.aaz9353
- 615 Cramer, J., Jørgensen, J.T., Hoffmann, B., Loft, S., Bräuner, E.V., Prescott, E., Ketzel, M.,
- 616 Hertel, O., Brandt, J., Jensen, S.S., Backalarz, C., Simonsen, M.K., Andersen, Z.J., 2020.
- 617 Long-Term Exposure to Air Pollution and Incidence of Myocardial Infarction: A Danish
- 618 Nurse Cohort Study. Environ Health Perspect 128, 057003.
- 619 https://doi.org/10.1289/EHP5818

620	Croft, D.P., Utell, M.J., Rich, D.Q., 2022. Can Long-Term Residential Air Cleaning Reduce
621	Cardiovascular Morbidity in COPD Patients? Am J Respir Crit Care Med rccm.202211-
622	2138ED. https://doi.org/10.1164/rccm.202211-2138ED
623	Danesh Yazdi, M., Wang, Y., Di, Q., Zanobetti, A., Schwartz, J., 2019. Long-term exposure to
624	PM2.5 and ozone and hospital admissions of Medicare participants in the Southeast USA.
625	Environment International 130, 104879. https://doi.org/10.1016/j.envint.2019.05.073
626	DeFilippis, E.M., Singh, A., Divakaran, S., Gupta, A., Collins, B.L., Biery, D., Qamar, A.,
627	Fatima, A., Ramsis, M., Pipilas, D., Rajabi, R., Eng, M., Hainer, J., Klein, J., Januzzi,
628	J.L., Nasir, K., Di, C.M.F., Bhatt, D.L., Blankstein, R., 2018. Cocaine and Marijuana Use
629	Among Young Adults With Myocardial Infarction. Journal of the American College of
630	Cardiology 71, 2540-2551. https://doi.org/10.1016/j.jacc.2018.02.047
631	Di, Q., Amini, H., Shi, L., Kloog, I., Silvern, R., Kelly, J., Sabath, M.B., Choirat, C., Koutrakis,
632	P., Lyapustin, A., Wang, Y., Mickley, L.J., Schwartz, J., 2020. Assessing NO2
633	Concentration and Model Uncertainty with High Spatiotemporal Resolution across the
634	Contiguous United States Using Ensemble Model Averaging. Environ. Sci. Technol. 54,
635	1372–1384. https://doi.org/10.1021/acs.est.9b03358
636	Di, Q., Amini, H., Shi, L., Kloog, I., Silvern, R., Kelly, J., Sabath, M.B., Choirat, C., Koutrakis,
637	P., Lyapustin, A., Wang, Y., Mickley, L.J., Schwartz, J., 2019. An ensemble-based model
638	of PM2.5 concentration across the contiguous United States with high spatiotemporal
639	resolution. Environment International 130, 104909.
640	https://doi.org/10.1016/j.envint.2019.104909

641	Druss, B.G., Bradford, W.D., Rosenheck, R.A., Radford, M.J., Krumholz, H.M., 2001. Quality
642	of Medical Care and Excess Mortality in Older Patients With Mental Disorders. Arch
643	Gen Psychiatry 58, 565. https://doi.org/10.1001/archpsyc.58.6.565
644	Feng, L., Wei, J., Liang, S., Sun, Z., Duan, J., 2020. miR-205/IRAK2 signaling pathway is
645	associated with urban airborne PM2.5-induced myocardial toxicity. Nanotoxicology 14,
646	1198–1212. https://doi.org/10.1080/17435390.2020.1813824
647	Frepe, I.B., ChB, S.K.M., Osornio-Vargas, A.R., 2006. The effects of air pollution on the health
648	of children 11.
649	Gandini, M., Scarinzi, C., Bande, S., Berti, G., Carnà, P., Ciancarella, L., Costa, G., Demaria,
650	M., Ghigo, S., Piersanti, A., Rowinski, M., Spadea, T., Stroscia, M., Cadum, E., 2018.
651	Long term effect of air pollution on incident hospital admissions: Results from the Italian
652	Longitudinal Study within LIFE MED HISS project. Environment International 121,
653	1087–1097. https://doi.org/10.1016/j.envint.2018.10.020
654	Gładka, A., Rymaszewska, J., Zatoński, T., 2018. Impact of air pollution on depression and
655	suicide. Int J Occup Med Environ Health 31, 711–721.
656	https://doi.org/10.13075/ijomeh.1896.01277
657	Gulati, R., Behfar, A., Narula, J., Kanwar, A., Lerman, A., Cooper, L., Singh, M., 2020. Acute
658	Myocardial Infarction in Young Individuals. Mayo Clinic Proceedings 95, 136–156.
659	https://doi.org/10.1016/j.mayocp.2019.05.001
660	Gwon, J.G., Park, J.H., Kim, J.S., Seo, HM., 2021. Exposure to Long-Term Air Pollution and
661	Incidence of Peripheral Arterial Disease in the General Population: A National
662	Population-Based Retrospective Cohort Study (preprint). In Review.
663	https://doi.org/10.21203/rs.3.rs-506193/v1

664	Hart, J.E., Puett, R.C., Rexrode, K.M., Albert, C.M., Laden, F., 2015. Effect Modification of
665	Long-Term Air Pollution Exposures and the Risk of Incident Cardiovascular Disease in
666	US Women. JAHA 4, e002301. https://doi.org/10.1161/JAHA.115.002301
667	Hartiala, J., Breton, C.V., Tang, W.H.W., Lurmann, F., Hazen, S.L., Gilliland, F.D., Allayee, H.,
668	2016. Ambient Air Pollution Is Associated With the Severity of Coronary Atherosclerosis
669	and Incident Myocardial Infarction in Patients Undergoing Elective Cardiac Evaluation.
670	JAHA 5, e003947. https://doi.org/10.1161/JAHA.116.003947
671	HCUP Databases, 2021. Healthcare Cost and Utilization Project (HCUP).
672	Hu, T., Zhu, P., Liu, Y., Zhu, H., Geng, J., Wang, B., Yuan, G., Peng, Y., Xu, B., 2021. PM2.5
673	induces endothelial dysfunction via activating NLRP3 inflammasome. Environmental
674	Toxicology 36, 1886–1893. https://doi.org/10.1002/tox.23309
675	Hystad, P., Larkin, A., Rangarajan, S., AlHabib, K.F., Avezum, Á., Calik, K.B.T., Chifamba, J.,
676	Dans, A., Diaz, R., du Plessis, J.L., Gupta, R., Iqbal, R., Khatib, R., Kelishadi, R., Lanas,
677	F., Liu, Z., Lopez-Jaramillo, P., Nair, S., Poirier, P., Rahman, O., Rosengren, A., Swidan,
678	H., Tse, L.A., Wei, L., Wielgosz, A., Yeates, K., Yusoff, K., Zatoński, T., Burnett, R.,
679	Yusuf, S., Brauer, M., 2020. Associations of outdoor fine particulate air pollution and
680	cardiovascular disease in 157 436 individuals from 21 high-income, middle-income, and
681	low-income countries (PURE): a prospective cohort study. The Lancet Planetary Health
682	4, e235–e245. https://doi.org/10.1016/S2542-5196(20)30103-0
683	Inserte, J., Barrabés, J.A., Aluja, D., Otaegui, I., Bañeras, J., Castellote, L., Sánchez, A.,
684	Rodríguez-Palomares, J.F., Pineda, V., Miró-Casas, E., Milà, L., Lidón, RM., Sambola,
685	A., Valente, F., Rafecas, A., Ruiz-Meana, M., Rodríguez-Sinovas, A., Benito, B., Buera,
686	I., Delgado-Tomás, S., Beneítez, D., Ferreira-González, I., 2021. Implications of Iron

687	Deficiency in STEMI Patients and in a Murine Model of Myocardial Infarction. JACC
688	Basic Transl Sci 6, 567–580. https://doi.org/10.1016/j.jacbts.2021.05.004
689	Ito, K., Thurston, G.D., Silverman, R.A., 2007. Characterization of PM2.5, gaseous pollutants,
690	and meteorological interactions in the context of time-series health effects models. J
691	Expo Sci Environ Epidemiol 17, S45–S60. https://doi.org/10.1038/sj.jes.7500627
692	Kelly, F.J., Dunster, C., Mudway, I., 2003. Air pollution and the elderly: oxidant/antioxidant
693	issues worth consideration. European Respiratory Journal 21, 70s-75s.
694	https://doi.org/10.1183/09031936.03.00402903
695	Klepeis, N.E., Nelson, W.C., Ott, W.R., Robinson, J.P., Tsang, A.M., Switzer, P., Behar, J.V.,
696	Hern, S.C., Engelmann, W.H., 2001. The National Human Activity Pattern Survey
697	(NHAPS): a resource for assessing exposure to environmental pollutants. J Expo Sci
698	Environ Epidemiol 11, 231–252. https://doi.org/10.1038/sj.jea.7500165
699	Langrish, J.P., Bosson, J., Unosson, J., Muala, A., Newby, D.E., Mills, N.L., Blomberg, A.,
700	Sandström, T., 2012. Cardiovascular effects of particulate air pollution exposure: time
701	course and underlying mechanisms. Journal of Internal Medicine 272, 224–239.
702	https://doi.org/10.1111/j.1365-2796.2012.02566.x
703	Larsen, B.A., Christenfeld, N.J.S., 2009. Cardiovascular Disease and Psychiatric Comorbidity:
704	The Potential Role of Perseverative Cognition. Cardiovascular Psychiatry and Neurology
705	2009, 1-8. https://doi.org/10.1155/2009/791017
706	Leogrande, S., Alessandrini, E.R., Stafoggia, M., Morabito, A., Nocioni, A., Ancona, C.,
707	Bisceglia, L., Mataloni, F., Giua, R., Mincuzzi, A., Minerba, S., Spagnolo, S., Pastore, T.,
708	Tanzarella, A., Assennato, G., Forastiere, F., 2019. Industrial air pollution and mortality

709	in the Taranto area, Southern Italy: A difference-in-differences approach. Environment
710	International 132, 105030. https://doi.org/10.1016/j.envint.2019.105030
711	Li, J., Liu, F., Liang, F., Huang, K., Yang, X., Xiao, Q., Chen, J., Liu, X., Cao, J., Chen, S.,
712	Shen, C., Yu, L., Lu, F., Wu, Xianping, Zhao, L., Wu, Xigui, Li, Y., Hu, D., Huang, J.,
713	Liu, Y., Lu, X., Gu, D., 2020. Long-Term Effects of High Exposure to Ambient Fine
714	Particulate Matter on Coronary Heart Disease Incidence: A Population-Based Chinese
715	Cohort Study. Environ. Sci. Technol. 54, 6812-6821.
716	https://doi.org/10.1021/acs.est.9b06663
717	Lipsett, M.J., Ostro, B.D., Reynolds, P., Goldberg, D., Hertz, A., Jerrett, M., Smith, D.F., Garcia,
718	C., Chang, E.T., Bernstein, L., 2011. Long-Term Exposure to Air Pollution and
719	Cardiorespiratory Disease in the California Teachers Study Cohort. Am J Respir Crit
720	Care Med 184, 828-835. https://doi.org/10.1164/rccm.201012-2082OC
721	Madrigano, J., Kloog, I., Goldberg, R., Coull, B.A., Mittleman, M.A., Schwartz, J., 2013. Long-
722	term Exposure to PM2.5 and Incidence of Acute Myocardial Infarction. Environmental
723	Health Perspectives 121, 192–196. https://doi.org/10.1289/ehp.1205284
724	Malik, A.O., Jones, P.G., Chan, P.S., Peri-Okonny, P.A., Hejjaji, V., Spertus, J.A., 2019.
725	Association of Long-Term Exposure to Particulate Matter and Ozone With Health Status
726	and Mortality in Patients After Myocardial Infarction. Circ: Cardiovascular Quality and
727	Outcomes 12, e005598. https://doi.org/10.1161/CIRCOUTCOMES.119.005598
728	Marchini, T., Magnani, N., Garces, M., Kelly, J., Paz, M., Caceres, L., Calabro, V., Lasagni
729	Vitar, R., Caltana, L., Contin, M., Reynoso, S., Lago, N., Vico, T., Vanasco, V., Wolf,
730	D., Tripodi, V., Gonzalez Maglio, D., Alvarez, S., Buchholz, B., Berra, A., Gelpi, R.,
731	Evelson, P., 2022. Chronic exposure to polluted urban air aggravates myocardial

732	infarction by impaired cardiac mitochondrial function and dynamics. Environmental
733	Pollution 295, 118677. https://doi.org/10.1016/j.envpol.2021.118677
734	McEwan, P.J., 2010. Empirical Research Methods in the Economics of Education, in: Peterson,
735	P., Baker, E., McGaw, B. (Eds.), International Encyclopedia of Education (Third
736	Edition). Elsevier, Oxford, pp. 187–192. https://doi.org/10.1016/B978-0-08-044894-
737	7.01211-2
738	McGuinn, L.A., Ward-Caviness, C.K., Neas, L.M., Schneider, A., Diaz-Sanchez, D., Cascio,
739	W.E., Kraus, W.E., Hauser, E., Dowdy, E., Haynes, C., Chudnovsky, A., Koutrakis, P.,
740	Devlin, R.B., 2016. Association between satellite-based estimates of long-term PM2.5
741	exposure and coronary artery disease. Environmental Research 145, 9-17.
742	https://doi.org/10.1016/j.envres.2015.10.026
743	Miller, K.A., Siscovick, D.S., Sheppard, L., Shepherd, K., Sullivan, J.H., Anderson, G.L.,
744	Kaufman, J.D., 2007. Long-Term Exposure to Air Pollution and Incidence of
745	Cardiovascular Events in Women. N Engl J Med 356, 447-458.
746	https://doi.org/10.1056/NEJMoa054409
747	Mohamed, M.O., Rashid, M., Farooq, S., Siddiqui, N., Parwani, P., Shiers, D., Thamman, R.,
748	Gulati, M., Shoaib, A., Chew-Graham, C., Mamas, M.A., 2019. Acute Myocardial
749	Infarction in Severe Mental Illness: Prevalence, Clinical Outcomes, and Process of Care
750	in U.S. Hospitalizations. Canadian Journal of Cardiology 35, 821-830.
751	https://doi.org/10.1016/j.cjca.2019.04.021
752	Mustafić, H., Jabre, P., Caussin, C., Murad, M.H., Escolano, S., Tafflet, M., Périer, MC.,
753	Marijon, E., Vernerey, D., Empana, JP., Jouven, X., 2012. Main Air Pollutants and

- 754 Myocardial Infarction: A Systematic Review and Meta-analysis. JAMA 307, 713–721.
- 755 https://doi.org/10.1001/jama.2012.126
- Nawrot, T.S., Perez, L., Künzli, N., Munters, E., Nemery, B., 2011. Public health importance of
- 757 triggers of myocardial infarction: a comparative risk assessment. The Lancet 377, 732–
- 758 740. https://doi.org/10.1016/S0140-6736(10)62296-9
- 759 Newcomer, J.W., Hennekens, C.H., 2007. Severe Mental Illness and Risk of Cardiovascular
- 760 Disease. JAMA 298, 1794. https://doi.org/10.1001/jama.298.15.1794
- 761 Ojha, N., Dhamoon, A.S., 2022. Myocardial Infarction, in: StatPearls. StatPearls Publishing,
   762 Treasure Island (FL).
- 763 Padda, J., Khalid, K., Hitawala, G., Batra, N., Pokhriyal, S., Mohan, A., Cooper, A.C., Jean-
- Charles, G., 2021. Acute Anemia and Myocardial Infarction. Cureus 13, e17096.
  https://doi.org/10.7759/cureus.17096
- 766 Palasubramaniam, J., Wang, X., Peter, K., 2019. Myocardial Infarction—From Atherosclerosis
- to Thrombosis. Arteriosclerosis, Thrombosis, and Vascular Biology 39, e176–e185.
- 768 https://doi.org/10.1161/ATVBAHA.119.312578
- Pereira, A.A., Sarnak, M.J., 2003. Anemia as a risk factor for cardiovascular disease. Kidney
  International 64, S32–S39. https://doi.org/10.1046/j.1523-1755.64.s87.6.x
- 771 Phillips, K., Luk, A., Soor, G.S., Abraham, J.R., Leong, S., Butany, J., 2009. Cocaine
- 772 Cardiotoxicity. Am J Cardiovasc Drugs 9, 177–196. https://doi.org/10.1007/BF03256574
- 773 Puett, R.C., Hart, J.E., Yanosky, J.D., Paciorek, C., Schwartz, J., Suh, H., Speizer, F.E., Laden,
- F., 2009. Chronic Fine and Coarse Particulate Exposure, Mortality, and Coronary Heart
- 775 Disease in the Nurses' Health Study. Environmental Health Perspectives 117, 1697–
- 776 1701. https://doi.org/10.1289/ehp.0900572

777	Qiu, X., Wei, Y., Amini, H., Wang, C., Weisskopf, M., Koutrakis, P., Schwartz, J., 2022. Fine
778	particle components and risk of psychiatric hospitalization in the U.S. Science of The
779	Total Environment 849, 157934. https://doi.org/10.1016/j.scitotenv.2022.157934
780	Qiu, X., Wei, Y., Weisskopf, M., Spiro, A., Shi, L., Castro, E., Coull, B., Koutrakis, P.,
781	Schwartz, J., 2023. Air pollution, climate conditions and risk of hospital admissions for
782	psychotic disorders in U.S. residents. Environmental Research 216, 114636.
783	https://doi.org/10.1016/j.envres.2022.114636
784	Rajagopalan, S., Al-Kindi, S.G., Brook, R.D., 2018. Air Pollution and Cardiovascular Disease:
785	JACC State-of-the-Art Review. Journal of the American College of Cardiology 72,
786	2054–2070. https://doi.org/10.1016/j.jacc.2018.07.099
787	Reich, B.J., Yang, S., Guan, Y., Giffin, A.B., Miller, M.J., Rappold, A., 2021. A Review of
788	Spatial Causal Inference Methods for Environmental and Epidemiological Applications.
789	International Statistical Review 89, 605-634. https://doi.org/10.1111/insr.12452
790	Rich, D.Q., Kipen, H.M., Zhang, J., Kamat, L., Wilson, A.C., Kostis, J.B., null, null, 2010.
791	Triggering of Transmural Infarctions, but Not Nontransmural Infarctions, by Ambient
792	Fine Particles. Environmental Health Perspectives 118, 1229–1234.
793	https://doi.org/10.1289/ehp.0901624
794	Rich, D.Q., Özkaynak, H., Crooks, J., Baxter, L., Burke, J., Ohman-Strickland, P., Thevenet-
795	Morrison, K., Kipen, H.M., Zhang, J., Kostis, J.B., Lunden, M., Hodas, N., Turpin, B.J.,
796	2013. The Triggering of Myocardial Infarction by Fine Particles Is Enhanced When
797	Particles Are Enriched in Secondary Species. Environ. Sci. Technol. 47, 9414–9423.
798	https://doi.org/10.1021/es4027248

799 Ros	vall, N.,	Raaschou-Nielsen,	O.,	Ketzel, M.,	Gammelmark,	A.,	Overvad,	K.,	Olsen, A.,	
---------	-----------	-------------------	-----	-------------	-------------	-----	----------	-----	------------	--

- 800 Sørensen, M., 2017. Long-term residential road traffic noise and NO2 exposure in
- 801 relation to risk of incident myocardial infarction A Danish cohort study. Environmental

802 Research 156, 80–86. https://doi.org/10.1016/j.envres.2017.03.019

- 803 Schneider, A., Neas, L.M., Graff, D.W., Herbst, M.C., Cascio, W.E., Schmitt, M.T., Buse, J.B.,
- 804 Peters, A., Devlin, R.B., 2010. Association of cardiac and vascular changes with ambient
- 805 PM2.5 in diabetic individuals. Part Fibre Toxicol 7, 14. https://doi.org/10.1186/1743-
- 806 8977-7-14
- 807 Schwartz, B.G., Rezkalla, S., Kloner, R.A., 2010. Cardiovascular Effects of Cocaine. Circulation
- 808 122, 2558–2569. https://doi.org/10.1161/CIRCULATIONAHA.110.940569
- Schwartz, J., Wei, Y., Yitshak-Sade, M., Di, Q., Dominici, F., Zanobetti, A., 2021. A national
  difference in differences analysis of the effect of PM2.5 on annual death rates.
- 811 Environmental Research 194, 110649. https://doi.org/10.1016/j.envres.2020.110649
- 812 Schwartz, J.D., Di, Q., Requia, W.J., Dominici, F., Zanobetti, A., 2021. A Direct Estimate of the
- 813 Impact of PM2.5, NO2, and O3 Exposure on Life Expectancy Using Propensity Scores.
- Epidemiology 32, 469–476. https://doi.org/10.1097/EDE.00000000001354
- 815 Shi, L., Kloog, I., Zanobetti, A., Liu, P., Schwartz, J.D., 2015. Impacts of Temperature and its

816 Variability on Mortality in New England. Nat Clim Chang 5, 988–991.

- 817 https://doi.org/10.1038/nclimate2704
- 818 Shi, L., Wu, X., Danesh Yazdi, M., Braun, D., Abu Awad, Y., Wei, Y., Liu, P., Di, Q., Wang,
- 819 Y., Schwartz, J., Dominici, F., Kioumourtzoglou, M.-A., Zanobetti, A., 2020. Long-term
- 820 effects of PM2.5 on neurological disorders in the American Medicare population: a

- 821 longitudinal cohort study. The Lancet Planetary Health 4, e557–e565.
- 822 https://doi.org/10.1016/S2542-5196(20)30227-8
- 823 Shumake, K.L., Sacks, J.D., Lee, J.S., Johns, D.O., 2013. Susceptibility of older adults to health
- 824 effects induced by ambient air pollutants regulated by the European Union and the United
- 825 States. Aging Clin Exp Res 25, 3–8. https://doi.org/10.1007/s40520-013-0001-5
- 826 Steiner, C., Elixhauser, A., Schnaier, J., 2002. The healthcare cost and utilization project: an
- 827 overview. Effective Clinical Practice 5.
- 828 Stewart, S., Keates, A.K., Redfern, A., McMurray, J.J.V., 2017. Seasonal variations in
- 829 cardiovascular disease. Nat Rev Cardiol 14, 654–664.
- 830 https://doi.org/10.1038/nrcardio.2017.76
- Valli, H., 2001. Gaseous pollutants in particulate matter epidemiology: confounders or
  surrogates? Environmental Health Perspectives 109.
- 833 Vodonos, A., Awad, Y.A., Schwartz, J., 2018. The concentration-response between long-term
- 834 PM2.5 exposure and mortality; A meta-regression approach. Environmental Research
- 835 166, 677–689. https://doi.org/10.1016/j.envres.2018.06.021
- 836 Wang, G., Jiang, R., Zhao, Z., Song, W., 2013. Effects of ozone and fine particulate matter
- 837 (PM2.5) on rat system inflammation and cardiac function. Toxicology Letters 217, 23–
- 838 33. https://doi.org/10.1016/j.toxlet.2012.11.009
- 839 Wang, R., Xiao, X., Cao, L., Shen, Z., Lei, Y., Cao, Y., 2016. Airborne fine particulate matter
- 840 induces an upregulation of endothelin receptors on rat bronchi. Environmental Pollution
- 841 209, 11–20. https://doi.org/10.1016/j.envpol.2015.10.043

842	Wang, Y., Kloog, I., Coull, B.A., Kosheleva, A., Zanobetti, A., Schwartz, J.D., 2016. Estimating
843	Causal Effects of Long-Term PM2.5 Exposure on Mortality in New Jersey. Environ
844	Health Perspect 124, 1182–1188. https://doi.org/10.1289/ehp.1409671
845	Wang, Y., Shi, L., Lee, M., Liu, P., Di, Q., Zanobetti, A., Schwartz, J.D., 2017. Long-term
846	exposure to PM2.5 and mortality among older adults in the Southeastern US.
847	Epidemiology 28, 207–214. https://doi.org/10.1097/EDE.0000000000000614
848	Weaver, A.M., McGuinn, L., Neas, L., Mirowsky, J., Devlin, R.B., Dhingra, R., Ward-Caviness,
849	C., Cascio, W.E., Kraus, W.E., Hauser, E.R., Di, Q., Schwartz, J., Diaz-Sanchez, D.,
850	2019. Neighborhood Sociodemographic Effects on the Associations Between Long-term
851	PM2.5 Exposure and Cardiovascular Outcomes and Diabetes Mellitus. Environ
852	Epidemiol 3, e038. https://doi.org/10.1097/EE9.000000000000038
853	Wei, Y., Qiu, X., Yazdi, M.D., Shtein, A., Shi, L., Yang, J., Peralta, A.A., Coull, B.A., Schwartz,
854	J.D., 2022. The Impact of Exposure Measurement Error on the Estimated Concentration-
855	Response Relationship between Long-Term Exposure to PM2.5 and Mortality. Environ
856	Health Perspect 130, 077006. https://doi.org/10.1289/EHP10389
857	Wei, Y., Wang, Y., Wu, X., Di, Q., Shi, L., Koutrakis, P., Zanobetti, A., Dominici, F., Schwartz,
858	J.D., 2020. Causal Effects of Air Pollution on Mortality Rate in Massachusetts. Am J
859	Epidemiol 189, 1316–1323. https://doi.org/10.1093/aje/kwaa098
860	Weichenthal, S., Hoppin, J.A., Reeves, F., 2014. Obesity and the cardiovascular health effects of
861	fine particulate air pollution. Obesity 22, 1580–1589. https://doi.org/10.1002/oby.20748
862	Weichenthal, S., Pinault, L.L., Burnett, R.T., 2017. Impact of Oxidant Gases on the Relationship
863	between Outdoor Fine Particulate Air Pollution and Nonaccidental, Cardiovascular, and
864	Respiratory Mortality. Sci Rep 7, 16401. https://doi.org/10.1038/s41598-017-16770-y

42

- 865 Weisskopf, M.G., Webster, T.F., 2017. Trade-offs of Personal Versus More Proxy Exposure
- 866 Measures in Environmental Epidemiology: Epidemiology 28, 635–643.
- 867 https://doi.org/10.1097/EDE.00000000000686
- 868 Wynands, Q., Cazorla, O., Gouzi, F., Grillet, P., Bourdin, A., 2022. Impact of air pollution and
- 869 emphysema on cardiovascular and hemodynamic features. European Respiratory Journal
- 870 60. https://doi.org/10.1183/13993003.congress-2022.2866
- Xu, M., Sbihi, H., Pan, X., Brauer, M., 2020. Modifiers of the effect of short-term variation in
- PM2.5 on mortality in Beijing, China. Environmental Research 183, 109066.
- 873 https://doi.org/10.1016/j.envres.2019.109066
- 874 Yang, J., Biery, D.W., Singh, A., Divakaran, S., DeFilippis, E.M., Wu, W.Y., Klein, J., Hainer,
- J., Ramsis, M., Natarajan, P., Januzzi, J.L., Nasir, K., Bhatt, D.L., Di Carli, M.F.,
- 876 Blankstein, R., 2020. Risk Factors and Outcomes of Very Young Adults Who Experience
- 877 Myocardial Infarction: The Partners YOUNG-MI Registry. The American Journal of

878 Medicine 133, 605-612.e1. https://doi.org/10.1016/j.amjmed.2019.10.020

- Yang, L., Zhang, Y., Qi, W., Zhao, T., Zhang, L., Zhou, L., Ye, L., 2022. Adverse effects of
- 880 PM2.5 on cardiovascular diseases. Reviews on Environmental Health 37, 71–80.
- 881 https://doi.org/10.1515/reveh-2020-0155
- Yitshak-Sade, M., Kloog, I., Zanobetti, A., Schwartz, J.D., 2019. Estimating the causal effect of
  annual PM2.5 exposure on mortality rates in the Northeastern and mid-Atlantic states.
- 884 Environmental Epidemiology 3, e052. https://doi.org/10.1097/EE9.000000000000052
- 885 Yu, B., Akushevich, I., Yashkin, A.P., Kravchenko, J., 2021. Epidemiology of Geographic

886 Disparities of Myocardial Infarction Among Older Adults in the United States: Analysis

887 of 2000–2017 Medicare Data. Frontiers in Cardiovascular Medicine 8.

- Zhao, J., Gao, Z., Tian, Z., Xie, Y., Xin, F., Jiang, R., Kan, H., Song, W., 2013. The biological
- 889 effects of individual-level PM2.5 exposure on systemic immunity and inflammatory
- response in traffic policemen. Occup Environ Med 70, 426–431.
- 891 https://doi.org/10.1136/oemed-2012-100864
- 892 Zhu, W., Cai, J., Hu, Y., Zhang, H., Han, X., Zheng, H., Wu, J., 2021. Long-term exposure to
- 893 fine particulate matter relates with incident myocardial infarction (MI) risks and post-MI
- 894 mortality: A meta-analysis. Chemosphere 267, 128903.
- 895 https://doi.org/10.1016/j.chemosphere.2020.128903

1

# 897 Supplementary Materials

898

# 899 Long-term Exposure to Ambient PM<sub>2.5</sub> and Hospitalizations for Myocardial Infarction

### 900 among U.S. Residents: A Difference-in-Differences Analysis

- 901 Yichen Wang<sup>a\*</sup>, Xinye Qiu<sup>a</sup>, Yaguang Wei<sup>a</sup>, Joel D. Schwartz<sup>a,b</sup>
- <sup>a</sup> Department of Environmental Health, Harvard T.H. Chan School of Public Health, Boston, MA,
- 903 USA
- 904 <sup>b</sup> Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA, USA
- 905
- 906 \*Corresponding author: Yichen Wang, Department of Environmental Health, Harvard T.H. Chan
- 907 School of Public Health, 401 Park Drive, Boston, MA 02215 USA. Email:
- 908 yichenwang@hsph.harvard.edu
- 909
- 910

### 911 Data S1. Supplemental Methods

HCUP data are developed and managed through a Federal-State-Industry partnership sponsored by the Agency for Healthcare Research and Quality. By providing uniformly formatted datasets for multiple states, the data are aimed to facilitate health services research and improve health care delivery. The SIDs in HCUP data contain the inpatient discharge records from community hospitals in 49 participating states, which account for about 97% of all U.S. community hospital discharges. The coverage of SIDs reaches all patients regardless of payer, including individuals covered by Medicare, Medicaid, or private insurance, and those uninsured.

2

919	In the present study, we included the SIDs for the following ten states that were allowed
920	by our current funding budget for research use from HCUP: Arizona (AZ), Michigan (MI), North
921	Carolina (NC), New York (NY), Rhode Island (RI), Washington (WA), New Jersey (NJ),
922	Maryland (MD), Georgia (GA), and Wisconsin (WI). For GA, we only have access to the data
923	after 2010. For WI, we only have access to the data after 2012. For MD, residential address ZIP
924	code information is only available after 2008 and admission year information is only available
925	after 2009. For the other included states, we have the complete records from 2002 to 2016.
926	

927

928 **Table S1.** Percent change in hospitalization rate for MI per 1-µg/m<sup>3</sup> increase in long-term

929 exposure to PM<sub>2.5</sub> across different age groups.

Age (years)	% Change (95% CI)
p for trend <sup>*</sup>	<0.001
0-34	2.40 (1.62, 3.20)
35-44	1.32 (0.93, 1.70)
45-54	0.87 (0.59, 1.16)
55-64	0.95 (0.68, 1.22)
65-74	1.01 (0.75, 1.28)
≥75	1.94 (1.69, 2.20)

<sup>930</sup> p for age trend was estimated from the adjusted model using age as a continuous term.

931

3

933 **Table S2.** Percent change in hospitalization rate for MI per 1-µg/m<sup>3</sup> increase in long-term

SES factors	10th percentile	90th percentile	<i>p</i> for interaction
Poverty	1.16 (0.86, 1.46)	1.42 (1.17, 1.67)	0.0497
Household income	1.31 (1.04, 1.57)	1.44 (1.07, 1.80)	0.5190
Population density	1.07 (0.82, 1.32)	1.19 (0.95, 1.43)	< 0.0001
High school or less	1.30 (0.97, 1.63)	1.37 (1.10, 1.65)	0.6640

934 exposure to PM<sub>2.5</sub> across different levels of SES factors.

935

- 936
- 937

938 **Table S3.** Interaction Odds Ratio (OR) for hospitalization for MI per 1-µg/m<sup>3</sup> increase in long-

939 term exposure to PM<sub>2.5</sub> comparing individuals with and without comorbidities.

Comorbidity	Cases (%)*	OR (95% CI)	р
Hypertension	1067896 (69.4)	1.037 (1.036, 1.039)	< 0.001
DM	448683 (29.2)	1.039 (1.037, 1.041)	< 0.001
COPD	324202 (21.1)	1.004 (1.002, 1.006)	< 0.001
Renal failure	290739 (18.9)	1.042 (1.039, 1.040)	< 0.001
IDA	238974 (15.5)	1.056 (1.053, 1.058)	< 0.001
Obesity	193034 (12.5)	1.007 (1.004, 1.011)	< 0.001
PVD	169422 (11.0)	1.014 (1.011, 1.016)	< 0.001
Depression	105388 (6.8)	0.995 (0.991, 0.998)	< 0.001
Other neurological disorders	93109 (6.1)	1.017 (1.041, 1.055)	< 0.001
Psychoses	31566 (2.1)	1.048 (1.041, 1.054)	< 0.001

- 940 \* The proportion was calculated as the case count divided by the MI hospitalizations from 2005
- 941 to 2015. DM, diabetes mellitus; COPD, chronic obstructive pulmonary disease; IDA, iron
- 942 deficiency anemia; PVD, peripheral vascular disorders.