

25 **Abstract**

26 **Introduction:** Adolescence is a stage of physical, psychological, and social changes,
27 which determine the personality of the individual, which can be influenced by physical
28 appearance because of alterations in the skin texture of adolescents. Patients with
29 dermatosis have a higher risk of developing depression, anxiety and suicidal ideas. Acne
30 is an important factor for quality of life and affects adolescents both physically and
31 psychosocially. **Objective :** To determine the factors associated with the repercussions of
32 acne in adolescents in Honduras. **Methodology:** Analytical cross-sectional study. The
33 Cardiff Acne Disability Index (CADI) scale was used through a questionnaire in Google
34 Forms. Descriptive and analytical statistics were obtained. **Results:** Of the 3,272 young
35 participants, with respect to quality of life, 25.8% (845) had medium or high
36 repercussions, 5.3% were very depressed by the appearance of their skin in the month
37 prior to filling out the questionnaire. When multivariate analysis was performed, it was
38 found that those who had medium or high repercussions due to acne were women (RPa:
39 1.33; 95%CI: 1.20-1.48; p-value<0.001), those of indigenous ethnicity (RPa: 1.36;
40 95%CI: 1.05-1.75; p-value=0.019), those who had scars on the face (RPa: 13.08; 95%CI:
41 10.02-17.05; p-value<0.001) or those who had the father (RPa: 1.13; 95%CI: 1.01-1.25;
42 p-value=0.031) or siblings with acne (RPa: 1.16; 95%CI: 1.04-1.28; p-value=0.005).
43 **Conclusion:** It was found that gender, belonging to an indigenous ethnicity, having scars
44 on the face or that having a close relative who had suffered from acne were associated
45 with medium or high repercussions on the quality of life; it is important to take care of
46 acne in this stage of life to avoid repercussions in adulthood.

47

48 **Keywords:** Acne Vulgaris, Adolescent, Central America, Quality of Life, Scar,
49 Honduras.

50 **Introduction**

51 Acne is a chronic inflammatory disease of the pilosebaceous unit, involving
52 increased androgen-induced sebum production, abnormal keratinization of the
53 pilosebaceous duct and an immune response to colonization by *Cutibacterium acnes*
54 (formerly *Propionibacterium acnes*) (1, 2). Acne is estimated to affect 9.4% of the world's
55 population, making it the eighth most prevalent disease worldwide, affecting 90% of
56 adolescents and young adults (3, 4).

57 It is known that adolescence is a crucial period for the development of social and
58 emotional skills important for mental well-being, such as adopting healthy sleep patterns,
59 exercising regularly, developing skills to maintain interpersonal relationships, coping
60 with difficult situations, solving problems and learning to manage emotions. Therefore,
61 it is important to have a favorable and protective environment in the family, school and
62 community in general (5). The transition from childhood to adolescence entails
63 physiological and psychological changes in the child (6), to which is now added the
64 mental challenges generated by the COVID-19 pandemic, and it is important that
65 adolescents receive the physical and mental care they need to develop, grow and thrive
66 (7).

67 The visibility, chronicity and recurrent course of many skin lesions contribute to
68 various psychosocial disorders, consequences of skin diseases. Patients with dermatoses
69 are at increased risk of developing depression, anxiety, and suicidal ideation (4, 8, 9). In
70 our society today, if certain social parameters are not met in physical appearance,
71 adolescents suffer from bullying by their peers, acne can lead to emotional stress and the
72 impact on quality of life can be significant (10).

73 There are studies that show the importance of acne, blemishes, scars and others
74 that could have sequelae on the skin in adolescents (11), however, most of these studies
75 have been conducted in Europe (12), the United States (13), India (14) and developed
76 countries, in which a very important role is given to mental health and its influence on
77 public health in general. On the other hand, in Latin American countries there are few
78 studies that report this situation, as in Argentina and Mexico, but they were carried out in
79 the pre-pandemic period (15, 16).

80 Among the few studies published in Honduras on the subject, in 2016 "Prevalence
81 of dermatoses in school-aged children" was published, finding a prevalence of acne of
82 1.2% (17), to date no studies have been published in adolescent population related to
83 acne. Therefore, the general objective is to determine the factors associated with the
84 repercussions of acne in adolescents in Honduras.

85 **METHODOLOGY**

86 The Scientific Research Unit (UIC) of the Faculty of Medical Sciences (FMC) -
87 UNAH, implemented the methodology of national research on priority health problems
88 in low-resource settings (18). The physicians who participated in the project were 65
89 eighth year medical students (MSS), who were trained in research methodology by the
90 UIC-FCM, in dermatology with emphasis on acne and in quality of life by professors of
91 the postgraduate courses of Dermatology and Psychiatry UNAH. After training, the
92 surveyors were distributed in 16 departments of the country; this distribution was done
93 by sampling in hospitals and primary care units, and information was collected from July
94 to August 2020. Each interviewer provided an average of 58 questionnaires, each form
95 had an average response time of 10 minutes.

96

97 To determine whether the number of respondents was sufficient to determine the
98 association between the variables, the statistical power calculation was performed, where
99 the only crosses that did not reach adequate power were the age of 13 years (power: 6%),
100 for white ethnicity (power: 69%) and for whether the siblings had suffered from acne
101 (power: 4%); therefore, only these 3 crosses should be taken with caution.

102 The study design was cross-sectional, analytical and multicenter, with a non-
103 random snowball sample, which included 3325 young people aged 12-17 years; in the
104 process of purging the database, 53 records were eliminated, 51 adolescents did not sign
105 the informed consent form, and 2 records were not filled out in their entirety. The
106 adolescents were children or family members within the staff of the care centers at where
107 there were doctors doing. They were provided with the Google Forms link, and they
108 socialized it with their peers. This type of survey was conducted due to the onset of the
109 pandemic and considering the biosecurity measures, the restrictions of movement adopted
110 by the country, the protocol, the safety of the surveyors and respondents. Therefore, it
111 was adapted to online questionnaire and informed assent was obtained from each
112 participant.

113 Before starting the project, ethical care was ensured, so the protocol was submitted
114 to the FCM Biomedical Research Ethics Committee (CEIB), approval code: 010-2020,
115 which was approved in the session of March 20, 2020. As part of their training, the MSS
116 completed the online course on Good Clinical Practices The Global Health Network
117 (www.tghn.org).

118 An instrument was prepared with the following sections: sociodemographic data
119 including general data (age, sex, schooling, origin, ethnicity), as well as related factors
120 and personal pathological history. To help determine whether they suffered from acne,

121 two images were presented for the adolescent to identify their type of acne, inflammatory
122 or non-inflammatory (comedogenic), and the Dermatological Questionnaire of Quality of
123 Life in Adolescents (The Cardiff Acne Disability Index) was included (19).

124 Motley and Finlay described a brief questionnaire designed to assess disability
125 caused by acne. The questionnaire adds a patient-oriented dimension to medical records,
126 highlights patients with unusually high levels of disability, and increases the relevant
127 information on which we base therapeutic decisions (20).

128 The Cardiff Acne Disability Index (CADI) is a questionnaire designed to measure
129 the quality of life of adolescents and young adults with acne, we used the Spanish version
130 validated in 2018, which was validated by Cardiff University. (19). As of October 2021,
131 a change in wording was made (21).

132 When all the information was available, the database was cleaned and the
133 variables were labeled and ready for analysis, which was performed with the statistical
134 program Stata (version 16). First, we proceeded to generate three tables describing the
135 population and their condition of acne, where we obtained the frequencies and
136 percentages of the categorical variables, as well as the mean and interquartile range of
137 age. Then the last tables were elaborated, where the RPc (crude prevalence ratios), RPa
138 (adjusted prevalence ratios), 95% confidence intervals (95% confidence intervals) and p-
139 values were obtained; all were obtained with the generalized linear models (Poisson
140 family, log link function and models for robust variances). For a variable to pass from the
141 crude model to the adjusted model, it had to have a p value <0.05; this same cut-off point
142 was used to determine the final statistical significance.

143

144 **RESULTS**

145 Of the 3272 respondents, 60.0% (1960) were female, the most frequent age was
146 16 years (25.1%), the median age was 15 years (interquartile range: 14-16 years),
147 mestizos were the majority (74.4%) and they stated that their mother (20.2%), father
148 (16.9%), brothers (37.7%), cousins (39.6%) and uncles (20.4%) had suffered from acne
149 in their family.

150 **Table 1**

151 **Table 1.** Characteristics of adolescents surveyed in Honduras, n=3272.

Variable	Frequency	Percentage
Sex		
Male	1306	40,0
Female	1960	60,0
Age (years old)		
12 years	190	5,8
13 years	353	10,8
14 years	454	13,9
15 years	683	20,9
16 years	820	25,1
17 years	772	23,6
Median and interquartile range	15 years	14-16 years
Educational level		
Pre-basic or basic	1415	45,7
Intermediate level	1681	54,3
Ethnic group		
Mestizos	2391	74,4
White	342	10,6
Other	378	11,7
Indigenous	105	3,3
Family with acne*		

Mother	658	20,2
Father	553	16,9
Brothers	1229	37,7
Cousins	1293	39,6
Uncles	666	20,4
You have scars on your face		
No	1701	52,0
yes	1571	48,0

152

153

154

*This section does not add up to 100% since the questions were asked in a way that could have multiple answers.

155

156

157

158

When asked if they currently had acne, 52.5% (1717) responded that, if they did, of those who had acne, 44.3% (761) had it already more than 6 months ago, 19.7% (339) had inflammatory type acne and as for severity, 4.8% (82) and 25.7% (441) had severe and moderate acne, respectively. **Table 2**

159

160

Table 2. Current acne sufferers among adolescent respondents in Honduras: characteristics and impact, n=3272.

Variable	Frequency	Percentage
Currently suffering from acne		
No	1555	47,5
Yes	1717	52,5
Months suffering from acne*		
< 1 month	224	13,1
1-3 months	452	26,3
>3 months and up to 6 months	280	16,3
> 6 months	761	44,3
It is inflammatory acne*.		
No	1378	80,3
Yes	339	19,7
Severity of acne*		
Light	1194	69,5

Mild	441	25,7
Severe	82	4,8

161

162 *These questions have only been answered by those who mentioned that they had acne.

163

164 According to the impact test questions, 4.4% said that they felt very much
 165 aggressive, frustrated or embarrassed in the past month, and 3.4% thought that the acne
 166 they had in the past month interfered with their daily social life, social events or their
 167 relationship with people of the opposite sex. In addition, 5.3% were very depressed about
 168 the appearance of their skin in the past month and 25.8% (845) had medium or high
 169 repercussions from acne. Table 3

170 **Table 3.** Disaggregated and global responses to the impact of acne among adolescent
 171 respondents in Honduras, n=3272.

Questions	Answers
1. As a result of having acne, have you been aggressive, frustrated or embarrassed during the past month?	In fact, very much = 4.4%. Very much= 8.5%. A little = 21.0%. Not at all = 66.1%.
2. Do you think the acne you had last month interfered with your daily social life, social events or your relationship with people of the opposite sex?	Very much; affected all activities=3.4%. Moderately; in most activities=7.5%. Sometimes or only in some activities=19.4% Not at all=69.7%
3. In the past month, have you stopped changing clothes or wearing your bathing suit in public places because of acne?	All the time=3.2% Most of the time=5,1%. Sometimes=12.4% Not at all=79.2%
4. How did the appearance of your skin make you feel last month?	Very depressed and sad=5.3%. Generally worried=12,0%. Sometimes worried=22.3% Not bothered=60.4%
5. Tell us how bad you think your acne is now:	The worst it could be=2.5%. A major problem=17,2%. A minor problem=23.1% No problem=57,2%

Degree of acne affect
None=47.5%
Low=26.7%
Medium=18.9%
High=6.9%

172

173 When bivariate analysis was performed, it was found that those who had medium or high
174 repercussions due to acne were females ($p<0.001$), those who were 14 years old
175 ($p=0.022$), 15 years old ($p<0.001$), 16 years old ($p<0.001$) or 17 years old ($p<0.001$),
176 those of middle level ($p<0, 001$), those who had scars from acne ($p<0.001$) or those who
177 had mother ($p<0.001$), father ($p<0.001$) or siblings with acne ($p<0.001$), on the contrary,
178 those of other ethnicities had less impact from acne ($p<0.001$). **Table 4**

179 **Table 4.** Bivariate analysis of factors associated with the impact of acne in adolescents in
180 Honduras, n=3272.

Variable	Acne repercussions		Bivariate analysis
	None or low	Medium or high	RPc (95%CI) p-value
Gender			
Male	1017 (77,9)	289 (22,1)	Comparison category
Female	1405 (71,7)	555 (28,3)	1,27 (1,13-1,45) <0,001
Age (years old)			
12 years old	165 (86,4)	25 (13,2)	Comparison category
13 years old	305 (86,4)	48 (13,6)	1,03 (0,66-1,62) 0,886
14 years old	358 (78,9)	96 (21,1)	1,61 (1,07-2,41) 0,022
15 years old	468 (68,5)	215 (31,5)	2,39 (1,63-3,50) <0,001
16 years old	582 (71,0)	238 (29,0)	2,21 (1,51-3,23) <0,001
17 years old	549 (71,1)	223 (28,9)	2,20 (1,50-3,21) <0,001
Levels of education			
Pre-basic or basic	1112 (78,6)	303 (21,4)	Comparison category
Medium level	1191 (70,9)	490 (29,1)	1,36 (1,20-1,54) <0,001
Ethnicity			
Mestizos	1737 (72,7)	654 (27,3)	Comparison category
White	258 (75,4)	84 (24,6)	0,90 (0,74-1,09) 0,284

Other	310 (82,0)	68 (18,0)	0,66 (0,53-0,82) <0,001
Indigenous	72 (68,6)	33 (31,4)	1,15 (0,86-1,54) 0,348
Family with acne*			
Mother	409 (62,2)	249 (37,8)	1,65 (,147-1,87) <0,001
Father	323 (58,4)	230 (41,6)	1,83 (1,62-2,07) <0,001
Brothers	761 (61,9)	468 (38,1)	2,06 (1,83-2,31) <0,001
Cousins	935 (72,3)	358 (27,7)	1,12 (0,99-1,26) 0,057
Uncles	480 (72,1)	186 (27,9)	1,10 (0,96-1,26) 0,174
Scars on face			
No	1644 (96,7)	57 (3,3)	Comparison category
Yes	783 (49,8)	788 (50,2)	14,97 (11,54-19,41) <0,001

181

182 *Only the responses of those who, if they suffered, are shown; for the bivariate analysis, they
 183 are always compared with family members who did not suffer. RPc (crude prevalence ratios),
 184 95% CI95% (95% confidence intervals) and p-values were obtained with generalized linear
 185 models (Poisson family, log link function and models for robust variances).

186 When performing the multivariate analysis, it was found that those who had medium or
 187 high repercussions due to acne were women (RPa: 1.33; 95%CI: 1.20-1.48; p-
 188 value<0.001), those of indigenous ethnicity (RPa: 1.36; 95%CI: 1.05-1.75; p-
 189 value=0.019), those who had scars on the face (RPa: 13.08; 95%CI: 10.02-17.05; p-
 190 value<0.001) or those who had the father (RPa: 1.13; 95%CI: 1.01-1.25; p-value=0.031)
 191 or siblings with acne (RPa: 1.16; 95%CI: 1.04-1.28; p-value=0.005). Table 5

192 **Table 5.** Multivariate analysis of factors associated with acne repercussions in adolescents in
 193 Honduras, n=3272.

Variable	RPc (95%CI) p-value
Gender	
Male	Comparison category
Female	1,33 (1,20-1,48) <0,001

Age (years old)	
12 years old	Comparison category
13 years old	0,83 (0,57-1,22) 0,345
14 years old	1,05 (0,75-1,48) 0,768
15 years old	1,37 (0,99-1,89) 0,055
16 years old	1,25 (0,91-1,72) 0,171
17 years old	1,25 (0,91-1,73) 0,170
Ethnicity	
Mestizos	Comparison category
White	0,99 (0,84-1,17) 0,914
Others	0,86 (0,71-1,04) 0,112
Indigenous	1,36 (1,05-1,75) 0,019
Family with acne*	
Mother	1,05 (0,93-1,17) 0,419
Father	1,13 (1,01-1,25) 0,031
Brothers	1,16 (1,04-1,28) 0,005
Scars on face	
No	Comparison category
Yes	13,08 (10,02-17,05) <0,001

194

195 The RPa (adjusted prevalence ratios), 95% confidence intervals (95% confidence intervals) and
196 p-values were obtained with the generalized linear models (Poisson family, log link function
197 and models for robust variances). Educational level was not included in the final model
198 because it expressed the same as age, and variables that did not have a $p < 0.05$ in the bivariate
199 model were not included.

200

201 **Discussion**

202 The objective of the study was to determine the factors associated with the
203 repercussions of acne in adolescents aged 12-17 years in Honduras, since there is no
204 national data available. The only study on the subject found acne in eighth place among

205 the most frequent dermatoses, with 1.2% in 59/15002 school-age children (17). A
206 systematic review that included the opinions and experiences of people with acne vulgaris
207 in 6 countries reported that the physical, psychological and social impact was common
208 and often generated problems in forming new relationships and maintaining current ones
209 (22).

210 Acne is an important factor for quality of life and affects adolescents both
211 physically and psychosocially, they may have difficulties in establishing social
212 relationships during this period, especially when so much importance is given to external
213 appearance (23). In this study we found that one out of four young people perceived
214 medium to high repercussions from acne on their face, knowing that skin conditions of
215 varying severity can affect the quality of life of patients and have psychiatric
216 consequences (24), Hazarika and Archana found facial acne as a single site involvement,
217 this being the most common type (60%), but also with multi-site involvement (face, chest
218 and back together) in 37% of cases, 75% had varying degrees of acne scarring, while
219 post-acne hyperpigmentation was observed in 79% of cases (25). In Europe through an
220 online survey on perceptions and management of acne, most people considered their acne
221 (all stages of severity) to be either not a problem or a minor problem, although 29.7%
222 considered it a major problem or burden (26).

223 We found in this study that females had more repercussions due to acne, a study
224 in high school adolescents in Egypt, which included 994 students, found an overall
225 prevalence of acne vulgaris of 33.5%, the mean age was 16.84 ± 0.87 , this being more
226 common among females, with the mild form predominating, followed by moderate and
227 severe with 6%. The CADI score was significantly related to disease grade and was
228 highest among those with severe grade, followed by moderate and finally mild disease
229 grade (27). Bosio et al. observed that the DLQI was higher in males than in females (15),

230 contrasting our results with those found in this study, where women have a greater impact
231 on quality of life due to acne. Therefore, this indicates that the suffering of this disease
232 may vary according to the population, so studies should be done for each country or
233 region.

234 Singh reported that "feelings of embarrassment and interference with social
235 activities" scored significantly higher for women in a study of 1392 subjects with acne,
236 using the Dermatologic Life Quality Index (DLQI) and the Cardiff Acne Disability Index
237 (CADI) (28), the strongest association was facial scarring, which resulted in the highest
238 frequency of acne-related repercussions, knowing that acne vulgaris can cause permanent
239 physical scarring, negatively affect quality of life and self-image, and has been associated
240 with higher rates of anxiety, depression and suicidal ideation (29). In Egypt, a study on
241 the severity and impact on quality of life and self-esteem among adolescents according
242 to the CADI score showed that 48.96% of students with clinically confirmed acne
243 experienced a mild level of disability, while 11.46% of students with acne had a severe
244 level of disability, low self-esteem was significantly more prevalent among females than
245 males (30).

246 Acne vulgaris appears to share the same pathogenesis regardless of race or
247 ethnicity but has different clinical presentations. In darker skin phototypes, there appears
248 to be a greater subclinical inflammatory response, even in noninflammatory lesions,
249 which is thought to trigger an exaggerated post inflammatory hyperpigmentation (PIH)
250 response and keloid scarring (31), this study found that adolescents of indigenous
251 ethnicity had fewer repercussions from acne. Among the adolescents studied aged 12- 17
252 years, moreover, those who had a parent or siblings with acne also had more acne-related
253 repercussions, knowing that positive family history is a strong predisposing factor in
254 influencing acne presentation, severity and scarring (32). Parental acne was significantly

255 associated with acne presentation and moderate/severe acne, while sibling acne was
256 significantly associated with grade 3/4 scarring (33). A systematic review of the
257 epidemiology of acne vulgaris showed a strong association between family history, age,
258 BMI, and skin type and acne presentation or severity in multiple studies (34). A cross-
259 sectional online survey study of individuals aged 15-24 years in Belgium, Czech and
260 Slovak Republic, France, Italy, Poland, and Spain (n = 10,521) showed that a history of
261 maternal or paternal acne was associated with an increased likelihood of acne (35).
262 Among the risk factors correlated with an increased likelihood of scarring were acne
263 severity, time between acne onset, first effective treatment, recurrent acne, and male sex
264 (36). A study in a group of 15- to 35-year-olds reported that acne outbreaks located on
265 the face were problematic for 81% of the study population (regardless of sex). Females
266 experienced psychological distress more frequently than males (37). Jerome Kaikati et al.
267 in a study conducted in Lebanon reported that acne creates a social barrier and is
268 intensified in women of childbearing age (38). Therefore, people need support to
269 understand the long-term management of acne, to develop control over acne and its
270 treatments, to recognize the impact (22).

271 **Limitations:** The collection of information was designed to be done in person in public
272 middle schools throughout the country, however, due to the quarantine for the COVID-
273 19 pandemic, it was modified and a Google Forms instrument was used, this through a
274 non-random snowball sampling. It is in this context that the schools were forced to be at
275 home, with restricted mobility and other characteristics of the pandemic. All this together
276 could have had a bias at the time of recruitment and how they understood the questions,
277 however, by using validated tests, the support of trained interviewers, the use of
278 supporting images so that they could answer about the type of acne they suffered from
279 and other ways that ensured a better approach to the respondents, all this together shows

280 that we tried to minimize the bias to its minimum proportion. In addition, many
281 adolescents were recruited, and although 3 crosstabs did not reach the necessary statistical
282 power, only two of them had no association at the end of the analysis, so that the number
283 of respondents was sufficient for the vast majority of the variable crosstabs that were
284 raised.

285 **Conclusion:** Based on the above, it is concluded that those who had a medium or high
286 repercussion due to acne were women, those of indigenous ethnicity, those who had scars
287 on their face or those who had a parent or siblings affected by acne.

288 Author Contributions Conceptualization: EET, CASM and KCC

289 Data curation: CM, JA, EET, LIZ, and HNCR.

290 Formal analysis: CM, LIZ and JA.

291 Funding acquisition: KCC.

292 Methodology: EET, CASM and KCC

293 Writing – original draft: EET, CASM, LIZ, KCC, CM, and JA.

294 Writing – review & editing: all authors have read and agreed to the published version of
295 the manuscript.

296 **Funding:** The current article processing charges (publication fees) were funded by the
297 Facultad de Ciencias Médicas (FCM) (2-03-01-01), Universidad Nacional Autónoma de
298 Honduras (UNAH), Tegucigalpa, MDC, Honduras, Central America (granted to Dr.
299 Chinchilla).

300 **Institutional Review Board Statement:** The study was conducted under the Declaration
301 of Helsinki. This research's preparation and execution fully complied with the
302 fundamental ethical principles of autonomy, justice, beneficence, and non-maleficence.
303 The Act Number (2020010), approved by the Ethics Committee in Biomedical Research

304 (CEIB) of the National Autonomous University of Honduras (UNAH), meeting of March
305 20, 2020.

306 **Data Availability Statement:** The data presented in this study are available on request
307 from the corresponding author.

308 **Acknowledgments:** All the MSS of the cohort October 2019 - October 2020 who
309 accepted to participate in the project, to the directors of the healthcare centers in the areas
310 that participated in the study, to the teachers of the UIC, for all the support provided in
311 the review of the MSS assigned to each one. To Ovidio Padilla. who designed the online
312 instrument and created the link for it, and Mauricio Gonzales for the data curation of the
313 study.

314 **Conflicts of Interest:** The authors declare no conflict of interest.

315 [Literature Cited](#)

316 1. Williams HC, Dellavalle RP, Garner S. Acne vulgaris. *Lancet* 2012; 379(9813):361–
317 72.

318 2. Layton AM, Thiboutot D, Tan J. Reviewing the global burden of acne: how could we
319 improve care to reduce the burden?*. *Br J Dermatol* 2021; 184(2):219–25.

320 3. Tan JKL, Bhate K. A global perspective on the epidemiology of acne. *Br J Dermatol*
321 2015; 172 Suppl 1:3–12.

322 4. Marta Szepietowska,¹ Alicja Dąbrowska,¹ Bernadetta Nowak,¹ Katarzyna
323 Skinderowicz,¹ Bartosz Wilczyński,¹ Piotr K. Krajewski,^{2,*} and Alina Jankowska-
324 Konsur². Alexithymia in Adolescents with Acne: Association with Quality of Life
325 Impairment and Stigmatization. *J Clin Med*. Feb 2022. Available from: URL:
326 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8837167/>.

327 5. WHO. Salud mental del adolescente; 2021 (consultado 2023). Available from: URL:
328 <https://www.who.int/es/news-room/fact-sheets/detail/adolescent-mental-health>.

329 6. Gniewosz G, Gniewosz B. Psychological Adjustment During Multiple Transitions
330 Between Childhood and Adolescence. *The Journal of Early Adolescence* 2019;
331 40(4):566–98.

- 332 7. Jones EAK, Mitra AK, Bhuiyan AR. Impact of COVID-19 on Mental Health in
333 Adolescents: A Systematic Review. *Int J Environ Res Public Health* 2021; 18(5).
- 334 8. Roosta N, Black DS, Peng D, Riley LW. Skin disease and stigma in emerging
335 adulthood: impact on healthy development. *J Cutan Med Surg* 2010; 14(6):285–90.
- 336 9. Ritvo E, Del Rosso JQ, Stillman MA, La Riche C. Psychosocial judgements and
337 perceptions of adolescents with acne vulgaris: A blinded, controlled comparison of adult
338 and peer evaluations. *BioPsychoSocial Med* 2011; 5(1):11. Available from: URL:
339 <https://bpsmedicine.biomedcentral.com/articles/10.1186/1751-0759-5-11>.
- 340 10. Leung AK, Barankin B, Lam JM, Leong KF, Hon KL. Dermatology: how to manage
341 acne vulgaris. *Drugs Context* 2021; 10.
- 342 11. Gallitano SM, Berson DS. How Acne Bumps Cause the Blues: The Influence of Acne
343 Vulgaris on Self-Esteem. *Int J Womens Dermatol* 2018; 4(1):12–7.
- 344 12. Gieler U, Gieler T, Kupfer JP. Acne and quality of life - impact and management. *J*
345 *Eur Acad Dermatol Venereol* 2015; 29 Suppl 4:12–4.
- 346 13. Jerry Tan,¹ Stefan Beissert,² Fran Cook-Bolden,³ Rajeev Chavda,⁴ Julie Harper,⁵
347 Adelaide Hebert,⁶ Edward Lain,⁷ Alison Layton,⁸ Marco Rocha,⁹ Jonathan Weiss,¹⁰
348 and Brigitte Drénocorresponding author¹¹. Impact of Facial Atrophic Acne Scars on
349 Quality of Life: A Multi-country Population-Based Survey. *Am J Clin Dermatol* 2022.
350 Available from: URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8776674/>.
- 351 14. Gudiya, M S. Cecilia, Sujata Satapathy,¹ and M Ramam². Assessment of Body Image
352 Disturbance, Self-Esteem and Quality of Life among Adolescents and Young Adults with
353 Acne in a Tertiary Care Facility of India. *Indian J Dermatol* Jan-feb 2022. Available from:
354 URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9154134/>.
- 355 15. Bosio Bonet M, Corball MV, Garay IS, Valente E, Ruiz-Lascano A. Epidemiología e
356 impacto en la calidad de vida de los pacientes con acné juvenil. *Piel* 2022. Available from:
357 URL: <https://www.sciencedirect.com/science/article/pii/S0213925122002192>.
- 358 16. Pantoja-Villa LF, Medina-Castillo DE, Avilés-Sánchez ME. Depresión y calidad de
359 vida en pacientes con acné al inicio y al final del tratamiento. *Dermatología Revista*
360 *Mexicana* 2019; 63(2):123–43.
- 361 17. Cerrato Hernández K, Enamorado B, Delgado L, Sierra M, Espinoza I, Martel B.
362 Prevalencia de dermatosis en niños escolares de Honduras. *Medicina cutánea ibero-*

- 363 latino-americana 2016; 44:177–82. Available from: URL:
364 <https://www.medigraphic.com/cgi-bin/new/resumen.cgi?IDARTICULO=70628>.
- 365 18. Espinoza-Turcios E, Gonzales-Romero RM, Sosa-Mendoza C, Sierra-Santos M,
366 Castro-Ramos HN, Zambrano LI et al. Factors associated with hopelessness, depression
367 and anxiety in the Honduran-Central America population during the COVID-19
368 pandemic. *Front. Psychiatry* 2023; 14.
- 369 19. Cardiff University. Índice de discapacidad del acné de Cardiff; 2018. Available from:
370 URL: [https://www.cardiff.ac.uk/medicine/resources/quality-of-life-](https://www.cardiff.ac.uk/medicine/resources/quality-of-life-questionnaires/cardiff-acne-disability-index)
371 [questionnaires/cardiff-acne-disability-index](https://www.cardiff.ac.uk/medicine/resources/quality-of-life-questionnaires/cardiff-acne-disability-index).
- 372 20. Motley RJ, Finlay AY. Practical use of a disability index in the routine management
373 of acne. *Clin Exp Dermatol* 1992; 17(1):1–3.
- 374 21. Abdelrazik YT, Ali FM, Salek MS, Finlay AY. Clinical experience and psychometric
375 properties of the Cardiff Acne Disability Index (CADI). *Br J Dermatol* 2021; 185(4):711–
376 24.
- 377 22. Ip A, Muller I, Geraghty AWA, Platt D, Little P, Santer M. Views and experiences of
378 people with acne vulgaris and healthcare professionals about treatments: systematic
379 review and thematic synthesis of qualitative research. *BMJ Open* 2021; 11(2):e041794.
- 380 23. Eyüboğlu M, Kalay I, Eyüboğlu D. Evaluation of Adolescents Diagnosed with Acne
381 Vulgaris for Quality of Life and Psychosocial Challenges. *Indian J Dermatol* 2018;
382 63(2):131–5.
- 383 24. Baker N, Billick SB. Psychiatric Consequences of Skin Conditions: Multiple Case
384 Study Analysis with Literature Review. *Psychiatr Q* 2022; 93(3):841–7.
- 385 25. Hazarika N, Archana M. The Psychosocial Impact of Acne Vulgaris. *Indian J*
386 *Dermatol* 2016; 61(5):515–20.
- 387 26. Szepletowski JC, Wolkenstein P, Veraldi S, Tennstedt D, Machovcová A, Delarue A.
388 Acne across Europe: an online survey on perceptions and management of acne. *J Eur*
389 *Acad Dermatol Venereol* 2018; 32(3):463–6.
- 390 27. El-Hamd MA, Nada EE-DA-A, Moustafa MA-K, Mahboob-Allah RA. Prevalence of
391 acne vulgaris and its impact of the quality of life among secondary school-aged
392 adolescents in Sohag Province, Upper Egypt. *J Cosmet Dermatol* 2017; 16(3):370–3.

- 393 28. Singh IP, Poonia K, Bajaj K. Quality of life in young adults with acne: Results of a
394 cross-sectional study. *J Cosmet Dermatol* 2021; 20(12):4017–23.
- 395 29. Eichenfield DZ, Sprague J, Eichenfield LF. Management of Acne Vulgaris: A
396 Review. *JAMA* 2021; 326(20):2055–67.
- 397 30. Tayel K, Attia M, Agamia N, Fadl N. Acne vulgaris: prevalence, severity, and impact
398 on quality of life and self-esteem among Egyptian adolescents. *Journal of the Egyptian*
399 *Public Health Association* 2020; 95(1):30.
- 400 31. Pathmarajah P, Peterknecht E, Cheung K, Elyoussfi S, Muralidharan V, Bewley A.
401 Acne Vulgaris in Skin of Color: A Systematic Review of the Effectiveness and
402 Tolerability of Current Treatments. *J Clin Aesthet Dermatol* 2022; 15(11):43–68.
- 403 32. Say Y-H, Heng AHS, Reginald K, Wong YR, Teh KF, Rawanan Shah SM et al.
404 Modifiable and non-modifiable epidemiological risk factors for acne, acne severity and
405 acne scarring among Malaysian Chinese: a cross-sectional study. *BMC Public Health*
406 2021; 21(1):601.
- 407 33. Heng AHS, Say Y-H, Sio YY, Ng YT, Chew FT. Epidemiological Risk Factors
408 Associated with Acne Vulgaris Presentation, Severity, and Scarring in a Singapore
409 Chinese Population: A Cross-Sectional Study. *Dermatology* 2022; 238(2):226–35.
- 410 34. Heng AHS, Chew FT. Systematic review of the epidemiology of acne vulgaris. *Sci*
411 *Rep* 2020; 10(1):5754.
- 412 35. Wolkenstein P, Machovcová A, Szepietowski JC, Tennstedt D, Veraldi S, Delarue A.
413 Acne prevalence and associations with lifestyle: a cross-sectional online survey of
414 adolescents/young adults in 7 European countries. *J Eur Acad Dermatol Venereol* 2018;
415 32(2):298–306.
- 416 36. Tan J, Kang S, Leyden J. Prevalence and Risk Factors of Acne Scarring Among
417 Patients Consulting Dermatologists in the USA. *J Drugs Dermatol* 2017; 16(2):97–102.
- 418 37. Kostecka M, Kostecka J, Szwed-Gułaga O, Jackowska I, Kostecka-Jarecka J. The
419 Impact of Common Acne on the Well-Being of Young People Aged 15-35 Years and the
420 Influence of Nutrition Knowledge and Diet on Acne Development. *Nutrients* 2022;
421 14(24).

422 38. Kaikati J, Zoghaib S, Kechichian E, Stephan F, Helou J, Sleilaty G et al. The impact
423 of acne treatment on quality of life and self-esteem: A prospective cohort study from
424 Lebanon. *Int J Womens Dermatol* 2021; 7(4):415–21.

425