**Looking under the lamp-post: quantifying the performance of contact tracing in the United**

**States during the SARS-CoV-2 pandemic**

**Authors:** Henry Bayly[1], Madison Stoddard[2], Debra Van Egeren[3], Ellie Murray[4], Julia Raifman[5],

Arijit Chakravarty[2+], Laura F White[1*+]

**Affiliations:**

[1] Department of Biostatistics, Boston University School of Public Health, Boston, MA, USA

[2] Fractal Therapeutics, Cambridge, MA, USA.

[3] Stanford University, Palo Alto, CA, USA.

[4] Department of Epidemiology, Boston University School of Public Health, Boston, MA, USA

[5] Department of Health Law, Policy and Management, Boston University School of Public

Health, Boston, MA, USA

* Corresponding author

+ contributed equally

**Word count:** 3498

**Abstract word count:** 215

**Abstract:**

Contact tracing forms a crucial part of the public-health toolbox in mitigating and understanding emergent pathogens and nascent disease outbreaks. Contact tracing in the United States was conducted during the pre-Omicron phase of the ongoing COVID-19 pandemic. This tracing relied on voluntary reporting and responses, often using rapid antigen tests (with a high false negative rate) due to lack of accessibility to PCR tests. These limitations, combined with SARS-CoV-2's propensity for asymptomatic transmission, raise the question "how reliable was contact tracing for COVID-19 in the United States"? We answered this question using a Markov model to examine the efficiency with which transmission could be detected based on the design and response rates of contact tracing studies in the United States. Our results suggest that contact tracing protocols in the U.S. are unlikely to have identified more than 1.65% (95% uncertainty interval: 1.62%-1.68%) of transmission events with PCR testing and 0.88% (95% uncertainty interval 0.86%-0.89%) with rapid antigen testing. When considering an optimal scenario, based on compliance rates in East Asia with PCR testing, this increases to 62.7% (95% uncertainty interval: 62.6%-62.8%). These findings highlight the limitations in interpretability for studies of SARS-CoV-2 disease spread based on U.S. contact tracing and underscore the vulnerability of the population to future disease outbreaks, for SARS-CoV-2 and other pathogens.

**Introduction**

The management and control of infectious disease has been a signal modern achievement.

Advances in epidemiological techniques pioneered during the 19[th] century established public

health as a discipline. Overlapping with, but distinct from the medical establishment and the

biopharmaceutical industry, modern public health organizations have sought to control disease

using nonpharmaceutical interventions (NPIs).

Contact tracing is a cornerstone of the public-health response, particularly with emergent

pathogens and nascent disease outbreaks [1]. Effective contact tracing facilitates estimates of

epidemiological parameters describing disease spread. In the current COVID-19 pandemic,

rigorous early studies relying on contact tracing revealed key epidemiological features of SARS-

CoV-2 such as asymptomatic transmission [2,3], superspreading [4], and aerosol transmission

[5–9]. This provided a basis for projecting the future course of the outbreak and designing a

public health response.

Effective contact tracing is also critical for limiting onward spread through the deployment of

test-and-trace and isolation protocols. Many Asia-Pacific countries effectively limited SARS-CoV-

2 community spread for the first two years of the pandemic, relying on contact tracing with

isolation of contacts, including strict testing and isolation efforts at their borders. For example,

South Korea used methods such as tracking credit card transactions and using closed circuit

televisions to link contacts together [10]. In China, specifically Hubei, suspected contacts were

placed under monitored house arrest throughout their quarantine period [11]. This strategy

permitted high levels of withing country contact and mobility while keeping case counts low

[12–18].

In the U.S., contact tracing was primarily performed in the pre-Omicron era, and largely

abandoned in early 2022 [19]. It has been widely recognized that contact tracing in the U.S. has

not slowed disease transmission [20]. Part of the challenge has been that the process varied

from state to state and relied on individual initiative and access to testing [19]. This meant that

an individual typically must be symptomatic, voluntarily seek testing, and have their positive

result reported to initiate contact tracing [21]. Public health officials initiated an investigation

by asking the index case to identify their contacts, who in turn would be interviewed. The

exposed contacts were monitored for symptoms and could choose to test for SARS-CoV-2 five

days after exposure. If positive, the contact (now a secondary case) would be asked to name

their contacts.

The process was largely voluntary, allowing for selection bias and many missed transmission

chains. There was often no system for identifying close contacts whom the index case did not

know personally. Many published papers noted that many named contacts were not

successfully traced [22–24] and not all symptomatic contacts were willing to undergo testing

[25]. A systematic surveillance-based cross-sectional study in the U.S. showed that 2 out of 3

index cases of COVID-19 were either not reached by tracers or declined to share contacts. Only

70% of named contacts agreed to be interviewed, and only 50% of those contacts were

monitored, leading to an average of less than one contact per index case being monitored [26].

Additionally, the CDC-recommended 15 minutes of contact within six feet over a 24-hour

period was somewhat arbitrary and never updated, even as evidence emerged indicating that

COVID-19 could be transmitted through brief interactions.

The implications of these limitations in contact tracing are significant. The relatively high

reproductive number for SARS-CoV-2 [27] would suggest that many transmission chains

generated from a single index case went undetected. Additionally, asymptomatic transmission

and superspreading behavior would also impact the efficacy of contact tracing for infection

control and the generalizability of inferences made about transmission dynamics [28,29].

In keeping with this voluntary and symptom-gated approach to contact tracing, there are many

examples of minimally observed onward transmission in settings where transmission would be

expected. This includes studies involving children with strong implications for policies related to

schools. The results of studies investigating children and COVID-19 transmission have

documented limited forward transmission, but this is often in context of significant mitigation

strategies being in place or incomplete contact tracing [30–32]. During the initial omicron surge,

when contact tracing was limited, schools struggled to remain open, reported high absenteeism

rates, and in some cases, relied on the national guard to teach courses and due to incomplete

contact tracing it was unclear what role children in schools played in transmission [33,34]. In

one another case, two COVID-19 positive hairdressers in Missouri saw 139 clients over a ten-

day period, with no reported onward transmission [24]. Notably, of the exposed clients, only

~75% (n=104) responded to contact tracers' requests for interviews, and only ~50% (n=67)

agreed to be tested. Biases in willingness to respond to interviews or agree to testing may have

concealed many onward transmission events.

Another example, demonstrated the challenges in identifying both primary and secondary

infections, was the Sturgis motorcycle rally in August 2020. Following this 10-day event in

Meade County, South Dakota (attended by approximately 460,000 persons [35] without [36–

38] any mask-wearing requirements or other mitigating policies [39]), there was a wave of

COVID-19 cases in Meade County and South Dakota. The counties outside of South Dakota that

contributed the highest inflows of rally attendees experienced a 6.4-12.5% increase in COVID-

19 cases relative to counties without inflows [40]. Despite evidence of population-level changes

in COVID-19 case counts following the rally, the CDC and Minnesota Department of Health were

able to identify only 21 person-to-person transmission events [41]. Out of the 86 positive cases,

only 41 reported being in close contact (defined as being within 6 feet of another person for

≥15 minutes) with other people, and they reported an average of 2.5 close contacts. Both

statistics are implausible for a 10-day motorcycle rally featuring indoor dining and concerts [42–

44]. The CDC's report does not specify how many of the 102 secondary contacts were tested,

consistent with other U.S. contact-tracing studies [45,46].

Examples such as these, coupled with the unique features of SARS-CoV-2 contact tracing in the

U.S. during the early part of the pandemic raise the question "what was the efficiency of

contact tracing, as it was implemented in the U.S."?  We answered this question using Markov

Chain modeling to synthesize data from multiple sources of information on testing and contact

tracing completeness to estimate the efficiency with which onward transmission could be

detected. Our model-based approach sought to quantify two metrics of performance for

contact tracing: 1) the percentage of all transmission pairs identified in a disease

cluster/outbreak, and 2) the percentage of onward transmission events identified from a

known index case. These metrics correspond broadly to the two contact-tracing scenarios

described above, the Sturgis motorcycle rally study (seeking all transmission pairs) and the

Missouri hairdressers (seeking onward transmission from a known index case). Our results

suggest that, as may be expected, contact tracing protocols in the United States are unlikely to

have identified more than a vanishingly small fraction of transmission events. We contrast this

with a similar model run that incorporates data from Asian countries with more comprehensive

contact tracing protocols in place, which yields a larger fraction of identified transmission

events.

**Methods**

*Model*

We created a Markov Model to represent the sequence of events in contact tracing depicted in

Figure 1. The model captures the steps in contact tracing beginning with identifying a primary

infectious individual through testing and then engaging that person in contact tracing by

accurately identifying their contacts. The final steps of the model details engaging the infected

contact and completing testing. We focus on estimating the probability of identifying infected

contacts only. We parameterize the model steps through literature review. We capture

uncertainty in the transition parameters by sampling over a literature-informed uncertainty

range of the parameters 10,000 times. When no data was available on a parameter value, we

sweep over the range of [0,1].

*Literature search and model parameterization*

We reviewed the literature to produce estimates for each of the parameters of our model

(Figure 1). We used the search term "(covid 19 or covid-19 or covid19) AND (case investigation

or contact tracing) AND (united states or US)" on PubMed to inform parameters that define the

likelihood of individuals naming their close contacts and of those contacts responding to a

contact tracing encounter and being tested. We collected data from contact tracing

investigations that reported the proportion of positive cases that named contacts, the

proportion of named contacts reached, the proportion of contacts that cooperated with

tracers, and/or the proportion of contacts tested.

We also parameterized our model to represent a setting where stringent contact tracing was

implemented. For this, we derived estimates from Taiwan and South Korea, which had rigorous

contact tracing protocols during the initial phase of the pandemic using the search terms '(covid

19 or covid-19 or covid19) AND (case investigation or contact tracing) AND (taiwan)' and '(covid

19 or covid-19 or covid19) AND (case investigation or contact tracing) AND (korea or south

korea)'. We run the model separately for RAT and PCR tests.

The model can be divided into two distinct categories of parameters that contribute to the

overall effectiveness of contact tracing: 1) efficiency of testing and 2) efficiency of contact

tracing. Efficiency of testing includes the proportion of symptomatic people receiving testing,

test sensitivity (RAT or PCR) [47,48], and the proportion of contacts tested. Efficiency of contact

tracing aggregates the probabilities that tracers contact a positive index case, a positive index

case names contacts, and named contacts are traced. Both aggregate parameters take values

between [0,1]. We plot all possible combinations of these parameters on a heatmap and

identify the estimates obtained for the U.S.-based and ideal scenarios.

*Code Availability.*

The Markov Model was implemented in Python, and code for running the simulations and

plotting the results are available in a Jupyter notebook on Github (https://github.com/Henry-

Bayly/ContactTracingMarkovModel).

**Results**

*Literature search*

Our literature search of U.S. studies yielded 1,355 papers. Of these, the first 350 were reviewed

to represent a random sample of the total papers found, as our goal was to conduct a

representative and not comprehensive literature review. We excluded 325 papers that did not

contain data relevant to the parameters required for our model, leaving twenty-five papers

with information on contact tracing parameters. When multiple papers had values for the same

parameter, for instance the probability of a case naming contacts, we assumed that the true

value was uniformly distributed in the range of the reported values across the papers. All

parameter values are shown in Table 1; sources of the parameters are shown in Tables S1-S6.

We were unable to identify precise parameter estimates for the probability of a symptomatic

person receiving testing and assume a uniform distribution over [0,1].

We reviewed 225 papers from South Korea and Taiwan. However, we were unable to

find studies quantifying contact tracing parameters related to the completeness of tracing. This

appeared to be due to a much more comprehensive approach to contact tracing in these

countries, negating the need to report on these parameters since it was assumed that reporting

was nearly complete. For example, South Korea used traditional shoe-leather epidemiology

along with large databases (global positioning system, credit card transactions, and closed-

circuit television) [10] and in one study of 5,706 index cases an average of 9.9 contacts per

index case were reported [10]. Taiwan similarly reported an average of 27.61 contacts per index

case [49]. This contrasts with the US where the average number of non-household contacts

reported in a large study was one for every three index cases [50]. Therefore, we hypothesized

a realistic range of [0.9, 1.0] for all parameters not associated with testing sensitivity in our

ideal model setting.

*Estimates of contact tracing efficacy*

We examined the efficiency of contact tracing along the two dimensions described previously:

1) the percentage of all transmission pairs identified in a disease cluster/outbreak, and 2) the

percentage of onward transmission events identified from a known index case.

We estimate a 0.88% chance (95% uncertainty range: 0.86%-0.89%) of identifying any

transmission pair in the U.S. when RAT are the primary testing modality. More specifically, we

estimate a 20.2% chance of identifying a positive index case, a 20.0% chance of identifying

contacts given that the index case has been identified, and a 21.2% chance of identifying

positive secondary cases given the index cases and contacts were identified. Using more

sensitive, but less available, PCR tests, we estimate a 1.65% (95% uncertainty range: 1.62%-

1.68%) chance of identifying a transmission pair (Figure 2, Table 2). Our model estimates a

27.7% chance of identifying a positive index case, a 19.8% chance of identifying contacts given

that the index case has been identified, and a 29.2% chance of identifying positive cases given

that the index cases and its contacts were correctly identified. By contrast, when we use an

idealized scenario, based on data from East Asia, we estimate a 62.7% (95% uncertainty range:

62.6%-62.8%) chance of identifying a given transmission pair when using PCR testing and 33.5%

(95% uncertainty range: 33.4%-33.6%) when using RAT (Figure 2, Table 2).

When an index case has been identified, we remove the associated model steps. In our U.S.

based example, we estimate a 2.62% (95% uncertainty interval: 2.59%-2.65%) and 3.62% (95%

uncertainty interval: 3.58%-3.66%) chance of identifying a positive contact of a known index

case using RAT and PCR testing, respectively. In our idealized tracing setting, these increase to

53.3% (95% uncertainty interval: 53.2%-53.4%) for RAT and 73.3% (95% uncertainty interval:

73.2%-73.4%) for PCR testing.

**Discussion**

Since its emergence in humans more than three years ago, SARS-CoV-2 has overwhelmed public

health institutions globally. The virus still exerts an enormous mortality and morbidity burden

worldwide, with nearly 7 million reported deaths thus far [51]. Underlying this is the systematic

failure of contact tracing, which was abandoned by most states in the United States by early

2022 [19] and contraindicated in CDC guidance for communities with "sustained ongoing

transmission" of COVID-19 [21]. The Lancet Commission report on COVID-19 [52] and several

reviews [20,53,54] have highlighted this shortcoming in pandemic response.

Our work takes this a step further by quantifying the extent to which contact tracing failed to

identify transmission events in the U.S. and in an idealized setting base on East Asian data. Our

demonstrate that the voluntary steps in contact tracing, i.e. seeking testing and interacting with

contact tracers, reduced the efficiency dramatically, with fewer than 2% of transmission events

identified compared to 62.7% in a setting where testing and compliance with tracing was

higher.

Contact tracing formed the basis of modern epidemiological practice, dating back to the

investigation of the 1854 Broad Street Cholera outbreak in Britain [55] that led to both a

mechanistic understanding of cholera transmission [56] and successful control of the outbreak.

A more recent example of highly successful and proactive contact tracing by public health

authorities was the effective suppression of monkeypox in the Western US in 2003 [57].

Notably, during the current pandemic, many other countries (such as China, Japan, South

Korea, Taiwan, Vietnam and Singapore) were successful at implementing contact tracing in the

first two years of the pandemic [14–18].

The impact of poor contact tracing in the U.S. has undermined our understanding of the

transmission potential of SARS-CoV-2. For example, the argument that schools do not

contribute to SARS-CoV-2 transmission was based in part on the lack of detection of

transmission chains in a school setting. Numerous publications showed a lack of contact-traced

chains of transmission in a school setting [58–62], while in effect lacking a positive control for

the ability to identify onward chains of transmission [63]. It is now clear that SARS-CoV-2 is

readily transmitted in schools [64–72], particularly when robust mitigation measures are not in

place [73,74]. Indeed, dramatic increases in case detection rates have been observed in studies

that relied on surveillance testing, rather than contact tracing [70]. Additionally, this led to the

conclusion that the most common source of transmission was gatherings in the home, but it is

unclear if this is a consequence of household contacts being easiest to identify or a result of

many transmission studies being conducted in settings with strict shut downs, where

households were one of the few places where transmission could occur [75]. Also, reports from

the West have pointed to a lack of detected transmission chains in air travel [76–78]. These

reports are contradicted by careful contact tracing studies from other countries, which have

clearly demonstrated person-to-person transmission in flight [79–81], even when robust

mitigation measures were in place [82].

Our work has several limitations. We have assumed instantaneous contact tracing, ignoring the

impact of tracing delays on infection control which has a significant impact on contact tracing

effectiveness against transmission [83,84] due to short incubation period of SARS-CoV-2 [85].

Instead, our estimates for contact tracing effectiveness apply to the informativeness of contact

tracing studies, and they form an upper bound for the effectiveness of contact tracing as a

transmission prevention measure. We do not account for asymptomatic transmission, again

making our estimates an upper bound. The percent of asymptomatic COVID-19 cases is

estimated to be anywhere between 1.6% and 56.5% [90–96], with a these cases having a

relative reduced infectiousness of 0 to 62% [90-97]. This would mean our estimated 1.65% of

transmission pairs identified with PCR testing could be as low as 0.9%, assuming no

asymptomatic index cases are identified. We also have not accounted for superspreading,

which has been estimated to be a significant feature in COVID-19 transmission [94–96]. This

implies that missing a superspreading index case would have tremendous impact on

downstream contact tracing efforts and that there is significant stochasticity [97]. We only

consider the probability of identifying infected contacts, but it is ideal to also identify

uninfected contacts accurately. Finally, we estimate the probability of naming an infected

contact using data describing the probability of naming any contacts at all (rather than the

probability of naming any given contact). This means that our final estimated probabilities are

upper bounds of the true values. Despite representing an upper bound, our contact tracing

estimates suggest that U.S. contact tracing studies fail to identify the vast majority of

transmission pairs and onward transmission events. This severely limits the inferences that can

be drawn from such studies.

Our work points to several key lessons for future public health efforts. First, compliance is a key

driver of contact tracing effectiveness. Methods to improve compliance will be crucial for future

contact tracing efforts- whether using technological approaches (such as mobile phone or

surveillance-camera based tracing) or by making changes to the legal framework around public

health efforts (see Supplementary Information S2 for a more on this topic).

Second, there is a pressing need for innovation, to develop contact tracing methodologies that

are more resistant to noncompliance. One such approach may be backward contact tracing,

which seeks to identify who infected the detected case. Here when contact tracing is executed

backward to identify the source of infection (parent), the more offspring (infections) a parent

has produced, the more frequently the parent shows up as a contact. Model-based analysis

suggests that a backwards contact tracing approach does not require sampling a network at

such a large scale as forward tracing [98,99] to understand transmission dynamics, and

addresses the problem of low compliance. This approach has been proposed by others for

COVID-19 [100–102], and has been empirically shown to be effective, particularly in identifying

superspreading events [103], however this is unlikely to be as helpful in reducing transmission.

Third, public health responses to future outbreaks must include educate the public about

behaviors with health outcomes, including creating a normative framework around contact

tracing compliance. Consistent messaging about limiting transmission and contact tracing are

key as has been noted by the Lancet Commission [52], among others [104,105]. This could

include reframing messaging to reduce stigma that has often been associated with contact

tracing [106] and which undermines contact tracing efficacy [107]. During the HIV epidemic,

contact tracers emphasized the index case's personal responsibility towards the health of their

sexual partners [1]. It also includes addressing misinformation, which led many to believe that

COVID-19 was a "hoax" [108] and public health measures were overreactions [109].

Finally, we have shown that testing availability and accuracy create a critical gap in contact

tracing efforts. Considering only the steps for testing accuracy and cases/contacts receiving

testing in our model we find that only 12.4% of possible cases *could* be identified with RAT, the

most available testing modality.  To effectively manage future outbreaks, tests need to be

sensitive, provide rapid results, and be readily available.

The work presented here adds to the growing body of literature [26,110,111] highlighting the

poor performance of contact tracing in the West during the ongoing pandemic and suggests

practical fixes for this problem, as we have described. In its absence, public health is forced to

rely on population-wide measures for disease spread and will not be able to fine-tune its

responses to match the situation. If we are to improve our response to the current crisis, or to

others in the future, we must improve our ability to deliver this key function.

**TABLES**

**Table 1.** Table of parameters derived from literature that were used in the model.

| Parameter (Step in the Model) | U.S. Probability Range | Ideal Contact Tracing Probability Range |
|---|---|---|
| Symptomatic case receives testing* | [0,1] | [0.90, 1.0] |
| COVID-19 test gives true positive* | [0.57, 0.73] (RAT) [48] and [0.88, 0.92] (PCR) [47] | [0.57, 0.73] (RAT) and [0.88, 0.92] (PCR) |
| Tracers make contact with a positive case** [22,26,45,46,50,112–116] | [0.41, 0.82] | [0.90, 1.0] |
| Positive case names any contacts**,a [22,26,45,112,113,116,117] | [0.17, 0.52] | [0.90, 1.0] |
| Tracers make contact with infected contacts of positive case**[22,26,46,112,113,115–117] | [0.28, 0.85] | [0.90, 1.0] |
| Infected contacts of positive case get tested* [113,117–121] | [0.19, 0.45] | [0.90, 1.0] |
| Contact's COVID-19 test gives a true positive* | [0.57, 0.73] (RAT) [48] and [0.88, 0.92] (PCR) [47] | [0.57, 0.73] (RAT) and [0.88, 0.92] (PCR) |

<sup>a</sup>These values represent the probability that a positive case names any contacts at all. We use

this as a proxy for the given step. Thus, our model represents an overestimate of the true

efficacy of contact tracing.

* parameter related to testing

**parameter related to tracing

**Table 2.** Probabilities of correctly identifying positive contacts of a given index case stratified by

contact tracing style and by use of RAT/PCR tests. Values in parenthesis represent the 95%

uncertainty intervals.

| | United States | | Ideal Contact Tracing | |
|---|---|---|---|---|
| | RAT | PCR Tests | RAT | PCR Tests |
| Percent of all transmission pairs in an outbreak | 0.88% (0.86%-0.89%) | 1.65% (1.62%-1.68%) | 33.5% (33.4%-33.6%) | 62.7% (62.6%-62.8%) |
| Percent of onward transmission from a known index case | 2.62% (2.59%-2.65%) | 3.62% (3.58%-3.66%) | 53.3% (53.2%-53.4%) | 73.3% (73.2%-73.4%) |

365 **FIGURES**

372    5 coincide with phase 2 (identifying contacts of positive cases). Finally, steps 6 and 7 coincide

373    with Phase 3 (identifying positive cases among contacts).

**Figure 2. Impact of contact tracing and testing on the probability of identifying a positive**

**contact of an infected individual with COVID-19.** Quality of testing refers to parts of the

process relating to testing and aggregates the following probabilities: symptomatic people

receiving testing, symptomatic index cases receive true positive test result (i.e. test sensitivity),

contacts receive testing, contacts receive a true positive test result. Quality of contact tracing

aggregates the probability of: tracers contact a positive index case, a positive index case names

contacts, and tracers contact the contacts of the index case. This shows that if testing and

contact tracing are done perfectly, we can expect to identify all contacts of infected individuals

(illustrated through the colors of the heat map). The blue circles correspond to our simulations

using RAT while the white circles correspond to PCR testing use. The two circles in the lower left

hand corner correspond to our United States estimates while the two in the upper half

correspond to our simulations using estimates from South Korea and Taiwan, where there was

stricter contact tracing.

**References**

1.   Brandt AM. The History of Contact Tracing and the Future of Public Health. Am J Public
Health. American Public Health Association; **2022**; 112(8):1097–1099.

2.   He X, Lau EHY, Wu P, et al. Temporal dynamics in viral shedding and transmissibility of
COVID-19. Nat Med. **2020**; 26(5):672–675.

3.   Sun K, Wang W, Gao L, et al. Transmission heterogeneities, kinetics, and controllability of
SARS-CoV-2. Science. American Association for the Advancement of Science; **2021**;
371(6526):eabe2424.

4.   Endo A, Centre for the Mathematical Modelling of Infectious Diseases COVID-19 Working
Group, Abbott S, Kucharski AJ, Funk S. Estimating the overdispersion in COVID-19
transmission using outbreak sizes outside China. Wellcome Open Res. **2020**; 5:67.

5.   Lu J, Gu J, Li K, et al. COVID-19 Outbreak Associated with Air Conditioning in Restaurant,
Guangzhou, China, 2020. Emerg Infect Dis. **2020**; 26(7):1628–1631.

6.   Eichler N, Thornley C, Swadi T, et al. Transmission of Severe Acute Respiratory Syndrome
Coronavirus 2 during Border Quarantine and Air Travel, New Zealand (Aotearoa). Emerg
Infect Dis. **2021**; 27(5):1274–1278.

7.   Kwon K-S, Park J-I, Park YJ, Jung D-M, Ryu K-W, Lee J-H. Evidence of Long-Distance Droplet
Transmission of SARS-CoV-2 by Direct Air Flow in a Restaurant in Korea. J Korean Med Sci.
**2020**; 35(46):e415.

8.   Wang J, Du G. COVID-19 may transmit through aerosol. Ir J Med Sci. **2020**; 189(4):1143–
1144.

9.   Shen Y, Li C, Dong H, et al. Airborne Transmission of COVID-19: Epidemiologic Evidence
from Two Outbreak Investigations. SSRN Electron J. **2020**; .

10.  Park YJ, Choe YJ, Park O, et al. Contact Tracing during Coronavirus Disease Outbreak, South
Korea, 2020. Emerg Infect Dis. **2020**; 26(10):2465–2468.

11.  Maier BF, Brockmann D. Effective containment explains subexponential growth in recent
confirmed COVID-19 cases in China. Science. **2020**; 368(6492):742–746.

12.  Lu N, Cheng K-W, Qamar N, Huang K-C, Johnson JA. Weathering COVID-19 storm: Successful
control measures of five Asian countries. Am J Infect Control. **2020**; 48(7):851–852.

13.  Zhou Y, Jiang H, Wang Q, Yang M, Chen Y, Jiang Q. Use of contact tracing, isolation, and
mass testing to control transmission of covid-19 in China. The BMJ. **2021**; 375:n2330.

14. El Guerche-Séblain C, Chakir L, Nageshwaran G, et al. Experience from five Asia-Pacific
countries during the first wave of the COVID-19 pandemic: Mitigation strategies and
epidemiology outcomes. Travel Med Infect Dis. **2021**; 44:102171.

15. Jefferies S, French N, Gilkison C, et al. COVID-19 in New Zealand and the impact of the
national response: a descriptive epidemiological study. Lancet Public Health. **2020**;
5(11):e612–e623.

16. Van Nguyen H, Lan Nguyen H, Thi Minh Dao A, et al. The COVID-19 pandemic in Australia:
Public health responses, opportunities and challenges. Int J Health Plann Manage. **2022**;
37(1):5–13.

17. Chen S-C. Taiwan's experience in fighting COVID-19. Nat Immunol. **2021**; 22(4):393–394.

18. Van Nguyen Q, Cao DA, Nghiem SH. Spread of COVID-19 and policy responses in Vietnam:
An overview. Int J Infect Dis IJID Off Publ Int Soc Infect Dis. **2021**; 103:157–161.

19. Feliciano M. State Approaches to Contact Tracing during the COVID-19 Pandemic [Internet].
Natl. Acad. State Health Policy. 2022 [cited 2022 Nov 26]. Available from:
https://www.nashp.org/state-approaches-to-contact-tracing-covid-19/

20. Clark E, Chiao EY, Amirian ES. Why Contact Tracing Efforts Have Failed to Curb Coronavirus
Disease 2019 (COVID-19) Transmission in Much of the United States. Clin Infect Dis Off
Publ Infect Dis Soc Am. **2021**; 72(9):e415–e419.

21. CDC. Health Departments [Internet]. Cent. Dis. Control Prev. 2020 [cited 2022 Nov 26].
Available from: https://www.cdc.gov/coronavirus/2019-ncov/php/contact-
tracing/contact-tracing-plan/contact-tracing.html

22. Lash RR, Donovan CV, Fleischauer AT, et al. COVID-19 Contact Tracing in Two Counties -
North Carolina, June-July 2020. MMWR Morb Mortal Wkly Rep. **2020**; 69(38):1360–1363.

23. Nadeem R. The Challenges of Contact Tracing as U.S. Battles COVID-19 [Internet]. Pew Res.
Cent. Internet Sci. Tech. 2020 [cited 2022 Nov 26]. Available from:
https://www.pewresearch.org/internet/2020/10/30/the-challenges-of-contact-tracing-
as-u-s-battles-covid-19/

24. Hendrix MJ, Walde C, Findley K, Trotman R. Absence of Apparent Transmission of SARS-
CoV-2 from Two Stylists After Exposure at a Hair Salon with a Universal Face Covering
Policy - Springfield, Missouri, May 2020. MMWR Morb Mortal Wkly Rep. **2020**;
69(28):930–932.

25. Doyle T, Kendrick K, Troelstrup T, et al. COVID-19 in Primary and Secondary School Settings
During the First Semester of School Reopening - Florida, August-December 2020. MMWR
Morb Mortal Wkly Rep. **2021**; 70(12):437–441.

26. Lash RR, Moonan PK, Byers BL, et al. COVID-19 Case Investigation and Contact Tracing in the
US, 2020. JAMA Netw Open. **2021**; 4(6):e2115850.

27. Alimohamadi Y, Taghdir M, Sepandi M. Estimate of the Basic Reproduction Number for
COVID-19: A Systematic Review and Meta-analysis. J Prev Med Pub Health. Korean
Society for Preventive Medicine; **2020**; 53(3):151–157.

28. Nakajo K, Nishiura H. Exploring secondary SARS-CoV-2 transmission from asymptomatic
cases using contact tracing data. Theor Biol Med Model. **2021**; 18(1):12.

29. Kinoshita R, Anzai A, Jung S-M, et al. Containment, Contact Tracing and Asymptomatic
Transmission of Novel Coronavirus Disease (COVID-19): A Modelling Study. J Clin Med.
**2020**; 9(10):3125.

30. Viner RM, Mytton OT, Bonell C, et al. Susceptibility to SARS-CoV-2 Infection Among Children
and Adolescents Compared With Adults: A Systematic Review and Meta-analysis. JAMA
Pediatr. **2021**; 175(2):143–156.

31. Kim J, Choe YJ, Lee J, et al. Role of children in household transmission of COVID-19. Arch Dis
Child. **2021**; 106(7):709–711.

32. Yung CF, Kam K, Nadua KD, et al. Novel Coronavirus 2019 Transmission Risk in Educational
Settings. Clin Infect Dis. **2021**; 72(6):1055–1058.

33. Schwartz S, Gewertz C. Omicron Is Making a Mess of Instruction, Even Where Schools Are
Open. Educ Week [Internet]. **2022** [cited 2023 Feb 20]; . Available from:
https://www.edweek.org/teaching-learning/omicron-is-making-a-mess-of-instruction-
even-where-schools-are-open/2022/01

34. Long C. Omicron Exacerbating School Staff Shortages | NEA [Internet]. [cited 2023 Feb 20].
Available from: https://www.nea.org/advocating-for-change/new-from-nea/omicron-
exacerbating-school-staff-shortages

35. 2020 Sturgis Rally Final Vehicle Counts [Internet]. [cited 2022 Nov 26]. Available from:
https://news.sd.gov/newsitem.aspx?id=27174

36. Blistein J. "Freedom-Loving People": Behind the Scenes at That Controversial Smash Mouth
Show in South Dakota [Internet]. Roll. Stone. 2020 [cited 2022 Nov 26]. Available from:
https://www.rollingstone.com/music/music-news/sturgis-covid-19-biker-fest-smash-
mouth-1043040/

37. Even the Official Motorcycle Brand of the Sturgis Rally Thinks the Mass Gathering Is Too
Risky [Internet]. [cited 2022 Nov 26]. Available from: https://news.yahoo.com/even-
official-motorcycle-brand-sturgis-
091043595.html?guce_referrer=aHR0cHM6Ly9lbi53aWtpcGVkaWEub3JnLw&guce_referr
er_sig=AQAAAE-Mbd02M7hUYmmufo9DeJ5BTNfsZOc1Z4B0b2yq9vvRmYegzua-

r6eYCg3WWyq8Hfffeq2WkWB6sR3jX1eeVKlMA_74t-hd-
vLk798wTDPvXiizPqVBjiNMviJblOYBHfn-7EGUdE_4PeQ7N-
eYEcch3PJ4C_8Tsi26OG7TPWAT&guccounter=2

38. "It's literally impossible to stop": Sturgis, South Dakota, braces as hundreds of thousands of
bikers arrived in the middle of a pandemic [Internet]. [cited 2022 Nov 26]. Available from:
https://news.yahoo.com/literally-impossible-stop-sturgis-south-231030345.html

39. Sturgis Motorcycle Rally [Internet]. Wikipedia. 2022 [cited 2022 Nov 26]. Available from:
https://en.wikipedia.org/w/index.php?title=Sturgis_Motorcycle_Rally&oldid=112059155
8

40. Dave D, McNichols D, Sabia JJ. The contagion externality of a superspreading event: The
Sturgis Motorcycle Rally and COVID-19. South Econ J. **2021**; 87(3):769–807.

41. Firestone MJ. COVID-19 Outbreak Associated with a 10-Day Motorcycle Rally in a
Neighboring State — Minnesota, August–September 2020. MMWR Morb Mortal Wkly
Rep [Internet]. **2020** [cited 2022 Nov 26]; 69. Available from:
https://www.cdc.gov/mmwr/volumes/69/wr/mm6947e1.htm

42. Harleys everywhere, masks nowhere: Sturgis expects crowd of 250,000 [Internet]. Chic.
Sun-Times. 2020 [cited 2022 Nov 26]. Available from:
https://chicago.suntimes.com/coronavirus/2020/8/7/21359318/sturgis-motorcycle-
harleys-everywhere-masks-nowhere

43. Rapier G. "If I die from the virus, it was just meant to be": 250,000 descend upon tiny South
Dakota town for world-famous motorcycle rally [Internet]. Bus. Insid. [cited 2022 Nov
26]. Available from: https://www.businessinsider.com/sturgis-motorcycle-rally-kicks-off-
despite-surging-coronavirus-cases-2020-8

44. 2020 Sturgis Motorcycle Rally attracts thousands with no mask requirements amid
pandemic [Internet]. [cited 2022 Nov 26]. Available from:
https://www.usatoday.com/picture-gallery/news/nation/2020/08/09/2020-sturgis-
motorcycle-rally-draws-thousands-no-mask-requirements-covid-19-
coronavirus/3331908001/

45. Bonacci RA, Manahan LM, Miller JS, et al. COVID-19 Contact Tracing Outcomes in
Washington State, August and October 2020. Front Public Health. **2021**; 9:782296.

46. Shelby T, Schenck C, Weeks B, et al. Lessons Learned From COVID-19 Contact Tracing During
a Public Health Emergency: A Prospective Implementation Study. Front Public Health.
**2021**; 9:721952.

47. Kortela E, Kirjavainen V, Ahava MJ, et al. Real-life clinical sensitivity of SARS-CoV-2 RT-PCR
test in symptomatic patients. PLoS ONE. **2021**; 16(5):e0251661.

48. S J, F S-R, P J, P B, M N. Diagnostic accuracy of a SARS-CoV-2 rapid antigen test in real-life
clinical settings. Int J Infect Dis IJID Off Publ Int Soc Infect Dis [Internet]. Int J Infect Dis;
**2021** [cited 2022 Nov 26]; 109. Available from:
https://pubmed.ncbi.nlm.nih.gov/34242764/

49. Cheng H-Y, Jian S-W, Liu D-P, et al. Contact Tracing Assessment of COVID-19 Transmission
Dynamics in Taiwan and Risk at Different Exposure Periods Before and After Symptom
Onset. JAMA Intern Med. **2020**; 180(9):1156–1163.

50. Hood JE, Kubiak RW, Avoundjian T, et al. A Multifaceted Evaluation of a COVID-19 Contact
Tracing Program in King County, Washington. J Public Health Manag Pract JPHMP. **2022**;
28(4):334–343.

51. WHO Coronavirus (COVID-19) Dashboard [Internet]. [cited 2023 Jan 15]. Available from:
https://covid19.who.int

52. The Lancet Commission on lessons for the future from the COVID-19 pandemic - The Lancet
[Internet]. [cited 2022 Nov 26]. Available from:
https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(22)01585-9/fulltext

53. Lewis D. Why many countries failed at COVID contact-tracing — but some got it right.
Nature. **2020**; 588(7838):384–387.

54. Kwon O. Evidence of the importance of contact tracing in fighting COVID-19. Epidemiol
Health. **2022**; 44:e2022006.

55. Tulchinsky TH. John Snow, Cholera, the Broad Street Pump; Waterborne Diseases Then and
Now. Case Stud Public Health. **2018**; :77–99.

56. Competing Theories of Cholera [Internet]. [cited 2022 Nov 27]. Available from:
https://www.ph.ucla.edu/epi/snow/choleratheories.html

57. CDC. Monkeypox in the U.S. [Internet]. Cent. Dis. Control Prev. 2022 [cited 2022 Nov 27].
Available from: https://www.cdc.gov/poxvirus/monkeypox/outbreak/us-outbreaks.html

58. COVID-19 in children and the role of school settings in transmission - second update
[Internet]. Eur. Cent. Dis. Prev. Control. 2021 [cited 2022 Nov 26]. Available from:
https://www.ecdc.europa.eu/en/publications-data/children-and-school-settings-covid-
19-transmission

59. Hershow RB. Low SARS-CoV-2 Transmission in Elementary Schools — Salt Lake County,
Utah, December 3, 2020–January 31, 2021. MMWR Morb Mortal Wkly Rep [Internet].
**2021** [cited 2022 Nov 26]; 70. Available from:
https://www.cdc.gov/mmwr/volumes/70/wr/mm7012e3.htm

60. Falk A, Benda A, Falk P, Steffen S, Wallace Z, Høeg TB. COVID-19 Cases and Transmission in
17 K-12 Schools - Wood County, Wisconsin, August 31-November 29, 2020. MMWR Morb
Mortal Wkly Rep. **2021**; 70(4):136–140.

61. Gandini S, Rainisio M, Iannuzzo ML, Bellerba F, Cecconi F, Scorrano L. No evidence of
association between schools and SARS-CoV-2 second wave in Italy [Internet]. medRxiv;
2020 [cited 2022 Nov 26]. p. 2020.12.16.20248134. Available from:
https://www.medrxiv.org/content/10.1101/2020.12.16.20248134v1

62. Zimmerman KO, Akinboyo IC, Brookhart MA, et al. Incidence and Secondary Transmission of
SARS-CoV-2 Infections in Schools. Pediatrics. **2021**; 147(4):e2020048090.

63. Johnson KE, Lachmann M, Stoddard M, et al. Detecting in-school transmission of SARS-CoV-
2 from case ratios and documented clusters. MedRxiv Prepr Serv Health Sci. **2021**;
:2021.04.26.21256136.

64. White LF, Murray EJ, Chakravarty A. The role of schools in driving SARS-CoV-2 transmission:
Not just an open-and-shut case. Cell Rep Med. **2022**; 3(3):100556.

65. Manica M, Poletti P, Deandrea S, et al. Estimating SARS-CoV-2 transmission in educational
settings: A retrospective cohort study. Influenza Other Respir Viruses [Internet]. [cited
2022 Nov 29]; n/a(n/a). Available from:
https://onlinelibrary.wiley.com/doi/abs/10.1111/irv.13049

66. Meuris C, Kremer C, Geerinck A, et al. Transmission of SARS-CoV-2 After COVID-19
Screening and Mitigation Measures for Primary School Children Attending School in
Liège, Belgium. JAMA Netw Open. **2021**; 4(10):e2128757.

67. Torres JP, Piñera C, De La Maza V, et al. Severe Acute Respiratory Syndrome Coronavirus 2
Antibody Prevalence in Blood in a Large School Community Subject to a Coronavirus
Disease 2019 Outbreak: A Cross-sectional Study. Clin Infect Dis Off Publ Infect Dis Soc
Am. **2021**; 73(2):e458–e465.

68. Gold JAW, Gettings JR, Kimball A, et al. Clusters of SARS-CoV-2 Infection Among Elementary
School Educators and Students in One School District - Georgia, December 2020-January
2021. MMWR Morb Mortal Wkly Rep. **2021**; 70(8):289–292.

69. Fontanet A, Tondeur L, Grant R, et al. SARS-CoV-2 infection in schools in a northern French
city: a retrospective serological cohort study in an area of high transmission, France,
January to April 2020. Euro Surveill Bull Eur Sur Mal Transm Eur Commun Dis Bull. **2021**;
26(15):2001695.

70. Crowe J, Schnaubelt AT, SchmidtBonne S, et al. Assessment of a Program for SARS-CoV-2
Screening and Environmental Monitoring in an Urban Public School District. JAMA Netw
Open. **2021**; 4(9):e2126447.

71. Buja A, Zabeo F, Cristofori V, et al. Opening Schools and Trends in SARS-CoV-2 Transmission
in European Countries. Int J Public Health. **2021**; 66:1604076.

72. Llupià A, Borràs-Santos A, Guinovart C, Utzet M, Moriña D, Puig J. SARS-CoV-2 transmission
in students of public schools of Catalonia (Spain) after a month of reopening. PloS One.
**2021**; 16(5):e0251593.

73. Theuring S, Thielecke M, Loon W van, et al. SARS-CoV-2 infection and transmission in school
settings during the second COVID-19 wave: a cross-sectional study, Berlin, Germany,
November 2020. Euro Surveill Bull Eur Sur Mal Transm Eur Commun Dis Bull. **2021**;
26(34):2100184.

74. Cowger TL, Murray EJ, Clarke J, et al. Lifting Universal Masking in Schools — Covid-19
Incidence among Students and Staff. N Engl J Med. Massachusetts Medical Society; **2022**;
387(21):1935–1946.

75. Lei H, Xu X, Xiao S, Wu X, Shu Y. Household transmission of COVID-19-a systematic review
and meta-analysis. J Infect. Elsevier; **2020**; 81(6):979–997.

76. Nir-Paz R, Grotto I, Strolov I, et al. Absence of in-flight transmission of SARS-CoV-2 likely due
to use of face masks on board. J Travel Med. **2020**; 27(8):taaa117.

77. Saretzki C, Bergmann O, Dahmann P, et al. Are small airplanes safe with regards to COVID-
19 transmission? J Travel Med. **2021**; 28(7):taab105.

78. Pombal R, Hosegood I, Powell D. Risk of COVID-19 During Air Travel. JAMA. **2020**;
324(17):1798.

79. Guo Q, Wang J, Estill J, et al. Risk of COVID-19 Transmission Aboard Aircraft: An
Epidemiological Analysis Based on the National Health Information Platform. Int J Infect
Dis IJID Off Publ Int Soc Infect Dis. **2022**; 118:270–276.

80. Khanh NC, Thai PQ, Quach H-L, et al. Transmission of SARS-CoV 2 During Long-Haul Flight.
Emerg Infect Dis. **2020**; 26(11):2617–2624.

81. Yang N, Shen Y, Shi C, et al. In-flight transmission cluster of COVID-19: a retrospective case
series. Infect Dis Lond Engl. **2020**; 52(12):891–901.

82. Bae SH, Shin H, Koo H-Y, Lee SW, Yang JM, Yon DK. Asymptomatic Transmission of SARS-
CoV-2 on Evacuation Flight. Emerg Infect Dis. **2020**; 26(11):2705–2708.

83. Kretzschmar ME, Rozhnova G, Bootsma MCJ, Boven M van, Wijgert JHHM van de, Bonten
MJM. Impact of delays on effectiveness of contact tracing strategies for COVID-19: a
modelling study. Lancet Public Health. **2020**; 5(8):e452–e459.

84. Johnson KE, Stoddard M, Nolan RP, White DE, Hochberg N, Chakravarty A. This time is
different: model-based evaluation of the implications of SARS-CoV-2 infection kinetics for
disease control [Internet]. medRxiv; 2020 [cited 2022 Nov 29]. p. 2020.08.19.20177550.
Available from: https://www.medrxiv.org/content/10.1101/2020.08.19.20177550v2

85. Quantifying SARS-CoV-2 transmission suggests epidemic control with digital contact tracing
| Science [Internet]. [cited 2022 Nov 26]. Available from:
https://www.science.org/doi/10.1126/science.abb6936?cookieSet=1

86. Gao Z, Xu Y, Sun C, et al. A systematic review of asymptomatic infections with COVID-19. J
Microbiol Immunol Infect. **2021**; 54(1):12–16.

87. Johansson MA, Quandelacy TM, Kada S, et al. SARS-CoV-2 Transmission From People
Without COVID-19 Symptoms. JAMA Netw Open. **2021**; 4(1):e2035057.

88. Ma Q, Liu J, Liu Q, et al. Global Percentage of Asymptomatic SARS-CoV-2 Infections Among
the Tested Population and Individuals With Confirmed COVID-19 Diagnosis: A Systematic
Review and Meta-analysis. JAMA Netw Open. **2021**; 4(12):e2137257.

89. Mahmood M, Ilyas N-U-A, Khan MF, Hasrat MN, Richwagen N. Transmission frequency of
COVID-19 through pre-symptomatic and asymptomatic patients in AJK: a report of 201
cases. Virol J. **2021**; 18(1):138.

90. Tan J, Ge Y, Martinez L, et al. Transmission roles of symptomatic and asymptomatic COVID-
19 cases: a modelling study. Epidemiol Infect. **2022**; 150:e171.

91. Just how contagious is asymptomatic Covid-19? [Internet]. [cited 2023 Mar 8]. Available
from: https://www.advisory.com/Daily-Briefing/2022/06/02/covid-transmission

92. Moghadas SM, Fitzpatrick MC, Sah P, et al. The implications of silent transmission for the
control of COVID-19 outbreaks. Proc Natl Acad Sci U S A. **2020**; 117(30):17513–17515.

93. Muller CP. Do asymptomatic carriers of SARS-COV-2 transmit the virus? Lancet Reg Health –
Eur [Internet]. Elsevier; **2021** [cited 2023 Mar 8]; 4. Available from:
https://www.thelancet.com/journals/lanepe/article/PIIS2666-7762(21)00059-4/fulltext

94. Chen PZ, Koopmans M, Fisman DN, Gu FX. Understanding why superspreading drives the
COVID-19 pandemic but not the H1N1 pandemic. Lancet Infect Dis. Elsevier; **2021**;
21(9):1203–1204.

95. Lakdawala SS, Menachery VD. Catch Me if You Can: Superspreading of COVID-19. Trends
Microbiol. **2021**; 29(10):919–929.

96. Chopping the tail: How preventing superspreading can help to maintain COVID-19 control -
ScienceDirect [Internet]. [cited 2022 Dec 1]. Available from:
https://www.sciencedirect.com/science/article/pii/S1755436520300487

97. Gupta M, Parameswaran GG, Sra MS, et al. Contact tracing of COVID-19 in Karnataka, India:
Superspreading and determinants of infectiousness and symptomatic infection. PLoS
ONE. **2022**; 17(7):e0270789.

98. Kojaku S, Hébert-Dufresne L, Mones E, Lehmann S, Ahn Y-Y. The effectiveness of backward
contact tracing in networks. Nat Phys. **2021**; 17:652–658.

99. A National Plan to Enable Comprehensive COVID-19 Case Finding and Contact Tracing in the
US. :16.

100.    Empirical evidence on the efficiency of backward contact tracing in COVID-19 | Nature
Communications [Internet]. [cited 2022 Nov 29]. Available from:
https://www.nature.com/articles/s41467-022-32531-6

101.    Bradshaw WJ, Alley EC, Huggins JH, Lloyd AL, Esvelt KM. Bidirectional contact tracing
could dramatically improve COVID-19 control. Nat Commun. Nature Publishing Group;
**2021**; 12(1):232.

102.    S0523_Oxford_-_Backwards_contact_tracing.pdf [Internet]. [cited 2022 Nov 29].
Available from:
https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachme
nt_data/file/1048768/S0523_Oxford_-_Backwards_contact_tracing.pdf

103.    Endo A, Centre for the Mathematical Modelling of Infectious Diseases COVID-19 Working
Group, Leclerc QJ, et al. Implication of backward contact tracing in the presence of
overdispersed transmission in COVID-19 outbreaks. Wellcome Open Res. **2020**; 5:239.

104.    Amara PS, Platt JE, Raj M, Nong P. Learning about COVID-19: sources of information,
public trust, and contact tracing during the pandemic. BMC Public Health. **2022**;
22(1):1348.

105.    Samuel G, Lucivero F, Johnson S, Diedericks H. Ecologies of Public Trust: The NHS COVID-
19 Contact Tracing App. J Bioethical Inq. **2021**; 18(4):595–608.

106.    Brandt AM. No Magic Bullet: A Social History of Venereal Disease in the United States
Since 1880- 35th Anniversary Edition. Oxford University Press; 2020.

107.    El-Sadr WM, Platt J, Bernitz M, Reyes M. Contact Tracing: Barriers and Facilitators. Am J
Public Health. American Public Health Association; **2022**; 112(7):1025–1033.

108.    Tanase L-M, Kerr J, Freeman ALJ, Schneider CR. COVID-19 risk perception and hoax
beliefs in the US immediately before and after the announcement of President Trump's
diagnosis. R Soc Open Sci. Royal Society; 9(8):212013.

109.    Nadeem R. Lack of Preparedness Among Top Reactions Americans Have to Public Health
Officials' COVID-19 Response [Internet]. Pew Res. Cent. Sci. Soc. 2022 [cited 2022 Nov

26]. Available from: https://www.pewresearch.org/science/2022/10/05/lack-of-
preparedness-among-top-reactions-americans-have-to-public-health-officials-covid-19-
response/

110.   Using a household-structured branching process to analyse contact tracing in the SARS-
712            CoV-2 pandemic | Philosophical Transactions of the Royal Society B: Biological Sciences
713            [Internet]. [cited 2022 Nov 29]. Available from:
714            https://royalsocietypublishing.org/doi/10.1098/rstb.2020.0267

715    111.   Davis EL, Lucas TCD, Borlase A, et al. Contact tracing is an imperfect tool for controlling
716            COVID-19 transmission and relies on population adherence. Nat Commun. Nature
717            Publishing Group; **2021**; 12(1):5412.

718    112.   Miller JS, Bonacci RA, Lash RR, et al. COVID-19 Case Investigation and Contact Tracing in
719            Central Washington State, June-July 2020. J Community Health. **2021**; 46(5):918–921.

720    113.   Stargel A, Taylor MM, Zansky S, Spencer K, Hogben M, Shultz A. Case Investigation and
721            Contact Tracing Efforts from Health Departments in the United States, November 2020–
722            December 2021. Clin Infect Dis Off Publ Infect Dis Soc Am. **2022**; :ciac442.

723    114.   Kalyanaraman N, Fraser MR. Containing COVID-19 Through Contact Tracing. Public Health
724            Rep. **2020**; 136(1):32–38.

725    115.   Spencer KD, Chung CL, Stargel A, et al. COVID-19 Case Investigation and Contact Tracing
726            Efforts from Health Departments - United States, June 25-July 24, 2020. MMWR Morb
727            Mortal Wkly Rep. **2021**; 70(3):83–87.

728    116.   Kanu FA, Smith EE, Offutt-Powell T, Hong R, Dinh T-H, Pevzner E. Declines in SARS-CoV-2
729            Transmission, Hospitalizations, and Mortality After Implementation of Mitigation
730            Measures— Delaware, March–June 2020. Morb Mortal Wkly Rep. **2020**; 69(45):1691–
731            1694.

732    117.   Sachdev DD, Brosnan HK, Reid MJA, et al. Outcomes of Contact Tracing in San Francisco,
733            California—Test and Trace During Shelter-in-Place. JAMA Intern Med. **2021**; 181(3):381–
734            383.

735    118.   Sachdev DD, Chew Ng R, Sankaran M, et al. Contact-Tracing Outcomes Among Household
736            Contacts of Fully Vaccinated Coronavirus Disease 2019 (COVID-19) Patients: San
737            Francisco, California, 29 January-2 July 2021. Clin Infect Dis Off Publ Infect Dis Soc Am.
738            **2022**; 75(1):e267–e275.

739    119.   Jones A, Fialkowski V, Prinzing L, Trites J, Kelso P, Levine M. Assessment of Day-7
740            Postexposure Testing of Asymptomatic Contacts of COVID-19 Patients to Evaluate Early
741            Release from Quarantine - Vermont, May-November 2020. MMWR Morb Mortal Wkly
742            Rep. **2021**; 70(1):12–13.

120. Matthias J. Notes from the Field: COVID-19 Case Investigation and Contact Tracing
Program — Spirit Lake Tribe, North Dakota, September–November 2020. MMWR Morb
Mortal Wkly Rep [Internet]. **2021** [cited 2022 Nov 27]; 70. Available from:
https://www.cdc.gov/mmwr/volumes/70/wr/mm7014a4.htm

121. Atherstone C, Siegel M, Schmitt-Matzen E, et al. SARS-CoV-2 Transmission Associated
with High School Wrestling Tournaments - Florida, December 2020-January 2021.
MMWR Morb Mortal Wkly Rep. **2021**; 70(4):141–143.