Subgrouping multimorbid patients with ischemic heart disease by means of unsupervised clustering: A cohort study of 72,249 patients defined by 3,046 diagnoses

Short title: Unsupervised clustering of patients with ischemic heart disease

Amalie D. Haue[1,2,¶], Peter C. Holm[1,¶], Karina Banasik[1], Agnete T. Lundgaard[1], Victorine P. Muse[1], Timo Röder[1], David Westergaard[1], Piotr J. Chmura[1], Alex H. Christensen[2,3], Peter EK. Weeke[2], Erik Sørensen[4], Ole BV. Pedersen[4,5], Sisse R. Ostrowski[4,6], Kasper K. Iversen[3], Lars V. Køber[2,6], Henrik Ullum[7], Henning Bundgaard[2,5*], Søren Brunak, PhD[1,8*]

[1]Novo Nordisk Foundation Center for Protein Research, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark
[2]Department of Cardiology, The Heart Center, Rigshospitalet, Copenhagen, Denmark
[3]Department of Cardiology, Copenhagen University Hospital, Herlev, Denmark
[4]Department of Clinical Immunology, Copenhagen University Hospital, Copenhagen, Denmark
[5]Department of Clinical Immunology, Zealand University Hospital, Køge, Denmark
[6]Department of Clinical Medicine, University of Copenhagen, Rigshospitalet, Copenhagen, Denmark
[7]Statens Serum Institut, Copenhagen, Denmark
[8]Copenhagen University Hospital, Rigshospitalet, Copenhagen, Denmark

*E-mail: soren.brunak@cpr.ku.dk (SB)

¶These authors contributed equally to this work.

Abstract

Background: There are no methods for classifying multimorbid patients with ischemic heart disease (IHD), although such methods might be clinically useful due to the marked differences in presentation and disease-course.

Methods: A population-based cohort study from a Danish secondary care setting of patients with IHD (2004-2016) and subjected to a coronary angiography (CAG) or coronary computed tomography angiography (CCTA). Data sources were The Danish National Patient Registry, in-hospital laboratory data, and genetic data from Copenhagen Hospital Biobank. Comorbidities included diagnoses assigned prior to presentation of IHD. Patients were clustered my means of the Markov Clustering Algorithm based on the entire spectrum of registered multimorbidity. The two prespecified outcomes were: New ischemic events (including death from IHD causes) and death from non-IHD causes. Patients were followed from date of CAG/CCTA until one of the two outcomes occurred or end of follow-up, whichever came first. Biological and clinical appropriateness of clusters was assessed by comparing risks (estimated from Cox proportional hazard models) in clusters and by phenotypic and genotypic enrichment analyses, respectively.

Findings: In a cohort of 72,249 patients with IHD (mean age 63.9 years, 63.1% males), 31 distinct clusters (C1-31, 67,136 patients) were identified. Comparing each cluster to the 30 others, eight clusters (9,590 patients) had statistically significantly higher (five clusters) or lower (three clusters) risk of new ischemic events; 18 clusters (35,982 patients) had a higher (11 clusters) or lower (seven clusters) risk of death from non-IHD causes. All clusters at increased risk of new ischemic events, associated with risk of death from non-IHD causes as well. Cardiovascular or inflammatory diseases were the commonly enriched in clusters (13), and distributions for 24 laboratory test results differed significantly across clusters. Polygenic risk scores for atrial fibrillation and diabetes were increased in x and y clusters respectively.

50   **Conclusions:** Clustering of patients with IHD based on comorbidities identified subgroups of

51   patients with significantly different clinical outcomes. This novel approach may support

52   differentiation of treatment intensity dependent on expected outcomes.

Non-standard abbreviations

CAG: Coronary arteriography

CCTA: Coronary computed tomography angiography

ICD-10: International Statistical Classification of Diseases and Related Health Problems 10th

Revision

IHD: Ischemic heart disease

MCL: Markov clustering

NPR: Danish National Patient Registry (NPR)

O/E-ratio: Observed-expected-ratio

PRS: Polygenic risk score

Introduction

Ischemic heart disease (IHD) is a common, chronic, complex disease and mode of onset,

disease burden and disease progression vary considerably between patients(1–3). This

heterogeneity relates to several factors, but a major contribution is multimorbidity as more

than 85% of IHD patients are diagnosed with other chronic diseases; a phenomenon coined

cardiometabolic multimorbidity(4,5). The increased mortality in patients with

cardiometabolic multimorbidity is generally only related to single disease states, such as

obstructive lung disease, diabetes, or stroke, although it is known that the risk of

cardiovascular diseases is increased in many chronic, inflammatory disorders(6,7). As more

patients at older age and with more and more co-morbidities are seen, new methods for

characterizing and studying cardiometabolic multimorbidity are needed(8–11).

Unsupervised clustering algorithms can systematically reveal structures in large, feature-rich

datasets and may be used to identify distinct patient subgroups within a heterogenous

population(12). Proof-of-concept analyses of cardiovascular phenotypes, including IHD,

heart failure, diabetes, and atrial fibrillation have already been performed(13–19). While

these studies successfully identify subgroups resembling those from traditional analyses, they

often fail to demonstrate that clustering analysis leads to novel understanding of a given

dataset. Rather, they are typically restricted to characterize high-, medium-, and low-risk

subgroups which by and large resemble more conservative approaches from an earlier, less

data-rich, epoch(20).

For decades, Danish healthcare registries have had a strong position within epidemiological

research(20–22). Given the opportunities for using clinical data more extensively, we carried

out an unsupervised clustering analysis of 72,249 patients with IHD based on their entire

disease history until IHD onset. Explicitly, we wanted to classify IHD based on the entire

spectrum of multimorbidity. We identified distinct patient subgroups derived from a total of

3,046 diagnoses assigned prior to IHD onset. The biological and clinical appropriateness

of the patient subgroups identified by unsupervised clustering analysis was asserted by

assessments of their associations with clinical outcomes and clinical characteristics,

laboratory data, and genetics (Figure 1).

Fig 1

95   **Fig 1: Graphical overview of study.** Conceptual figure displaying the cohort of patients

96   with IHD. A.: Assemblage of patient-specific vectors and an *f* x *p* matrix, where *f*

97   corresponds to the number of diagnoses and *p* corresponds to the number of included patients

98   B: Unsupervised clustering of IHD patients using the MCL algorithm. that were the basis for

99 construction of a matrix. C: which was the basis for performing unsupervised clustering to

100 identify distinct clusters. D: Cluster-based risk stratification of cohort. Red: Increased risk of

101 both outcomes. Orange: Increased risk of death from non-IHD causes, only. Yellow: No

102 statistically significant associations. Light green: Reduced risk of death from non-IHD

103 causes. Dark green: Reduced risk of new ischemic events or non-IHD causes. E: Phenotypic

104 and genetic characterization of clusters. IHD: Ischemic heart disease. MCL: Markov

105 Clustering.

Methods

Data sources, study population, and outcomes

Data from the Danish National Patient Registry (NPR) and the Danish Registry for Causes of Death were linked to in-hospital electronic health data covering the two Danish healthcare regions in Eastern Denmark (~2.9 mil inhabitants), and the Copenhagen Hospital Biobank Cardiovascular Disease Cohort(21,23,24). Linkage of different healthcare data sources was obtained via the personal identification number and only patients admitted to a hospital in Eastern Denmark in years 2004 to 2016 were considered(25). We identified all patients in NPR who were assigned an ICD-10 code for IHD(26). To increase the positive predictive value of IHD diagnoses and align included patients in time, we further required that patients had been subjected to coronary arteriography (CAG) or coronary computed tomography angiography (CCTA). To qualify that CAG/CCTAs were conclusive for IHD, patients were only included if the CAG/CCTA was performed during a contact where patients were assigned an ICD-10 code for IHD. We set the earliest CAG/CCTA fulfilling this criterium as the index date and excluded patients with an index date before year 2004 or after 2016 (Fig 2).

Fig 2

Fig 2: Flowchart: Data sources, study population, and outcomes. Gray: Identification.

Blue: Screening. Red: Eligibility. Green: Inclusion and outcomes. AMI: Acute myocardial

infarction. UAP: Unstable angina pectoris. NPR: The Danish National Patient Registry. IHD:

ischemic heart disease (ICD-10 codes I20-I25). CAG: Coronary arteriography. CCTA:

Coronary computed tomography angiography. ICD-10: International Statistical Classification

of Diseases and Related Health Problems 10th Revision. SKS: Sundhedsvæsenets

Klassifikationssystem (The Danish Health Authority Classification System).

There were two predefined outcomes: 1) New ischemic events and 2) Death from other

causes than IHD (non-IHD causes). The outcome "new ischemic event" was a composite

outcome of a) hospitalization minimum 30 days after index for myocardial infarction or

unstable angina pectoris (i.e., hospitalization with myocardial infarction or unstable angina

pectoris as the primary diagnosis), b) revascularization not related to the index date and c)

any death where IHD was listed as the primary or secondary cause. Outcomes were obtained

from NPR and Danish Registry for Causes of Death. Eligible codes for inclusion, outcomes

and specific cutoffs are available in S1 Fig and S1 Table.

Data preprocessing and application of the Markov cluster algorithm

We performed a clustering analysis of included patients based on their multimorbidity prior

to their IHD diagnosis (index) using the Markov cluster (MCL) algorithm(27).

Multimorbidity was represented as patient-specific vectors using diagnoses assigned prior to

or at index. ICD-10 codes assigned to less than five patients (n=1,673) were excluded from

the analysis owing to legal regulations in Denmark. As we focused the studies on

multimorbidity in IHD, ICD-10 codes for IHD (I20-I25) were excluded from patients-specific

vectors. Thus, a total of 3,046 ICD-10 codes were the basis for constructing a patient

similarity network that was used as MCL algorithm input. Patient-specific vectors consisted

of length 3,046 with integers indicating the number of times a patient had been assigned a

particular ICD-10 code. The length of the vectors corresponded to the number of input

features (ICD-10 codes). By combining the patient-specific vectors from all included

patients, a matrix of size p x f was constructed, where p indicates the number of included

patients and f indicates the number of input features (ICD-10 codes). Following a series of

preprocessing steps, a patient similarity network was created based on the p x f matrix and

used as input for the MCL algorithm(28). Resulting clusters were denoted C followed by an

integer indicating the rank of the clusters with respect to cluster size (number of patients).

Thus, C1 denotes the largest cluster and cluster-membership was used to denote a cluster as a

covariate in subsequent analyses. Robustness of clustering was assessed by generating a

series of diluted and shuffled versions of the resulting clusters (reference clustering), and

their similarity was quantified using the variance of information measure as previously

described(29). Explicitly, a series of diluted and shuffled versions of the input graph were

generated(30). In total, 500 variations of the input graph were constructed by shuffling and

deleting edges, respectively. The variation in the graphs was then quantified by means of

variation of information parameter. Details regarding the MCL settings and a description of

cluster robustness assessment are available in the S1 Appendix.

Preprocessing of laboratory and genetic data

Clusters were characterized by laboratory and genetic data based on the subset of patients

where these data types were available. A panel of 25 different lab parameters was included in

the analyses. Only tests taken up to 90 days before index or at the day of index were included.

Included lab tests were plasma levels of potassium, sodium, hemoglobin, estimated

glomerular filtration rate (eGFR), creatinine, carbamide, glucose, troponin (I/T), HDL

cholesterol, LDL cholesterol, total cholesterol, leukocytes, C-reactive protein, lymphocytes,

monocytes, neutrophils, basophils, platelets, INR, alanine transaminase, albumin, alkaline

phosphatase, bilirubin, and triglyceride. For every cluster, a *score* was computed based on the

number of patients with a lab test below, within, or above the standard reference value,

indicated by -1, 0 and 1, respectively. *Score* was defined as the mean of the summarized

values per cluster.

Autosomal genotype data were obtained by identifying included patients who were also

among the study participants in the Copenhagen Hospital Biobank – Cardiovascular Disease

Cohort(24). For included patients with genetic data available, we calculated polygenic risk

scores (PRSs) for 14 traits, obtained from nine GWAS meta-analyses (atrial fibrillation, BMI-

adjusted non-insulin diabetes, chronic kidney disease, HDL cholesterol levels, heart failure,

LDL cholesterol levels, stroke, total cholesterol levels, triglyceride levels)(31–36) and five

GWAS (acute myocardial infarction, coronary artery disease, diastolic blood pressure, non-

alcoholic fatty liver disease, systolic blood pressure)(37–40). PRSs were calculated using the

"LDpred2-auto" algorithm, implemented in the R package "bigsnpr" (version 1.11.6) with R

version 4.0.0 and the workflow management system Snakemake(41–43). Each trait's PRS

distribution was scaled to a mean of zero and a standard deviation of one.

Statistical analyses of clusters identified by the MCL algorithm

As the study was designed to identify patient subgroups and not individual variation, clusters

of size < 500 were excluded from the remaining analyses. Mean age at IHD onset in each

cluster was compared to the mean age at onset in all the other clusters using Tukey's Honest

Significant Difference (HSD) method. Significance level was set to 0.05 and P-values were

adjusted assuming 465 tests (adj. P-val.).

To investigate the association between cluster-membership and the competing risks of new

ischemic events and death from non-IHD causes, we used Cox proportional-hazards models

(Cox models). Patients were followed from index until occurrence of either of the two

outcomes, or end of follow-up (year 2018), whichever came first. The dependent variable was

either risk of new ischemic events or death from non-IHD causes, and the independent

variables were cluster, sex, and age at index. To age-adjust the models, analyses were

performed using restricted cubic spline with three knots for age at index. Follow-up time was

truncated to a maximum of five years. For each cluster, hazard ratios (HRs) and 95%

confidence intervals (CIs) were estimated by comparing HRs for the members of the cluster

with the HRs with that of non-members.

Further characterization of clusters consisted of: (1) phenotypic enrichment analysis, (2)

characterization of clusters with respect to their laboratory profiles and (3) a test for genetic

enrichment. The phenotypic enrichment analysis was carried out based on ratios between

Observed (O) and Expected (E) frequencies of diagnoses in the clusters (O/E-ratios). That is,

ratios between the frequencies of ICD-10 codes in each cluster (observed frequencies) and

the frequencies of ICD-10 codes in the entire population (expected frequencies) were

calculated and expressed as O/E-ratios(44). In subsequent characterization of clusters,

enrichment denoted O/E-ratios > 2, and clusters were characterized as having little

enrichment if the sum of the ten largest O/E-ratios < 50. Inverse changes were used to denote

O/E-ratios between 0 and 1.

Hierarchical clustering was applied to estimate the cluster similarity with respect to the

laboratory tests using the Euclidean distance between the *score* of each cluster for each test.

For each of the fourteen traits we calculated PRSs for, we used Wilcoxon rank-sum tests to

compare the PRS distribution of each cluster to the combined PRS distribution of PRSs in all

other clusters. Resulting P-values were converted to the false discovery rate (FDR) to account

for multiple testing, with a total of 434 tests. We report effect sizes as calculated by the

"wilcox.test" function built into R version 4.0.0. Level of significance was set to FDR < 0.05,

assuming 434 tests.

Further details regarding preprocessing and analyses of laboratory and genetic data are

available in the S2 Appendix.

Results

Cohort demographics and co-morbidities

A total of 72,249 patients (63.1% males, mean age 63.9 years) were included (Table 1).

Angina pectoris (I20) was the most common IHD diagnosis (38,239 patients, 52.9%),

followed by acute myocardial infarction (I21) (33,229 patients, 46.0 %) and chronic IHD

(I25) (22,750 patients, 31.5%). The most common co-morbidity prior to the IHD index was

hypertension (I10.9) (24,818 patients, 34.4%) followed by dyslipidemia (E78.0) (12,780

patients, 17.7%) and non-insulin dependent diabetes (E11.9) (7,551 patients, 10,5%). Prior to

index, the mean number of diagnoses per patient was 8.1. A total of 68,103 patients (94.3%)

were co-morbid at index. The overall incidence (new ischemic events and death from non-

IHD causes) was 94 events per 1000 person-years (Table 1).

Table 1: Patient demographics, comorbidities, and outcomes

Cohort demographics	Total	Males	Females
Number of patients (%)	72,249	45,576 (63.1)	26,673 (36,1)
Mean age at index (SD)	63.9 (11.9)	62.9 (11.6)	65.6 (12.1)
IHD manifestations (ICD-10)	**Total**	**Males**	**Females**
Angina pectoris (I20)	38,239	22,628	15,611
Acute myocardial infarction (I21)	33,299	27,720	10,579
Subsequent myocardial infarction (I22)	61	34	27
Certain current complications following acute myocardial infarction (I23)	138	92	46
Other acute ischemic heart diseases (I24)	1,341	814	527
Chronic ischemic heart disease (I25)	22,750	14,589	8,152
Common comorbidities (ICD-10)	**Total**	**Males**	**Females**
Primary (essential) hypertension (I10.9)	24,818	14,508	10,310
Hypercholesterolemia (E78.0)	12,780	7,842	4,938
Non-insulin dependent diabetes (E11.9)	7,551	4,891	2,660
Atrial fibrillation and atrial flutter, unspecified (I48.9)	7,075	4,509	2,566
Heart failure, unspecified (I50.9)	6,160	4,059	2,101
Chest pain, unspecified (R07.9)	5,863	3,441	2,422
Senile cataract, unspecified (H25.9)	5,764	2,795	2,969
Pneumonia, unspecified (J18.9)	5,469	3,236	2,260
Hyperlipidemia, unspecified (E78.5)	5,002	3,306	1,696
Chronic obstructive pulmonary disease (J44.9)	4,621	2,449	2,172
Outcomes, number of cases	**Total**	**Males**	**Females**
New ischemic events (%)	14,679	10,152	4,527
■ *Myocardial infarction*	*5,833*	*3,709*	*2,124*
■ *Revascularization*	*6,282*	*4.718*	*2,124*
■ *Death caused by IHD*	*2,563*	*1.724*	*839*
Death from non-IHD causes (%)	10,684	6,710	3,974
Censored (%)	46,886	28,713	18,172
Outcomes, time to event	**Mean time to event in years (SD)**		
	Total	**Males**	**Females**
New ischemic events	1.48 (1.40)	1.49 (1.41)	1.48 (1.40)
■ Myocardial infarction	2.40 (1.87)	2.41 (1.89)	2.38 (1.85)
■ Revascularization	2.25 (1.88)	2.28 (1.89)	2.16 (1.84)
■ Death caused by IHD	1.92 (1.13)	1.95 (2.02)	1.88 (2.05)
Death from non-IHD causes	2.16 (1.50)	2.14 (1.49)	2.20 (1.51)
Censored	4.37 (1.08)	4.36 (1.09)	4.39 (1.06)
Total	3.72 (1.64)	3.67 (1.67)	3.81 (1.60)

 Unsupervised clustering of multimorbid patients with IHD

In the cohort, the MCL algorithm identified 36 distinct clusters based on the set of 3,046

ICD-10 codes assigned to the patients prior to or at index. The 36 clusters contained a total of

68,084 patients. The remaining 4,365 patients in the cohort (6,0%) that did not cluster were

primarily patients with no diagnoses prior to index (>99%). This observation served as a

negative control as the MCL algorithm correctly identified patients without registered co-

morbidity prior to index. Further, clusters robustness was assessed as described in Methods

with a reasonable variation of information. Next, the 31 of the 36 clusters with >500 patients

(67,136 patients) were characterized (Table 2). Using Tukey's HSD to compare the age at

index between all 31 clusters (a total of 466 combinations), we found significant differences

in 391 comparisons (S3 Table). For demographics of patients that did not cluster or were in

clusters of size < 500, see S4 Table.

Table 2: Cluster demographics, characteristics, and associations with outcomes

Cluster	Size	Mean age at index in years (SD)	Males	Females	New ischemic events		Death from non-IHD causes	
					HR	Adj. P-val.	HR	Adj. P-val.
C1	7,191	64.8 (11.3)	3,897	3,294	1.000	> 0.050	0.856	> 0.050
C2	5,990	58.6 (11.5)	2,862	3,127	0.825	< 0.001	0.600	< 0.001
C3	4,641	56.8 (11.4)	2,727	1,914	0.757	< 0.001	0.586	< 0.001
C4	4,401	69.6 (10.2)	2,853	1,548	0.920	> 0.050	1.461	< 0.001
C5	4,290	63.9 (10.7)	2,803	1,487	1.402	< 0.001	1.629	< 0.001
C6	3,589	59.7 (10.9)	2.388	1,201	0.969	> 0.050	0.675	< 0.001
C7	3,309	63.8 (11.0)	2,025	1,284	0.889	> 0.050	0.611	< 0.001
C8	2,802	71.1 (10.9)	1,867	935	0.943	> 0.050	0.842	> 0.050
C9	2,581	63.7 (11.8)	1,803	778	1.314	< 0.001	1.789	> 0.050
C10	2,562	74.2 (9.6)	1,225	1,337	0.978	> 0.050	0.928	> 0.050
C11	2,292	66.1 (11.0)	2,186	106	0.926	> 0.050	0.650	< 0.001
C12	2,213	70.3 (10.2)	2,068	145	0.920	> 0.050	0.805	> 0.050
C13	2,070	58.6 (10.2)	1,348	722	0.946	> 0.050	0.577	< 0.050
C14	2,070	68.2 (9.6)	1,030	1,010	1.146	> 0.050	3.390	< 0.001
C15	2,040	63.9 (10.1)	1,208	805	1.031	> 0.050	0.784	> 0.050
C16	1,654	64.1 (12.1)	1,013	641	1.107	> 0.050	1.761	< 0.001
C17	1,281	65.3 (9.9)	714	567	1.001	> 0.050	1.761	< 0.001
C18	1,251	68.2 (9.8)	802	449	1.790	< 0.001	3.421	< 0.001
C19	1,168	58.5 (9.7)	995	173	0.752	< 0.050	1.571	> 0.050
C20	1,119	71.5 (11.3)	713	406	1.213	> 0.050	1.782	< 0.001
C21	1,000	61.0 (11.0)	769	231	1.116	> 0.050	0.890	> 0.050
C22	988	69.2 (10.4)	516	472	1.023	> 0.050	0.978	> 0.050
C23	935	58.7 (12.2)	588	347	1.609	< 0.001	2.275	< 0.001
C24	932	67.9 (10.1)	28	904	0.787	> 0.050	1.589	< 0.001
C25	860	56.2 (9.9)	664	196	0.978	> 0.050	2.691	< 0.001
C26	852	58.7 (12.1)	391	461	0.939	> 0.050	1.108	> 0.050
C27	823	65.1 (10.9)	532	291	1.201	> 0.050	1.289	> 0.050
C28	686	71.7 (8.0)	673	13	0.866	> 0.050	1.786	< 0.001
C29	550	57.2 (11.1)	435	115	0.906	> 0.050	0.985	> 0.050
C30	533	61.2 (11.7)	391	172	1.874	< 0.001	5.364	< 0.001
C31	520	64.4 (11.2)	213	307	1.052	> 0.050	1.484	> 0.050
NA*	5,113	60.1 (11.1)	3,878	1,235	NA	NA	NA	NA

*Patients that did not cluster or were in clusters of size < 500

Clusters, clinical outcomes, and phenotypic enrichment

To assess if the unsupervised clustering identified patient subgroups at different risks of

disease progression, we used cluster-membership (C1-C31) as a covariate in a series of Cox

models. A total of 14,679 patients experienced a new ischemic event during follow-up and

10,684 patients died from other causes than IHD. Mean follow-up times was 3.72 years

(Table 1). Risks for new ischemic events and death from non-IHD causes in each cluster were

compared to the pooled risk for patients in the remaining 30 clusters. The survival analysis

demonstrated that the MCL algorithm stratified patients according to risk of new ischemic

events and death from non-IHD causes (Fig 3). Comparing each cluster to all the others

(n=30), seven clusters (20,221 patients) had a statistically significantly higher (five clusters)

or lower (two clusters) risk of new ischemic events; and 18 clusters (43,173 patients) had a

higher (14 clusters) or lower (four clusters) risk of death from non-IHD causes. All clusters at

increased risk of new ischemic events, associated with risk of death from non-IHD causes as

well. Thus, 11 clusters (23,963 patients) did not have altered risk of the two outcomes, when

compared to the other clusters (Fig 1D).

Fig 3

Fig 3: Risk of new ischemic events and non-IHD causes stratified by cluster. Forest
plots where clusters are shown against HR for new ischemic events (left) and death from non-
IHD causes (right). X-axis: HR for a single cluster relative to mean HR of the 30 other
clusters. Y-axis: Clusters arranged by risk of new ischemic events, increasing risk from top to
bottom. IHD: Ischemic heart disease. HR: Hazzard ratio.

The distribution of O/E-ratios was heavily left-skewed as less than 99% (n=101) of all O/E-

ratios were >10 and roughly 7% (n=887) of all O/E-ratios were >2. About 60% of all O/E-

ratios (n=8,056) were in the range of 0 and 1 corresponding to inverse changes. Generally,

clusters that had high risk of new ischemic events or death from non-IHD causes were also

characterized by large, summarized O/E-values corresponding to a high degree of

multimorbidity (S5 Table). To obtain an overview of the results of the risk stratification in

conjunction with the enrichment analysis, clusters were described according to three

categories: (1) clusters enriched for cardiometabolic or -vascular diseases, (2) clusters

enriched for degenerative or inflammatory diseases and (3) clusters enriched characterized by

little enrichment and inverse changes.

Clusters enriched for cardiometabolic and -vascular diseases

Four of the five clusters at increased risk of new ischemic events (and death from non-IHD

causes) were enriched for diabetes (C5, C18, C23, and C30). In these four clusters, HRs

ranged from 1.40 (C5, 95%CI: 1.30;1.50, adj. P-val. < 0.001) to 1.88 (C30, 95%CI:

1.60;2.00, adj. P-val. < 0.001) with a significant difference in age at index (C5: 63.9 years,

C30: 61.2 years, Adj. P-val. < 0.001, TukeyHSD). C18 and C23 were only enriched for

insulin-dependent diabetes, but differed in that C18 was also enriched for insulin-dependent

diabetes with vascular complications and periphery atherosclerosis. In contrast, C5 was only

enriched for non-insulin dependent diabetes and included diabetes with as well as without

complications. Lastly, C30 was only enriched for diabetes with complications (insulin and

non-insulin dependent) and was the diabetes cluster enriched for chronic kidney disease and

bacterial infections, as well (S5 Table).

Other cardiac diseases that displayed enrichment were supraventricular arrythmias (C4),

cardiomyopathies (C9), and valve diseases (C20). Of the three clusters, only C9 had

increased risk of new ischemic events (HR: 1.31 (C9, 95%CI: 1.20;1.44, adj. P-val.: < 0.001).

Risk of death from non-IHD causes was 1.79 (95%CI: 1.60;2.00, adj. P-val. < 0.001). In

contrast, C4 and C20 only had increased risk of death from non-IHD causes with HRs of 1.49

(C4, 95%CI: 1.34;1.59, adj. P-val. < 0.001) and 1.78 (C20, 95%CI: 1.54;2.04, adj. P-val. <

0.001). Interestingly, the cluster enriched for cerebrovascular diseases (C27) did not have

altered risk of any of the two outcomes. In sum, all clusters that had increased risk of new

ischemic events were enriched for cardiometabolic diseases, albeit not all clusters enriched

for cardiometabolic and -vascular diseases had increased risk of new ischemic events (Table

2 and S5 Table).

Clusters enriched for degenerative or inflammatory diseases

Six clusters (C7, C13, C14, C22, C26, and C31) were enriched for diagnoses describing

degenerative or inflammatory diseases, i.e., osteoarthritis (C7), degenerative spine disease

(C13 and C22), chronic obstructive pulmonary disease (C14), asthma (C26), and rheumatoid

arthritis (C31). Remarkably, none of the four clusters had increased risk of new ischemic

events and only one cluster (C14) had increased risk of death from non-IHD causes (HR:

3.39, 95%CI: 3.09;3.71, adj. P-val. < 0.001). Conversely, C7 and C13 had reduced risk of

death from non-IHD causes (C7, HR: 0.61, 95%CI: 0.52;0.72, adj. P-val. < 0.001 and C13,

HR: 0.58, 95%CI: 0.45;0.74, adj. P-val. < 0.001). Age at index for the clusters enriched for

degenerative or inflammatory diseases range between 58.6 years (C13) and 69.2 years (C22)

(Table 2). Taken together, these findings hint to the dual nature of inflammation as a potential

disease modifier as well as a risk factor.

Clusters characterized by little enrichment and inverse changes

Six clusters (C1, C2, C3, C6, C15, and C17) were characterized by little enrichment, which

included the two clusters with reduced risk of new ischemic events (C2, HR: 0.82, 95%CI:

0.76;0.89, adj. P-val. < 0.001 and C3, HR: 0.76, 95%CI: 0.52;0.69, adj. P-val. < 0.001). Not

surprisingly, none of these six clusters had increased risk of either of the two outcomes, but

three clusters (C2, C3, and C6) had reduced risk of death from non-IHD causes (C2, HR:

0.60, 95%CI: 0.52;0.69, adj. P-val. < 0.001, C3, HR: 0.59, 95%CI: 0.59;0.69, adj. P-val. <

0.001 and C6, HR: 0.68, 95%CI: 0.57;0.79, adj. P-val. < 0.001) (Table 2). It was a common

attribute of the clusters without altered risk of any of the two outcomes that O/E-ratios for

hypertension and dyslipidemia were among the largest. In contrast, diabetes, heart failure,

and chronic obstructive pulmonary disease frequently displayed inverse changes (O/E-ratios

< 1) in these clusters (S5 Table). Taken together, these observations indicate that risk of

disease progression in this populations necessitates a more sophisticated analysis of

multimorbidity.

For a list with results of the enrichment analysis for all clusters, including the 13 clusters not

described above, S5 Table.

Clusters and their association with laboratory measurements and genetic data

Clusters were also characterized by means of datatypes not included among the MCL

algorithm input features. For patients in the 31 clusters, we had laboratory measurements on

30,755 (49.5%) and genetic data on 19,422 (31.3%). To assess if the phenotypic differences

captured by the MCL algorithm were also reflected in laboratory measurements, we tested if

the distributions of test results within and out of reference ranges differed significantly. There

were significantly different distributions of tests within and out of reference ranges in clusters

for the 24 most frequent tests. Overall, this indicates that the phenotypic patterns within the

entire spectrum of cardiovascular multimorbidity registered before index correlate with

results of clinical laboratory tests (S6 Table). Thus, these findings are a strong indicator that

the patterns captured by the MCL algorithm are biologically relevant. For a graphical

summary of the laboratory scores in each cluster, see S4 Figure.

Finally, we identified 41 cases (out of 434 tests) where the PRS distribution for a specific

trait in cluster was significantly different from that trait's combined PRS distribution of the

other 30 clusters. Among these cases, we found the largest effects size to be a higher genetic

risk for atrial fibrillation in cluster C4 (0.57, FDR < 0.001) as well as a higher genetic risk for

non-insulin dependent diabetes in cluster C5 (0.55, FDR < 0.001). These findings are

congruent with the results of the enrichment analysis for C4 and C5, respectively. In contrast,

C1 (phenotypically characterized by inverse changes) had relatively large, positive effect

sizes for systolic as well as diastolic blood pressure (0.20 and 0.16, FDR < 0.001). Similarly,

there were positive effect sizes for total cholesterol and triglycerides in C6, which was also

characterized by little phenotypic enrichment as well as a high degree of inverse changes. A

list of significant effect sizes for the 41 significant cases, see S7 Table.

Discussion

In this study, we developed a novel, data-driven method for structuring the entire spectrum of

multimorbidity by means of an unsupervised clustering analysis. In a cohort of 72,249

patients with IHD patients, we identified 31 distinct clusters (67,136 patients) based on 3,046

diagnoses assigned prior to or at index. By comparing risk of new ischemic events and death

from non-IHD causes across clusters and then performing an enrichment analysis, we found

that clusters at increased risk of new ischemic events were enriched for diabetes (four

clusters) or cardiomyopathies (one cluster). Neither the cluster enriched for supraventricular

arrythmias, nor valve diseases had increased risk of new ischemic events. Degenerative and

inflammatory diseases were enriched in a total of six clusters and displayed no clear trend in

their relation to the outcomes. The results of the enrichment analysis were supported by

trends in laboratory test results and clusters enriched for supraventricular arrythmias and non-

insulin diabetes also had congruently, higher genetic risks.

The results of the study agree with common knowledge on risk of IHD, while also adding

insights to the disease-diseases associations, which are currently underappreciated in the

literature. The fact that clusters enriched for diabetes were generally the most high-risk

clusters serves as a methodological reality check(6). Added value of the study lies in the fact

that the method allows for a more sophisticated description of such associations, as the

method allows to study the entire spectrum of multimorbidity. For example, four clusters

were enriched for diabetes, which is in line with the current paradigm that a single term is

insufficient to describe a multifactorial disease, such as diabetes(16,29). By integrating

different data types, the findings indicate how phenotypic and genetic data complement each

other, by exemplifying (1) that clustering analysis facilitates stronger genetic signals in

patient subgroups and (2) that genetic data may unveil patterns not captured by phenotypic

data alone.

In addition, the method developed in this study and subsequent findings add perspective to

the relatively limited body of literature regarding associations between chronic inflammatory

and cardiovascular diseases(7). While previous studies have concluded that the risk of

cardiovascular diseases is increased in most chronic inflammatory disorders, the results of

our study indicate that pre-existing degenerative or inflammatory disorders in patients with

IHD do not increase the risk of new ischemic events.

The pre-selected outcomes in the present study are also a unique aspect of the study, as

previous clustering analyses within the cardiovascular domain studies have mainly analyzed

all-cause mortality(17,18). This aspect of the study allows to distinguish between risk of

progression related to IHD and risk of progression that is related to comorbidity drawing

attention to important aspects of multimorbidity in this domain. For example, clusters

enriched for supraventricular arrythmias and chronic obstructive pulmonary disease,

respectively, only had increased risk of death from non-IHD causes. The study design,

including the enrichment analysis, also revealed that classical risk factors for IHD (e.g.,

hypertension and dyslipidemia) did not drive the clustering. This finding agrees with

previously published comorbidity phenotypes in patients with IHD(18). We argue that the

present study displays that continuous exploration and characterization of multimorbidity in

IHD are key elements in optimizing the exploit the full potential of continuously developing

treatment strategies.

Previous clustering analyses within the cardiovascular domain have typically included either

thousands of patients or hundreds of input features, but not both(14,15). For example, Hall et

al. defined multimorbidity using only eight different chronic conditions, whereas Crowe et al.

defined multimorbidity with reference to 20 predefined conditions(17,18). Thus, the scale of

our study exceeds that of previous work, as it includes more than 70,000 patients and more

than 3,000 input features. And further, we limited the risk of introducing bias by not exerting

feature selection prior to clustering.

The two main limitations with respect to the data foundation are that (1) owing to the novelty

of the method, there were no standardized way of assessing the representation of

multimorbidity and (2) it was only a subset for which laboratory and genetic data were

available. These challenges are naturally overcome in clustering analyses based on data from

randomized controlled trials, such as the studies by Inohara et al, and Karwath et al.(15,19)

However, in the present, data-rich era, we argue that it is highly important to develop

methods for structuring and studying other data than what is being collected for trials. Ideally,

the two approaches, based on nationwide data and randomized controlled trials, respectively,

will complement each other; and will facilitate more precise identification of patients who are

likely to benefit from different treatment options as well as guide optimized selection of

patients for randomized controlled trials .

In sum, the study further showcases the strengths of a more fine-grained analysis of patient

subgroups, which, in turn, may pave the way for successful implementation of precision

medicine. Owing to its flexibility, the comprehensive, data-driven analysis of cardiovascular

multimorbidity represents a novel method for characterizing multimorbidity in IHD with

great potential of applying it to other diseases of interest or other clinical data. Such trends

may guide clinical decision making in cases, where for example it is not obvious how to

manage the angiographic findings or the combination of drugs that a specific patient will

benefit most from.

In conclusion, the present study cements the complexity of multimorbid patients with IHD

and exemplifies the clinical relevance of a more fine-grained patient subgrouping by carrying

out a cluster-based risk-stratifying the cohort. Further, owing to its flexibility, the

comprehensive, data-driven method of cardiovascular multimorbidity presented here

represents a novel method for characterizing multimorbidity in IHD with great potential.

Improved patient subgrouping may be critical guide future clinical decision making in cases,

where it is non-trivial how to manage the angiographic findings or to find the optimal

combination of drugs for a given patient.

Acknowledgement

The authors would like to thank (1) research programmer, Troels Siggaard, Novo Nordisk Foundation Center for Research, University of Copenhagen, Denmark for continuous and reliable infrastructure support, and (2) Head of Cardiovascular Research, Hilma Hólm, deCODE genetics, Iceland for insightful comments.

References

1. Antman EM, Braunwald E. Managing Stable Ischemic Heart Disease. N Engl J Med. 2020 Apr 9;382(15):1468–70.

2. Ferraro R, Latina JM, Alfaddagh A, Michos ED, Blaha MJ, Jones SR, et al. Evaluation and Management of Patients With Stable Angina: Beyond the Ischemia Paradigm. J Am Coll Cardiol. 2020 Nov 10;76(19):2252–66.

3. Nabel EG, Braunwald E. A tale of coronary artery disease and myocardial infarction. N Engl J Med. 2012 Jan 5;366(1):54–63.

4. Forman DE, Maurer MS, Boyd C, Brindis R, Salive ME, Horne FM, et al. Multimorbidity in Older Adults With Cardiovascular Disease. J Am Coll Cardiol. 2018 May 15;71(19):2149–61.

5. Afilalo J, Alexander KP, Mack MJ, Maurer MS, Green P, Allen LA, et al. Frailty Assessment in the Cardiovascular Care of Older Adults. J Am Coll Cardiol. 2014 Mar 4;63(8):747–62.

6. The Emerging Risk Factors Collaboration. Association of Cardiometabolic Multimorbidity With Mortality. JAMA. 2015 Jul 7;314(1):52–60.

7. Dregan A, Charlton J, Chowienczyk P, Gulliford MC. Chronic Inflammatory Disorders and Risk of Type 2 Diabetes Mellitus, Coronary Heart Disease, and Stroke. Circulation. 2014 Sep 2;130(10):837–44.

8. Glynn LG. Multimorbidity: another key issue for cardiovascular medicine. The Lancet. 2009 Oct 24;374(9699):1421–2.

9. Joshi A, Rienks M, Theofilatos K, Mayr M. Systems biology in cardiovascular disease: a multiomics approach. Nat Rev Cardiol. 2021 May;18(5):313–30.

10. Khera Amit V., Kathiresan Sekar. Is Coronary Atherosclerosis One Disease or Many? Circulation. 2017 Mar 14;135(11):1005–7.

11. Rahimi K, Lam CSP, Steinhubl S. Cardiovascular disease and multimorbidity: A call for interdisciplinary research and personalized cardiovascular care. PLOS Med. 2018 Mar 27;15(3):e1002545.

12. Kaufman L, Rousseeuw PJ. Finding Groups in Data: An Introduction to Cluster Analysis. John Wiley & Sons; 2009. 369 p.

13. Shah RV, Yeri AS, Murthy VL, Massaro JM, D'Agostino R Sr, Freedman JE, et al. Association of Multiorgan Computed Tomographic Phenomap With Adverse Cardiovascular Health Outcomes: The Framingham Heart Study. JAMA Cardiol. 2017 Nov 1;2(11):1236–46.

14. Ahmad T, Pencina MJ, Schulte PJ, O'Brien E, Whellan DJ, Piña IL, et al. Clinical implications of chronic heart failure phenotypes defined by cluster analysis. J Am Coll Cardiol. 2014 Oct 28;64(17):1765–74.

15. Inohara T, Shrader P, Pieper K, Blanco RG, Thomas L, Singer DE, et al. Association of
of Atrial Fibrillation Clinical Phenotypes With Treatment Patterns and Outcomes: A
Multicenter Registry Study. JAMA Cardiol. 2018 Jan 1;3(1):54–63.

16. Ahlqvist E, Storm P, Käräjämäki A, Martinell M, Dorkhan M, Carlsson A, et al. Novel
subgroups of adult-onset diabetes and their association with outcomes: a data-driven
cluster analysis of six variables. Lancet Diabetes Endocrinol [Internet]. 2018 Mar;0(0).
Available from: http://www.ncbi.nlm.nih.gov/pubmed/29503172

17. Hall M, Dondo TB, Yan AT, Mamas MA, Timmis AD, Deanfield JE, et al.
Multimorbidity and survival for patients with acute myocardial infarction in England and
Wales: Latent class analysis of a nationwide population-based cohort. PLOS Med. 2018
Mar 6;15(3):e1002501.

18. Crowe F, Zemedikun DT, Okoth K, Adderley NJ, Rudge G, Sheldon M, et al.
Comorbidity phenotypes and risk of mortality in patients with ischaemic heart disease in
the UK. Heart. 2020 Jun 1;106(11):810–6.

19. Karwath A, Bunting KV, Gill SK, Tica O, Pendleton S, Aziz F, et al. Redefining β-
blocker response in heart failure patients with sinus rhythm and atrial fibrillation: a
machine learning cluster analysis. The Lancet. 2021 Oct 16;398(10309):1427–35.

20. Bowman L, Baras A, Bombien R, Califf RM, Chen Z, Gale CP, et al. Understanding the
use of observational and randomized data in cardiovascular medicine. Eur Heart J. 2020
Jul 14;41(27):2571–8.

21. Schmidt M, Schmidt SAJ, Adelborg K, Sundbøll J, Laugesen K, Ehrenstein V, et al. The
Danish health care system and epidemiological research: from health care contacts to
database records. Clin Epidemiol. 2019;11:563–91.

22. Hemingway H, Asselbergs FW, Danesh J, Dobson R, Maniadakis N, Maggioni A, et al.
Big data from electronic health records for early and late translational cardiovascular
research: challenges and potential. Eur Heart J. 2018 Apr 21;39(16):1481–95.

23. Helweg-Larsen K. The Danish Register of Causes of Death. Scand J Public Health. 2011
Jul;39(7 Suppl):26–9.

24. Sørensen E, Christiansen L, Wilkowski B, Larsen MH, Burgdorf KS, Thørner LW, et al.
Data Resource Profile: The Copenhagen Hospital Biobank (CHB). Int J Epidemiol
[Internet]. 2020 Nov 10 [cited 2020 Dec 13];(dyaa157). Available from:
https://doi.org/10.1093/ije/dyaa157

25. Schmidt M, Pedersen L, Sørensen HT. The Danish Civil Registration System as a tool in
epidemiology. Eur J Epidemiol. 2014 Aug 1;29(8):541–9.

26. Sundbøll J, Adelborg K, Munch T, Frøslev T, Sørensen HT, Bøtker HE, et al. Positive
predictive value of cardiovascular diagnoses in the Danish National Patient Registry: a
validation study. BMJ Open [Internet]. 2016 Nov 1 [cited 2020 Jan 15];6(11). Available
from: https://bmjopen.bmj.com/content/6/11/e012832

27. Enright AJ, Van Dongen S, Ouzounis CA. An efficient algorithm for large-scale
detection of protein families. Nucleic Acids Res. 2002 Apr;30(7):1575–84.

28. MCL - a cluster algorithm for graphs [Internet]. [cited 2023 Jan 30]. Available from:
http://micans.org/mcl/

29. Kirk IK, Simon C, Banasik K, Holm PC, Haue AD, Jensen PB, et al. Linking glycemic
dysregulation in diabetes to symptoms, comorbidities, and genetics through EHR data
mining. Valencia A, Barkai N, editors. eLife. 2019 Dec 10;8:e44941.

30. Meilă M. Comparing clusterings—an information based distance. J Multivar Anal. 2007
May 1;98(5):873–95.

31. Nielsen JB, Thorolfsdottir RB, Fritsche LG, Zhou W, Skov MW, Graham SE, et al.
Biobank-driven genomic discovery yields new insight into atrial fibrillation biology. Nat
Genet. 2018 Sep;50(9):1234–9.

32. Mahajan A, Taliun D, Thurner M, Robertson NR, Torres JM, Rayner NW, et al. Fine-
mapping type 2 diabetes loci to single-variant resolution using high-density imputation
and islet-specific epigenome maps. Nat Genet. 2018 Nov;50(11):1505–13.

33. Wuttke M, Li Y, Li M, Sieber KB, Feitosa MF, Gorski M, et al. A catalog of genetic loci
associated with kidney function from analyses of a million individuals. Nat Genet. 2019
Jun;51(6):957–72.

34. Surakka I, Horikoshi M, Mägi R, Sarin AP, Mahajan A, Lagou V, et al. The impact of
low-frequency and rare variants on lipid levels. Nat Genet. 2015 Jun;47(6):589–97.

35. Shah S, Henry A, Roselli C, Lin H, Sveinbjörnsson G, Fatemifar G, et al. Genome-wide
association and Mendelian randomisation analysis provide insights into the pathogenesis
of heart failure. Nat Commun. 2020 Jan 9;11(1):163.

36. Malik R, Chauhan G, Traylor M, Sargurupremraj M, Okada Y, Mishra A, et al.
Multiancestry genome-wide association study of 520,000 subjects identifies 32 loci
associated with stroke and stroke subtypes. Nat Genet. 2018 Apr;50(4):524–37.

37. Jiang L, Zheng Z, Fang H, Yang J. A generalized linear mixed model association tool for
biobank-scale data. Nat Genet. 2021 Nov;53(11):1616–21.

38. van der Harst P, Verweij N. Identification of 64 Novel Genetic Loci Provides an
Expanded View on the Genetic Architecture of Coronary Artery Disease. Circ Res. 2018
Feb 2;122(3):433–43.

39. Hoffmann TJ, Ehret GB, Nandakumar P, Ranatunga D, Schaefer C, Kwok PY, et al.
Genome-wide association analyses using electronic health records identify new loci
influencing blood pressure variation. Nat Genet. 2017 Jan;49(1):54–64.

40. Anstee QM, Darlay R, Cockell S, Meroni M, Govaere O, Tiniakos D, et al. Genome-wide
association study of non-alcoholic fatty liver and steatohepatitis in a histologically
characterised cohort☆. J Hepatol. 2020 Sep 1;73(3):505–15.

41. Privé F, Arbel J, Vilhjálmsson BJ. LDpred2: better, faster, stronger. Bioinformatics. 2020
Dec 1;36(22–23):5424–31.

42. R Core Team. R: A Language and Environment for Statistical Computing [Internet].
Vienna, Austria: R Foundation for Statistical Computing; 2020. Available from:
https://www.R-project.org/

43. Mölder F, Jablonski KP, Letcher B, Hall MB, Tomkins-Tinch CH, Sochat V, et al.
Sustainable data analysis with Snakemake [Internet]. F1000Research; 2021 [cited 2023
Mar 16]. Available from: https://f1000research.com/articles/10-33

44. Violán C, Roso-Llorach A, Foguet-Boreu Q, Guisado-Clavero M, Pons-Vigués M, Pujol-
Ribera E, et al. Multimorbidity patterns with K-means nonhierarchical cluster analysis.
BMC Fam Pract. 2018 Jul 3;19(1):108.

Supporting information captions

Supporting information captions

S1 Fig: Classification of new ischemic events.

S1 Table: Eligible codes for inclusion and outcomes

S1 Appendix: MCL algorithm settings and assessment of cluster robustness

• **S2 Fig: Selection of number of components.**

• **S3 Fig: Limiting edge-density and average node degree in sex-specific similarity**

networks.

S2 Appendix: Preprocessing of laboratory data

• **S2 Table: Laboratory codes included in assessment of data quality and completeness**

S3 Appendix: Calculation of polygenetic risk scores for 14 traits

S3 Table: Comparison of mean age at index in 31 cluster using Tukey's HSD

S4 Table: Demographics for patients not cluster or were in clusters of size < 500

S5 Table: Cluster-wise summarized O/E-ratios, 10 largest O/E-ratios and 10 lowest

O/E-ratios.

S6 Table: Chi-squared test for distribution laboratory values in clusters

S7 Table: Traits with significantly different PGS distributions in clusters