Delineating disorder-general and disorder-specific dimensions of psychopathology from

functional brain networks in a developmental clinical sample

Authors:

Irene Voldsbekk[1,2], Rikka Kjelkenes[1,2], Andreas Dahl[1,2], Madelene C. Holm[1,2], Martina J.

Lund[1], Tobias Kaufmann[1,3], Christian K. Tamnes[1,4,5], Ole A. Andreassen[1,6], Lars T.

Westlye[1,2,6], Dag Alnæs[1,7]

[1] Norwegian Centre for Mental Disorders Research (NORMENT), Institute of Clinical

Medicine, University of Oslo, & Division of Mental Health and Addiction, Oslo University

Hospital, Oslo, Norway

[2] Department of Psychology, University of Oslo, Oslo, Norway

[3] Department of Psychiatry and Psychotherapy, University of Tübingen, Germany

[4] Department of Psychiatric Research, Diakonhjemmet Hospital, Oslo, Norway

[5] PROMENTA Research Center, Department of Psychology, University of Oslo, Oslo,

Norway

[6] KG Jebsen Centre for Neurodevelopmental Disorders, University of Oslo, & Department of

Neurology, Oslo University Hospital, Oslo, Norway

[7] Kristiania University College, Oslo, Norway

Corresponding author:

Full name: Irene Voldsbekk

Address: NORMENT, Oslo universitetssykehus HF

Klinikk psykisk helse og avhengighet

Seksjon for psykoseforskning/TOP

Ullevål sykehus, bygg 49

Postboks 4956 Nydalen

0424 Oslo

Phone: +47 41495202

Email: irene.voldsbekk@psykologi.uio.no

Keywords: Brain; Multivariate; Functional connectivity; Psychopathology; Developmental

Abstract

Background: The interplay between functional brain network maturation and psychopathology during development remains elusive. In pursuit of the structure of psychopathology and its underlying neurobiological mechanisms, mapping of both shared and unique functional connectivity patterns across developmental clinical populations is needed.

Methods: We investigated shared associations between resting-state functional connectivity patterns and psychopathology in children and adolescents aged 5-21 (n=1880, 62% male). Specifically, we used partial least squares (PLS) to identify latent variables (LV) between functional connectivity and both symptom scores and diagnostic information. We also investigated associations between functional connectivity and each diagnosis specifically, controlling for other diagnosis categories.

Results: PLS identified several statistically significant LVs between functional brain networks and symptoms, mapping onto the psychopathology hierarchy. The first LV resembled a general psychopathology factor, followed by dimensions of externalising-internalising, neurodevelopment, somatic complaints, and thought problems. Another PLS with diagnostic data revealed one significant LV, resembling a cross-diagnostic case-control pattern. The diagnosis-specific PLS identified a unique connectivity pattern for autism spectrum disorder (ASD). All LVs were associated with distinct patterns of functional connectivity.

Conclusions: Covariance in developmental functional brain networks supports transdiagnostic and symptom-based dimensions of psychopathology.

1 Introduction

Establishing the structure of psychopathology and its underlying neurobiological mechanisms is a critical step towards personalised approaches in mental health research and care. The high rate of comorbidity between diagnoses challenges the utility of traditional case-control designs and motivates novel strategies for clinical phenotyping such as transdiagnostic assessment of psychopathology dimensions (Caspi et al., 2014; Insel et al., 2010). Existing diagnostic categories do not map onto disease-specific neurobiological substrates (Insel & Cuthbert, 2015), and many of the detected abnormalities in both genetics (Hindley et al., 2022; Lahey et al., 2011; Pettersson et al., 2016; Roelfs et al., 2021), brain structure (Goodkind et al., 2015; Opel et al., 2020a) and brain function (Elliott et al., 2018; McTeague et al., 2017; McTeague et al., 2020; Sha et al., 2019) are shared across disorders. Investigation into the neurobiological substrates of distinct symptom dimensions may therefore elucidate the brain-based underpinnings of mental disorders.

Childhood and adolescence is characterised by large scale reorganisation and maturation of the brain and its functional networks (Paus et al., 2008; Power et al., 2010). Given that mental disorders often first manifest during this time (Caspi et al., 2020; Kessler et al., 2007), aberrant functional network development may represent a key aetiological component in mental disorders (Casey et al., 2014; Paus et al., 2008). Indeed, while sensory and motor regions and their associated functional networks typically are fully developed by late childhood, the association cortex, and implicated functional networks such as the default mode network (DMN), take longer to mature. This might leave these brain regions vulnerable to emerging psychopathology during neurodevelopment (Sydnor et al., 2021). To identify biologically informed dimensions of psychopathology, investigating associations between functional brain networks and psychopathology during childhood and adolescence is imperative.

Psychopathology is increasingly conceptualised as a hierarchical structure (Caspi et

al., 2014; Kotov et al., 2017; Lahey et al., 2017). This hierarchy consists of a general

psychopathology factor as the highest order, reflecting a general vulnerability to

psychopathology, followed by increasingly narrow dimensions, such as internalising and

externalising. These reflect anxious and depressive symptoms, and aggressive, rule-breaking,

and hyperactive symptoms, respectively. A neurodevelopmental factor is also often included

to reflect autistic-like traits and symptoms of attention deficit hyperactivity disorder

(ADHD). For example, recent work in the Adolescent Brain Cognitive Development (ABCD)

cohort (Casey et al., 2018) derived five dimensions of psychopathology (i.e. internalising,

externalising, neurodevelopmental, detachment, and somatoform) using exploratory factor

analysis on symptom data (Michelini et al., 2019). Similar psychopathology dimensions have

been derived from both symptom data (Karcher et al., 2021) and diagnostic data (Lees et al.,

2021) and then associated with patterns of functional connectivity obtained from resting-state

functional magnetic resonance imaging (rs-fMRI). However, these studies derived

dimensions of psychopathology from symptom data or diagnostic information in isolation,

and only afterwards associated them with functional connectivity. To identify brain-based

dimensions of psychopathology, the functional brain networks should inform the estimation

of psychopathology dimensions per se.

Doing exactly this, studies have identified symptom dimensions by finding their

maximal correlation with functional connectivity in youth aged 8-22 (Xia et al., 2018) and

preadolescents aged 9-11 (i.e. the ABCD cohort) (Kebets et al., 2023). In youth, dimensions

of mood, psychosis, fear, and externalisation symptoms exhibited both unique and a shared

pattern of connectivity. In ABCD, dimensions derived from structural and functional brain

patterns simultaneously resembled a general psychopathology factor along with internalising-

externalising, neurodevelopmental, somatoform, and detachment dimensions. Although this

work is promising with respect to identifying biologically informed dimensions of

psychopathology, the investigation of mental health symptoms in population-based studies

may not generalise to clinical populations (Vanes & Dolan, 2021). While investigation in

young, representative cohorts is essential to understand putative developmental mechanisms

relevant for psychopathological vulnerability, it is equally important to map the relevance of

these findings to individuals already diagnosed with a mental disorder. Mapping of disorder-

general and disorder-specific patterns in clinical populations is needed to elucidate the

underlying neurobiological mechanisms of psychopathology.

Patterns of connectivity related to symptoms of anxiety, irritability, and ADHD were

replicated across two independent clinical samples of children and adolescents (Linke et al.,

2021). Specifically, this study identified one dimension consisting of all three domains, while

the second dimension captured shared aspects of irritability and ADHD, and the third was

specific to anxiety. This indicates clinically relevant disorder-general (i.e. shared across

disorders) and disorder-specific effects in functional networks of children and adolescents.

Moreover, it points to the possibility of decomposing irritability, a symptom shared between

anxiety and ADHD, into disorder-specific and common components based on patterns of

brain connectivity. However, this study did not investigate connectivity patterns related to

broad transdiagnostic symptom dimensions but maintained a focus limited to anxiety,

irritability, and attention problems. Moreover, the degree of overlap between functional

networks linked to symptom dimensions and those related to diagnosis remain to be

determined.

In the current study, we aimed to investigate dimensions linking functional

connectivity and psychopathology in a clinical sample of children and adolescents. We used

partial least squares (PLS) (Krishnan et al., 2011), a multivariate technique that identifies

shared associations across two high-dimensional matrices. This enables identification of

dimensions of psychopathology derived from connectivity patterns in functional brain

networks. Specifically, we wanted to highlight similarities and differences across dimensions

derived from symptom data vs diagnostic classifications in the same sample. To do this, we

investigated associations between functional connectivity and a) symptom scores, and b)

diagnostic information. In addition, we investigated associations between functional

connectivity and each diagnosis specifically, controlling for other diagnosis categories.

2 Material and methods

2.1 Sample

The sample was recruited from New York City, USA to participate in the Healthy Brain

Network (HBN) (Alexander et al., 2017), a cohort consisting of children and adolescents

aged 5-21. Participants were recruited through community sampling where children with

clinical concerns were encouraged to participate. The participants underwent a

comprehensive assessment of biological and socio-environmental factors, in addition to

diagnostic evaluation by qualified health personnel. After quality control and data cleaning

(see below), the final sample for our analyses consisted of 1880 participants (721 females).

Sample demographics are provided in Figure 1.

2.2 Clinical measures

Symptom scores were obtained from the Child Behaviour Checklist (CBCL) (Achenbach &

Rescorla, 2001), which assesses emotional, behavioural, and social problems in children by

parent report. Parents scored their children on 113 items as either 0 ("Not true"), 1

("Somewhat or sometimes true") or 2 ("Very true or often true"). The responses to these

items result in eight syndrome measures, previously found to be the best-fitting model for

data obtained from both general and clinical populations (Achenbach & Rescorla, 2001;

Ivanova et al., 2007): Anxious/Depressed, Withdrawn/Depressed, Somatic Complaints,

Social Problems, Thought Problems, Attention Problems, Rule-Breaking Behaviour, and

Aggressive Behaviour.

Diagnostic information was obtained by a computerised version of the Schedule for

Affective Disorders and Schizophrenia – Children´s version (KSADS) (Kaufman et al.,

1997), which is a clinician-administered semi-structured psychiatric interview based on

DSM-5. Based on this interview and revision of all other collected materials, a consensus

regarding clinical diagnosis was made by a team of licensed clinicians. We then categorised

diagnoses as either "ADHD", "ASD", "anxiety disorders", "mood disorders", "other

neurodevelopmental disorders", "other disorders" or "no diagnosis". Most participants had

more than one diagnosis.

Figure 1. Sample distributions. **A.** Diagnosis categories (more than one per individual possible). **B.**

Clinical symptoms. **C.** Comorbidity. **D.** Age and sex. The lines indicate mean age for each sex.

CBCL; child behaviour checklist. ADHD; attention-deficit hyperactivity disorder. ASD; autism

spectrum disorder. ND; neurodevelopmental.

Of those with complete MRI data (see below), 1992 participants had available both

diagnostic data and symptom data. Participants with more than 10% missing symptom data

were excluded (n=112). For the remaining participants (n=1880), missing values were

imputed with knnimpute in MATLAB (MathWorks, 2020).

2.3 MRI pre-processing

We accessed rs-fMRI and T_1-weighted structural MRI for the current study. MRI data were

acquired at four different sites: a mobile scanner at Staten Island (SI), Rutgers University

Brain Imaging Centre, Citigroup Biomedical Imaging Centre (CBIC) and Harlem CUNY

Advanced Science Research Centre. A detailed overview of the MRI protocol is available

elsewhere

(http://fcon_1000.projects.nitrc.org/indi/cmi_healthy_brain_network/MRI%20Protocol.html).

T1-weighted MRI data (n=3334) were processed using FreeSurfer v. 7.1.0 (Fischl, 2012) and

quality controlled using the MRIQC classifier (Esteban et al., 2017). For participants with

more than one T_1-weighted scan, we selected the sequence with the best estimated quality, as

previously described (Voldsbekk et al., 2023).

For individuals with sufficient structural MRI image quality (n=3213), we submitted

rs-fMRI images for pre-processing along the following pipeline. We applied FSL MCFLIRT

(Jenkinson et al., 2002) for motion correction, high-pass temporal filtering (cut-off: 100),

spatial smoothing (FWHM: 6) and distortion correction as part of FEAT (Woolrich et al.,

2001). The rs-fMRI images were also registered to the structural image using FLIRT

(Jenkinson et al., 2002) and boundary-based registration (Greve & Fischl, 2009). Next, for

additional removal of artefacts and noise, we performed non-aggressive ICA-AROMA

(Pruim, Mennes, Buitelaar, et al., 2015; Pruim, Mennes, van Rooij, et al., 2015) and FIX

(Griffanti et al., 2014; Salimi-Khorshidi et al., 2014) with a threshold of 20. During this

procedure, 595 participants were excluded due to missing data or insufficient image quality.

As an additional step, quality control of the raw rs-fMRI images was performed using

MRIQC. Estimations of temporal signal-to-noise ratio (tSNR) and mean framewise

displacement (FD), as calculated by MRIQC, were used as covariates in subsequent analyses.

2.4 Network analysis

To increase reproducibility, nodes were estimated from the Schaefer parcellation with 100

parcels and 7 networks (Schaefer et al., 2018). These networks include visual A, visual B,

visual C, auditory, somatomotor A, somatomotor B, language, salience A, salience B, control

A, control B, control C, default A, default B, default C, dorsal attention A and dorsal

attention B. As an additional quality check of the estimated parcels, participants with data in

less than 60% of voxels for each parcel were excluded (n=290). An overview of percentage

missing data in each parcel is shown in Figure S1. To check that 60% is a reasonable

threshold, balancing exclusion of participants vs completeness of voxel data, time series

correlations were computed in the subset of participants with no missing data between the

full parcel time series (i.e., from 100% of voxels) and parcel time series based on 60% of the

voxels of that parcel (removing those voxels most frequently missing). The correlation

between the full 100% parcel and the 60% parcel time series was high for every parcel (all

higher than .87, see Figure S2). Parcel timeseries were then imported to FSLNets

(https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/FSLNets), as implemented in MATLAB (MathWorks,

2020), for estimation of edges (n=2328). In this step, we calculated the partial correlations

between nodes using L2-norm ridge regression, as these are considered a better measure of

direct connectivity strength (Marrelec et al., 2006). Finally, edges were z-transformed using

Fisher's transformation and we extracted the upper triangle of the correlation matrix for

further analysis, yielding 4,950 unique edges reflecting the connection strength between

nodes for each participant.

2.5 Partial-least squares

To assess shared associations between edges (i.e. functional connectivity strength between

two brain regions) and symptom data, we used rotated behavioural PLS in PLS Application

(Krishnan et al., 2011), entering z-transformed symptom data as behavioural variables. This

approach yields latent variables (LV) reflecting maximal covariance across both matrices.

See figure 2 for an overview of the PLS analysis pipeline. Briefly, the PLS analysis estimates

a cross-covariance matrix between imaging and behavioural data. This matrix is then inputted

to singular value decomposition, yielding a total number of LVs corresponding to the number

of behavioural variables. For each of these LVs, we get a singular value and the weights of

each imaging and behavioural variable, as well as for each subject. The significance of LVs

was assessed using permutation testing (n=5000). Then, the stability of edges for each

significant LV was estimated using bootstrapping with replacement (n=1000), thresholding at

$|pseudo-z|>3|$ (McIntosh & Lobaugh, 2004) for significance. Symptom loadings onto each LV

was extracted as the correlation between weights on each LV and the original symptom data.

Next, entering diagnostic information (one column for each diagnosis: 1 as having the

diagnosis, 0 as not – more than one possible per participant) as behavioural variables, we then

used rotated behavioural PLS to decompose data into putative specific and shared disorder

dimensions (i.e. diagnosis-based patterns). Then, to test for diagnosis-specific patterns

explicitly, we ran non-rotated behavioural PLS, in which associations between edges and

each diagnostic category was tested. This test was run for each diagnosis category separately,

Partial Least Squares (PLS)

Figure 2. Illustration of the PLS analysis pipeline. The cross-covariance matrix between imaging data

and behavioural data is estimated. This matrix is then inputted to singular value decomposition,

yielding singular values for each LV, as well as imaging, behavioural, and subject weights. Then,

loadings onto each LV are calculated as the correlation between subject weights and the original

imaging and behavioural data, respectively. PLS; partial least squares. LV; latent variable.

while controlling for all other diagnosis categories (see Table S1 for an overview of

contrasts).

Prior to running PLS, edges were adjusted for sex, age, tSNR, FD, and scanner site.

PLS was run using Spearman´s rank correlation. Significant LVs were then plotted using R

version 4.1.2 (https://cran.r-project.org). To aid in the visualisation and interpretation of the

high dimensional connectivity patterns, edges were summarised across networks for

significant LVs. To investigate the loading of nodes, we also estimated the nodal loading

strength across the connectivity matrix for each connectivity pattern identified by PLS using

the Brain Connectivity Toolbox (Rubinov & Sporns, 2010) in MATLAB. To obtain a more

detailed overview of each connectivity pattern, we plotted nodal strength and edge strength

using BrainNet Viewer (Xia et al., 2013).

2.6 Consistency across age, sex, ethnicity, socio-economic status, and intelligence

Given that aberrant brain development may represent a key aetiological component in mental

disorders (Casey et al., 2014; Paus et al., 2008), we assessed whether the shared associations

between edges and symptom data differed as a function of age or sex. To do this, we reran

the symptom-based PLS without adjusting edges for age and sex. These results revealed

similar patterns of covariation (Figure S3), with highly correlated feature weights across

overlapping dimensions for all LVs (Table S2). To examine whether the shared associations

identified were generalisable across ethnic groups, we plotted the correlations between edges

and symptoms by ethnic group (Figure S4). Similarly, to examine whether the shared

associations identified were generalisable across levels of socioeconomic status (SES), we

plotted the correlations by median-split of household income, as a proxy for SES (Figure S5).

To examine whether the shared associations identified were generalisable across levels of

intelligence, we plotted the correlations by full scale IQ split into ± 70 (Figure S6). Finally,

symptom loadings and connectivity loadings from the symptom-based PLS were regressed

against each diagnosis category separately, with "no diagnosis" as a reference group. We also

regressed loadings against number of diagnoses, interpreting the latter as a proxy of cross-

diagnostic vulnerability. All associations were adjusted for age, age^2, and sex.

3 Results

3.1 Symptom-based dimensions

Based on the scree plot of percent cross-block covariance explained (Figure S7), we selected

the first six LVs in the symptom-based PLS for further analysis. Of these, five were

significant (r=.69, p=.01; r=.69, p=.038; r=.70, p=.012; r=.69, p=.031; r=68, p=.012,

respectively; Figure S8). Each LV represents a distinct pattern that relates a weighted set of

symptoms to a weighted set of functional brain network connections. Inspection of the most

heavily weighted symptoms for each LV revealed that they resemble the psychopathology

hierarchy: the first LV resembled a general psychopathology factor (see Figure 3A), while

the remaining four represented increasingly narrow dimensions (see Figure 4). Specifically,

LV2 was related to externalising-internalising, LV3 to neurodevelopment, LV4 to somatic

complaints, and LV6 to thought problems.

These psychopathology dimensions were identified by their shared associations with

specific patterns of connectivity. For each dimension, these patterns were widely distributed

across functional networks (see Figure 3 and S9). LV1 was related to weaker connectivity

Figure 3. The first dimension of shared associations between functional connectivity and clinical

symptoms resembled a general psychopathology factor. **A.** The highest loading symptoms of this

dimension. Loadings reflect correlations between LV weights and original data. **B.** Strength of edges

and nodes that contributed to this dimension. Edges are coloured red for higher connectivity and blue

for lower connectivity. Nodes are coloured based on network membership. **C.** Both increased and

reduced connectivity in specific edges contributed. Magnitude in this plot reflects summarised edge

strength across each network. **E-F.** Connectivity and symptom loadings across diagnostic categories

(left) and number of comorbidities (right). In these plots, the data is centred around the mean of no

diagnosis. LV; latent variable. ADHD; attention-deficit hyperactivity disorder. ASD; autism spectrum

disorder. ND; neurodevelopmental. Vis; visual network. SomMot; somatomotor network.

SalVentAttn; salience network. Cont; control network. Default; default mode network. DorsAttn;

dorsal attention network.

Figure 4. Dimensions of shared associations between functional connectivity and clinical symptoms

map onto the hierarchical structure of psychopathology. **A.** LV2 map onto symptoms of higher

externalisation and lower internalisation. **B.** LV3 map onto symptoms of neurodevelopmental

problems. **C.** LV4 map onto symptoms of higher somatic complaints and lower withdrawn/depressive

symptoms. **D.** LV6 map onto symptoms of higher thought problems. LV; latent variable.

between the salience network and limbic and visual network (Figure 3B-C). In addition, LV1

was related to stronger connectivity between the limbic and visual network and between the

salience network and DMN. The nodes with strongest loading on this pattern implicated the

control network, DMN, and salience network. LV2 was related to stronger within- and

between-network connectivity in the visual network and between the salience network and

DMN, with the strongest nodes in the DMN and visual network (Figure S9). LV3 was related

to stronger connectivity between the DA and limbic network and between the DMN and

visual network. The strongest nodes were distributed across the DMN, DA, control, and

visual network. LV4 was related to weaker connectivity between the DA and control

network, stronger connectivity between the visual and limbic network, and within the control

network. The strongest nodes were in the control network. LV6 was related to weaker

connectivity between the limbic and DA and control network, as well as within the DMN.

The strongest nodes were in the somatomotor network and the limbic network.

Figure 5. One dimension of shared associations between functional connectivity and diagnosis

categories, resembling a cross-diagnostic case-control difference. **A.** The diagnosis dimension

reflected a pattern across all diagnostic categories vs no diagnosis. **B.** Strength of edges and nodes that

contributed to this dimension. Edges are coloured red for higher connectivity and blue for lower

connectivity. Nodes are coloured based on network membership. **C.** Both increased and reduced

connectivity in specific edges contributed. Magnitude in this plot reflects summarised edge strength

across each network. ADHD; attention-deficit hyperactivity disorder. ND; neurodevelopmental. ASD;

autism spectrum disorder. Vis; visual network. SomMot; somatomotor network. SalVentAttn;

salience network. Cont; control network. Default; default mode network. DorsAttn; dorsal attention

network.

3.2 Diagnosis-based dimensions

The rotated diagnosis-based PLS identified one significant LV (r=.69, p=.009). As shown in

Figure 5A, this pattern resembled a general case vs control pattern across all diagnoses.

Partly consistent with the general psychopathology pattern of the symptom-based LV1, a

higher loading on this LV (i.e. having a diagnosis) entailed stronger connectivity between the

limbic and visual network, in addition to weaker connectivity within the somatomotor

network and between this network and the salience network and DMN (Figure 5B-C). The

strongest nodes were distributed across all these networks, as well as the DA network.

Figure 6. ASD was the only diagnosis category exhibiting a unique pattern of connectivity **A.**

Strength of edges and nodes specific to ASD. Edges are coloured red for higher connectivity and blue

for lower connectivity. Nodes are coloured based on network membership. **B.** Both increased and

reduced connectivity in specific edges contributed. Magnitude in this plot reflects summarised edge

strength across each network. ASD; autism spectrum disorder. Vis; visual network. SomMot;

somatomotor network. SalVentAttn; salience network. Cont; control network. Default; default mode

network. DorsAttn; dorsal attention network.

3.3 Diagnosis-specific patterns

The non-rotated diagnosis-specific PLS, which tested each diagnosis category separately

while controlling for all other diagnosis categories, identified a unique connectivity pattern

for ASD (r=.67, p=.012). As shown in Figure 6, the ASD-specific pattern was widely

distributed, including weaker connectivity within the somatomotor network and between the

DA network and visual network. The strongest nodes implicated the salience, visual, and

control network. In addition, the non-rotated diagnosis-specific PLS identified a unique

pattern of no diagnosis vs all diagnoses (Figure S10). This pattern implicated increased

connectivity between the somatomotor, and salience and DMN, in addition to lower

connectivity between the visual and limbic network, as well as between the DMN and DA

network. The strongest nodes implicated in this pattern were in the DA network. The

remaining diagnosis categories did not show disorder-specific patterns of functional

connectivity.

3.4 Symptom-based pattern evident across diagnostic boundaries

To understand the distribution of symptom dimensions in more detail, we plotted them

against diagnosis categories (Figure S11). As shown in Figure 3E-F, there was a consistency

between a higher degree of comorbidity and higher loading on LV1. We also observed some

expected variation in symptom loading with respect to specific diagnoses, such as patients

with ADHD and ASD loading more highly on the neurodevelopmental dimension (LV3;

Figure S11). On the externalising-internalising dimension (LV2), mood disorder, ASD, and

anxiety disorder loaded more negatively, consistent with increasing symptoms of

internalising being typical for these diagnosis categories. In addition, linear models revealed

that all diagnosis categories showed higher symptom and connectivity loadings on LV1

compared to having no diagnosis (Table S3). There was also a significant linear association

with the number of diagnoses for both symptom and connectivity loadings on LV1 (Table S4

and S5). This was true when including "no diagnosis" in the model or not, suggesting that

this effect was not driven by case-control effects.

4 Discussion

Through shared associations between mental health data and functional connectivity, the current study delineated shared and unique patterns in child and adolescent functional brain networks. We found that dimensions of clinical symptoms map onto specific patterns of brain connectivity, implicating both a general psychopathology factor and dimensions overlapping with subscale syndromes of CBCL. The rotated decomposition of diagnostic data (i.e. the diagnosis-based PLS) revealed one significant dimension, implicating a cross-diagnostic pattern only. The disorder-specific tests revealed specific patterns of connectivity related to ASD and no diagnosis (i.e., a disorder-general effect), but not for any other diagnosis. For the symptom-based dimensions, we found that increased comorbidity was consistently related to both increased symptom burden and increased connectivity aberrations. Taken together, these results indicate that compared to diagnostic classifications in isolation, transdiagnostic and symptom-based dimensions of psychopathology are more closely mapped to the functional networks of the brain during the formative years of childhood and adolescence.

The clinical dimensions revealed by shared associations between functional connectivity and symptoms in the current study adhere to the hierarchical structure of psychopathology, implicating a general psychopathology factor, followed by dimensions of externalising-internalising, neurodevelopment, somatic complaints, and thought problems. PLS derives orthogonal LVs, leaving dimensions independent. Capturing externalisation-internalisation as the second latent pattern is consistent with previous work (Kebets et al., 2023; Linke et al., 2021). Next, we identified a neurodevelopmental dimension, which was also captured in Kebets and colleagues (Kebets et al., 2023). In fact, the dimensions this study detected in the ABCD sample were the same as the symptom-based dimensions we captured here in a clinical sample. This degree of overlap across samples is striking, suggesting generalisable patterns of functional connectivity-psychopathology associations,

and strongly supporting the conceptualisation of general population vs clinical population as existing on a continuum. Interestingly, given the overlap between the current functional connectivity-derived symptom dimensions and CBCL subscale syndromes derived from symptom data alone, it also appears that the psychopathology hierarchy is represented in functional networks during development.

In contrast to our findings, Linke and colleagues (Linke et al., 2021) identified a dimension specific to anxiety symptoms, which did not emerge in the current study. Neither did we detect a pattern specific to having an anxiety diagnosis. This discrepancy may reflect diversity in the range of symptoms and diagnostic groups included. Indeed, in the current study, the diagnostic range was broader than in the study by Linke and colleagues (Linke et al., 2021), while the symptom domains assessed by Xia and colleagues (Xia et al., 2018) were more closely mapped to adult psychopathology than to child and adolescent symptomology, which may explain their differences.

Alterations in functional connectivity of the DMN have previously been implicated in several neurological and mental disorders (van den Heuvel & Sporns, 2019). In addition, DMN connectivity has been linked to general psychopathology (Elliott et al., 2018; Karcher et al., 2021; Kebets et al., 2023; Sato et al., 2018). In line with this, DMN nodes were some of the strongest loading nodes of the general psychopathology factor (i.e. symptom-based LV1) in the current study. This factor was further characterised by a distributed pattern involving weaker connectivity between the salience network and limbic and visual network. It also implicated stronger connectivity between DMN and salience network, and between the control network and somatomotor network. In line with this, the limbic, salience, fronto-parietal and sensorimotor networks are also implicated in general psychopathology (Vanes & Dolan, 2021).

The connectivity pattern related to a general psychopathology factor in the ABCD

sample (Kebets et al., 2023) was only weakly consistent with current findings, implicating

increased connectivity between the DMN and salience network. Partly consistent with a

general psychopathology dimension derived from diagnostic data (Lees et al., 2021) and

symptom scores (Karcher et al., 2021) also in ABCD, the current study identified

hypoconnectivity within the DA network. Importantly, the methodology used to derive a

general psychopathology factor in these studies differed from the approach used here, which

may explain the observed variation in detected connectivity patterns across studies. This may

also explain the weak overlap between other symptom-based psychopathology dimensions

between current and previous work (Karcher et al., 2021; Kebets et al., 2023). In addition,

previous work has shown that while connectivity-informed clinical dimensions were

replicable across two samples, the connectivity patterns themselves were less so (Linke et al.,

2021). The authors attributed this to a "many-to-one" mapping between neural and clinical

variables, which may also explain the lack of overlap in specific connectivity patterns

identified in the current work compared to previous work.

The diagnosis-based analysis revealed only one significant dimension, resembling a

cross-diagnostic case-control pattern. This pattern was characterised by no diagnosis

exhibiting the highest loading, with all the other diagnosis categories exhibiting smaller

associations in the opposite direction. If, instead, connectivity patterns specific to each

diagnosis were detectable, we would have expected this test to reveal several dimensions

(i.e., LVs), each consisting of loadings from one (or a few) diagnoses. In addition, we

identified a no diagnosis vs all diagnoses specific pattern, representing an inverse cross-

diagnostic case-control pattern. Although the weighting of each diagnosis category differed

between these two analyses, the overarching connectivity patterns for these dimensions

revealed inverse overlap. Indeed, although the symptom-based general psychopathology

dimension exhibited a more distributed connectivity pattern, there was also some overlap

between this pattern and cross-diagnostic case-control pattern. Together, these patterns

implicate connectivity between the visual and limbic networks in separating no diagnosis

from having a diagnosis.

ASD was the only diagnosis category exhibiting a detectable unique pattern of

connectivity. This pattern was widely distributed, implicating altered connectivity within

several brain networks. Hyperconnectivity within several large-scale brain networks has

previously been implicated in ASD (Uddin et al., 2019). The distributed nature of the ASD-

specific pattern identified in the current study, including both increased and reduced

connectivity within and between several networks, is in line with the notion that both

hyperconnectivity and hypoconnectivity may underlie ASD (Kana et al., 2011). Importantly,

the finding that ASD was the only diagnosis group exhibiting a unique connectivity pattern

has implications for our understanding of the neurobiological substrates of ASD, but also for

our understanding of psychopathology and ASD in this landscape more broadly. The current

work supports the understanding that rather than belonging in the general psychopathology

domain, ASD likely represent a separate neurodevelopmental dimension (Opel et al., 2020b;

Ronald, 2019).

Although our findings provide several new insights into the link between functional

brain connectivity and the structure of childhood psychopathology, some limitations should

be noted. First, functional connectivity results are known to be influenced by methodological

choices (Sala-Llonch et al., 2019; Shirer et al., 2015), complicating the identification of

robust and replicable results. To increase replicability of the current work, we relied on an

established parcellation scheme (Schaefer et al., 2018). Second, several functional

connectivity patterns identified implicated the limbic network, a network known to be

sensitive to susceptibility artefacts and reduced signal (Khatamian et al., 2016). Although we

did additional measures to reduce the influence of reduced signal on our analysis, we cannot

completely rule out that our results are influenced by this confound. Third, as the sample

consisted of mainly children, and most of them with at least one mental disorder, motion was

an issue. To ameliorate this influence, we used the MRIQC classifier to exclude participants

with insufficient image quality, cleaned data using FIX and AROMA, and regressed out

measures of image quality and motion from the data. Fourth, the cross-sectional nature of the

study design prohibits any conclusion to be drawn with respect to the within-person temporal

dynamics of any identified pattern. Fifth, given that the symptom data was continuous, and

the diagnostic analysis separated the sample into groups, the power to extract maximal

covariance between symptoms and functional connectivity was better than that for the

diagnostic analysis. As such, these results cannot be directly compared. Finally, the sample

was enriched with children diagnosed with ADHD and other neurodevelopmental conditions.

Although this may be representative of a developmental clinical sample, it may not generalise

to other clinical populations. For example, this has implications for the comparison of the

current results to other studies investigating a derived general psychopathology factor.

However, given that we do see a great degree of overlap between the current results and other

studies, our findings may translate to other populations.

Conclusions

The current work found that dimensions of psychopathology derived from clinical symptoms

were associated with specific patterns of functional connectivity in the developing brain,

while ASD was the only diagnostic category to exhibit such a specific pattern. This

contributes to a growing body of evidence in favour of dimensional and transdiagnostic

classifications of psychopathology (Vanes & Dolan, 2021). In this classification,

neurodevelopmental conditions such as ASD may possess specific abnormalities in

functional connectivity networks above and beyond those related to general psychopathology.

This has implications for the pursuit of individualised brain-based surrogate markers in

mental health research and care, which in turn may lead to improved prevention and

intervention of mental disorders.

Acknowledgements

This project was funded by research grants from the Research Council of Norway (Grant

Nos. L.T.W: 249795, T.K: 276082, 323961. C.K.T: 288083, 323951), the South-Eastern

Norway Regional Health Authority (Grant Nos. L.T.W: 2014097, 2015073, 2016083,

2018076, 2019101. C.K.T: 2019069, 2021070, 2023012, 500189. D.A: 2019107, 2020086.

O.A.A: 223273), the Norwegian ExtraFoundation for Health and Rehabilitation (L.T.W:

Grant No. 2015/FO5146), KG Jebsen Stiftelsen, ERA-Net Cofund through the ERA PerMed

project IMPLEMENT, and the European Research Council under the European Union s

Horizon 2020 research and Innovation program (L.T.W: ERC StG Grant No. 802998. O.A.A:

H2020 RIA grant # 847776).

The work was performed on the Service for Sensitive Data (TSD) platform, owned by

the University of Oslo, operated, and developed by the TSD service group at the University

of Oslo IT-Department (USIT). Computations were also performed using resources provided

by UNINETT Sigma2—the National Infrastructure for High Performance Computing and

Data Storage in Norway.

Declaration of Interest

O.A.A. is a consultant to cortechs.ai and has received speaker's honorarium from Janssen,

Lundbeck, Sunovion.

References

Achenbach, T. M., & Rescorla, L. A. (2001). Manual for the ASEBA school-age forms &
   profiles: an integrated system of multi-informant assessment Burlington, VT:
   University of Vermont. *Research Center for Children, Youth, & Families*, *1617*.

Alexander, L. M., Escalera, J., Ai, L., Andreotti, C., Febre, K., Mangone, A., Vega-Potler,
   N., Langer, N., Alexander, A., & Kovacs, M. (2017). An open resource for
   transdiagnostic research in pediatric mental health and learning disorders. *Scientific
   data*, *4*(1), 1-26.

Casey, B. J., Cannonier, T., Conley, M. I., Cohen, A. O., Barch, D. M., Heitzeg, M. M.,
   Soules, M. E., Teslovich, T., Dellarco, D. V., & Garavan, H. (2018). The adolescent
   brain cognitive development (ABCD) study: imaging acquisition across 21 sites.
   Developmental Cognitive Neuroscience, *32*, 43-54.

Casey, B. J., Oliveri, M. E., & Insel, T. (2014). A neurodevelopmental perspective on the
   research domain criteria (RDoC) framework. *Biol Psychiatry*, *76*(5), 350-353.
   https://doi.org/10.1016/j.biopsych.2014.01.006

Caspi, A., Houts, R. M., Ambler, A., Danese, A., Elliott, M. L., Hariri, A., Harrington, H.,
   Hogan, S., Poulton, R., Ramrakha, S., Rasmussen, L. J. H., Reuben, A., Richmond-
   Rakerd, L., Sugden, K., Wertz, J., Williams, B. S., & Moffitt, T. E. (2020).
   Longitudinal Assessment of Mental Health Disorders and Comorbidities Across 4
   Decades Among Participants in the Dunedin Birth Cohort Study. *JAMA network
   open*, *3*(4), e203221-e203221. https://doi.org/10.1001/jamanetworkopen.2020.3221

Caspi, A., Houts, R. M., Belsky, D. W., Goldman-Mellor, S. J., Harrington, H., Israel, S.,
   Meier, M. H., Ramrakha, S., Shalev, I., Poulton, R., & Moffitt, T. E. (2014). The p
   Factor: One General Psychopathology Factor in the Structure of Psychiatric

Disorders? *Clin Psychol Sci*, *2*(2), 119-137.

https://doi.org/10.1177/2167702613497473

Elliott, M. L., Romer, A., Knodt, A. R., & Hariri, A. R. (2018). A Connectome-wide

Functional Signature of Transdiagnostic Risk for Mental Illness. *Biological*

Psychiatry, *84*(6), 452-459.

https://doi.org/https://doi.org/10.1016/j.biopsych.2018.03.012

Esteban, O., Birman, D., Schaer, M., Koyejo, O. O., Poldrack, R. A., & Gorgolewski, K. J.

(2017). MRIQC: Advancing the automatic prediction of image quality in MRI from

unseen sites. *PLoS ONE*, *12*(9), e0184661.

https://doi.org/10.1371/journal.pone.0184661

Fischl, B. (2012). FreeSurfer. *NeuroImage*, *62*(2), 774-781.

https://doi.org/https://doi.org/10.1016/j.neuroimage.2012.01.021

Goodkind, M., Eickhoff, S. B., Oathes, D. J., Jiang, Y., Chang, A., Jones-Hagata, L. B.,

Ortega, B. N., Zaiko, Y. V., Roach, E. L., Korgaonkar, M. S., Grieve, S. M., Galatzer-

Levy, I., Fox, P. T., & Etkin, A. (2015). Identification of a Common Neurobiological

Substrate for Mental Illness. *JAMA Psychiatry*, *72*(4), 305-315.

https://doi.org/10.1001/jamapsychiatry.2014.2206

Greve, D. N., & Fischl, B. (2009). Accurate and robust brain image alignment using

boundary-based registration. *NeuroImage*, *48*(1), 63-72.

https://doi.org/10.1016/j.neuroimage.2009.06.060

Griffanti, L., Salimi-Khorshidi, G., Beckmann, C. F., Auerbach, E. J., Douaud, G., Sexton, C.

E., Zsoldos, E., Ebmeier, K. P., Filippini, N., Mackay, C. E., Moeller, S., Xu, J.,

Yacoub, E., Baselli, G., Ugurbil, K., Miller, K. L., & Smith, S. M. (2014). ICA-based

artefact removal and accelerated fMRI acquisition for improved resting state network

imaging. *NeuroImage*, *95*, 232-247.

https://doi.org/https://doi.org/10.1016/j.neuroimage.2014.03.034

Hindley, G., Frei, O., Shadrin, A. A., Cheng, W., O'Connell, K. S., Icick, R., Parker, N.,

Bahrami, S., Karadag, N., Roelfs, D., Holen, B., Lin, A., Fan, C. C., Djurovic, S.,

Dale, A. M., Smeland, O. B., & Andreassen, O. A. (2022). Charting the Landscape of

Genetic Overlap Between Mental Disorders and Related Traits Beyond Genetic

Correlation. *American Journal of Psychiatry*, *179*(11), 833-843.

https://doi.org/10.1176/appi.ajp.21101051

[Record #1022 is using a reference type undefined in this output style.]

Insel, T. R., & Cuthbert, B. N. (2015). Brain disorders? precisely. *Science*, *348*(6234), 499-

500.

Ivanova, M. Y., Achenbach, T. M., Dumenci, L., Rescorla, L. A., Almqvist, F., Weintraub,

S., Bilenberg, N., Bird, H., Chen, W. J., Dobrean, A., Döpfner, M., Erol, N.,

Fombonne, E., Fonseca, A. C., Frigerio, A., Grietens, H., Hannesdóttir, H.,

Kanbayashi, Y., Lambert, M., Larsson, B., Leung, P., Liu, X., Minaei, A., Mulatu, M.

S., Novik, T. S., Oh, K. J., Roussos, A., Sawyer, M., Simsek, Z., Steinhausen, H.-C.,

Metzke, C. W., Wolanczyk, T., Yang, H.-J., Zilber, N., Zukauskiene, R., & Verhulst,

F. C. (2007). Testing the 8-Syndrome Structure of the Child Behavior Checklist in 30

Societies. *Journal of Clinical Child & Adolescent Psychology*, *36*(3), 405-417.

https://doi.org/10.1080/15374410701444363

Jenkinson, M., Bannister, P., Brady, M., & Smith, S. (2002). Improved optimization for the

robust and accurate linear registration and motion correction of brain images.

NeuroImage, *17*(2), 825-841. https://doi.org/10.1016/S1053-8119(02)91132-8

Kana, R. K., Libero, L. E., & Moore, M. S. (2011). Disrupted cortical connectivity theory as

an explanatory model for autism spectrum disorders. *Physics of Life Reviews*, *8*(4),

410-437. https://doi.org/https://doi.org/10.1016/j.plrev.2011.10.001

Karcher, N. R., Michelini, G., Kotov, R., & Barch, D. M. (2021). Associations Between

Resting-State Functional Connectivity and a Hierarchical Dimensional Structure of

Psychopathology in Middle Childhood. *Biological Psychiatry: Cognitive*

Neuroscience and Neuroimaging, *6*(5), 508-517.

https://doi.org/https://doi.org/10.1016/j.bpsc.2020.09.008

Kaufman, J., Birmaher, B., Brent, D., Rao, U. M. A., Flynn, C., Moreci, P., Williamson, D.,

& Ryan, N. (1997). Schedule for Affective Disorders and Schizophrenia for School-

Age Children-Present and Lifetime Version (K-SADS-PL): Initial Reliability and

Validity Data. *Journal of the American Academy of Child & Adolescent Psychiatry*,

36(7), 980-988. https://doi.org/https://doi.org/10.1097/00004583-199707000-00021

Kebets, V., Piguet, C., Chen, J., Ooi, L. Q. R., Kirschner, M., Siffredi, V., Misic, B., Yeo, B.

T., & Bernhardt, B. (2023). Multimodal neural correlates of childhood

psychopathology. *bioRxiv*, 2023.2003. 2002.530821.

Kessler, R. C., Angermeyer, M., Anthony, J. C., De Graaf, R., Demyttenaere, K., Gasquet, I.,

De Girolamo, G., Gluzman, S., Gureje, O., & Haro, J. M. (2007). Lifetime prevalence

and age-of-onset distributions of mental disorders in the World Health Organization's

World Mental Health Survey Initiative. *World psychiatry*, *6*(3), 168.

Khatamian, Y. B., Golestani, A. M., Ragot, D. M., & Chen, J. J. (2016). Spin-Echo Resting-

State Functional Connectivity in High-Susceptibility Regions: Accuracy, Reliability,

and the Impact of Physiological Noise. *Brain Connectivity*, *6*(4), 283-297.

https://doi.org/10.1089/brain.2015.0365

Kotov, R., Krueger, R. F., Watson, D., Achenbach, T. M., Althoff, R. R., Bagby, R. M.,

Brown, T. A., Carpenter, W. T., Caspi, A., & Clark, L. A. (2017). The Hierarchical

Taxonomy of Psychopathology (HiTOP): A dimensional alternative to traditional

nosologies. *Journal of abnormal psychology*, *126*(4), 454.

Krishnan, A., Williams, L. J., McIntosh, A. R., & Abdi, H. (2011). Partial Least Squares

(PLS) methods for neuroimaging: A tutorial and review. *NeuroImage*, *56*(2), 455-475.

https://doi.org/https://doi.org/10.1016/j.neuroimage.2010.07.034

Lahey, B. B., Krueger, R. F., Rathouz, P. J., Waldman, I. D., & Zald, D. H. (2017). A

hierarchical causal taxonomy of psychopathology across the life span. *Psychological*

Bulletin, *143*(2), 142-186. https://doi.org/10.1037/bul0000069

Lahey, B. B., Van Hulle, C. A., Singh, A. L., Waldman, I. D., & Rathouz, P. J. (2011).

Higher-Order Genetic and Environmental Structure of Prevalent Forms of Child and

Adolescent Psychopathology. *Archives of general psychiatry*, *68*(2), 181-189.

https://doi.org/10.1001/archgenpsychiatry.2010.192

Lees, B., Squeglia, L. M., McTeague, L. M., Forbes, M. K., Krueger, R. F., Sunderland, M.,

Baillie, A. J., Koch, F., Teesson, M., & Mewton, L. (2021). Altered Neurocognitive

Functional Connectivity and Activation Patterns Underlie Psychopathology in

Preadolescence. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*,

6(4), 387-398. https://doi.org/https://doi.org/10.1016/j.bpsc.2020.09.007

Linke, J. O., Abend, R., Kircanski, K., Clayton, M., Stavish, C., Benson, B. E., Brotman, M.

A., Renaud, O., Smith, S. M., Nichols, T. E., Leibenluft, E., Winkler, A. M., & Pine,

D. S. (2021). Shared and Anxiety-Specific Pediatric Psychopathology Dimensions

Manifest Distributed Neural Correlates. *Biological Psychiatry*, *89*(6), 579-587.

https://doi.org/10.1016/j.biopsych.2020.10.018

662 Marrelec, G., Krainik, A., Duffau, H., Pélégrini-Issac, M., Lehéricy, S., Doyon, J., & Benali,

663   H. (2006). Partial correlation for functional brain interactivity investigation in

664   functional MRI. *NeuroImage*, *32*(1), 228-237.

665 MathWorks. (2020). *MATLAB*. In

666 McIntosh, A. R., & Lobaugh, N. J. (2004). Partial least squares analysis of neuroimaging

667   data: applications and advances. *NeuroImage*, *23*, S250-S263.

668 McTeague, L. M., Huemer, J., Carreon, D. M., Jiang, Y., Eickhoff, S. B., & Etkin, A. (2017).

669   Identification of Common Neural Circuit Disruptions in Cognitive Control Across

670   Psychiatric Disorders. *American Journal of Psychiatry*, *174*(7), 676-685.

671   https://doi.org/10.1176/appi.ajp.2017.16040400

672 McTeague, L. M., Rosenberg, B. M., Lopez, J. W., Carreon, D. M., Huemer, J., Jiang, Y.,

673   Chick, C. F., Eickhoff, S. B., & Etkin, A. (2020). Identification of common neural

674   circuit disruptions in emotional processing across psychiatric disorders. *American*

675   *Journal of Psychiatry*, *177*(5), 411-421.

676 Michelini, G., Barch, D. M., Tian, Y., Watson, D., Klein, D. N., & Kotov, R. (2019).

677   Delineating and validating higher-order dimensions of psychopathology in the

678   Adolescent Brain Cognitive Development (ABCD) study. *Translational Psychiatry*,

679   *9*(1), 261. https://doi.org/10.1038/s41398-019-0593-4

680 Opel, N., Goltermann, J., Hermesdorf, M., Berger, K., Baune, B. T., & Dannlowski, U.

681   (2020a). Cross-Disorder Analysis of Brain Structural Abnormalities in Six Major

682   Psychiatric Disorders: A Secondary Analysis of Mega- and Meta-analytical Findings

683   From the ENIGMA Consortium. *Biological Psychiatry*, *88*(9), 678-686.

684   https://doi.org/https://doi.org/10.1016/j.biopsych.2020.04.027

685 Opel, N., Goltermann, J., Hermesdorf, M., Berger, K., Baune, B. T., & Dannlowski, U.

686   (2020b). Cross-disorder analysis of brain structural abnormalities in six major

psychiatric disorders: a secondary analysis of mega-and meta-analytical findings from the ENIGMA consortium. *Biological Psychiatry*, *88*(9), 678-686.

Paus, T., Keshavan, M., & Giedd, J. N. (2008). Why do many psychiatric disorders emerge during adolescence? *Nat Rev Neurosci*, *9*(12), 947-957. https://doi.org/10.1038/nrn2513

Pettersson, E., Larsson, H., & Lichtenstein, P. (2016). Common psychiatric disorders share the same genetic origin: a multivariate sibling study of the Swedish population. *Molecular Psychiatry*, *21*(5), 717-721. https://doi.org/10.1038/mp.2015.116

Power, J. D., Fair, D. A., Schlaggar, B. L., & Petersen, S. E. (2010). The development of human functional brain networks. *Neuron*, *67*(5), 735-748.

Pruim, R. H. R., Mennes, M., Buitelaar, J. K., & Beckmann, C. F. (2015). Evaluation of ICA-AROMA and alternative strategies for motion artifact removal in resting state fMRI. *NeuroImage*, *112*, 278-287. https://doi.org/https://doi.org/10.1016/j.neuroimage.2015.02.063

Pruim, R. H. R., Mennes, M., van Rooij, D., Llera, A., Buitelaar, J. K., & Beckmann, C. F. (2015). ICA-AROMA: A robust ICA-based strategy for removing motion artifacts from fMRI data. *NeuroImage*, *112*, 267-277. https://doi.org/https://doi.org/10.1016/j.neuroimage.2015.02.064

Roelfs, D., Alnæs, D., Frei, O., van der Meer, D., Smeland, O. B., Andreassen, O. A., Westlye, L. T., & Kaufmann, T. (2021). Phenotypically independent profiles relevant to mental health are genetically correlated. *Transl Psychiatry*, *11*(1), 202. https://doi.org/10.1038/s41398-021-01313-x

Ronald, A. (2019). Editorial: The psychopathology p factor: will it revolutionise the science and practice of child and adolescent psychiatry? [https://doi.org/10.1111/jcpp.13063].

Journal of Child Psychology and Psychiatry, *60*(5), 497-499.

https://doi.org/https://doi.org/10.1111/jcpp.13063

Rubinov, M., & Sporns, O. (2010). Complex network measures of brain connectivity: uses and interpretations. *NeuroImage*, *52*(3), 1059-1069.

Sala-Llonch, R., Smith, S. M., Woolrich, M., & Duff, E. P. (2019). Spatial parcellations, spectral filtering, and connectivity measures in fMRI: Optimizing for discrimination. *Human Brain Mapping*, *40*(2), 407-419.

Salimi-Khorshidi, G., Douaud, G., Beckmann, C. F., Glasser, M. F., Griffanti, L., & Smith, S. M. (2014). Automatic denoising of functional MRI data: Combining independent component analysis and hierarchical fusion of classifiers. *NeuroImage*, *90*, 449-468. https://doi.org/https://doi.org/10.1016/j.neuroimage.2013.11.046

Sato, J. R., Biazoli, C. E., Salum, G. A., Gadelha, A., Crossley, N., Vieira, G., Zugman, A., Picon, F. A., Pan, P. M., Hoexter, M. Q., Amaro, E., Anés, M., Moura, L. M., Del'Aquilla, M. A. G., McGuire, P., Rohde, L. A., Miguel, E. C., Jackowski, A. P., & Bressan, R. A. (2018). Association between abnormal brain functional connectivity in children and psychopathology: A study based on graph theory and machine learning. *The World Journal of Biological Psychiatry*, *19*(2), 119-129. https://doi.org/10.1080/15622975.2016.1274050

Schaefer, A., Kong, R., Gordon, E. M., Laumann, T. O., Zuo, X.-N., Holmes, A. J., Eickhoff, S. B., & Yeo, B. T. T. (2018). Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic Functional Connectivity MRI. *Cerebral Cortex*, *28*(9), 3095-3114. https://doi.org/10.1093/cercor/bhx179

Sha, Z., Wager, T. D., Mechelli, A., & He, Y. (2019). Common Dysfunction of Large-Scale Neurocognitive Networks Across Psychiatric Disorders. *Biological Psychiatry*, *85*(5), 379-388. https://doi.org/https://doi.org/10.1016/j.biopsych.2018.11.011

Shirer, W. R., Jiang, H., Price, C. M., Ng, B., & Greicius, M. D. (2015). Optimization of rs-

fMRI pre-processing for enhanced signal-noise separation, test-retest reliability, and

group discrimination. *NeuroImage*, *117*, 67-79.

Sydnor, V. J., Larsen, B., Bassett, D. S., Alexander-Bloch, A., Fair, D. A., Liston, C.,

Mackey, A. P., Milham, M. P., Pines, A., & Roalf, D. R. (2021). Neurodevelopment

of the association cortices: Patterns, mechanisms, and implications for

psychopathology. *Neuron*, *109*(18), 2820-2846.

Uddin, M. N., Figley, T. D., Solar, K. G., Shatil, A. S., & Figley, C. R. (2019). Comparisons

between multi-component myelin water fraction, T1w/T2w ratio, and diffusion tensor

imaging measures in healthy human brain structures. *Scientific reports*, *9*(1), 1-17.

van den Heuvel, M. P., & Sporns, O. (2019). A cross-disorder connectome landscape of brain

dysconnectivity. *Nature Reviews Neuroscience*, *20*(7), 435-446.

https://doi.org/10.1038/s41583-019-0177-6

Vanes, L. D., & Dolan, R. J. (2021). Transdiagnostic neuroimaging markers of psychiatric

risk: A narrative review. *NeuroImage: Clinical*, *30*, 102634.

https://doi.org/https://doi.org/10.1016/j.nicl.2021.102634

Voldsbekk, I., Kjelkenes, R., Wolfers, T., Dahl, A., Lund, M. J., Kaufmann, T., Fernandez-

Cabello, S., de Lange, A.-M. G., Tamnes, C. K., Andreassen, O. A., Westlye, L. T., &

Alnæs, D. (2023). Shared pattern of impaired social communication and cognitive

ability in the youth brain across diagnostic boundaries. *Developmental Cognitive*

Neuroscience, *60*, 101219. https://doi.org/https://doi.org/10.1016/j.dcn.2023.101219

Woolrich, M. W., Ripley, B. D., Brady, M., & Smith, S. M. (2001). Temporal

Autocorrelation in Univariate Linear Modeling of FMRI Data. *NeuroImage*, *14*(6),

1370-1386. https://doi.org/https://doi.org/10.1006/nimg.2001.0931

Xia, C. H., Ma, Z., Ciric, R., Gu, S., Betzel, R. F., Kaczkurkin, A. N., Calkins, M. E., Cook,

P. A., García de la Garza, A., Vandekar, S. N., Cui, Z., Moore, T. M., Roalf, D. R.,

Ruparel, K., Wolf, D. H., Davatzikos, C., Gur, R. C., Gur, R. E., Shinohara, R. T.,

Bassett, D. S., & Satterthwaite, T. D. (2018). Linked dimensions of psychopathology

and connectivity in functional brain networks. *Nature Communications*, *9*(1), 3003.

https://doi.org/10.1038/s41467-018-05317-y

Xia, M., Wang, J., & He, Y. (2013). BrainNet Viewer: A Network Visualization Tool for

Human Brain Connectomics. *PLoS ONE*, *8*(7), e68910.

https://doi.org/10.1371/journal.pone.0068910