Study of the Effect of Intestinal Microbes on Obesity: A Bibliometric Analysis

Zehao Su[1,2], Chenyu Tian[1,2], Guan Wang[1*], Jingjing Guo[1,2*], Xiaoyan Yang[1,2*]

[1] Department of Endocrinology & Metabolism and West China Biomedical Big Data Center, West
China Hospital, Sichuan University/West China School of Nursing, Chengdu 610041, Sichuan,
China.

[2]Med-X Center for Informatics, Sichuan University, Chengdu 610041, China.

Keywords: obesity[1], intestinal microbiota[2], bibliometric analysis[3], visualization[4], co-occurrence
analysis[5].

Abstract

Background: Obesity is a serious public health problem. According to statistics, there are millions
obese people worldwide. The intestinal microbiota is the microbiota that lives in the gastrointestinal
tract of the human body and can interact with the external environment through diet. Probing and
summarizing the relationship between intestinal microbes and obesity has important guiding
significance for the accurate control of the research direction and expanding the choice of obesity
treatment methods. We aimed to use bibliometrics to qualitatively and quantitatively analyze the
published literature and to reveal the research hotspots and development trends in the effects of
intestinal microbes on obesity research.

Methods: We obtained documents from the core collection of Web of Science (WoSCC) from 2013–
2022 on February 2nd, 2023. Microsoft Excel, Biblioshiny R language software packages and
CiteSpace were employed to collect publication data, analyze publication trends, and visualize relevant
results.

Results: We identified 8888 original research articles on the effects of intestinal microbes on obesity
published between 2013 and 2022. China made the highest number of publications on the topic, with
3036 publications (34.16%), and the United States with 87065 citations made the greatest contribution.
University of Copenhagen was the most prolific institution (209 publications). Patrice D. Cani
published the most related articles (97 publications), whereas Peter J. Turnbaugh was cited the most
frequently (3307 citations). Nutrients was the journal with the most studies (399 publications, 4.49%),
and Nature was the most commonly co-cited journal (6446 citations). The burst keywords "obese
patient" and "serum" recently appeared as research frontiers.

Conclusion: Although China has the largest number of publications, the United States, depending on
its largest citations, has become the leader in the effects of intestinal microbes on obesity research.
Current research hotspots focus on related mechanisms of the effects of intestinal microbes on obesity
and therapeutic methods for obesity with intestinal microbes. Through diet, probiotic preparations and
regulation of the intestinal flora, fecal microbe transplantation may have a significant impact on
reducing the obesity epidemic. Considering the great potential and application prospects, intestinal
microbe applications in obesity therapy will remain a research hotspot in the future.

1   Introduction

Obesity is a serious public health problem. According to statistics, there are approximately 600 million obese people and 1.9 billion overweight individuals worldwide (El-Sayed Moustafa and Froguel, 2013). It is predicted that the number of obese people in the world will reach 1.12 billion by 2030 (KellyYang et al., 2008). As far as we know, obesity is not only a change in appearance but also associated with disorders of lipid and glucose metabolism, chronic inflammation, oxidative stress and increased risk of a variety of diseases. According to research, obesity is one of the nonnegligible reasons for cardiovascular disease, diabetes and malignant tumors (BoccellinoDi Domenico et al., 2018; Scherer and Hill, 2016), and it is estimated that more than 80% of the global burden of diabetes and cardiovascular disease will occur in developing countries such as China and India in 2025 (MohanGupta et al., 2016). However, the etiology of obesity is not yet completely clear. The increasing prevalence of obesity will not be explained by changes in energy balance alone. Therefore, it is vital for the whole world to determine the formation mechanism of obesity and then explore effective methods to treat it and reduce the incidence and mortality of obesity-related diseases.

The intestinal microbiota is a microbiota that lives in the gastrointestinal tract of the human body and can interact with the external environment through diet (AlessandroMarcello et al., 2017). These microbes colonize the intestinal tract through mother-to-child transmission from the prenatal period. The colonization of human intestinal microbes continues after birth and is regulated by factors including gestational age, mode of delivery (natural delivery or cesarean section), diet (breastfeeding or baby food), hygiene and antibiotics. The environment and diet of the first three years of life are essential for the acquisition of adult-like microbiota and the establishment of bacteria-host symbiosis that affect immune and nervous system development. At the age from 2 to 5, the intestinal microbiota will gradually stabilize, and there is no significant difference with intestinal microbiota in adults (RodríguezMurphy et al., 2015). Under normal circumstances, the intestinal microbiota maintains a symbiotic relationship with the human body, and a good symbiotic relationship between them is very important for the maintenance of human health (JandhyalaTalukdar et al., 2015), but instantaneous changes in the intestinal ecosystem can lead to the destruction of the symbiotic relationship between microorganisms and hosts, such as aging, obesity, sedentary lifestyles, and dietary patterns, and the effects of antibiotics can change the intestinal microbiota (NazliYang et al., 2004). Because of the important role of the intestinal ecosystem in maintaining host physiology, its changes can also cause a variety of physiological disorders in the human body, including low-grade inflammation, metabolic disorders, excessive lipid accumulation and loss of insulin sensitivity, in turn leading to changes in the composition of the intestinal microbiota and decreasing the diversity of flora and metabolic pathways (Halmos and Suba, 2016; Sidhu and van der Poorten, 2017), thus increasing the risk of metabolic diseases such as obesity and diabetes (TangKitai et al., 2017). Probing and summarizing the relationship between intestinal microbes and obesity has important guiding significance for the accurate control of the research direction and expanding the choice of obesity treatment methods.

Bibliometrics is a discipline that uses mathematics and statistics to quantitatively analyze literature and information. Through quantitative analysis of the subject literature, we can construct its knowledge structure and explore its development trend. At present, it has been widely used in many research fields (GaoDou et al., 2021; LiuGao et al., 2022; WangMa et al., 2020). Overall, this analysis method and the traditional literature review are both based on previous research, summarize the current situation and shortcomings of the research, and finally guide further research. Literature review emphasizes the content of the article, that is, to summarize the aspects and shortcomings of the existing research, and most of them quote the representative papers in the literature according to a preset research context.

On the other hand, bibliometric analysis places more emphasis on the analysis of "quantity". There is
no need to analyze the research content of each document in detail but to analyze the number of
published documents, the distribution of authors and the relationship of citations in the target field.
Most of the included documents are highly quoted. Compared with the traditional literature review,
bibliometric analysis can make a more intuitive systematic analysis of all the literature in this field in
a visual way, which is helpful for researchers entering a new field to grasp the overall trend of the field.
There have been many review articles in the field of intestinal microbes and obesity, but the quantitative
analysis in the field is insufficient. Therefore, in addition to briefly summarizing the hotpots in review
articles, we use bibliometrics to qualitatively and quantitatively analyze the published literature and
intends to reveal the research hotspots and development trends in this field.

2   Materials and methods

2.1   Data sources and search strategies

Data were obtained from the core collection of Web of Science (WoSCC), a database of Clarivate
Analytics. The following search strategies were presented: ((overweight OR obesity OR obes* OR
over$weight) AND (gastrointestin* OR gastro-intestin* OR gut OR intestin*) AND (bacteria OR
prebiotic OR probiotic OR microbiot* OR microbiome* OR flora OR microflora)). In the end, a total
of 8888 original articles in English were retrieved. All the documents were downloaded as "Full Record
and Cited References" and saved as "Bibtex" and "plain text file". The above data were obtained on
2nd February 2023.

2.2   Data analysis and graph acquisition

The corresponding files retrieved from WoSCC were imported into Microsoft Excel (16.66.1),
Biblioshiny and CiteSpace (6.1. R6) to perform a bibliometric and visual analysis. Annual publications,
country annual publications, and publisher annual publications were analyzed with Microsoft Excel.
The Biblioshiny platform provides a web-based graphical interface in the Bibliometrix package, which
was used to conduct total citation, average article cited, publications of journal and country
collaboration map. CiteSpace is interactive visualization analysis software developed by Professor
Chen Chaomei to conduct different types of network analyses, such as country collaboration networks,
institution collaboration networks, author collaboration networks, co-cited author collaboration
networks, co-cited journal collaboration networks, and keyword co-occurrence, which aid in visually
analyzing the knowledge domain and emerging trends. A dual-map overlay was constructed for
journals. The parameters of CiteSpace were as follows: time slicing (2013 to 2022), years per slice (1),
links (strength: cosine, scope: within slices), selection criteria (g-index: k=5 in country, institution,
author collaboration network analysis and keyword co-occurrence), pruning (pathfinder, pruning sliced
networks). Cluster labels used keywords by log likelihood ratio (LLR). To identify emerging topics,
we detected keywords and references with a strong citation burst.

Ethical approval was waived by the local ethical board because the data came from public databases,
and no human or animal subjects were involved.

3   Results

3.1   An overview of the annual growth trend

As shown in Figure 1A, from 2013 to 2022, a total of 8888 English-language articles met our
inclusion criteria. The number of annual publications began to exceed 500 in 2016, and only after 3

years did the annual publications exceed 1000. The annual number of publications increased from 267 in 2013 to 1768 in 2022, with an amazing increase of over sixfold. We further analyzed the annual national output of the 10 most productive countries and regions (Figure 1B). The different colors represent different countries, and the slope represents the speed of annual publication number increase. China ranked first in the number of annual publications, followed by the United States and Spain. The top 10 publishers according to their contribution to the total number of articles about the effect of intestinal microbes on obesity are shown in Figure 1C. The different colors represent different years' publications. Regarding the number of publications, Elsevier (1685) and Springer Nature (1388) far surpassed other publishers, confirming their unsurpassable international position in the publishing industry.

3.2 Country influence and collaboration

In the ten years from 2013-2022, 124 countries or regions participated in the domain of intestinal microbes and obesity. China ranked first with 3036 publications, accounting for 34.16% of all publications, followed by the United States (2356, 26.51%) and Spain (480, 5.40%) (Figure 2A, Table 1). Nevertheless, the United States tops the ranking of the most cited countries with a frequency (number of documents) of 87065, followed by China (65250), France (13809), and Canada (13496) (Supplementary Table S1). The country collaboration map shows the overall perspective of countries' and regions' academic cooperation (Supplementary Figure S1). The shade of blue represents the number of articles published, dark blue indicates more publications, and as the color becomes lighter, the publications decreased. The red line represents that there is academic collaboration between the linked two countries. The thicker the line is, the more cooperation two countries have. The United States had the broadest academic associations with other countries or regions, while its collaboration with China was the tightest, followed by Canada. The United States and China both had a frequent connection with European countries. Moreover, among European countries, there was also tight collaboration, but communication among the other countries or regions still needs to be strengthened.

There were 179 institutions included in the results of institution analysis. The top 10 institutions mostly came from China (5) and the USA (2), followed by Denmark (1), Belgium (1), and Spain (1). Chinese academic of sciences, the University of Copenhagen and Shanghai Jiao Tong University had the highest total link strength, revealing close cooperation with other institutions (Figure 2B, Table 2).

3.3 Distribution and co-authorship of authors

In this analysis, 16 authors had published at least 21 publications. The 10 most productive authors are listed in Table 3. Most of the authors were from Europe, with Belgium (n=3), followed by China (n=3), Canada (n=2), Sweden (n=1), and France (n=1). Patrice D. Cani (97 publications) and Nathalie M. Delzenne (73 publications), both from Université catholique de Louvain, contributed the top two most publications, followed by Reimer, Raylene A (51 publications) from the University of Calgary. The top two authors maintained the most contact with other research labs, and the cooperation between Patrice D. Cani and Nathalie M. Delzenne was close (Figure 3A). Co-cited authors are shown in Figure 3B and Table 4. Peter J. Turnbaugh, and Patrice D. Cani rank the top two, which manifests their centrality in the research field.

3.4   Analysis of journals and co-cited academic journals

As shown in Table 5, the top 10 most productive journals published 2155 publications, accounting for 24.25% of all publications. Nutrients (impact factor (IF) 2021, 6.706) published the most articles (399 publications, 4.49%), followed by Scientific Reports (IF 2021, 4.997; 339 publications, 3.81%) and Food & Function (IF 2021, 6.317; 281 publications, 3.16%). The top three highest co-cited journals were Nature (IF 2021, 69.504; 6446 citations), PLOS ONE (IF 2021, 3.752; 6247 citations), and Proceedings of the National Academy of Sciences of the United States of America (IF 2021, 12.779; 5329 citations). The co-citation relationship among different journals was visualized in a co-citation network (Figure 4, Table 6). Nature had the highest centrality (0.18), followed by PLOS ONE (0.17), which means that the published articles in the two journals were widely recognized and cited by researchers.

Supplementary Figure S2 shows the dual-map overlay of journals. The yellow, green, and orange spline waves represent citations made by the source articles. Each spline curve starts with the citing map on the left and points to the cited map on the right. The label represents the subject covered by the journal. The dual-map overlay shows seven main citation paths. The published articles were mostly focused on journals in the fields of medicine, medical, clinical, molecular, biology, and immunology and partly focused on veterinary, animal and science. Cited articles were mostly published in journals in the fields of molecular biology and genetics, and many articles were published in journals in the fields of health, nursing, medicine, environmental, toxicology and nutrition.

3.5   Analysis of keywords

3.5.1 Co-occurrence analysis

Keywords could accurately reflect the main research point of an article, which has a high condensation in a research field and can directly point to the center of the text. Therefore, a high frequency of keywords represents hot issues in a research field and research hotspots. The keyword co-occurrence graph for the effects of intestinal microbes on obesity is shown in Figure 5. The density value was 0.0457. The ten most frequent keywords were gut microbiota, obesity, inflammation, insulin resistance, intestinal microbiota, health, metabolism, diet, gut microbiome, and metabolic syndrome (Table 7).

3.5.2 Cluster analysis

The clustering analysis of the keywords yielded 288 nodes and 1083 links, with a Q value of 0.4123 (>0.3) and an S value of 0.7178 (>0.5), which are two important indicators to evaluate the significance of the clustering effect. There were 8 clusters, including diversity, insulin resistance, short-chain fatty acids, activation, food intake, bariatric surgery, body composition, and nonalcoholic fatty liver disease (Figure 6). "diversity" # 0 was the largest cluster, followed by "insulin resistance" # 1 and "short-chain fatty acids" # 2. Timeline View analysis was conducted to further analyze the keywords of the effects of intestinal microbes on obesity (Supplementary Figure S3).

3.5.3 Burst detection.

CiteSpace was further used to perform burst detection with high frequency and therefore revealed research frontiers and hotspot trends over a period. As shown in Supplementary Table S2, the first 25 keywords were sorted by strengths of burst to discover the research hotspots about the effects of intestinal microbes on obesity. Keywords such as "ecology" (strength 28.33), "human gut microbiota" (strength 27.34), "microbiota" (strength 23.44), "diet induced obesity" (strength 21.12), and "irritable

203 bowel disease" (strength 16) were the strongest keywords leading the research boom from 2013-2018.
204 Keywords such as "obese" (strength 9.3) and "serum" (9.69) were the most recent trending keywords.
205 Supplementary Table S3 shows the top 25 references with the strongest citation bursts, which were
206 recognized as a critical milestone and led the development direction of the field for a while.

4   Discussion

208   In this study, we performed a bibliometric analysis of the effects of intestinal microbes on obesity-
209 related studies from 2013 to 2022 using the core collection of WoSCC to comprehensively understand
210 global research trends and hotspots and provide references for researchers in this field or those who
211 want to become involved in the field. Based on the above analysis, it is certain that obesity is closely
212 influenced by intestinal microbes.

4.1   Global trends in effects of intestinal microbes on obesity

The amount of annual scientific production is an important indicator of development in an academic
field. In our study, a total of 8888 original articles from 1392 journals met the inclusion criteria. The
article annual growth rate reached 22.93%, and the annual growth rate from 2019 to 2020 was the
highest. An article published by Lawrence A. David et al., which was the most co-cited, found that an
altered diet could lead to a rapid gut microbiome response. The results showed that short-term
consumption of diets composed entirely of animal or plant products alters microbial community
structure and overwhelms interindividual differences in microbial gene expression (DavidMaurice et
al., 2014). The second most cited article was published by Amandine Everard et al. They isolated a
mucin-degrading bacterium that resides in the mucus layer, named *Akkermansia muciniphila*. The
results showed that the abundance of *A. muciniphila* decreased in obese and type 2 diabetic mice, but
prebiotic feeding could normalize *A. muciniphila* abundance. Moreover, high-fat diet-induced
metabolic disorders, including fat-mass gain, metabolic endotoxemia, adipose tissue inflammation, and
insulin resistance, could be reversed by *A. muciniphila* treatment (EverardBelzer et al., 2013). Although
these two articles are about different research, they both proved the correlations between intestinal
microbes and the host, which have greatly driven the development of the field.

Among the 10 most influential journals in effects of intestinal microbes on obesity research, four
were also among the top 10 co-cited journals (Nature, Proceedings of the national academy of sciences
(PNAS), Gut, Science). Among these, the IF value of Nature was 69.504 (2021), followed by Science
(63.832), Gut (31.795) and PNAS (12.779), indicating their outstanding contributions in the field of
effects of intestinal microbes on obesity research. The most frequently co-cited journals were Nature,
PLOS ONE, PNAS, and Gut. Notably, three of the top 10 co-cited references were published in Nature,
one of them was published in PNAS, and one of them was published in Gut. The main research hotspots
were about the mechanism of dietary intervention or human genetics to intestinal microbes. Moreover,
richness or kinds of intestinal microbes can also modulate the host's metabolism, such as obesity or
type 2 diabetes. These findings have laid the foundation for a deeper understanding of the regulatory
relationship between intestinal microbes and obesity or type 2 diabetes.

4.2   Co-authorship networks in effects of intestinal microbes on obesity research

The closeness of collaboration between countries/regions, institutions, and authors was assessed,
which can help find the laws of scientific research cooperation, guiding more effective scientific

research activities, and promoting potential collaborative opportunities for other groups. According to
Table 1, China, with 3036 articles, ranks first among the top 10 total publication countries, followed
by the United States, with 2356 publications. Notably, the total number of publications in the USA
(87065) was greater than that in China (65250) and far greater than that in France (13809), Canada
(13496) and the other top 10 most cited countries. The total citations of the United States and China
are far more than several other top 10 most cited countries citations combined. Without doubt, the
United States and China demonstrate their position as leaders in the field. Notably, Israel with 49
publications was the most cited country per article (average article citations, 124.18), suggesting that
the quality of research on effects of intestinal microbes on obesity in Israel is very high, the similar
story is unfolding in many European countries such as Belgium (average article citations, 107.85),
Sweden (average article citations, 103.95), etc., which sets a model for other countries/regions in this
research field (Supplementary Table S1, S4). To improve the quality of articles, the reliability of the
data source and the rigor of the experimental design may be the first factors affecting the quality of
publications, and more attention should be given to these aspects in future studies. According to Figure
2B, the bottom right is most dominated by universities from China, while the upper left is most
dominated by schools in North America or Europe. Nevertheless, more cooperation between the two
clusters could be conducted to catalyze breakthrough progress in research on the effects of intestinal
microbes on obesity.

Four of the top 10 institutions are based in China, followed by the United States with three, which
are thereby maximizing regional advantages and demonstrating the dominance of the United States
and China in the field. This may partly explain why China and the United States consistently maintain
a high quantity of publications. University of Copenhagen in Denmark was the most productive
institution worldwide, followed by Chinese academic of sciences in China, indicating that these two
institutions participated in the most collaborations with other institutions worldwide. Although Spain
and Canada ranked third and fourth in terms of total publications, only one of the Spanish research
institutions ranked in the top 10, indicating a lack of institutions with professional and research stature
in terms of the effects of intestinal microbes on obesity research (Table 1-2). The most effective
organizations and groups are leading the trends in the effects of intestinal microbes on obesity research;
thus, further study at these institutions will ensure continuous future development in this field.

272　4.3　Basic knowledge and hotspots in effects of intestinal microbes on obesity research

273　4.3.1 Basic knowledge

The most frequently cited references possess pivotal academic position in the field. Either positive
or negative conclusions from the documents influence others' research direction. In our study, the top
co-cited references were used to investigate the knowledge base for the effects of intestinal microbes
on obesity research. Among the top 10 co-cited references, we found that second-generation
sequencing was a vital technology that relies on its high-throughput characteristic, making it easy to
sequence the transcriptome or genome of a species (Schuster, 2008). With the development of
sequencing technology, a large amount of data is generated. To make more accurate and full use of
sequencing data, software and data analysis platforms have also been developed. An open-source
software package named DADA2 is used for modeling and correcting Illumina-sequenced amplicon
errors. With the help of this software, researchers could accurately reconstruct amplicon-sequenced
communities at the highest resolution, which ensured the accuracy of the research to the greatest extent
(CallahanMcMurdie et al., 2016). Another utility tool named QIIME 2 could serve not only as a
marker-gene analysis tool but also as a multidimensional and powerful data science platform that can
be rapidly adapted to analyze diverse microbiome features (BolyenRideout et al., 2019). These tools
help drive rapid development in microbiome research. With the help of sequencing, recent research

found that the richness of the human gut microbiome correlated with human metabolic markers. Individuals with a low bacterial richness were characterized by more marked overall adiposity, insulin resistance and dyslipidemia and a more pronounced inflammatory phenotype when compared with high bacterial richness individuals (Le ChatelierNielsen et al., 2013). Correspondingly, with higher gut microbiome gene richness and *A. muciniphila* abundance could exhibit the healthiest metabolic status, particularly in fasting plasma glucose, plasma triglycerides and body fat distribution (DaoEverard et al., 2016). The obesity-associated gut microbial species *Bacteroides thetaiotaomicron,* a glutamate-fermenting commensal, is linked to changes in circulating metabolites, and the abundance of *Bacteroides thetaiotaomicron* is markedly decreased in obese individuals and inversely correlated with serum glutamate concentration (LiuHong et al., 2017). Human obesity is a heterogeneous condition in the context of pathogenesis, pathophysiology and therapeutic responsiveness. Studies of alterations in the genome—the microbial gut metagenome—may define subsets of adult individuals with different metabolic risk profiles, which could contribute to resolving some of the heterogeneity associated with adiposity-related phenotypes (Le ChatelierNielsen et al., 2013). All these findings suggest that gut microbiome richness is a key factor in maintaining the homeostasis of human health.

The gut microbiome can rapidly respond to an altered diet, and a dynamic balance can be achieved between intestinal microbes and diet immediately, indicating that not only a long-term diet but also a short-term diet can influence the gut microbiome (DavidMaurice et al., 2014). Weight-loss intervention by bariatric surgery partially reversed obesity-associated microbial and metabolic alterations in obese individuals (LiuHong et al., 2017). This means that external interventions can affect intestinal microbes. Research also proved that transmissible and modifiable interactions between diet and microbiota influence host biology. The transformation correlated with invasion of members of *Bacteroidales* from Ln into Ob microbiota that prevented development of increased adiposity and body mass in Ob cage mates and transformed their microbiota's metabolic profile to a leanlike state. Therefore, it may be possible to intervene in obesity by targeting the gut microbiota (RidauraFaith et al., 2013), and there is already evidence proving this speculation. *Akkermansia muciniphila* is a mucin-degrading bacterium that resides in the mucus layer. Research found that *A. muciniphila* decreased in obese and type 2 diabetic mice, and *A. muciniphila* treatment reversed high-fat diet-induced metabolic disorders, including fat-mass gain, metabolic endotoxemia, adipose tissue inflammation, and insulin resistance (EverardBelzer et al., 2013). Pasteurized *A. muciniphila* could enhance its capacity to reduce fat mass development, insulin resistance and dyslipidemia. Moreover, Amuc_1100, a purified membrane protein from *Akkermansia muciniphila*, could improve the gut barrier and partly recapitulate the beneficial effects of the bacterium (PlovierEverard et al., 2017). All these findings are of great driving significance, indicating that it is feasible to use microbes to treat obesity, but more research is needed to support this hypothesis.

In addition to the underlying effect of intestinal microbes on obesity, substantial evidence has suggested that intestinal microbes also have an effect on type 2 diabetes, which is also a major public health issue throughout the world. A metagenome-wide association study (MGWAS) was used to identify and validate type-2-diabetes-associated markers and to establish the concept of a metagenomic linkage group, which enabled taxonomic species-level analyses. Type 2 diabetes is accompanied by a decrease in the abundance of some universal butyrate-producing bacteria and an increase in various opportunistic pathogens (QinLi et al., 2012). Conversely, high bacterial richness was related to a reduction in insulin resistance (Le ChatelierNielsen et al., 2013), and *A. muciniphila* treatment could also reverse insulin resistance (EverardBelzer et al., 2013) (Table 8).

4.3.2 Research hotspots

Keywords with high frequency are usually used to accurately reveal the main research interests and hotspots in the field within a certain period, which help researchers to precisely catch up to the research trends from numerous studies. By analyzing the co-occurrence keywords, "gut microbiota" and "obesity" had the highest frequency and centrality ranking the top two, indicating that the main research interests focused on the relationship between "gut microbiota" and "obesity". The effects of intestinal microbes on obesity have been supported by several trials, which have all shown that intestinal microbes modulate the metabolism of obesity (EverardBelzer et al., 2013; Le ChatelierNielsen et al., 2013; RidauraFaith et al., 2013). The clustering function was used to show the whole network map. The 8 clusters of keywords presented in this paper illustrate 8 research topics, which to some extent represent 8 significant aspects of the research. Studies in the same cluster might have similar research themes. As shown in Figure 6, the top two largest clusters, "diversity" # 0 and "insulin resistance" # 1, showed that the two topics were still the main themes in the effect of intestinal microbes on obesity research. Simultaneously, a timeline view analysis was conducted to further observe the evolution pace of each cluster and recognize the pivotal research direction through a microperspective. We can clearly observe in Supplementary Figure S3 that the research focus has shifted with time; by 2020, the research on several topics was more balanced. After 2020, the research in the "shot-chain fatty acids" #2 and "activation" #3 directions gradually weakened. The results indicated that gut microbiome research and the treatment of obesity and obesity-related diseases have become the main themes.

Furthermore, burst detection was an analytic tool that could effectively capture the dramatic increases in references or keywords in one research field within a specified period; therefore, it served as an important indicator of research hotspots or research frontiers over time. The evolution of the burst keywords over the last 10 years is shown in Supplementary Table S2. According to the ranking by burstiness strength, we observed that the top 15 keywords started to burst in 2013, and "ecology" remained the longest with an end time in 2018. More importantly, the "obese patient" and "serum" burst separately in 2019 and 2020 and continue to burst until now, which indicated that the two topics have received continuous attention in recent years and might be the main trends of research on the effect of intestinal microbes on obesity. Notably, there was a sudden and rapid growth in the number of articles in the field, which might be related to the term "serum", which was only a new burst in that year.

Bariatric surgery is considered the only effective and sustainable weight loss method for obese patients. There is a 50–70% decrease in body weight and fat mass after surgical procedures such as Roux-Y gastric bypass (RYGB) and sleeve gastrectomy (SG) (PaganelliLuyer et al., 2019). Nevertheless, obese patients undergoing bariatric surgery may experience overgrowth of small intestinal bacteria (SabateCoupaye et al., 2017), a condition that snags with weight loss and increases the risk of micronutrient deficiency that appears to be harmful for the configuration and composition of intestinal microbiota (HibberdWu et al., 2017; Mach and Clark, 2017). To maintain the effect of surgery and avoid weight rebound, it is important for obese patients to correct the microbial balance and improve microbiota-host interactions with specific interventions (Aron-WisnewskyPrifti et al., 2019). Research has also shown that the elevated pH resulting from RYGB could ensure the survival of probiotic bacteria, making it possible for surgical patients to receive probiotic therapy (80). Probiotics are a kind of active microorganism beneficial to the host that colonizes the human intestinal tract and reproductive system, and they can improve the host microecological balance and play a beneficial role. Bariatric surgery is often followed by an increase in *Streptococcaceae* and a decline in *Bifidobacteriaceae* (PaganelliLuyer et al., 2019). Proper supplementation with probiotics can compensate for the intestinal microbial imbalance caused by surgery. Moreover, recent research showed that probiotic intervention could increase the levels of peptide YY and GLP-1 in mice (TianLi et al., 2016), reduce the level of intestinal inflammation (TsaiLin et al., 2019), and induce the

production of anti-inflammatory cytokines (FolignéParayre et al., 2016). Meanwhile, intestinal peptide
signals activate the gut-brain axis, which, in turn, exerts endocrine effects on other organ systems,
especially the brain, regulating appetite, metabolism, and other dietary behaviors(Bliss and Whiteside,
2018; Hussain and Bloom, 2013; NeyrinckVan Hée et al., 2012; RastelliVan Hul et al., 2020).
Consequently, weight loss and reduced insulin resistance occur. However, due to the heterogeneity of
experimental techniques, methods and objects, the effectiveness and safety of microbial product
intervention in improving obesity against intestinal microbiota need to be further verified, and the
mechanism is not clear. Few studies have focused on the development of new functional
microbiological products, and most of them have been carried out in mice. Before microbial products
are reasonably and effectively treated for obesity, a large number of studies are needed, especially
randomized controlled clinical trials.

Serum samples are a less invasive biomatrix that reflects the dynamic changes in the metabolome
of the whole organism (DunnBroadhurst et al., 2011). The serum metabolome has emerged as a
technique that focuses on defining the functional status of host-microbial relationships in biological
specimens, which can reflect the dynamic changes in metabolites and explore disease-related
metabolites or dysregulated metabolic pathways (metabolome analysis for investigating host-gut
microbiota interactions). A top 10 co-cited reference showed the gut microbiome and serum
metabolome alterations in obesity that patients after bariatric surgery not only appeared partially
reversed obesity-associated microbial but also accompanied by metabolic alterations, including the
decreased abundance of *B. thetaiotaomicron* and the elevated serum glutamate concentration,
suggesting that it may be possible to intervene in obesity by targeting the gut microbiota (LiuHong et
al., 2017). There are complex interactions between the host and intestinal microbes in carbohydrate,
amino acid, lipid and nucleic acid metabolism. Intestinal microbes can use their respective metabolites
to maintain intestinal viability while affecting the development, homeostasis and function of the host
immune system through nutrition- and metabolite-dependent mechanisms (Brestoff and Artis, 2013).
Small molecules, such as vitamins, fatty acids, amino acids, and bile acids, regulate host-intestinal
metabolic homeostasis by binding to specific host membranes or nuclear receptors (HustedTrauelsen
et al., 2017; PutignaniDel Chierico et al., 2016). The references with the highest citation bursts are
unique contributions to the field of study. We found that articles related to bile acids have become the
most cited articles in recent years (KohDe Vadder et al., 2016; WangLiao et al., 2019). Bile acids
directly and rapidly affect the metabolism of bacteria, including membrane damage and disruption of
amino acid, nucleotide and carbohydrate metabolism, and short-term exposure to bile acids
significantly affects host metabolism by altering the bacterial community structure (TianGui et al.,
2020). Another study showed that the gut microbiota also produced many other metabolites with
important functions, such as short-chain fatty acids (SCFAs) and bile acids (KrishnanDing et al., 2018).
Short-chain fatty acids can directly activate G-coupled receptors, inhibit histone deacetylases, and
serve as energy substrates (KohDe Vadder et al., 2016). They thus affect various physiological
processes and may contribute to health and disease. According to the above analysis, although notable
advances in the effects of intestinal microbes on obesity have been made, our understanding of the
interrelationships between them remains descriptive, and we still have numerous gaps to fill.

4.4   Strengths and Limitations

The bibliometric study conducted a systematic analysis of the basic situation, research hotspots,
and trends in effects of intestinal microbes on obesity from a visualization perspective. The results of
the bibliometric study were objective and accurate, which could provide a comprehensive guide for

academics who are already or wish to work in this field. Nevertheless, there are still some limitations
in our study. First, owing to the nature of the CiteSpace software, our data are filtered only from the
WoSCC database, which is not sufficiently comprehensive and may lead to data omission. Second, our
results were processed by CiteSpace software with certain algorithms, which could lead to bias in some
of the results. Finally, only English articles were included from the database and analysis, potentially
leading to a source bias.

5 Conclusion

Intestinal microbes possess essential research value and application prospects in obesity therapy.
Compared with other review articles, the contribution of the study is evident in its visualized ways to
reveal the effect of intestinal microbes on obesity. Research on the effects of intestinal microbes on
obesity is increasing rapidly, highlighted by the outstanding contributions of China, the United States
and Spain to the development of this field. Current research hotspots focus on related mechanisms of
the effects of intestinal microbes on obesity and therapeutic methods for obesity with intestinal
microbes. Diet, probiotic preparations and regulation of the intestinal flora, fecal microbe
transplantation may have significant impacts on reducing the obesity epidemic. "Obese patient" and
"serum" have become the latest burst that have attracted attention in recent years and will serve as the
focus of future research. More cooperation and communication between countries and institutions
should be strengthened to promote development in this field and benefit more patients with
obesity.Data availability statement

The original contributions presented in the study are included in the article, and further inquiries
can be directed to the corresponding author.

Author Contributions

447 ZS and JG conceived the study. XY guided the methodological process. ZS, CT participated in
448 data collection and analysis. ZS wrote the manuscript. GW, JG and XY revised the manuscript. All
449 authors contributed to the article and approved the submitted version.

Funding

451 This study was funded by West China Hospital, Sichuan University (Project No. 339180262).

Acknowledgments

453 The authors are indebted to Prof. Zhiyong Zong, who contributed his time, knowledge, and
454 energy to developing this document.

Conflict of interest

456 The authors declare that the research was conducted in the absence of any commercial or financial
457 relationships that could be construed as a potential conflict of interest.

Publisher's note

459 All claims expressed in this article are solely those of the authors and do not necessarily represent
460 those of their affiliated organizations or those of the publisher, the editors and the reviewers. Any
461 product that may be evaluated in this article or claim that may be made by its manufacturer is not

462   guaranteed or endorsed by the publisher.

Supplementary Material

464   The Supplementary Material for this article can be found online at:

References

Alessandro, F., Marcello, D., Rosa, D. S., Valentina, G., and Luca, M. 2017. Gut microbiota, obesity and metabolic
disorders. *Minerva gastroenterologica e dietologica* 63 （4）, 337-44. doi: 10.23736/S1121-421X.17.02376-5.

Aron-Wisnewsky, J., Prifti, E., Belda, E., Ichou, F., Kayser, B. D., Dao, M. C., et al. 2019. Major microbiota dysbiosis in
severe obesity: fate after bariatric surgery. *Gut* 68 （1）, 70-82. doi: 10.1136/gutjnl-2018-316103.

Bliss, E. S., and Whiteside, E. 2018. The Gut-Brain Axis, the Human Gut Microbiota and Their Integration in the
Development of Obesity. *Front Physiol* 9, 900. doi: 10.3389/fphys.2018.00900.

Boccellino, M., Di Domenico, M., Donniacuo, M., Bitti, G., Gritti, G., Ambrosio, P., et al. 2018. AT1-receptor blockade:
Protective effects of irbesartan in cardiomyocytes under hypoxic stress. *PLoS One* 13 （10）, e0202297. doi:
10.1371/journal.pone.0202297.

Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., et al. 2019. Reproducible,
interactive, scalable and extensible microbiome data science using QIIME 2. *Nat Biotechnol* 37 （8）, 852-857. doi:
10.1038/s41587-019-0209-9.

Brestoff, J. R., and Artis, D. 2013. Commensal bacteria at the interface of host metabolism and the immune system.
Nature immunology 14 （7）, 676-684. doi: 10.1038/ni.2640.

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J., and Holmes, S. P. 2016. DADA2: High-
resolution sample inference from Illumina amplicon data. *Nat Methods* 13 （7）, 581-3. doi: 10.1038/nmeth.3869.

Dao, M. C., Everard, A., Aron-Wisnewsky, J., Sokolovska, N., Prifti, E., Verger, E. O., et al. 2016. Akkermansia
muciniphila and improved metabolic health during a dietary intervention in obesity: relationship with gut microbiome
richness and ecology. *Gut* 65 （3）, 426-36. doi: 10.1136/gutjnl-2014-308778.

David, L. A., Maurice, C. F., Carmody, R. N., Gootenberg, D. B., Button, J. E., Wolfe, B. E., et al. 2014. Diet rapidly and
reproducibly alters the human gut microbiome. *Nature* 505 （7484）, 559-63. doi: 10.1038/nature12820.

Dunn, W. B., Broadhurst, D., Begley, P., Zelena, E., Francis-McIntyre, S., Anderson, N., et al. 2011. Procedures for
large-scale metabolic profiling of serum and plasma using gas chromatography and liquid chromatography coupled to
mass spectrometry. *Nature protocols* 6 （7）, 1060-1083. doi: 10.1038/nprot.2011.335.

El-Sayed Moustafa, J. S., and Froguel, P. 2013. From obesity genetics to the future of personalized obesity therapy. *Nat*
Rev Endocrinol 9 （7）, 402-13. doi: 10.1038/nrendo.2013.57.

Everard, A., Belzer, C., Geurts, L., Ouwerkerk, J. P., Druart, C., Bindels, L. B., et al. 2013. Cross-talk between
Akkermansia muciniphila and intestinal epithelium controls diet-induced obesity. *Proc Natl Acad Sci U S A* 110 （22）,
9066-71. doi: 10.1073/pnas.1219451110.

Foligné, B., Parayre, S., Cheddani, R., Famelart, M. H., Madec, M. N., Plé, C., et al. 2016. Immunomodulation properties
of multi-species fermented milks. *Food Microbiol* 53 （Pt A）, 60-9. doi: 10.1016/j.fm.2015.04.002.

Gao, K., Dou, Y., Lv, M., Zhu, Y., Hu, S., and Ma, P. 2021. Research hotspots and trends of microRNA in
periodontology and dental implantology: a bibliometric analysis. *Ann Transl Med* 9（14）, 1122. doi: 10.21037/atm-21-
726.

Halmos, T., and Suba, I. 2016. [Physiological patterns of intestinal microbiota. The role of dysbacteriosis in obesity,
insulin resistance, diabetes and metabolic syndrome]. *Orv Hetil* 157（1）, 13-22. doi: 10.1556/650.2015.30296.

Hibberd, M. C., Wu, M., Rodionov, D. A., Li, X., Cheng, J., Griffin, N. W., et al. 2017. The effects of micronutrient
deficiencies on bacterial species from the human gut microbiota. *Science translational medicine* 9（390）, eaal4069. doi:
10.1126/scitranslmed.aal4069.

Hussain, S. S., and Bloom, S. R. 2013. The regulation of food intake by the gut-brain axis: implications for obesity. *Int J*
Obes (Lond) 37（5）, 625-33. doi: 10.1038/ijo.2012.93.

Husted, A. S., Trauelsen, M., Rudenko, O., Hjorth, S. A., and Schwartz, T. W. 2017. GPCR-mediated signaling of
metabolites. *Cell metabolism* 25（4）, 777-796. doi: 10.1016/j.cmet.2017.03.008.

Jandhyala, S. M., Talukdar, R., Subramanyam, C., Vuyyuru, H., Sasikala, M., and Nageshwar Reddy, D. 2015. Role of
the normal gut microbiota. *World J Gastroenterol* 21（29）, 8787-803. doi: 10.3748/wjg.v21.i29.8787.

Kelly, T., Yang, W., Chen, C. S., Reynolds, K., and He, J. 2008. Global burden of obesity in 2005 and projections to
2030. *Int J Obes (Lond)* 32（9）, 1431-7. doi: 10.1038/ijo.2008.102.

Koh, A., De Vadder, F., Kovatcheva-Datchary, P., and Bäckhed, F. 2016. From Dietary Fiber to Host Physiology: Short-
Chain Fatty Acids as Key Bacterial Metabolites. *Cell* 165（6）, 1332-1345. doi: 10.1016/j.cell.2016.05.041.

Krishnan, S., Ding, Y., Saedi, N., Choi, M., Sridharan, G. V., Sherr, D. H., et al. 2018. Gut Microbiota-Derived
Tryptophan Metabolites Modulate Inflammatory Response in Hepatocytes and Macrophages. *Cell Rep* 23（4）, 1099-
1111. doi: 10.1016/j.celrep.2018.03.109.

Le Chatelier, E., Nielsen, T., Qin, J., Prifti, E., Hildebrand, F., Falony, G., et al. 2013. Richness of human gut microbiome
correlates with metabolic markers. *Nature* 500（7464）, 541-6. doi: 10.1038/nature12506.

Liu, R., Hong, J., Xu, X., Feng, Q., Zhang, D., Gu, Y., et al. 2017. Gut microbiome and serum metabolome alterations in
obesity and after weight-loss intervention. *Nat Med* 23（7）, 859-868. doi: 10.1038/nm.4358.

Liu, Z., Gao, K., Hai, Y., and Liu, T. 2022. Developments, Focuses, and Trends in Early-Onset Scoliosis From 2005 to
2020: A Systematic Bibliometric Analysis. *World Neurosurg* 158, e697-e710. doi: 10.1016/j.wneu.2021.11.044.

Mach, N., and Clark, A. 2017. Micronutrient deficiencies and the human gut microbiota. *Trends in microbiology* 25（8），
607-610. doi: 10.1016/j.tim.2017.06.004.

Mohan, I., Gupta, R., Misra, A., Sharma, K. K., Agrawal, A., Vikram, N. K., et al. 2016. Disparities in Prevalence of
Cardiometablic Risk Factors in Rural, Urban-Poor, and Urban-Middle Class Women in India. *PLoS One* 11（2），
e0149437. doi: 10.1371/journal.pone.0149437.

Nazli, A., Yang, P. C., Jury, J., Howe, K., Watson, J. L., Söderholm, J. D., et al. 2004. Epithelia under metabolic stress
perceive commensal bacteria as a threat. *Am J Pathol* 164（3）, 947-57. doi: 10.1016/s0002-9440(10)63182-3.

Neyrinck, A. M., Van Hée, V. F., Piront, N., De Backer, F., Toussaint, O., Cani, P. D., et al. 2012. Wheat-derived
arabinoxylan oligosaccharides with prebiotic effect increase satietogenic gut peptides and reduce metabolic endotoxemia
in diet-induced obese mice. *Nutr Diabetes* 2（1）, e28. doi: 10.1038/nutd.2011.24.

Paganelli, F. L., Luyer, M., Hazelbag, C. M., Uh, H. W., Rogers, M. R. C., Adriaans, D., et al. 2019. Roux-Y Gastric
Bypass and Sleeve Gastrectomy directly change gut microbiota composition independent of surgery type. *Sci Rep* 9（1），
10979. doi: 10.1038/s41598-019-47332-z.

Plovier, H., Everard, A., Druart, C., Depommier, C., Van Hul, M., Geurts, L., et al. 2017. A purified membrane protein
from Akkermansia muciniphila or the pasteurized bacterium improves metabolism in obese and diabetic mice. *Nat Med*
23（1）, 107-113. doi: 10.1038/nm.4236.

Putignani, L., Del Chierico, F., Vernocchi, P., Cicala, M., Cucchiara, S., Dallapiccola, B., et al. 2016. Gut microbiota
dysbiosis as risk and premorbid factors of IBD and IBS along the childhood–adulthood transition. *Inflammatory bowel*
diseases 22（2）, 487-504. doi: 10.1097/MIB.0000000000000602.

Qin, J., Li, Y., Cai, Z., Li, S., Zhu, J., Zhang, F., et al. 2012. A metagenome-wide association study of gut microbiota in
type 2 diabetes. *Nature* 490（7418）, 55-60. doi: 10.1038/nature11450.

Rastelli, M., Van Hul, M., Terrasi, R., Lefort, C., Régnier, M., Beiroa, D., et al. 2020. Intestinal NAPE-PLD contributes
to short-term regulation of food intake via gut-to-brain axis. *Am J Physiol Endocrinol Metab* 319（3）, E647-e657. doi:
10.1152/ajpendo.00146.2020.

Ridaura, V. K., Faith, J. J., Rey, F. E., Cheng, J., Duncan, A. E., Kau, A. L., et al. 2013. Gut microbiota from twins
discordant for obesity modulate metabolism in mice. *Science* 341（6150）, 1241214. doi: 10.1126/science.1241214.

Rodríguez, J. M., Murphy, K., Stanton, C., Ross, R. P., Kober, O. I., Juge, N., et al. 2015. The composition of the gut
microbiota throughout life, with an emphasis on early life. *Microb Ecol Health Dis* 26, 26050. doi:
10.3402/mehd.v26.26050.

Sabate, J.-M., Coupaye, M., Ledoux, S., Castel, B., Msika, S., Coffin, B., et al. 2017. Consequences of small intestinal
bacterial overgrowth in obese patients before and after bariatric surgery. *Obesity surgery* 27, 599-605. doi:

Scherer, P. E., and Hill, J. A. 2016. Obesity, Diabetes, and Cardiovascular Diseases: A Compendium. *Circ Res* 118（11）,
1703-5. doi: 10.1161/circresaha.116.308999.

Schuster, S. C. 2008. Next-generation sequencing transforms today's biology. *Nat Methods* 5（1）, 16-8. doi:
10.1038/nmeth1156.

Sidhu, M., and van der Poorten, D. 2017. The gut microbiome. *Australian family physician* 46（4）, 206-211. doi:
10.3316/informit.741870194405168.

Tang, W. H., Kitai, T., and Hazen, S. L. 2017. Gut Microbiota in Cardiovascular Health and Disease. *Circ Res* 120（7）,
1183-1196. doi: 10.1161/circresaha.117.309715.

Tian, P., Li, B., He, C., Song, W., Hou, A., Tian, S., et al. 2016. Antidiabetic (type 2) effects of Lactobacillus G15 and
Q14 in rats through regulation of intestinal permeability and microbiota. *Food Funct* 7（9）, 3789-3797. doi:
10.1039/c6fo00831c.

Tian, Y., Gui, W., Koo, I., Smith, P. B., Allman, E. L., Nichols, R. G., et al. 2020. The microbiome modulating activity of
bile acids. *Gut microbes* 11（4）, 979-996. doi: 10.1080/19490976.2020.1732268.

Tsai, Y. L., Lin, T. L., Chang, C. J., Wu, T. R., Lai, W. F., Lu, C. C., et al. 2019. Probiotics, prebiotics and amelioration
of diseases. *J Biomed Sci* 26（1）, 3. doi: 10.1186/s12929-018-0493-6.

Wang, K., Liao, M., Zhou, N., Bao, L., Ma, K., Zheng, Z., et al. 2019. Parabacteroides distasonis Alleviates Obesity and
Metabolic Dysfunctions via Production of Succinate and Secondary Bile Acids. *Cell Rep* 26（1）, 222-235.e5. doi:
10.1016/j.celrep.2018.12.028.

Wang, Z., Ma, D., Pang, R., Xie, F., Zhang, J., and Sun, D. 2020. Research progress and development trend of social
media big data (smbd): Knowledge mapping analysis based on citespace. *ISPRS International Journal of Geo-*
Information 9（11）, 632. doi: 10.3390/ijgi9110632.

FIGURE 1 Global publication trend and country or region analysis for the effect of intestinal microbes on obesity research. (A) Annual worldwide publication output. (B) Growth trends in publication output from the top 10 most productive countries. (C) Top 10 publishers according to their contribution to the total number of articles on the effect of intestinal microbes on obesity research.

FIGURE 2 CiteSpace network visualization map of countries/regions and institutions. (A) CiteSpace network visualization map of countries/regions involved in the effect of intestinal microbes on obesity research (node label: by citation, label font size: proportional). (B) CiteSpace network visualization map of institutions involved in the effect of intestinal microbes on obesity research (node label: by citation, label font size: proportional).

FIGURE 3 CiteSpace network visualization map of authors and co-cited authors. (A) CiteSpace network visualization map of authors involved in the effect of intestinal microbes on obesity research (node label: by citation, label font size: proportional). (B) CiteSpace network visualization map of co-cited authors involved in the effect of intestinal microbes on obesity research (node label: by centrality, label font size: proportional).

FIGURE 4 CiteSpace network visualization map of co-cited journals involved in the effect of intestinal microbes on
obesity research (node label: by centrality, label font size: proportional).

FIGURE 5 Keyword co-occurrence network for the effect of intestinal microbes on obesity research (node label: by
centrality, label font size: uniformed).

FIGURE 6 Keyword clusters for the effect of intestinal microbes on obesity research.

TABLE 1 Top 10 productive countries/regions related to the effect of intestinal microbes on obesity.

Rank	Countries	Articles	Centrality	Percentage(N/8888)
1	China	3036	0.02	34.16%
2	the United States	2356	0.25	26.51%
3	Spain	480	0.14	5.40%
4	Canada	466	0.07	5.24%
5	France	432	0.11	4.86%
6	Japan	406	0.02	4.57%
7	Germany	382	0.12	4.30%
8	Italy	372	0.12	4.19%
9	South Korea	364	0.05	4.10%
10	England	355	0.23	3.99%

597 TABLE 2 Top 10 productive institutions related to the effect of intestinal microbes on obesity.

Rank	Institutions	Articles	Centrality	Country
1	University of Copenhagen	209	0.28	Denmark

2	Chinese academic of sciences	202	0.18	China
3	Shanghai Jiao Tong university	174	0.09	China
4	China Agricultural University	129	0.04	China
5	Catholic University of Louvain	114	0.08	Belgium
6	Zhejiang University	107	0.14	China
7	Inst Salud Carlos III	94	0.10	Spain
8	University of California, Davis	93	0.11	the United States
9	University of Illinois	91	0.09	the United States
10	University of Chinese academic of sciences	90	0.02	China

599 TABLE 3 Top 10 productive authors related to the effect of intestinal microbes on obesity.

Rank	Name	Articles	Centrality	Country
1	Patrice D. Cani	97	0.03	Belgium
2	Nathalie M. Delzenne	73	0.03	Belgium
3	Reimer, Raylene A	51	0.00	Canada
4	Backhed, Fredrik	35	0.06	Sweden
5	Neyrinck, Audrey M	34	0.01	Belgium
6	Wang, Jing	33	0.00	China
7	Liu, Bin	32	0.00	China
8	Marette, Andre	28	0.01	Canada
9	Chen, Wei	28	0.01	China
10	Clement, Karine	27	0.01	France

601 TABLE 4 Top 10 co-cited authors related to the effect of intestinal microbes on obesity.

Rank	Cited authors	Articles	Centrality	Country
1	Peter J. Turnbaugh	3307	0.33	the United States
2	Patrice D. Cani	2879	0.27	Belgium
3	Ruth E. Ley	2508	0.11	the United States
4	Backhed, Fredrik	1660	0.09	Sweden
5	J. Gregory Caporaso	1448	0.1	the United States
6	Amandine Everard	1370	0.08	Belgium
7	Junjie Qin	1250	0.12	China
8	Robert C Edgar	935	0.01	the United States
9	Nicola Segata	806	0.01	Italy
10	Emmanuelle Le Chatelier	751	0.01	France

604   TABLE 5 Top 10 productive journals related to the effect of intestinal microbes on obesity.

Sources	Articles	Percentage(N/8888)	IF
Nutrients	399	4.49%	6.706
Scientific Reports	339	3.81%	4.997
Food & Function	281	3.16%	6.317
PLOS ONE	265	2.98%	3.752
Frontiers in microbiology	226	2.54%	6.064
Journal of functional foods	171	1.92%	5.223
Molecular nutrition & food research	136	1.53%	6.575
Frontiers in nutrition	116	1.31%	6.590
Journal of agricultural and food chemistry	114	1.28%	5.895
International journal of molecular sciences	108	1.22%	6.208

606   TABLE 6 Top 10 co-cited journals related to the effect of intestinal microbes on obesity.

Rank	Co-cited journals	Articles	Centrality	IF
1	Nature	6446	0.18	69.504
2	PLOS ONE	6247	0.17	3.752
3	Proceedings of the national academy of sciences (PNAS)	5329	0.02	12.779
4	Gut	4389	0.01	31.795
5	Science	4180	0.01	63.832
6	Diabetes	4032	0.02	9.305
7	Scientific reports	3979	0.01	4.997
8	Cell	3308	0.01	66.850
9	Nutrients	3250	0.05	6.706
10	Gastroenterology	3114	0.04	33.883

608   TABLE 7 Top 10 keywords related to the effect of intestinal microbes on obesity.

Rank	Keywords	Articles	Centrality
1	gut microbiota	4673	0.39
2	obesity	3587	0.23
3	inflammation	1578	0.04
4	insulin resistance	1368	0.12
5	intestinal microbiota	1038	0.01
6	health	869	0.02
7	metabolism	862	0.02

8	diet	848	0.01
9	gut microbiome	825	0.01
10	metabolic syndrome	714	0.01

610 TABLE 8 The top 10 co-cited references related to the effect of intestinal microbes on obesity.

Rank	Title	Author	Article type	Journal	DOI	Articles
1	Diet rapidly and reproducibly alters the human gut microbiome	Lawrence A. David et al., 2013	Letter	Nature	10.1038/nature12820	383
2	Cross-talk between *Akkermansia muciniphila* and intestinal epithelium controls diet-induced obesity	Amandine Everard et al.,2013	Article	PNAS	10.1073/pnas.1219451110	367
3	*Akkermansia muciniphila* and improved metabolic health during a dietary intervention in obesity-relationship with gut microbiome richness and ecology	Maria Carlota Dao et al., 2016	Article	Gut	10.1136/gutjnl-2014-308778	303
4	A metagenome-wide association study of gut microbiota in type 2 diabetes	Junjie Qin et al., 2012	Article	Nature	10.1038/nature11450	291
5	Richness of human gut microbiome correlates with metabolic markers	Emmanuelle Le Chatelier et al., 2013	Article	Nature	10.1038/nature12506	285
6	Gut microbiota from twins discordant for obesity modulate metabolism in mice	Vanessa K. Ridaura et al., 2013	Article	Science	10.1126/science.1241214	276
7	A purified membrane protein from *Akkermansia muciniphila* or the pasteurized bacterium improves metabolism in obese and diabetic mice	Hubert Plovier et al., 2017	Letters	Nature Medicine	10.1038/nm.4236	275
8	DADA2: High-resolution sample inference from Illumina amplicon data	Benjamin J Callahan et al., 2016	Brief communications	Nature methods	10.1038/nmeth.3869	273
9	Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2	Evan Bolyen et al., 2019	Correspondence	Nature biotechnology	10.1038/s41587-019-0209-9	265
10	Gut microbiome and serum metabolome alterations in obesity and after weight-loss intervention	Ruixin Liu et al., 2017	Article	Nature medicine	10.1038/nm.4358	264