Title: Determining prescriptions in electronic health care (EHR) data: methods for development of

standardised, reproducible drug codelists

Authors: Emily L Graul[1], Philip W Stone[1,2], Georgie M Massen[2], Sara Hatam[3], Alexander

Adamson[1,2], Spiros Denaxas[4,5], Nicholas S Peters[2], Jennifer K Quint[1,2]

Addresses:

1. School of Public Health, Imperial College London, London, UK

2. National Heart & Lung Institute, Imperial College London, London, UK

3. Usher Institute, University of Edinburgh, Edinburgh, UK

4. University College London, Institute of Health Informatics, London, UK

5. British Heart Foundation Data Science Centre, Health Data Research UK, London, UK

Corresponding Author: Emily L Graul

School of Public Health, Imperial College London – White City Campus

86 Wood Lane

London, UK W12 0BZ

Phone: 02075948821

Email: e.graul@imperial.ac.uk

Key words: Codelist, Prescriptions, Value Sets, Electronic Health Records, epidemiology

Word Count: 3919/4000

ABSTRACT

Objective

Epidemiological research using electronic healthcare records(EHR) uses combinations of codes to define diseases and prescriptions (or phenotypes), requiring transparency and reproducibility. Yet methodology to generate codelists varies, manifesting in misclassification bias. Therefore we designed methodology enabling codelist reproducibility and generalisability across study contexts.

Materials and Methods

We developed a process to generate drug codelists, testing this using the Clinical Practice Research Datalink (CPRD) Aurum database, accounting for missing data in 'attribute' variables searched to identify codes. We generated a 1) cardiovascular codelist and 2) codelist for inhaled Chronic Obstructive Pulmonary Disease (COPD) therapies, applying them to a sample cohort of 335,931 COPD patients. We compared searching on all search variables (A, "gold standard") to B) chemical and C) ontological information only.

Results

In our full search (A), within follow-up, we determined 165,150 patients (49.2% of cohort) prescribed drugs from the cardiovascular codelist. For the COPD inhalers codelist, we determined 317,963 patients (94.7% of cohort) prescribed. Considering output within each individual value set, Search C missed substantial prescriptions, including vasodilator anti-hypertensives (19,696 prescriptions for A and B; 1,145 for C), and for SAMA (35,310 for A and B; 564 for C).

Discussion

Regardless of database and study context, we recommend the full method (A) for comprehensiveness. Despite database used, there are special considerations when generating adaptable drug codelists, including fluctuating status, cohort-specific drug indications, underlying hierarchical ontology, and collinearity in covariate analyses.

Conclusions

Generating drug codelists must use standardisable and reproducible methodology based on underlying ontology, with end-to-end clinical input.

BACKGROUND AND SIGNIFICANCE

Health data research & codelist generation

Research using electronic health records (EHR) is increasingly used to inform patient care across a breadth of longitudinal data sources, including the U.S. Veterans EHR System, the INSIGHT Clinical Research Network (CRN) database, the National Patient-Centered Clinical Research Network (PCORnet), the National COVID Cohort Collaborative (N3C) Data Enclave, Duke Health Data Warehouse, the SAIL Databank, the Clinical Practice Research Datalink (CPRD), NHSDigital and Northwest London Whole Systems Integrated Care (WSIC), with some allowing linkage to mortality, socioeconomic, registry and audit data[1–11].

Determining exposures, outcomes and covariates is central to EHR research[12,13] through generation of medical and drug "codelists" for an overarching clinical definition. Unfortunately, both methodology and reporting vary, forming sources of misclassification bias when ascertaining conditions and prescriptions. Methodology may not involve clinician review, manifesting in exclusion of necessary codes alongside inclusion of inappropriate codes. The Reporting of studies Conducted using Observational Routinely-collected health Data (RECORD) statement calls for EHR studies to provide "a complete list of codes and algorithms used to classify exposures, outcomes, confounders, and effect modifiers…considering the risk of misclassification bias…authors should provide sufficient detail to make their research *reproducible* and to make the *risk of bias* apparent" (emphasis added)[14].

Calls regarding transparency and bias extend beyond making codelists freely accessible in repositories. It also requires a systematic and standardisable yet malleable approach incorporating clinical input, designed around proactively considering when bias can manifest upon subsequent application to cohorts. Malleability within codelist design can make reproducibility and generalisation to other contexts and databases possible, allowing consistent definition (harmonisation) of phenotypes[15]. Malleability facilitates researchers to adapt and reuse others' codelists for their study and, conversely, facilitates researchers to contribute their codelists to others' studies appropriately[16]. Literature on drug codelists has been primarily high-level[17,18], underlying steps for its generation of the codelists themselves not usually described.

Aim

Given unique considerations for drug codelists, we developed a standardisable, reproducible method

for creating drug codelists that incorporates end-to-end clinical expertise, considers missing data in

search 'attribute' variables, considers fluctuating status, and is adaptable to other studies and

databases. We then operationalise the methodology to generate a codelist for oral drugs for

hypertension and heart failure, and a disease-specific codelist, all inhaled therapies for COPD. We

also apply the codelists to a sample cohort, according to study considerations and clinical review.

MATERIALS AND METHODS

Defining phenotypes, value sets, and ontologies

Common to both medical and drug codelist generation is establishing the single overarching clinical

definition (analogous to a "phenotype" for medical codelists), premised by a database's underlying

drug hierarchical ontology (e.g., British National Formulary, BNF, Anatomic Classification System,

ATC, and RxNorm)[19–21]. Within this definition are "value sets", a "uniquely identifiable set of valid

concept representations" for "possible values of a coded data element in an information model"[22].

Depending on purpose and study context there may be one or multiple value sets. Ideally, searches

to identify all possible codes are conducted purely using the drugs' chemical terms or the database's

ontology, but missing data can prevent this.

Our methodology

Our methodology for generating drug codelists has seven steps (summarised in Figure 1) and is

available on our Github repository. It is based on our methodology for generating medical codelists,

also available in our Github repository.

Figure 1. Summary of methodology for generating a product codelist
Note: Steps in grey font may be database-dependent and modified as necessary given missing data.

Step 1: Defining purpose and value sets

The first stage is to identify value sets consisting of drug class(es) based on an underlying ontology

and within classes a list of drugs with corresponding chemical and potentially multiple proprietary

names. The route(s) of administration (e.g., oral, parenteral, ocular) should be specified. The aim may

be to produce a broader codelist permitting modification for various contexts, or to generate a study-

specific codelist.

We consider clinical input regarding chemistry to improve search efficiency. For example, searching

on common compounds, active or blocking groups, or side chains such as *-nitrate -arginine -*

hydrochloride -mesilate is not recommended. Whilst these suffixes may be listed as part of the drug

name, they are not the chemical-of-interest and may lead to inefficiently large search outputs.

Premising step 1 is confirming the database's underlying ontology to systematically collate a list of all

possible terms (i.e., the BNF for UK databases) and reliable resource to facilitate reproducibility of

collation. The most user-friendly BNF resource is OpenPrescribing [23], a interface utilizing raw data

from UK National Health Service Business Services Authority (NHSBSA), with an ability to easily

extract corresponding proprietary names and identify discontinued drugs.

Step 2: Conducting search

The second step is to search based on the collated list, using wildcard (*) characters to pick up terms

in any location within a string.

Step 2a: Search the database drug dictionary

Before the search, we import the database's drug 'dictionary' text file. We import all 'attribute'

variables searched upon as strings.

We search the browser in 2 stages:

(i) For each drug within value sets, search by the chemical and all proprietary names, for a string

match on the associated 'attribute' variables.

(ii)Search on underlying ontology, considering syntax with slashes(/) as medicines may be

indicated for multiple conditions and located in chapters not primarily of interest but still used for

the indication-of-focus (e.g., searching for sildenafil (parenteral route) may fall into section

ontologies for both hypertension and erectile dysfunction).

This automated search nests chemical and proprietary terms within each drug list, with corresponding

lists nested within broader value sets (Figure 2), in effect sorting output for (i) by value set.

149
Figure 2. Flow diagram showing the Step 2 search process for drug codes.
A main clinical definition for the drug codelist is established, premised by the underlying database ontology.
Within each given value set ('parent' list) are nested 'child' lists each corresponding to individual drugs, chemical
and proprietary names. The CPRD-specific search attributes are *termfromemis* and *productname* (containing
chemical and proprietary information) and *drugsubstancename* (chemical information). Due to missing data within
the search 'attribute' variables we search on all three variables in Step 2a(i), with an additional ontological code
search in Step 2a(ii), checking for search term completeness of the former by comparing the (i) and (ii) in Step 2b
iteratively. Therefore, Step 2a(ii) and 2b may be database dependent given missing data. In Stata, parent and
child lists take the form of local macros; in R a comparable step would be to name a list of vectors, and nesting
the lists as necessary. The full Stata and R code including all drug codelist generation steps is located on our
[Github repository](#).
161
162

Step 2b: Tag codes additionally identified by searching on (ii) underlying ontology; Repeat

2a-2b iteratively

We tag outstanding codes not found by searching on chemical and proprietary terms alone, allowing

researchers to check if they included all possible terms. Outstanding codes are identified and tagged

if the row total for columns with "1" denoting searches (i) and (ii) within 2a sum to least two. If

outstanding codes are present, one should add the additional names to the value sets, re-running

steps 2a to 2b. Ideally, upon multiple iterations of re-running, there should be an absence of tags,

indicating inclusion of all appropriate terms. This additional step may seem redundant but is most

important to check codelist completeness.

In place of this step we incorporated an expanded search using Systematized Nomenclature of

Medicine-Clinical Terms (SNOMED-CT, a commonly-used international reference terminology)

concept IDs to check for outsanding synonym codes fitting our value sets, as recommended for

medical codelists[24]. However, in CPRD, although for clinical events, a given SNOMED-CT Concept

ID will match with multiple SNOMED-CT description IDs, for prescription events, a given SNOMED-

CT Concept ID does not match with more than one SNOMED-CT Description ID. This is because the

UK SNOMED-CT drug extension is derived from the Dictionary of Medicines and Devices (dm+d),

with the SNOMED-CT Concept ID being identical to the dm_d code, and CPRD only utilises dm+d

codes, preventing this approach.

Step 3: Exclusions

Step three consists of, after manual review, excluding codes. This is distinguished from later

exclusion based on epidemiological and clinical considerations specific to study context. Elimination

may be based on information from drug name, route, and/or formulation. The broad search may pick

up different medications with the same active chemical but of an inappropriate route, i.e., for a

different medical indication corresponding to a different organ system.

We do not eliminate by product identifiers; it is a less transparent method. We do not eliminate by

ontology chapter either, not only due to missing data, but anecdotal evidence suggests some drugs

may have intended medical indication(s) corresponding to multiple chapters, which cannot be

assumed and is not reflected in its ontological classification code.

Step 4: Cleaning

Step 4a: Eliminate mutually exclusive codes across value sets

This step places a temporary tag on codes mutually exclusive across value sets, a possibility given

the broad search, e.g., even within a single BNF class chapter, active ingredients may overlap among

the individual sections.

This tag allows researchers to write code automating the re-sorting process to make these sets

exclusive.

Step 4b. Tagging mutually exclusive codes across value sets, with intentional overlap, for

clinician and/or epidemiologist

This step consists of proactively placing permanent tags on codes corresponding to fixed combination

drugs with 'intentional overlap.'

We define 'intentional overlap' when one code corresponds to a fixed combination drug consisting of

two drug classes (i.e., mechanisms of action) such that it resides in multiple ontological sections, e.g.,

hydrochlorothiazide/captopril, a single drug including both diuretic and Renin-angiotensin system

components.

Clinician input is considered to provide a list of possible suffixes for the tags, e.g., "*azide*" for

diuretics, or "*pril*" for angiotensin-converting enzyme (ACE) inhibitors and angiotensin receptor

blockers (ARBs).

Step 4c. Modifying value sets as necessary

This step allows for value set modification. For example, combining sets into a broader value set upon

study context or computational considerations (e.g., Stata has macro character limits).

Step 5. Comparing to previous lists or mapping ontologies

This step is for merging together and comparing current and previous codelist versions, and to merge

and map codes labelled under different ontologies(e.g., ATC-BNF mapping). Comparison facilitates

correct categorization and possible identification of outstanding codes. Beyond completeness,

mapping allows harmonisation and reproducibility to other database contexts[15].

This leads to the 'raw' codelist that is not study-specific, and ready for adaptation to a cohort through

clinical review.

Step 6. Sending 'raw' codelist for clinical review to generate study-specific codelist

Clinician(s) review the 'raw' codelist and each code (observation) is labelled as the following

'certainty' categories:

0 = "clear exclusion"

1 = "certainty" to be included

2 = "uncertainty" if to include for future sensitivity analyses

Review by one clinician trained in epidemiology and familiar with using the database is necessary, *at*

least. When placing certainty categories, clinicians may consider:

–   *cohort-of-interest* specific to their expertise/organ system. If multiple systems are involved,

243   i.e., multimorbidity, multiple clinicians are necessary

244   -   *operationalisation of codes in clinical settings* and *patient behaviour or prescription*

245   *commonality* within context

246   -   *database characteristics* (e.g., drug with substantially low number of issues may have low

247   frequency of prescription events when later applied to cohort)

An additional step of resolving discordances among 'certainty' categories is required given multiple

clinicians. Step 6 adapts a publication's methods on generating medical codelists[12] after which the

codelist is finalised for study context.

Step 7. Keeping a 'master' codelist

Researchers should keep a 'master' codelist with all versions and tags for reasons pertaining to

malleability: to allow adaptation for sensitivity analyses, and to allow generalisation and harmonisation

to other and across study contexts.

It should contain:

(i) raw codes – all codes not study-specific, initially generated by epidemiologist (broad and all-

encompassing) sorted by ontology

(ii) clinical input – columns per clinician (≥1 column with 0/1/2s)

(iii) study-specific codes – column for tailoring to context (1s 'certainty' only)

(iv) tags for individual codes for fixed combination drugs also classified within different ontology

chapters

Operationalising our methodology

The methodology was applied to generate two codelists, 1) drugs for hypertension and heart failure (BNF Chapter 2.5) and 2) all inhaled therapies for COPD (BNF Ch. 3.1.1-.2; 3.1.4; 3.2) in CPRD Aurum[23].

We estimated the number of respective prescriptions among a cohort with COPD within the Clinical Practice Research Datalink (CPRD) Aurum in England. Many individuals with COPD require inhaled therapies for reduced lung function and have co-morbidities, including cardiovascular disease[25]. In this open cohort, patients were included and started follow up at latest of 1 January 2010 if they 1) were diagnosed with validated codes for COPD[26], 2) were at least 40 years, 3) had continuous 1 year GP registration, and 4) data was deemed 'acceptable' quality. Follow-up ended on the earliest of: 31 December 2019, the last collection date, date of patient death, or transfer out of GP practice.

We compared output of the following searches, for the codelist and upon application to the cohort:

    A. Using our full 'gold-standard' methodology, searching on chemical and proprietary terms and BNF codes (chemical and proprietary names on *termfromemis*, *productname* variables, chemical names only for *drugsubstancename* due the nature of this variable, BNF codes for *bnfchapter*)

    B. Using our methodology, but searching on chemical names only (within *drugsubstancename*)

    C. Using our methodology, but searching on the BNF code only (within CPRD's *bnfchapter*)

Outcome measures were product codes, drug issues, and prescriptions in overall and by value set. Analyses were conducted using Stata v17 (Texas, USA). We wrote final scripts in Stata, translating into R v4.2.0. A summary of each codelist's purpose and operationalisation is in Supplementary Table S1. R and Stata scripts, as well as a full list of our terms for the value sets, are in located in our [Github repository](#).

RESULTS

Generating raw codelists

We operationalised the methodology in CPRD Aurum to generate two codelists, a cardiovascular codelist for hypertension and heart failure medication and a codelist for inhaled COPD therapies.

We designed value sets around the codelist purpose (e.g., repository or disease-focused). We nested

terms corresponding to each drug within 'child' lists, then nesting each drug list into 'parent'

lists(Figure 2). For example, for the indoramin child list (cardiovascular codelist), we searched for

chemical and proprietary terms, nesting this list within the BNF chapter 2.5.4 value set). In the COPD

inhalers codelist, because intentions were to distinguish disease-specific drugs, we separated value

by type even though chemical compositions overlapped, e.g., inhaled corticosteroids (ICS) vs. inhaled

corticosteroid-long-acting muscarinic antagonists (ICS-LAMA).

In the CPRD Aurum drug dictionary, a given unique identifier ('*prodcodeid*' variable) can include

missing data on its following 'attributes' including active chemical ingredients ('*drugsubstancename*'),

ontology ('*bnfchapter*'), and route ('*routeofadministration*'). (A description of missing data in the CPRD

drug dictionary is in Supplementary Table S2.) Ideally, searches to identify all possible codes would

be conducted purely using chemical terms or the database's ontology, but missing data prevented

this, while the most-complete *termfromemis* variable lists drugs by *either* chemical or proprietary

name. Therefore, in in Step 2a(i) we searched on multiple 'attribute' variables: *termfromemis*,

productname and *drugsubstancename* (Figure 2).

Prior to producing raw codelists, we excluded 26 codes from the cardiovascular codelist, and 206

codes from the COPD inhalers codelist. This was due to cases where composition was correct but

route was incorrect given indication (e.g., *cutaneous* minodoxil, *ocular* guanethidine monosulfate,

salbutamol *nebuliser solutions*), our string match inadvertently picked up codes for a different purpose

or distinct chemical compound (e.g., Glutenex prescription from searching "*tenex*"; apra-clonidine

from searching on "*clonidine*"), or the term was not part of value sets (e.g., ICS-salbutamol codes).

We tagged codes for fixed combination drugs also classified within other BNF chapters. For the

cardiovascular codelist, this comprised codes for Ch. 2.2 diuretics and 2.6 antianginal drugs, e.g.,

Lisinopril-Hydrochlorothiazide (diuretic and angiotensin-converting enzyme (ACE) inhibitor). For the

COPD inhalers codelist this comprised codes for Ch. 3.3, for Salbutamol-sodium cromoglycate.

After respective exclusions and tags, the raw cardiovascular codelist contained 601 codes of both oral

and parenteral routes, and for inhaled COPD therapies, 259 codes. For the COPD inhalers codelist,

after subsequently merging with a previous codelist mapped to NHS Business Services Authority

TRUD ATC-BNF ontology mapping files, this led to a final count of 472 codes. Of these codes, 77

were new codes not in the previous codelist, and 13 outstanding codes were from the previous

codelist not in the new codelist. Most new codes were ICS- or short acting beta-agonist (SABA)-

based (Supplementary Table S5).

Clinical review

The first clinician, a respiratory consultant and epidemiologist, removed codes for drugs not part of

value sets: for the cardiovascular codelist, nine for Selexipag and one for Sodium Nitroprusside from

searching on BNF ontology(Step 2a,ii). There was concern such prescriptions were rare given few

issues and it being less likely these infrequently-used, new, or discontinued drugs were prescribed in

the COPD cohort. For the COPD inhalers codelist, we removed three codes (0s) consisting of an

ambiguous term for route – "liquid" or "solution"– potentially corresponding to nebulised therapies. We

retained and tagged one code for paediatrics.

For the cardiovascular codelist, 33 codes of the parenteral route were given 0s as this route is not

typically prescribed in UK primary care, leaving oral medications. A second clinician, a cardiologist,

reviewed the 1s, agreeing on all codes.

Final codelists

The final cardiovascular codelist consisted of 568 codes for oral medications (Figure 3;Supplmentary

TableS3), including the 66 and 28 product codes tagged for overlap with Chapter 2.2 and 2.6. The

value set with the greatest count was for drugs targeting the Renin-aldosterone system (Ch. 2.5.5,

N=375), whereas 2.5.3 and 2.5.8 were the smallest (N=4, N=2 respectively).

The final COPD inhalers codelist consisted of 456 codes(Figure 3;Supplmentary TableS3). The

largest value sets were for ICS and long-acting beta-agonists (ICS-LABA)(N=201, N=71,

respectively).

Applying codelists to find prescriptions

We applied the codelists to a cohort of 335,931 patients diagnosed with COPD, according to study population considerations and clinical input.

For the cardiovascular codelist, within the decade follow-up, 165,150 patients (49.2% of cohort) were prescribed at least one of the drugs (Figure 3;Supplmentary TableS3). As in the case with count, the value set for chapter 2.5.5 had the greatest number of patients prescribed (N=151,225, 45.0% of cohort). Chapters 2.5.3 and 2.5.8 did not have prescriptions.

For the COPD inhalers codelist, we determined 317,963 patients (94.7% of cohort) prescribed at least one of the drugs (Figure 4;Supplmentary TableS4). Counts and prescriptions followed different patterns. Whilst ICS had greatest count (N=213 codes), SABA had the most prescriptions (N=297,966; 88.7% of cohort).

Comparing to restricted searches

We compared output of our full 'gold-standard' searches (A) to two restricted searches still using our methodology but searching on (B) chemical terms within *drugsubstancename* and (C) BNF ontology within *bnfchapter*.

For the cardiovascular codelist, search B identified 505 codes and 155,678 patients prescribed (46.3% of cohort) at least one of the drugs across value sets. Search C identified 267 codes and 150,669 patients (44.9% of cohort) prescribed at least one of the drugs across value sets (Figure 3; Supplementary TableS3).

For the COPD inhalers codelist, search B identified 351 codes, and 317,957 patients prescribed (95% of cohort) at least one of the drugs across value sets. Search C identified 185 codes and 315,749 patients (94% of cohort) prescribed at least one of the drugs across value sets (Figure 4; Supplementary TableS4).

The percent increase in output from searching on BNF ontology only (C) to the gold-standard (Search A) was the most pronounced (cardiovascular codelist: 113% and 9.6% increase in codes and

prescriptions, respectively; COPD codelist 147% and 0.7%). However, we observed marginal

increases when comparing Search A to searching solely on chemical information (B) (cardiovascular

codelist: 12.5%, 6.08% increase respectively; COPD codelist 29%, 0.0002% increase respectively).

Considering restricted searches by value set, there were absent or marginal increases in prescriptions

in some value sets upon using Search A (close to 0%), but remarkable increase in others, particularly

for C (up to 24802% BNF 2.5.4; up to 6161% for SAMA). Search B led to higher output compared with

C, except for Ch. 2.5.5. Here, greater counts in B vs. C did not translate to greater prescriptions

(N=343 counts, N=138,992 prescriptions for B; N=217 and N=150,117 for C).

Figure 3. Comparison of codelist output by search type for the cardiovascular codelist
Adapted from an UpSet plot design[27,28]. RAAS=renin-angiotensin-aldosterone system. Results refer to post-clinicians' input. Counts derive from codelist generation, prescribed patients determined after codelist applied to COPD cohort. Prescriptions by value set are mutually exclusive as some patients are prescribed drugs of different classes across value sets, e.g., prescribed drug falling into 'at least' one subsection. Partially shaded dots for Ch. 2.5.8 refer to presence of codes in codelist, but absence upon application to cohort to determine prescriptions. Search A refers to the use of our methodology, searching *termfromemis, productname, drugsubstancename*, and *bnfchapter* variables. Search B refers to the use of our methodology, but searching on *drugsubstancename* variable, only. Search C refers to the use of our methodology, but searching on *bnfchapter* variable, only (Search C). Refer to Supplementary Table S3 for full data.

413
Figure 4. Comparison of codelist output by search type for the COPD inhalers codelist
Adapted from an UpSet plot design[27,28]. Results refer to post-clinicians' input. Counts derive from codelist
generation, prescribed patients determined after codelist applied to COPD cohort. Prescriptions by value set are
mutually exclusive as some patients are prescribed drugs of different classes across value sets, e.g., prescribed
drug falling into 'at least' one subsection. Search A refers to the use of our methodology, searching
termfromemis, productname, drugsubstancename, and *bnfchapter* variables. Search B refers to the use of our
methodology, but searching on *drugsubstancename* variable, only. Search C refers to the use of our
methodology, but searching on *bnfchapter* variable, only (Search C). Refer to Supplementary Table S4 for full
data.

DISCUSSION

Summary

We developed a standardisable, reproducible method for creating comprehensive drug codelists

using a semi-automated process incorporating end-to-end clinician expertise, considering missing

data and fluctuating status that is adaptable to other studies and databases.

We applied the methodology to generate two codelists that were implemented on a sample cohort of

patients with COPD in CPRD Aurum, according to study considerations and clinical review.

Evaluation

Searching only on BNF ontology (C) was insufficient and missed the most prescriptions in CPRD.

Given similar results for searches A and B, there may be a marginal opportunity cost between

searching on all variable attributes versus chemical terms alone, depending on drugs desired (i.e.,

restricting to a portion of the value sets, e.g., a short-acting muscarinic antagonist (SAMA)-only

codelist; or drugs in 2.5.4 only), and cohort size (e.g., rare vs. common disease). Value set Ch. 2.5.5

results could be explained by, despite fewer counts, saturation of prescriptions among particular

codes, upon codelist application to the study cohort.

Although we focus on CPRD and the cohort-of-interest, methods centre around addition of new

information (e.g., drugs, proprietary names), completeness, and context-specific adaptations,

applicable to other databases and underlying ontologies.

Recommendations

We recommend the full, 'gold-standard' method (A) if comprehensiveness and reaching statistical

power (i.e. studies where codelist defines cohort or exposure) or when sample size (e.g., propensity

scoring on complete data) is of focus. . In our database context which contained missing data in the

drug dictionary, a comprehensive search on multiple 'attribute' variables was warranted, but this may

not be the case for other data sources with data completeness.

If the aim is to produce a broader codelist (i.e., higher specificity, lower sensitivity) permitting

modification for various contexts, we recommend limiting *a priori* exclusion criteria in step 1; rather,

designing sets around underlying taxonomy. If the aim is disease-specific, value set generation

should be clinician-led but still designed to permit malleability for different studies, i.e., single/fewer

classes (e.g., statins only). To ensure all possible terms are found we recommend steps 2a-2b

iteratively.

Comparisons to previous codelists and/or mapping files can pick up additional codes. In studies

incorporating multiple databases, we recommend merging with ontology mapping files to enable

codelist harmonisation[15,29], such as ATC-BNF mapping.

Current literature

Complexities of creating medical codelists are described elsewhere[12,24], and in some databases,

recorded prescriptions are treated separately from medical events and often recorded in separate

tables using different coding schemes[2,5,9]. Few studies outline methods for codelist development,

with the paucity of literature prominent for drug codelists where focus has been high-level, covering

challenges, assumptions, and principles in EHR research, e.g., software and analytical techniques,

data preparation, and defining periods for drug covariates[17,18], but not on underlying steps for

codelist generation. Of literature on medical codelists, the focus was on general guidelines for

researchers to follow to enhance reproducibility[12,30], including incorporating clinician review to

explore codes' uncertainty[12], exploring applications of codelists to sample cohorts[12], and

identifying and comparing disease phenotypes based on restricted versus expanded conceptual

definitions[16,24].

Other considerations

We recommend consideration of the database and context in all study stages, yet emphasize building

modifiability into codelist methodology to allow generalisation. Modifications may derive from study

nature, including the period (e.g., retrospective with discontinued drugs in-use during the study),

cohort-of-interest (e.g., patients with COPD where certain cardiovascular prescriptions are not

indicated), and subsequent statistical analyses. In studies with drug covariates, overlaps in class

could present collinearity, where exclusion of overlapping codes may be required. Our solution was

proactively tagging overlapping codes, drawing upon clinical expertise. Codelist tailoring may relate to

database factors, such as data type (e.g., medications recorded in primary care versus claims data)

and taxonomy (e.g., BNF or ATC codes).

Due to changes in new, existing, and discontinued drugs, and periods-of-interest for retrospective

studies, the same codelist may need to be updated based on older or newer database versions.

Using nested lists allows for maintained organisation despite these realities. Future methods may

consider adding an extra column to the codelist showing "in use" status, although anecdotal evidence

suggests applying codes for discontinued drugs to a newer cohort will not pick up prescriptions.

Results on number of patients prescribed may differ depending on the cohort. Before generating final

study-specific codelists, clinical input for exclusion may differ by disease, e.g., drugs contraindicated

in specific patients such the codelist would be adapted to exclude corresponding value set(s).

Prescriptions may change if the codelist is applied to separate disease cohorts, where prescription

commonality by disease varies (e.g., cohorts with COPD, diabetes, Chronic Kidney Disease, or

multimorbidity).

Results on number of prescriptions may differ depending on how codelists are operationalised, i.e., as

covariates or exposures, part of inclusion, or accounting factors such as duration and frequency[31]

and for combination, open therapies (e.g., for SAMA-SABA beyond fixed therapies determined

through codelist generation).

Regarding external overlap to other taxonomic sections, the act of tagging is information proactively

sought and built into initial steps, allowing for streamlined adaptability, e.g., should covariate

collinearity arise. When considering adjustment of confounding covariates, covariate collinearity could

distort observed exposure-outcome effects. Resolving collinearity through codelist adaptation should

include clinical input.

CONCLUSION

We designed a semi-automated process to generate drug codelists using standardisable and

reproducible methodology. Despite database identity, there are special considerations when

generating adaptable drug codelists, including in-flux status, cohort-specific drug indication and

exclusions, database-specific underlying hierarchical ontology, and operationalisation relating to

inclusion and covariate analysis. Regardless, many EHR researchers are not clinicians; supplemental

input is necessary.

AUTHOR CONTRIBUTIONS

EG devised the protocol, sought ethics approval, and supervised the project, with PS advising on methodology. PS, GM, and SH provided methodological and analytical support. EG drafted the manuscript. JQ provided respiratory clinical input when generating the two cohort-specific codelists. NP provided clinical input when generating the cardiovascular cohort-specific codelist. AA translated the STATA script into R. All authors contributed content, reviewed and approved the final manuscript.

FUNDING

No funding is reported for this study. This research was supported by the NIHR Imperial Biomedical Research Centre (BRC).

CONFLICTS OF INTEREST

JQ has received grants from MRC, HDR UK, GSK, BI, asthma+lung UK, and AZ and personal fees for advisory board participation, consultancy or speaking fees from GlaxoSmithKline, Evidera, AstraZeneca, Insmed. NP has received funding from Imperial Health Charity, SD is supported by the BHF Data Science Centre led by HDR UK (grant SP/19/3/34678), BigData@Heart Consortium, funded by the Innovative Medicines Initiative-2 Joint Undertaking under grant agreement 116074, the NIHR Biomedical Research Centre at University College London Hospital NHS Trust (UCLH BRC), a BHF Accelerator Award (AA/18/6/24223), E) the CVD-COVID-UK/COVID-IMPACT consortium and the Multimorbidity Mechanism and Therapeutic Research Collaborative (MMTRC, grant number MR/V033867/1). PS reports grants from asthma+lung UK and Gilead. EG, GM, AA, and SH have nothing to disclose.

PATIENT CONSENT FOR PUBLICATION

Not applicable.

ACKNOWLEDGEMENTS

The data used are from CPRD obtained under license from the UK Medicines and Healthcare products Regulatory Agency. The data is provided by patients and collected by the National Health Service as part of their care and support.

PEER REVIEW

Externally peer reviewed.

DATA AVAILABILITY STATEMENT

Data may be obtained from a third party and are not publicly available. Data are available on request from CPRD. CPRD data provision requires purchase of a license, and this license does not permit the authors to make them publicly available to all.

ETHICS APPROVAL

CPRD has NHS Health Research Authority (HRA) Research Ethics Committee (REC) approval to allow the collection and release of anonymised primary care data for observational research [NHS HRA REC reference number: 05/MRE04/87]. Each year CPRD obtains Section 251 regulatory support through the HRA Confidentiality Advisory Group (CAG), to enable patient identifiers, without accompanying clinical data, to flow from CPRD contributing GP practices in England to NHS Digital, for the purposes of data linkage [CAG reference number: 21/CAG/0008]. The protocol for this research was approved by CPRD's Research Data Governance (RDG) Process (protocol number: 22_002515) and the approved protocol is available upon request. Linked pseudonymised data was provided for this study by CPRD. Data is linked by NHS Digital, the statutory trusted third party for linking data, using identifiable data held only by NHS Digital. Select general practices consent to this process at a practice level with individual patients having the right to opt-out.

REFERENCES

1   Xu Z, Wang F, Adekkanattu P, *et al.* Subphenotyping depression using machine learning and electronic health records. *Learn Health Syst* 2020;**4**:e10241. doi:10.1002/lrh2.10241

2   NCATS National COVID Cohort Collaborative (N3C) Data Enclave. COVID-19 Clinical Data Warehouse Data Dictionary: Based on OMOP Common Data Model Specifications Version 5.3. https://ncats.nih.gov/files/OMOP_CDM_COVID.pdf

3   Justice AC, Gordon KS, Romero J, *et al.* Polypharmacy-associated risk of hospitalisation among people ageing with and without HIV: an observational study. *Lancet Healthy Longev* 2021;**2**:e639–50. doi:10.1016/S2666-7568(21)00206-3

4   WSIC Data Specification, v11. https://www.registerfordiscover.org.uk/uploads/files/1539001703datadictionary.pdf

5   CPRD Aurum Data Specification, v2.8. 2022.https://cprd.com/sites/default/files/2022-

08/CPRD%20Aurum%20Data%20Specification%20v2.8.pdf

6   VA Family of EHR Cohorts (VACo Family). https://medicine.yale.edu/intmed/vacs/ (accessed 31

Mar 2023).

7   Wood A, Denholm R, Hollings S, *et al.* Linked electronic health records for research on a

nationwide cohort of more than 54 million people in England: data resource. *BMJ* 2021;**373**:n826.

doi:10.1136/bmj.n826

8   Data. SAIL Databank. https://saildatabank.com/data/ (accessed 31 Mar 2023).

9   Common Data Model (CDM) Specification, Version 6.0. 2020.https://pcornet.org/wp-

content/uploads/2022/01/PCORnet-Common-Data-Model-v60-2020_10_221.pdf (accessed 31

Mar 2023).

10   Wolf A, Dedman D, Campbell J, *et al.* Data resource profile: Clinical Practice Research Datalink

(CPRD) Aurum. *Int J Epidemiol* 2019;**48**:1740–1740g. doi:10.1093/ije/dyz034

11   Healthcare Epidemiology. Duke Dep. Med. https://medicine.duke.edu/education-and-

training/fellowship-programs/infectious-diseases/training-and-curriculum/research-

pathways/healthcare-epidemiology (accessed 1 Apr 2023).

12   Watson J, Nicholson BD, Hamilton W, *et al.* Identifying clinical features in primary care electronic

health record studies: methods for codelist development. *BMJ Open* 2017;**7**:e019637.

doi:10.1136/bmjopen-2017-019637

13   Sydes MR, Barbachano Y, Bowman L, *et al.* Realising the full potential of data-enabled trials in

the UK: a call for action. *BMJ Open* 2021;**11**:e043906. doi:10.1136/bmjopen-2020-043906

14   Benchimol EI, Smeeth L, Guttmann A, *et al.* The REporting of studies Conducted using

Observational Routinely-collected health Data (RECORD) Statement. *PLOS Med*

2015;**12**:e1001885. doi:10.1371/journal.pmed.1001885

15   Abbasizanjani H, Torabi F, Bedston S, *et al.* Harmonising electronic health records for

reproducible research: challenges, solutions and recommendations from a UK-wide COVID-19

research collaboration. *BMC Med Inform Decis Mak* 2023;**23**:8. doi:10.1186/s12911-022-02093-0

16   Jayatunga W, Stone P, Aldridge RW, *et al.* Code sets for respiratory symptoms in electronic

health records research: a systematic review protocol. *BMJ Open* 2019;**9**:e025965.

doi:10.1136/bmjopen-2018-025965

17   Pye SR, Sheppard T, Joseph RM, *et al.* Assumptions made when preparing drug exposure data

for analysis have an impact on results: An unreported step in pharmacoepidemiology studies.

Pharmacoepidemiol Drug Saf 2018;**27**:781–8. doi:10.1002/pds.4440

18   Denaxas S, Direk K, Gonzalez-Izquierdo A, *et al.* Methods for enhancing the reproducibility of

biomedical research findings using electronic health records. *BioData Min* 2017;**10**:31.

doi:10.1186/s13040-017-0151-7

19   VANDF (National Drug File). UMLS Metathesaurus.

https://www.nlm.nih.gov/research/umls/sourcereleasedocs/current/VANDF/index.html (accessed

1 Feb 2023).

20   ATC (Anatomical Therapeutic Chemical Classification System). UMLS Metathesaurus.

https://www.nlm.nih.gov/research/umls/sourcereleasedocs/current/ATC/index.html (accessed 1

Feb 2023).

21   RxNorm Technical Documentation.

2021.https://www.nlm.nih.gov/research/umls/rxnorm/docs/techdoc.html#s1_0

22   SNOMED International. 2.2. Value Set. In: *Practical Guide to Reference Sets*. International

Health Terminology Standards Development Organisation 2023.

https://confluence.ihtsdotools.org/display/DOCRFSPG/2.2.+Value+Set (accessed 18 Mar 2023).

23   Bennett Institute for Applied Data Science, University of Oxford. All BNF sections.

OpenPrescribing. 2023.https://openprescribing.net/bnf/ (accessed 9 Feb 2023).

24   Elkheder M, Gonzalez-Izquierdo A, Qummer Ul Arfeen M, *et al.* Translating and evaluating

historic phenotyping algorithms using SNOMED CT. *J Am Med Inform Assoc JAMIA*

2023;**30**:222–32. doi:10.1093/jamia/ocac158

25   Morgan AD, Zakeri R, Quint JK. Defining the relationship between COPD and CVD: what are the

implications for clinical practice? *Ther Adv Respir Dis* 2018;**12**:175346581775052.

doi:10.1177/1753465817750524

26   Quint JK, Müllerova H, DiSantostefano RL, *et al.* Validation of chronic obstructive pulmonary

disease recording in the Clinical Practice Research Datalink (CPRD-GOLD). *BMJ Open*

2014;**4**:e005540. doi:10.1136/bmjopen-2014-005540

27   Conway JR, Lex A, Gehlenborg N. UpSetR: an R package for the visualization of intersecting

sets and their properties. *Bioinformatics* 2017;**33**:2938–40. doi:10.1093/bioinformatics/btx364

28   Lex A, Gehlenborg N, Strobelt H, *et al.* UpSet: Visualization of Intersecting Sets. *IEEE Trans Vis*

Comput Graph 2014;**20**:1983–92. doi:10.1109/TVCG.2014.2346248

29   NHS Digital. NHSBSA dm+d. TRUD.

https://isd.digital.nhs.uk/trud/users/guest/filters/0/categories/6/items/24/releases (accessed 1 Feb

2023).

30   Davé S, Petersen I. Creating medical and drug code lists to identify cases in primary care

databases. *Pharmacoepidemiol Drug Saf* 2009;**18**:704–7. doi:10.1002/pds.1770

31   Whittaker HR, Wing K, Douglas I, *et al.* Inhaled Corticosteroid Withdrawal and Change in Lung

Function in Primary Care Patients with Chronic Obstructive Pulmonary Disease in England. *Ann*

Am Thorac Soc 2022;**19**:1834–41. doi:10.1513/AnnalsATS.202111-1238OC