Resource allocation strategies for insecticide-treated bed nets to achieve malaria eradication

Nora Schmit[1]*, Hillary M Topazian[1]*, Matteo Pianella[1]*, Giovanni D Charles[1], Peter Winskill[1], Michael T White[2], Katharina Hauck[3], Azra C Ghani[1]

[1] MRC Centre for Global Infectious Disease Analysis, Imperial College London, London, United Kingdom

[2] Infectious Disease Epidemiology and Analytics G5 Unit, Department of Global Health, Institut Pasteur, Université de Paris, Paris, France

[3] MRC Centre for Global Infectious Disease Analysis, Jameel Institute, Imperial College London, London, United Kingdom

* These authors contributed equally to this work

Corresponding author:

Nora Schmit

MRC Centre for Global Infectious Disease Analysis

Imperial College London

St Mary's Campus

Norfolk Place, London W2 1PG

n.schmit17@imperial.ac.uk

ABSTRACT

Background: Large reductions in the global malaria burden have been achieved in the last

decades, but plateauing funding poses a challenge for progressing towards the ultimate goal of

malaria eradication. We aimed to determine the optimal strategy to allocate global resources to

achieve this goal.

Methods: Using previously published mathematical models of *Plasmodium falciparum* and

Plasmodium vivax transmission incorporating insecticide-treated nets (ITNs) as an illustrative

intervention, we sought to identify the global funding allocation that maximized impact under

defined objectives and across a range of global funding budgets.

Results: We found that the optimal strategy for case reduction closely mirrored an allocation

framework that prioritizes funding for high-transmission settings, resulting in total case

reductions of 76% (optimal strategy) and 66% (prioritizing high-transmission settings) at

intermediate budget levels. Allocation strategies that had the greatest impact on case reductions

were associated with lesser near-term impacts on the global population at risk, highlighting a

trade-off between reducing burden and "shrinking the map" through a focus on near-elimination

settings. The optimal funding distribution prioritized high ITN coverage in high-transmission

settings endemic for *P. falciparum* only, while maintaining lower levels in low-transmission

settings. However, at high budgets, 62% of funding was targeted to low-transmission settings

co-endemic for *P. falciparum* and *P. vivax*.

Conclusions: These results support current global strategies to prioritize funding to high-

burden *P. falciparum*-endemic settings in sub-Saharan Africa to minimize clinical malaria burden

and progress towards elimination but highlight competing goals of reducing the global

population at risk and addressing the burden of *P. vivax*.

INTRODUCTION

Global support for malaria eradication has fluctuated in response to changing health policies

over the past 75 years. From near global endemicity in the 1900's over 100 countries have

eliminated malaria, with 10 of these certified malaria-free by the World Health Organization

(WHO) in the last two decades (Feachem et al., 2010, Shretta et al., 2017, Weiss et al., 2019).

Despite this success, 41% and 57% of the global population in 2017 were estimated to live in

areas at risk of infection with *Plasmodium falciparum* and *Plasmodium vivax*, respectively

(Weiss et al., 2019, Battle et al., 2019). In 2021 there were an estimated 247 million new

malaria cases and over 600,000 deaths, primarily in children under 5 years of age (World Health

Organization, 2022b). Mosquito resistance to the insecticides used in vector control, parasite

resistance to both first-line therapeutics and diagnostics, and local active conflicts continue to

threaten elimination efforts (World Health Organization, 2020). Nevertheless, the global

community continues to strive towards the ultimate aim of eradication, which could save millions

of lives and thus offer high returns on investment (Chen et al., 2018, Strategic Advisory Group

on Malaria Eradication, 2020).

The global goals outlined in the Global Technical Strategy for Malaria (GTS) 2016-2030 include

reducing malaria incidence and mortality rates by 90%, achieving elimination in 35 countries,

and preventing re-establishment of transmission in all countries currently classified as malaria-

free by 2030 (World Health Organization, 2015). Various stakeholders have also set timelines

for the wider goal of global eradication, ranging from 2030 to 2050 (World Health Organization,

2020, Chen et al., 2018, Strategic Advisory Group on Malaria Eradication, 2020). However,

there remains a lack of consensus on how best to achieve this longer-term aspiration.

Historically, large progress was made in eliminating malaria mainly in lower-transmission

countries in temperate regions during the Global Malaria Eradication Program in the 1950s, with

the global population at risk of malaria reducing from around 70% of the world population in 1950 to 50% in 2000 (Hay et al., 2004). Renewed commitment to malaria control in the early 2000s with the Roll Back Malaria initiative subsequently extended the focus to the highly endemic areas in sub-Saharan Africa (Feachem et al., 2010). Whilst it is now widely acknowledged that the current tool set is insufficient in itself to eradicate the parasite, there continues to be debate about how resources should be allocated (Snow, 2015). Some advocate for a focus on high-burden settings to lower the overall global burden (World Health Organization, 2019), while others call for increased funding to middle-income low-burden countries through a 'shrink the map strategy' where elimination is considered a driver of global progress (Newby et al., 2016). A third set of policy options is influenced by equity considerations including allocating funds to achieve equal allocation per person at risk, equal access to bed nets and treatment, maximize lives saved, or to achieve equitable overall health status (World Health Organization, 2013, Raine et al., 2016).

Global strategies are influenced by international donors, which represent 68% of the global investment in malaria control and elimination activities (World Health Organization, 2022b). The Global Fund and the U.S. President's Malaria Initiative are two of the largest contributors to this investment. Their strategies pursue a combination approach, prioritizing malaria reduction in high-burden countries while achieving sub-regional elimination in select settings (The Global Fund, 2021, United States Agency for International Development and Centers for Disease Control and Prevention, 2021). Given that the global investment for malaria control and elimination still falls short of the 6.8 billion USD currently estimated to be needed to meet GTS 2016-2030 goals (World Health Organization, 2020), an optimized strategy to allocate limited resources is critical to maximizing the chance of successfully achieving the GTS goals and longer-term eradication aspirations.

In this study, we use mathematical modelling to explore the optimal allocation of limited global resources to maximize the long-term reduction in *P. falciparum* and *P. vivax* malaria. Our aim is to determine whether financial resources should initially focus on high-transmission countries, low-transmission countries, or a balance between the two across a range of global budgets. In doing so, we consider potential trade-offs between short-term gains and long-term impact. We use compartmental deterministic versions of two previously developed and tested individual-based model of *P. falciparum* and *P. vivax* transmission respectively (Griffin et al., 2010, White et al., 2018). Using the compartmental model structures allows us to fully explore the space of possible resource allocation decision using optimization, which would be prohibitively costly to perform using more complex individual-based models. Furthermore, to evaluate the impact of resource allocation options, we focus on a single intervention - insecticide-treated nets (ITNs). Whilst in reality national malaria elimination programs encompass a broad range of preventative and therapeutic tools alongside different surveillance strategies as transmission decreases, this simplification is made for computational feasibility, with ITNs chosen as they (a) provide both an individual protective effect and population-level transmission reductions (i.e. indirect effects); (b) are the most widely used single malaria intervention other than first-line treatment; and (c) extensive distribution and costing data are available that allow us to incorporate their decreasing technical efficiency at high coverage.

RESULTS

We identified 105 malaria-endemic countries based on 2000 *P. falciparum* and *P. vivax* prevalence estimates (before the scale-up of interventions), of which 44, 9, and 52 were endemic for *P. falciparum* only, *P. vivax* only, and co-endemic for both species, respectively. Globally, the clinical burden of malaria was focused in settings of high transmission intensity endemic for *P. falciparum* only, followed by low-transmission settings co-endemic for *P. falciparum* and *P. vivax* (**Figure 1A**). Conversely, 89% of the global population at risk of malaria

was located in co-endemic settings with very low and low transmission intensities (**Figure 1B**).

All 25 countries with high transmission intensity and 11 of 17 countries with moderate transmission intensity were in Africa, while almost half of global cases and population at risk in low-transmission co-endemic settings originated in India.

Deterministic compartmental versions of two previously published and validated mathematical models of *P. falciparum* and *P. vivax* malaria transmission dynamics (Griffin et al., 2010, Griffin et al., 2014, Griffin et al., 2016, White et al., 2018) were used to explore associations between ITN use and clinical malaria incidence. In model simulations, the relationship between ITN usage and malaria infection outcomes varied by the baseline entomological inoculation rate (EIR), representing local transmission intensity, and parasite species (**Figure 2**). The same increase in ITN usage achieved a larger relative reduction in clinical incidence in low-EIR than in high-EIR settings. Low levels of ITN usage were sufficient to eliminate malaria in low-transmission settings, whereas high ITN usage was necessary to achieve a substantial decrease in clinical incidence in high-EIR settings. At the same EIR value, ITNs also led to a larger relative reduction in *P. falciparum* than *P. vivax* clinical incidence. However, ITN usage of 80% was not sufficient to lead to full elimination of either *P. falciparum* or *P. vivax* in the highest transmission settings. In combination, the models projected that ITNs could reduce global *P. falciparum* and *P. vivax* cases by 83.6% from 252.0 million and by 99.9% from 69.3 million in 2000, respectively, assuming a maximum ITN usage of 80%.

We next used a non-linear generalized simulated annealing function to determine the optimal global resource allocation for ITNs across a range of budgets. We defined optimality as the funding allocation across countries which minimizes a given objective. We considered two objectives: first, reducing the global number of clinical malaria cases, and second, reducing both the global number of clinical cases and the number of settings not having yet reached a pre-elimination phase. The latter can be interpreted as accounting for an additional positive

contribution of progressing towards elimination on top of a reduced case burden (e.g. general health system strengthening through a reduced focus on malaria). To relate funding to impact on malaria, we incorporated a non-linear relationship between costs and ITN usage, resulting in an increase in the marginal cost of ITN distribution at high coverage levels (Bertozzi-Villa et al., 2021). We considered a range of fixed budgets, with the maximum budget being that which enabled achieving the lowest possible number of cases in the model. Low, intermediate and high budget levels refer to 25%, 50% and 75% of this maximum, respectively.

In our main analysis we ignored the time dimension over which funds are distributed, instead focusing on the endemic equilibrium reached for each level of allocation (sensitivity to this assumption is explored in a second analysis with dynamic re-allocation every 3 years). The optimal strategies were compared with three existing approaches to resource allocation: 1) prioritization of high-transmission settings, 2) prioritization of low-transmission (near-elimination) settings, and 3) proportional allocation by disease burden. Strategies prioritizing high- or low-transmission settings involved sequential allocation of funding to groups of countries based on their transmission intensity (from highest to lowest EIR or *vice versa*). The proportional allocation strategy mimics the current allocation algorithm employed by the Global Fund: budget shares are distributed according to malaria disease burden in the 2000-2004 period (The Global Fund, 2019). To allow comparison with this existing funding model, we also started allocation decisions from the year 2000.

We found that the optimal strategies for reducing total malaria cases (i.e. global burden) and for case reduction and pre-elimination to be similar to the strategy that prioritized funding for high-transmission settings. These three strategies achieved the largest reductions in global malaria cases at all budgets, including reductions of 76%, 73% and 66% at the intermediate budget level, respectively (**Figure 3A**, **Table 1**). At low to intermediate budgets, the proportional

allocation strategy also reduced malaria cases effectively by up to 53%. While these four

scenarios had very similar effects on malaria cases at low budgets, they diverged with

increasing funding, where the proportional allocation strategy did not achieve substantial further

reductions. Depending on the available budget, the optimal strategy for case reduction averted

up to 31% more cases than prioritization of high-transmission settings and 64% more cases

than proportional allocation, corresponding to respective differences of 37.9 and 74.5 million

cases globally.

We additionally found there to be a trade-off between reducing global cases and reducing the

global population at risk of malaria. Both the optimal strategies and the strategy prioritizing high-

transmission settings did not achieve substantial reductions in the global population at risk until

large investments were reached (**Figure 3B, Table 1**). Even at a high budget, the global

population at risk was only reduced by 19% under the scenario prioritizing high-transmission

settings, with higher reductions of 42-58% for the optimal strategies, while proportional

allocation had almost no effect on this outcome. Conversely, diverting funding to prioritize low-

transmission settings was highly effective at increasing the number of settings eliminating

malaria, achieving a 56% reduction in the global population at risk already at intermediate

budgets. However, this investment only led to a minimal reduction of 24% in total malaria case

load (**Figure 3**, **Table 1**). At high budget levels, prioritizing low-transmission settings resulted in

up to 3.8 times (a total of 159.4 million) more cases than the optimal allocation for case

reduction. Despite the population at risk remaining relatively large with the optimal strategy for

case reduction and pre-elimination, it nevertheless led to pre-elimination in more malaria-

endemic settings than all other strategies (**Figure S7**), in addition to close to minimum cases

across all budgets (**Figure 3**).

The allocation strategies also had differential impacts on *P. falciparum* and *P. vivax* cases, with case reductions generally occurring first for *P. falciparum* except when prioritizing low-transmission settings. *P. vivax* cases were not substantially affected at low global budgets for all other allocation strategies, and proportional allocation had almost no effect on reducing *P. vivax* clinical burden at any budget (**Figure 3C**), leading to a temporary increase in the proportion of total cases attributable to *P. vivax* relative to *P. falciparum*. The global population at risk remained high with the optimal strategy for case reduction even at high budgets, partly due to a large remaining population at risk of *P. vivax* infection (**Figure 3D**), which was not targeted when aiming to minimize total cases (**Figure 1**).

The optimized distribution of funding to minimize clinical burden depended on the available global budget and was driven by the setting-specific transmission intensity and the population at risk (**Figure 4**, **Figure 1**). With very low to low budget levels, as much as 85% of funding was allocated to moderate to high transmission settings (**Figure 4A, Figure S8A**). This allocation pattern led to the maximum ITN usage of 80% being reached in settings of high transmission intensity and smaller population sizes even at low budgets, while maintaining lower levels in low-transmission settings with larger populations (**Figure 4B, Figure S8B**). The proportion of the budget allocated to low and very low transmission settings increased with increasing budgets, and low transmission settings received the majority of funding at intermediate to maximum budgets. This allocation pattern remained very similar when optimizing for both case reduction and pre-elimination (**Figure S9**). Similar patterns were also observed for the optimized distribution of funding between settings endemic for only *P. falciparum* compared to *P. falciparum* and *P. vivax* co-endemic settings (**Figure 4C-D**), with the former being prioritized at low to intermediate budgets. At the maximum budget, 70% of global funding was targeted at low- and very low-transmission settings co-endemic for both parasite species.

To evaluate the robustness of the results, we conducted a sensitivity analysis on our

assumption on ITN distribution efficiency. Results remained similar when assuming a linear

relationship between ITN usage and distribution costs (**Figure S10**). While the main analysis

involves a single allocation decision to minimise long-term case burden (leading to a constant

ITN usage over time in each setting irrespective of subsequent changes in burden), we

additionally explored an optimal strategy with dynamic re-allocation of funding every 3 years to

minimise cases in the short term. At high budgets, capturing dynamic changes over time

through re-allocation of funding based on minimizing *P. falciparum* cases every 3 years led to

the same case reductions over time as a one-time optimization with allocation of a constant ITN

usage (**Figure S11**). At lower budgets, re-allocation every 3 years achieved a higher impact at

several timepoints, but total cases remained similar between the two approaches. Although

reallocation of resources from settings which achieved elimination to higher transmission

settings did not lead to substantially fewer cases, it reduced total spending over the 39-year

period in some cases (**Figure S11**).

DISCUSSION

Our study highlights the potential impact that funding allocation decisions could have on the

global burden of malaria. We estimated that optimizing ITN allocation to minimize global clinical

incidence could, at a high budget, avert 83% of clinical cases compared to no intervention. In

comparison, the optimal strategy to minimize clinical incidence and maximize the number of

settings reaching pre-elimination averted 82% of clinical cases, prioritizing high-transmission

settings 81%, proportional allocation 61%, and prioritizing low-transmission settings 37%. Our

results support initially prioritizing funding towards reaching high ITN usage in the high-burden

P. falciparum-endemic settings to minimize global clinical cases and advance elimination in

more malaria-endemic settings, but highlight the trade-off between this strategy and reducing

the global population at risk of malaria as well as addressing the burden of *P. vivax*.

Prioritizing low-transmission settings demonstrated how focusing on "shrinking the malaria map"

by quickly reaching elimination in low-transmission countries diverts funding away from the high-

burden countries with the largest caseloads. Prioritizing low-transmission settings achieved

elimination in 42% of settings and reduced the global population at risk by 56% when 50% of

the maximum budget had been spent, but also resulted in 3.2 times more clinical cases than the

optimal allocation scenario. Investing a larger share of global funding towards high-transmission

settings aligns more closely with the current WHO "high burden to high impact" approach, which

places an emphasis on reducing the malaria burden in the 11 countries which comprise 70% of

global cases (World Health Organization, 2019). Previous research supports this approach,

finding that the 20 highest burden countries would need to obtain 88% of global investments to

reach case and mortality risk estimates in alignment with GTS goals (Patouillard et al., 2017).

This is similar to the modelled optimized funding strategy presented here, which allocated up to

76% of very low budgets to settings of high transmission intensity located in sub-Saharan Africa.

An initial focus on high- and moderate-transmission settings is further supported by our results

showing that a balance can be found between achieving close to optimal case reductions while

also progressing towards elimination in the maximum number of settings. Even within a single

country, targeting interventions to local hot-spots has been shown to lead to higher cost savings

than universal application (Barrenho et al., 2017), and could lead to elimination in settings

where untargeted interventions would have little impact (Bousema et al., 2012).

Assessing optimal funding patterns is a global priority due to the funding gap between supply

and demand for resources for malaria control and elimination (World Health Organization,

2022b). However, allocation decisions will remain important even if more funding became

available, as some of the largest differences in total cases between the modelled strategies

occurred at intermediate to high budgets. Our results suggest that most of global funding should

only be focused in low-transmission settings co-endemic for *P. falciparum* and *P. vivax* at high

budgets once ITN use has already been maximized in high-transmission settings. Global

allocation decisions are likely to affect *P. falciparum* and *P. vivax* burden differently, which could

have implications for the future global epidemiology of malaria. For example, with a focus on

disease burden reduction, a temporary increase in the proportion of malaria cases attributable

to *P. vivax* was projected, in line with recent observations in near-elimination areas (Battle et al.,

2019, Price et al., 2020). Nevertheless, even when international funding for malaria increased

between 2007-2009, African countries remained the major recipients of financial support, while

P. vivax-dominant countries were not as well funded (Snow et al., 2010). This serves as a

reminder that achieving elimination of malaria from all endemic countries will ultimately require

targeting investments so as to also address the burden of *P. vivax* malaria.

Different priorities in resource allocation decisions greatly affect which countries receive funding

and what health benefits are achieved. The modelled strategies follow key ethical principles in

the allocation of scarce healthcare resources, such as targeting those of greatest need

(*prioritizing high-transmission settings*, *proportional allocation*) or those with the largest

expected health gain (*optimized for case reduction*, *prioritizing high-transmission settings*)

(World Health Organization, 2013). Allocation proportional to disease burden did not achieve as

great an impact as other strategies because the funding share assigned to settings was

constant irrespective of the invested budget and its impact. In modelling this strategy, we did

not reassign excess funding in high-transmission settings to other malaria interventions, as

would likely occur in practice. This illustrates the possibility that such an allocation approach can

potentially target certain countries disproportionally and result in further inequities in health

outcomes (Barrenho et al., 2017). From an international funder perspective, achieving vertical

equity might therefore also encompass higher disbursements to countries with lower

affordability of malaria interventions (Barrenho et al., 2017), as reflected in the Global Fund's

proportional allocation formula which accounts for the economic capacity of countries and

specific strategic priorities (The Global Fund, 2019). While these factors were not included in the

proportional allocation used here, the estimated impact of these two strategies was

nevertheless very similar (**Supplementary Figure S12**).

While our models are based on country patterns of transmission settings and corresponding

populations in 2000, there are several factors leading to heterogeneity in transmission dynamics

at the national and sub-national level which were not modelled and limit our conclusions.

Seasonality, changing population size, and geographic variation in *P. vivax* relapse patterns or

in mosquito vectors could affect the projected impact of ITNs and optimized distribution of

resources across settings. The two representative *Anopheles* species used in the simulations

are also both very anthropophagic, which may have led to overestimation of the effect of ITNs in

some settings. By using ITNs as the sole means to reduce mosquito-to-human transmission, we

did not capture the complexities of other key interventions that play a role in burden reduction

and elimination, the geospatial heterogeneity in cost-effectiveness and optimized distribution of

intervention packages on a sub-national level, or related pricing dynamics (Conteh et al., 2021,

Drake et al., 2017). For *P. vivax* in particular, reducing the global economic burden and

achieving elimination will depend on incorporation of hypnozoitocidal treatment and G6PD

screening into case management (Devine et al., 2021). Furthermore, for both parasites,

intervention strategies generally become more focal as transmission decreases, with targeted

surveillance and response strategies prioritized over widespread vector control. Therefore,

policy decisions should additionally be based on analysis of country-specific contexts, and our

findings are not informative for individual country allocation decisions. Results do however

account for non-linearities in the relationship between ITN distribution and usage to represent

changes in cost as a country moves from control to elimination: interventions that are effective

in malaria control settings, such as widespread vector control, may be phased out or limited in

favor of more expensive active surveillance and a focus on confirmed diagnoses and at-risk populations (Shretta et al., 2017). We also assumed that transmission settings are independent of each other, and did not allow for the possibility of re-introduction of disease, such as has occurred throughout the Eastern Mediterranean from imported cases (World Health Organization Regional Office for the Eastern Mediterranean, 2007). While our analysis presents allocation strategies to progress towards eradication, the results do not provide insight into allocation of funding to maintain elimination. In practice, the threat of malaria resurgence has important implications for when to scale back interventions.

Our analysis demonstrates the most impactful allocation of a global funding portfolio for ITNs to reduce global malaria cases. Unifying all funding sources in a global strategic allocation framework as presented here requires international donor allocation decisions to account for available domestic resources. National governments of endemic countries contribute 31% of all malaria-directed funding globally (World Health Organization, 2020), and government financing is a major source of malaria spending in near-elimination countries in particular (Haakenstad et al., 2019). Within the wider political economy which shapes the funding landscape and priority setting, there remains substantial scope for optimizing allocation decisions, including improving efficiency of within-country allocation of malaria interventions. Subnational malaria elimination in localized settings within a country can also provide motivation for continued elimination in other areas and friendly competition between regions to boost global elimination efforts (Lindblade and Kachur, 2020). Although more efficient allocation cannot fully compensate for projected shortfalls in malaria funding, mathematical modelling can aid efforts in determining optimal approaches to achieve the largest possible impact with available resources.

MATERIALS AND METHODS

Transmission models

We used deterministic compartmental versions of two previously published individual-based

transmission models of *P. falciparum* and *P. vivax* malaria to estimate the impact of varying ITN

usage on clinical incidence in different transmission settings. The *P. falciparum* model has

previously been fitted to age-stratified data from a variety of sub-Saharan African settings to

recreate observed patterns in parasite prevalence ($PfPR_{2-10}$), the incidence of clinical disease,

immunity profiles, and vector components relating rainfall, mosquito density, and the

entomological inoculation rate (EIR) (Griffin et al., 2016). We developed a deterministic version

of an existing individual-based model of *P. vivax* transmission, originally calibrated to data from

Papua New Guinea but also shown to reproduce global patterns of *P. vivax* prevalence and

clinical incidence (White et al., 2018). Models for both parasite species are structured by age

and heterogeneity in exposure to mosquito bites, and account for human immunity patterns.

They model mosquito transmission and population dynamics, and the impact of scale-up of

ITNs in identical ways. Full assumptions, mathematical details and parameter values can be

found in the Supplementary Material and in previous publications (Griffin et al., 2010, Griffin et

al., 2014, Griffin et al., 2016, White et al., 2018).

Data sources

We calibrated the model to baseline transmission intensity in all malaria-endemic countries

before scale-up of interventions, using the year 2000 as an indicator of these levels in line with

the current allocation approach taken by the Global Fund (The Global Fund, 2019). Annual EIR

was used as a measure of parasite transmission intensity, representing the rate at which people

are bitten by infectious mosquitoes. We simulated models to represent a wide range of EIRs for

P. falciparum and *P. vivax*. These transmission settings were matched to 2000 country-level

prevalence data resulting in EIRs of 0.001-80 for *P. falciparum* and 0.001-1.3 for *P. vivax*. *P.*

falciparum estimates came from parasite prevalence in children aged 2-10 years and *P. vivax*

prevalence estimates came from light microscopy data across all ages, based on standard

reporting for each species (Weiss et al., 2019, Battle et al., 2019). The relationship between

parasite prevalence and EIR for specific countries is shown in **Figures S5 and S6**. In each

country, the population at risk for *P. falciparum* and *P. vivax* malaria was obtained by summing

WorldPop gridded 2000 global population estimates (Tatem, 2017) within Malaria Atlas Project

transmission spatial limits using geoboundaries (Runfola et al., 2020) (Supplementary Materials:

Country-level data). The analysis was conducted on the national level, since this scale also

applies to funding decisions made by international donors (The Global Fund, 2019). As this

exercise represents a simplification of reality, population sizes were held constant, and

projected population growth is not reflected in the number of cases and the population at risk in

different settings. Seasonality was also not incorporated in the model, as EIRs are matched to

annual prevalence estimates and the effects of seasonal changes are averaged across the time

frame captured. For all analyses, countries were grouped according to their EIR, resulting in a

range of transmission settings compatible with the global distribution of malaria. Results were

further summarized by grouping EIRs into broader transmission intensity settings according to

WHO prevalence cut-offs of 0-1%, 1-10% 10-35% and ≥35% (World Health Organization,

2022a). This corresponded approximately to classifying EIRs of less than 0.1, 0.1 to 1, 1 to 7

and 7 or higher as very low, low, moderate and high transmission intensity, respectively.

Interventions

In all transmission settings, we simulated the impact of varying coverages of ITNs on clinical

incidence. While most countries implement a package of combined interventions, to reduce the

computational complexity of the optimization we considered the impact of ITN usage alone in

addition to 40% treatment of clinical disease. ITNs are a core intervention recommended for

large-scale deployment in areas with ongoing malaria transmission by WHO (Winskill et al.,

2019, World Health Organization, 2022a) and funding for vector control represents much of

global investments required for malaria control and elimination (Patouillard et al., 2017).

Modelled coverages represent population ITN usage between 0 and 80%, with the upper limit

reflective of common targets for universal access (Koenker et al., 2018). In each setting, the

models were run until clinical incidence stabilized at a new equilibrium with the given ITN usage.

Previous studies have shown that, as population coverage of ITNs increases, the marginal cost

of distribution increases as well (Bertozzi-Villa et al., 2021). We incorporated this non-linearity in

costs by estimating the annual ITN distribution required to achieve the simulated population

usage based on published data from across Africa, assuming that nets would be distributed on

a 3-yearly cycle and accounting for ITN retention over time (Supplementary Material). The cost

associated with a given simulated ITN usage was calculated by multiplying the number of nets

distributed per capita per year by the population size and by the unit cost of distributing an ITN,

assumed to be $3.50 (Sherrard-Smith et al., 2022).

Optimization

The optimal funding allocation for case reduction was determined by finding the allocation of

ITNs b across transmission settings that minimizes the total number of malaria cases at

equilibrium. Case totals were calculated as the sum of the product of clinical incidence $cinc_i$ and

the population p_i in each transmission setting i. Simultaneous optimization for case reduction

and pre-elimination was implemented with an extra weighting term in the objective function,

corresponding to a reduction in total remaining cases by a proportion w of the total cases

averted by the ITN allocation, C. This therefore represents a positive contribution for each

setting reaching the pre-elimination phase. The weighting on pre-elimination compared to case

reduction was 0 in the scenario optimized for case reduction, and varied between 0.5 and 100

times in the other optimization scenarios. Resource allocation must respect a budget constraint,

which requires that the sum of the cost of the ITNs distributed cannot exceed the initial budget

B, with b_i the initial number of ITNs distributed in setting i and c the cost of a single pyrethroid-

treated net. The second constraint requires that the ITN usage $b_i^\star$ must be between 0 and 80%

(Koenker et al., 2018), with ITN usage being a function of ITNs distributed, as shown in the

following equation.

$$\min_{b \in \mathbb{R}^n} \left[\sum_i^n cinc_i * p_i - w * C * \sum_{i=1}^n j_i \right]$$

$$\text{s.t.} \qquad \sum_{i=1}^n b_i * c \leq B$$

$$0 \leq b_i^\star \leq 0.8 \ \forall \ i = 1, \dots, n$$

$$C = Cases \ at \ baseline - \sum_i^n cinc_i * p_i$$

$$j_i = \begin{cases} 1, & cinc_i < 1/1000 \\ 0, & cinc_i \geq 1/1000 \end{cases}$$

$$b_i^\star = f(b_i)$$

$$\text{for all } i = 1, \dots, n$$

9
10
The optimization was undertaken using generalized simulated annealing (Xiang et al., 2013).

We included a penalty term in the objective function to incorporate linear constraints. Further

details can be found in the Supplementary Material.

The optimal allocation strategy for minimizing cases was also examined over a period of 39

years using the *P. falciparum* model, comparing a single allocation of a constant ITN usage to

minimize clinical incidence at 39 years, to reallocation every 3 years (similar to Global Fund

allocation periods (The Global Fund, 2016)) leading to varying ITN usage over time. At the

beginning of each 3-year period, we determined the optimized allocation of resources to be held

fixed until the next round of funding, with the objective of minimizing 3-year global clinical

incidence. Once *P. falciparum* elimination is reached in a given setting, ITN distribution is

discontinued, and in the next period the same total budget *B* will be distributed among the

remaining settings. We calculated the total budget required to minimize case numbers at 39

years and compared the impact of re-allocating each 3 years with a one-time allocation of 25%,

50%, 75% and 100% of the budget. To ensure computational feasibility, 39 years was used as it

was the shortest time frame over which the effect of re-distribution of funding from countries

having achieved elimination could be observed.

Analysis

We compared the impact of the two optimal allocation strategies (scenario 1A and 1B) and three

additional allocation scenarios on global malaria cases and the global population at risk.

Modelled scenarios are shown in **Table 2**. Scenarios 1C-1E represent existing policy strategies

that involve prioritizing high-transmission settings, prioritizing low-transmission (near-

elimination) settings, or resource allocation proportional to disease burden in the year 2000.

Global malaria case burden and the population at risk were compared between baseline levels

in 2000 and after reaching an endemic equilibrium under each scenario for a given budget.

Certification of malaria elimination requires proof that the chain of indigenous malaria

transmission has been interrupted for at least 3 years and a demonstrated capacity to prevent

return transmission (World Health Organization, 2018). In our analysis, transmission settings

were defined as having reached malaria elimination once less than one case remained per the

setting's total population. Once a setting reaches elimination, the entire population is removed

from the global total population at risk, representing a 'shrink the map' strategy. The pre-

elimination phase was defined as having reached less than 1 case per 1000 persons at risk in a

setting (Mendis et al., 2009).

All strategies were evaluated at different budgets ranging from 0 to the minimum investment

required to achieve the lowest possible number of cases in the model (noting that ITNs alone

are not predicted to eradicate malaria in our model). No distinctions were made between

national government spending and international donor funding, as the purpose of the analysis

was to look at resource allocation and not to recommend specific internal and external funding

choices.

All analyses were conducted in R v. 4.0.5 (R Foundation for Statistical Computing, Vienna,

Austria). The sf (v. 0.9-8, Pebesma 2018), raster (v. 3.4-10, Hijmans & van Etten 2012), and

terra (v.1.3-4, Hijmans 2021) packages were used for spatial data manipulation. The akima

package (v.0.6-2.2, Akima and Gebhardt 2021) was used for surface development, and the

GenSA package (v.1.1.7, Gubian et al.) for model optimization.

1 **FIGURES AND TABLES**

2
Figure 1. Global distribution of *P. falciparum* and *P. vivax* malaria burden in 2000 (in the

absence of insecticide-treated nets) obtained from the Malaria Atlas Project (Weiss et al., 2019,

Battle et al., 2019). A) The annual number of clinical cases and B) the population at risk of

malaria across settings with different transmission intensities and endemic for *P. falciparum*, *P.*

vivax or co-endemic for both species. The number of countries in each setting is indicated below

the figure.

2 **Figure 2.** Modelled impact of insecticide-treated net (ITN) usage on malaria epidemiology by

3 the setting-specific transmission intensity, represented by the baseline entomological inoculation

4 rate. The impact on the clinical incidence and prevalence of *P. falciparum* malaria (panels A and

5 B) and on the clinical incidence and prevalence of *P. vivax* malaria (panels C and D) is shown.

6 Panels A and C represent the clinical incidence for all ages.

Figure 3. Global clinical cases and population at risk of malaria under different allocation strategies at varying budgets. The impact on total malaria cases (panel A), total population at risk (panel B), individual *P. falciparum* and *P. vivax* cases (panel C) and population at risk of either species (panel D) are shown. Budget levels range from 0, representing no usage of insecticide-treated nets, to the budget required to achieve the maximum possible impact. Optimizing for case reduction generally leads to declining populations at risk as the budget increases, but this is not guaranteed due to the possibility of redistribution of funding between settings to minimize cases. The strategy optimizing case reduction and pre-elimination shown here places the same weighting (1:1) on reaching pre-elimination in a setting as on averting total cases, but conclusions were the same for weights of 0.5-100 on pre-elimination.

2 **Figure 4**. Optimal strategy for funding allocation across settings to minimize malaria case

3 burden at varying budgets. Panels show optimized allocation patterns across settings of

4 different transmission intensity (panels A and B) and different endemic parasite species (panels

5 C and D). The proportion of the total budget allocated to each setting (panels A and C) and the

6 resulting mean population usage of insecticide-treated nets (ITNs) (panels B and D) are shown.

Table 1. Relative reduction in malaria cases and population at risk under different allocation

strategies. Reductions are shown relative to the baseline of 321 million clinical cases and 4.1

billion persons at risk in the absence of interventions. Low, intermediate and high budget levels

represent 25%, 50% and 75% of the maximum budget, respectively. The strategy optimizing

case reduction and pre-elimination shown here places the same weighting (1:1) on reaching

pre-elimination in a setting as on averting total cases.

SCENARIO	Budget level	Clinical cases		Population at risk	
		Number (millions)	Relative reduction (%)	Number (billions)	Relative reduction (%)
OPTIMIZED FOR CASE REDUCTION	Low	136.1	58	4.1	0
	Intermediate	77.0	76	3.9	6
	High	53.9	83	2.4	42
	Maximum	41.5	87	0.4	91
OPTIMIZED FOR CASE REDUCTION & PRE-ELIMINATION	Low	161.3	50	4.0	3
	Intermediate	87.8	73	3.5	16
	High	58.8	82	1.8	58
	Maximum	41.5	87	0.4	91
PRIORITIZE HIGH-TRANSMISSION SETTINGS	Low	153.9	52	4.0	2
	Intermediate	109.5	66	3.4	17
	High	61.8	81	3.3	19
	Maximum	41.5	87	0.4	91
PROPORTIONAL ALLOCATION	Low	166.9	48	4.1	1
	Intermediate	150.4	53	4.0	4
	High	123.8	61	4.0	4
	Maximum	116.0	64	3.9	5
PRIORITIZE LOW-TRANSMISSION SETTINGS	Low	268.2	17	2.3	44
	Intermediate	245.2	24	1.8	56
	High	202.1	37	0.5	88
	Maximum	41.5	87	0.4	91

Table 2. Overview of modelled scenarios for allocation of funding to different transmission settings. Strategies 1A-1E compare resource allocation scenarios using clinical incidence values from each transmission setting at equilibrium after ITN coverage has been introduced. Strategies 2A-2B are compared as part of the allocation over time sub-analysis. EIR: entomological inoculation rate.

	STRATEGY	MODELLING APPROACH/ASSUMPTIONS
1A	Optimized for total malaria case reduction	Generalized simulated annealing is used to determine the optimal allocation of a given budget to minimize the total number of global malaria cases.
1B	Optimized for total malaria case reduction and pre-elimination	Generalized simulated annealing is used to determine the optimal allocation of a given budget to minimize the total number of global malaria cases while placing a premium on the pre-elimination phase being reached in a setting.
1C	Prioritize high-transmission settings	Funding is allocated to groups of countries according to transmission intensity (*P. falciparum* + *P. vivax* EIR). For a given budget, the transmission settings with highest EIR are prioritized, increasing ITN coverage in increments of 1% in each setting until malaria is eliminated or until an increase in coverage leads to no further decrease in cases, before allocating to the next-highest EIR setting.
1D	Prioritize low-transmission (near-elimination) settings	Funding is allocated to groups of countries according to transmission intensity (*P. falciparum* + *P. vivax* EIR). For a given budget, the transmission settings with lowest EIR are prioritized, increasing ITN coverage in increments of 1% in each setting until malaria is eliminated or until an increase in coverage leads to no further decrease in cases, before allocating to the next-lowest EIR setting.
1E	Proportional allocation	Funding is allocated to groups of countries in proportion to their disease burden. Budget shares are calculated using country data from the World Malaria Report (World Health Organization, 2020) and account for the country-specific total malaria cases (*P. falciparum* and *P. vivax*), deaths, incidence and mortality rate in 2000-2004, scaled by the subsequent increase in the population at risk (The Global Fund, 2019).
2A	One-time optimized allocation for *P. falciparum* case reduction	Generalized simulated annealing is used to determine the optimized allocation at a given budget, minimizing the total number of global *P. falciparum* cases after 39 years, resulting in constant ITN usage in each setting over this time period.
2B	Optimized allocation every three years for *P. falciparum* case reduction	Generalized simulated annealing is used to determine the optimized allocation at a given budget, minimizing the total number of global *P. falciparum* cases after every 3-year period for 39 years, allowing ITN usage to

vary in each setting every 3 years.

FUNDING

This work was supported by the Wellcome Trust [reference 220900/Z/20/Z]. NS, HMT, MP,

GDC, PW, KH and ACG also acknowledge funding from the MRC Centre for Global Infectious

Disease Analysis [reference MR/R015600/1], jointly funded by the UK Medical Research

Council (MRC) and the UK Foreign, Commonwealth & Development Office (FCDO), under the

MRC/FCDO Concordat agreement and is also part of the EDCTP2 programme supported by the

European Union. KH also acknowledges funding by Community Jameel. Disclaimer: "The views

expressed are those of the author(s) and not necessarily those of the NIHR, the UK Health

Security Agency or the Department of Health and Social Care." For the purpose of open access,

the authors have applied a 'Creative Commons Attribution' (CC BY) license to any Author

Accepted Manuscript version arising from this submission.

AUTHOR CONTRIBUTIONS

N. S, H. M. T., M. P., G. D. C., K. H. and A. C. G. conceived the study. N. S., H. M. T., and M. P.

processed the data, conducted the analysis, and created the visualizations. N. S., H. M. T., M.

P. and G. D. C. developed the computational infrastructure and methodology. P. W., M. T. W.,

K. H. and A. C. G. contributed domain knowledge. K. H. and A. C. G. acquired funding for the

study and provided guidance on analysis and interpretation. N. S. and H. M. T. wrote the first

draft of the manuscript. All authors contributed to interpreting the results, reviewing and revising

the manuscript, and approved the final version for submission.

CONFLICTS OF INTEREST

No reported conflicts of interest.

REFERENCES

BARRENHO, E., MIRALDO, M., SHAIKH, M. & ATUN, R. 2017. Vertical and horizontal equity of funding for malaria control: a global multisource funding analysis for 2006–2010. *BMJ Global Health,* 2**,** e000496-e000496.

BATTLE, K. E., LUCAS, T. C. D., NGUYEN, M., HOWES, R. E., NANDI, A. K., TWOHIG, K. A., PFEFFER, D. A., CAMERON, E., RAO, P. C., CASEY, D., GIBSON, H. S., ROZIER, J. A., DALRYMPLE, U., KEDDIE, S. H., COLLINS, E. L., HARRIS, J. R., GUERRA, C. A., THORN, M. P., BISANZIO, D., FULLMAN, N., HUYNH, C. K., KULIKOFF, X., KUTZ, M. J., LOPEZ, A. D., MOKDAD, A. H., NAGHAVI, M., NGUYEN, G., SHACKELFORD, K. A., VOS, T., WANG, H., LIM, S. S., MURRAY, C. J. L., PRICE, R. N., BAIRD, J. K., SMITH, D. L., BHATT, S., WEISS, D. J., HAY, S. I. & GETHING, P. W. 2019. Mapping the global endemicity and clinical burden of Plasmodium vivax, 2000–17: a spatial and temporal modelling study. *The Lancet,* 394**,** 332-343.

BERTOZZI-VILLA, A., BEVER, C. A., KOENKER, H., WEISS, D. J., VARGAS-RUIZ, C., NANDI, A. K., GIBSON, H. S., HARRIS, J., BATTLE, K. E., RUMISHA, S. F., KEDDIE, S., AMRATIA, P., ARAMBEPOLA, R., CAMERON, E., CHESTNUTT, E. G., COLLINS, E. L., MILLAR, J., MISHRA, S., ROZIER, J., SYMONS, T., TWOHIG, K. A., HOLLINGSWORTH, T. D., GETHING, P. W. & BHATT, S. 2021. Maps and metrics of insecticide-treated net access, use, and nets-per-capita in Africa from 2000-2020. *Nature Communications,* 12**,** 3589.

BOUSEMA, T., GRIFFIN, J. T., SAUERWEIN, R. W., SMITH, D. L., CHURCHER, T. S., TAKKEN, W., GHANI, A., DRAKELEY, C. & GOSLING, R. 2012. Hitting Hotspots: Spatial Targeting of Malaria for Control and Elimination. *PLOS Medicine,* 9**,** e1001165-e1001165.

CHEN, I., COONEY, R., FEACHEM, R. G. A., LAL, A. & MPANJU-SHUMBUSHO, W. 2018. The Lancet Commission on malaria eradication. *The Lancet,* 391**,** 1556-1558.

CONTEH, L., SHUFORD, K., AGBORAW, E., KONT, M., KOLACZINSKI, J. & PATOUILLARD, E. 2021. Costs and cost-effectiveness of malaria control interventions: a systematic literature review. *Value in Health,* 24**,** 1213-1222.

DEVINE, A., BATTLE, K. E., MEAGHER, N., HOWES, R. E., DINI, S., GETHING, P. W., SIMPSON, J. A., PRICE, R. N. & LUBELL, Y. 2021. Global economic costs due to vivax malaria and the potential impact of its radical cure: A modelling study. *PLOS Medicine,* 18**,** e1003614-e1003614.

DRAKE, T. L., LUBELL, Y., KYAW, S. S., DEVINE, A., KYAW, M. P., DAY, N. P. J., SMITHUIS, F. M. & WHITE, L. J. 2017. Geographic Resource Allocation Based on Cost Effectiveness: An Application to Malaria Policy. *Appl Health Econ Health Policy,* 15**,** 299-306.

FEACHEM, R. G. A., PHILLIPS, A. A., HWANG, J., COTTER, C., WIELGOSZ, B., GREENWOOD, B. M., SABOT, O., RODRIGUEZ, M. H., ABEYASINGHE, R. R., GHEBREYESUS, T. A. & SNOW, R. W. 2010. Shrinking the malaria map: progress and prospects. *The Lancet,* 376**,** 1566-1578.

GRIFFIN, J. T., BHATT, S., SINKA, M. E., GETHING, P. W., LYNCH, M., PATOUILLARD, E., SHUTES, E., NEWMAN, R. D., ALONSO, P. & CIBULSKIS, R. E. 2016. Potential for reduction of burden and local elimination of malaria by reducing Plasmodium falciparum malaria transmission: a mathematical modelling study. *The Lancet Infectious Diseases,* 16**,** 465-472.

GRIFFIN, J. T., FERGUSON, N. M. & GHANI, A. C. 2014. Estimates of the changing age-burden of Plasmodium falciparum malaria disease in sub-Saharan Africa. *Nature communications,* 5**,** 3136.

GRIFFIN, J. T., HOLLINGSWORTH, T. D., OKELL, L. C., CHURCHER, T. S., WHITE, M., HINSLEY, W., BOUSEMA, T., DRAKELEY, C. J., FERGUSON, N. M., BASÁÑEZ, M. G. & GHANI, A. C. 2010. Reducing Plasmodium falciparum malaria transmission in Africa: a model-based evaluation of intervention strategies. *PLoS Medicine,* 7.

HAAKENSTAD, A., HARLE, A. C., TSAKALOS, G., MICAH, A. E., TAO, T., ANJOMSHOA, M., COHEN, J., FULLMAN, N., HAY, S. I., MESTROVIC, T., MOHAMMED, S., MOUSAVI, M., NIXON, M. R., PIGOTT, D., TRAN, K., MURRAY, C. J. L. & DIELEMAN, J. L. 2019. Tracking spending on malaria by source in 106 countries, 2000-16: an economic modelling study. *The Lancet Infectious Diseases,* 19**,** 703-716.

HAY, S. I., GUERRA, C. A., TATEM, A. J., NOOR, A. M. & SNOW, R. W. 2004. The global distribution and population at risk of malaria: past, present, and future. *Lancet Infect Dis,* 4**,** 327-36.

KOENKER, H., ARNOLD, F., BA, F., CISSE, M., DIOUF, L., ECKERT, E., ERSKINE, M., FLOREY, L., FOTHERINGHAM, M., GERBERG, L., LENGELER, C., LYNCH, M., MNZAVA, A., NASR, S., NDIOP, M., POYER, S., RENSHAW, M., SHARGIE, E., TAYLOR, C., THWING, J., VAN HULLE, S., YE, Y., YUKICH, J. & KILIAN, A. 2018. Assessing whether universal coverage with insecticide-treated nets has been achieved: is the right indicator being used? *Malaria Journal,* 17**,** 1-11.

LINDBLADE, K. A. & KACHUR, S. P. 2020. Opportunities for Subnational Malaria Elimination in High-Burden Countries. *The American Journal of Tropical Medicine and Hygiene,* 103**,** 2153-2153.

MENDIS, K., RIETVELD, A., WARSAME, M., BOSMAN, A., GREENWOOD, B. & WERNSDORFER, W. H. 2009. From malaria control to eradication: The WHO perspective. *Trop Med Int Health,* 14**,** 802-9.

NEWBY, G., BENNETT, A., LARSON, E., COTTER, C., SHRETTA, R., PHILLIPS, A. A. & FEACHEM, R. G. A. 2016. The path to eradication: a progress report on the malaria-eliminating countries. *The Lancet,* 387**,** 1775-1784.

PATOUILLARD, E., GRIFFIN, J., BHATT, S., GHANI, A. & CIBULSKIS, R. 2017. Global investment targets for malaria control and elimination between 2016 and 2030. *BMJ global health,* 2**,** e000176.

PRICE, R. N., COMMONS, R. J., BATTLE, K. E., THRIEMER, K. & MENDIS, K. 2020. Plasmodium vivax in the Era of the Shrinking P. falciparum Map. *Trends Parasitol,* 36**,** 560-570.

RAINE, R., OR, Z., PRADY, S. & BEVAN, G. 2016. Challenges, solutions and future directions in the evaluation of service innovations in health care and public health. Essay 5 - Evaluating health-care equity. *Health Services and Delivery Research,* 4.

RUNFOLA, D., ANDERSON, A., BAIER, H., CRITTENDEN, M., DOWKER, E., FUHRIG, S., GOODMAN, S., GRIMSLEY, G., LAYKO, R., MELVILLE, G., MULDER, M., OBERMAN, R., PANGANIBAN, J., PECK, A., SEITZ, L., SHEA, S., SLEVIN, H., YOUNGERMAN, R. & HOBBS, L. 2020. geoBoundaries: A global database of political administrative boundaries. *PLOS ONE,* 15**,** e0231866-e0231866.

SHERRARD-SMITH, E., WINSKILL, P., HAMLET, A., NGUFOR, C., N'GUESSAN, R., GUELBEOGO, M. W., SANOU, A., NASH, R. K., HILL, A., RUSSELL, E. L., WOODBRIDGE, M., TUNGU, P., KONT, M. D., MCLEAN, T., FORNADEL, C., RICHARDSON, J. H., DONNELLY, M. J., STAEDKE, S. G., GONAHASA, S., PROTOPOPOFF, N., ROWLAND, M. & CHURCHER, T. S. 2022. Optimising the deployment of vector control tools against malaria: a data-informed modelling study. *The Lancet Planetary Health,* 6**,** e100-e109.

SHRETTA, R., LIU, J., COTTER, C. & AL, E. 2017. Malaria Elimination and Eradication. *In:* HOLMES, K. K., BERTOZZI, S., BLOOM, B. & AL, E. (eds.) *Major Infectious Diseases.*

3rd ed. Washington, DC: The International Bank for Reconstruction and Development / The World Bank.

SNOW, R. W. 2015. Global malaria eradication and the importance of Plasmodium falciparum epidemiology in Africa. *BMC Med,* 13**,** 23.

SNOW, R. W., OKIRO, E. A., GETHING, P. W., ATUN, R. & HAY, S. I. 2010. Equity and adequacy of international donor assistance for global malaria control: an analysis of populations at risk and external funding commitments. *The Lancet,* 376**,** 1409-1416.

STRATEGIC ADVISORY GROUP ON MALARIA ERADICATION 2020. Malaria eradication: benefits, future scenarios & feasibility. Geneva: World Health Organization.

TATEM, A. J. 2017. WorldPop, open data for spatial demography. *Scientific Data.* Nature Publishing Groups.

THE GLOBAL FUND 2016. The Global Fund Strategy 2017-2022: Investing to end epidemics. Geneva: The Global Fund.

THE GLOBAL FUND. 2019. *Description of the 2020-2022 Allocation Methodology* [Online]. Available: https://www.theglobalfund.org/media/9224/fundingmodel_2020-2022allocations_methodology_en.pdf [Accessed August 30, 2023].

THE GLOBAL FUND 2021. Approval of the Global Fund Strategy Narrative. 46th Board Meeting. GF/B46/03 Revision 1. Geneva: The Global Fund.

UNITED STATES AGENCY FOR INTERNATIONAL DEVELOPMENT & CENTERS FOR DISEASE CONTROL AND PREVENTION 2021. End Malaria Faster - U.S. President's Malaria Initiative Strategy 2021-2026. Washington, DC.

WEISS, D. J., LUCAS, T. C., NGUYEN, M., NANDI, A. K., BISANZIO, D., BATTLE, K. E., CAMERON, E., TWOHIG, K. A., PFEFFER, D. A. & ROZIER, J. A. 2019. Mapping the global prevalence, incidence, and mortality of Plasmodium falciparum, 2000–17: a spatial and temporal modelling study. *The Lancet,* 394**,** 322-331.

WHITE, M. T., WALKER, P., KARL, S., HETZEL, M. W., FREEMAN, T., WALTMANN, A., LAMAN, M., ROBINSON, L. J., GHANI, A. & MUELLER, I. 2018. Mathematical modelling of the impact of expanding levels of malaria control interventions on Plasmodium vivax. *Nature Communications,* 9**,** 3300.

WINSKILL, P., WALKER, P. G., CIBULSKIS, R. E. & GHANI, A. C. 2019. Prioritizing the scale-up of interventions for malaria control and elimination. *Malaria Journal,* 18**,** 1-11.

WORLD HEALTH ORGANIZATION 2013. Briefing note - How should funds for malaria control be spent when there are not enough? *Malaria Policy Advisory Committee to the WHO: March 2013 meeting.* Geneva: World Health Organization.

WORLD HEALTH ORGANIZATION 2015. Global Technical Strategy for Malaria 2016-2030. Geneva: World Health Organization.

WORLD HEALTH ORGANIZATION 2018. Update on the e-2020 initiative of 21 malaria-eliminating countries: report and country briefs. Geneva: World Health Organization.

WORLD HEALTH ORGANIZATION 2019. High burden to high impact: a targeted malaria response. Geneva: World Health Organization.

WORLD HEALTH ORGANIZATION 2020. World malaria report 2020. Geneva: World Health Organization.

WORLD HEALTH ORGANIZATION 2022a. WHO Guidelines for malaria, 31 March 2022. Geneva: World Health Organization.

WORLD HEALTH ORGANIZATION 2022b. World malaria report 2022. Geneva: World Health Organization.

WORLD HEALTH ORGANIZATION REGIONAL OFFICE FOR THE EASTERN MEDITERRANEAN 2007. Guidelines on prevention of the reintroduction of malaria. *EMRO Technical Publications Series 34.* Cairo: World Health Organization.

XIANG, Y., GUBIAN, S., SUOMELA, B. & HOENG, J. 2013. Generalized simulated annealing for global optimization: the GenSA package. *The R Journal,* 5**,** 13-28.