

1 Superior antibody immunogenicity of a RH5 blood-stage malaria vaccine in 2 Tanzanian infants as compared to adults

3
4 Sarah E. Silk^{1,2,3}, Wilmina F. Kalinga⁴, Ivanny M. Mtaka⁴, Nasoro S. Lilolime⁴, Maximillian Mpina⁴,
5 Florence Milando⁴, Saumu Ahmed⁴, Ababacar Diouf⁵, Fatuma Mkwepu⁴, Beatus Simon⁴, Thabit
6 Athumani⁴, Mohammed Rashid⁴, Latipha Mohammed⁴, Omary Lweno⁴, Ali M. Ali⁴, Gloria Nyaulingo⁴,
7 Bakari Mwalimu⁴, Sarah Mswata⁴, Tunu G. Mwamlima⁴, Jordan R. Barrett^{1,2,3}, Lawrence T. Wang^{1,3}, Yrene
8 Themistocleous², Lloyd D. W. King^{1,2,3}, Susanne H. Hodgson^{1,2,3}, Ruth O. Payne², Carolyn M. Nielsen^{1,2,3},
9 Alison M. Lawrie², Fay L. Nugent^{1,2}, Jee-Sun Cho^{1,2,3}, Carole A. Long⁵, Kazutoyo Miura⁵, Simon J.
10 Draper^{1,2,3,*}, Angela M. Minassian^{1,2,3,*†}, and Ally I. Olotu^{4,*}.

11
12 ¹ Department of Biochemistry, University of Oxford, Dorothy Crowfoot Hodgkin Building, Oxford, OX1
13 3QU, UK.

14 ² Centre for Clinical Vaccinology and Tropical Medicine, Jenner Institute, University of Oxford, Old
15 Road Campus, Oxford, OX3 7LE, UK.

16 ³ Kavli Institute for Nanoscience Discovery, Dorothy Crowfoot Hodgkin Building, University of Oxford,
17 Oxford, OX1 3QU, UK.

18 ⁴ Interventions and Clinical Trials Department, Ifakara Health Institute, P.O. Box 74, Bagamoyo,
19 Tanzania.

20 ⁵ Laboratory of Malaria and Vector Research, NIAID/NIH, Rockville, MD 20852, USA.

21 * These authors contributed equally.

22
23 † Corresponding author: Dr Angela M. Minassian; angela.minassian@bioch.ox.ac.uk

Silk SE *et al.*

- 24 This research was funded in part by the UK Medical Research Council (MRC) [Grant number:
25 MR/P020593/1]. For the purpose of Open Access, the author has applied a CC BY public copyright
26 licence to any Author Accepted Manuscript (AAM) version arising from this submission.

27 **Abstract**

28 **Background**

29 RH5 is the leading blood-stage candidate antigen for inclusion in a *Plasmodium falciparum* malaria
30 vaccine, however, its safety profile and ability to induce functional immune responses in a malaria-
31 endemic population are unknown. Characterising safety and immunogenicity is key to refine and progress
32 next-generation RH5-based blood-stage malaria vaccines to field efficacy assessment.

33 **Methods**

34 A Phase 1b, single-center, dose-escalation, age de-escalation, double-blind, randomized, controlled trial
35 was conducted in Bagamoyo, Tanzania. Healthy adults (18-35 years), young children (1-6 years) and
36 infants (6-11 months) were recruited to receive a priming dose of viral-vectored ChAd63 RH5 (or rabies
37 control vaccine) followed by a booster dose of MVA RH5 (or rabies control vaccine) 8 weeks later. The
38 primary outcomes were the number of solicited and unsolicited adverse events following vaccination and
39 the number of serious adverse events over the whole study period. Secondary outcomes included
40 quantitative and qualitative measures of the anti-RH5 immune response. All participants receiving at least
41 one dose of vaccine were included in the primary analyses.

42 **Findings**

43 Between 12th April and 25th October 2018 a total of 63 adults, children and infants were recruited and
44 primed and 60 of these were boosted, all completing six months of follow-up post-priming vaccination.
45 Vaccinations were well-tolerated with participants reporting predominantly mild reactogenicity, with
46 profiles comparable between ChAd63 RH5, MVA RH5 and rabies vaccine groups, and across the age
47 groups. No serious adverse events were reported during the study period. RH5-specific T cell, B cell and
48 serum antibody responses were induced by vaccination. Higher anti-RH5 serum IgG responses were
49 observed post-boost in the 1-6 year old children (median 93 µg/mL; range: 31-508 µg/mL) and infants
50 (median 149 µg/mL; range: 29-352 µg/mL) as compared to adults (median 14 µg/mL; range: 9-15
51 µg/mL). These contracted over time post-boost, but the same hierarchy of responses across the age groups

52 was maintained to end of follow-up at 16 weeks post-boost (day 168). Vaccine-induced anti-RH5
53 antibodies were functional showing growth inhibition activity (GIA) *in vitro* against *P. falciparum* blood-
54 stage parasites. The highest levels were observed in the 6-11 month old infants, with 6/11 showing >60%
55 GIA following dilution of total IgG to 2.5 mg/mL (median 61%; range: 41-78%).

56

57 **Interpretation**

58 The ChAd63-MVA RH5 vaccine regimen shows an acceptable safety and reactogenicity profile and
59 encouraging immunogenicity in children and infants residing in a malaria-endemic area. The levels of
60 functional GIA observed in the RH5 vaccinated 6-11 month old infants are the highest levels reported to-
61 date following human vaccination. These data support onward clinical development of RH5-based blood-
62 stage vaccines that aim to protect against clinical malaria in young African infants.

63

64 **Funding**

65 Medical Research Council, London, United Kingdom.

66

67 **Trial Registration**

68 ISRCTN registry: 47448832 and ClinicalTrials.gov: NCT03435874.

69 **Introduction**

70 Over the last two decades, expanded access to malaria prevention tools has led to a major reduction in the
71 global burden of this disease, however, progress has stalled with current data showing that cases and
72 deaths are now rising once again, likely magnified by the SARS-CoV-2 pandemic ¹. Notably, over 80%
73 of deaths occur in children under the age of 5 in sub-Saharan Africa and a highly effective vaccine
74 remains urgently needed. Encouragingly, subunit vaccine strategies targeting the invasive sporozoite
75 stage of *Plasmodium falciparum* are now showing moderate levels of efficacy in field trials ^{2,3}, however,
76 durability of protection remains a key challenge. These vaccines also necessitate sterilizing immunity,
77 with only a single break-through sporozoite leading to the subsequent pathogenic blood-stage of infection.
78 An alternative and complementary approach is to vaccinate against the blood-stage merozoite to inhibit
79 erythrocyte invasion, thus, leading to control and/or clearance of blood-stage parasitemia, minimizing
80 morbidity and mortality and reducing transmission ⁴. Indeed, combining a blood-stage vaccine component
81 with existing anti-sporozoite vaccines is now widely regarded as the most promising strategy to achieve a
82 high efficacy intervention.

83
84 While a number of factors have long-stalled progress in the design of vaccines that can impact the blood-
85 stage of the parasite's lifecycle ⁴, the identification of an essential, highly conserved, antibody-
86 susceptible, protein complex used by *P. falciparum* merozoites to invade erythrocytes has propelled the
87 field over the last 10 years ⁵. Vaccine development efforts are most advanced for one component of this
88 invasion complex: reticulocyte-binding protein homolog 5 (**RH5**) ⁶. This vaccine target binds basigin on
89 the erythrocyte surface ⁷, a receptor-ligand interaction that is critical for parasite invasion and its human
90 host tropism ⁸. Significant *in vivo* protection with RH5-based vaccines was first demonstrated against a
91 stringent blood-stage *P. falciparum* challenge in an *Aotus* monkey model ⁹. Here, protection was
92 correlated with functional antibody activity measured using an *in vitro* assay of growth inhibition activity

93 (GIA); an immune mechanism subsequently validated by passive transfer of anti-RH5 monoclonal
94 antibody ¹⁰.

95
96 This paved the way for the first RH5-based vaccine to enter Phase 1a clinical testing in healthy UK adults
97 in 2014 (VAC057; ClinicalTrials.gov: NCT02181088). This vaccine utilized recombinant replication-
98 deficient chimpanzee adenovirus serotype 63 (ChAd63) and the attenuated orthopoxvirus modified
99 vaccinia virus Ankara (MVA), delivered in an 8-week prime-boost regimen, to enable *in situ* expression
100 of RH5 by virally infected cells (ChAd63-MVA RH5). The vaccine was well-tolerated and induced
101 functional human antibodies that exhibited cross-strain *in vitro* GIA ¹¹. However, although the levels of
102 anti-RH5 serum immunoglobulin G (IgG) in the UK adult participants greatly exceeded the levels
103 observed in African adults following years of natural malaria exposure, they fell below the protective
104 immunological threshold predicted by the *Aotus* model. In parallel and in a series of other clinical trials,
105 immunization with the same ChAd63-prime MVA-boost viral-vectored delivery platform recombinant for
106 the liver-stage malaria antigen ME-TRAP, reported ten-fold higher antibody levels in West African
107 infants as compared to UK adults and West African adults ¹². Given deployment trials of the WHO pre-
108 qualified RTS,S/AS01 vaccine are focussed on infants starting from 5 months of age, and given that a ten-
109 fold higher anti-RH5 IgG antibody response in African infants could potentially translate into protection
110 against clinical malaria, we proceeded to further assess the ChAd63-MVA RH5 in this target population.

111
112 We therefore conducted a single-center, dose-escalation, age de-escalation, double-blind, randomized,
113 controlled Phase 1b trial to explore the tolerability, safety and immunogenicity of an RH5-based vaccine
114 in a malaria-endemic area for the first time, to inform crucial decisions on progress to field efficacy
115 studies and/or iterative vaccine refinement.

116 **Results**

117 **Study recruitment and vaccinations**

118 In total 125 participants were screened and 63 of these were enrolled into the VAC070 Phase 1b trial
119 (**Figure 1**). Vaccinations began on 12th April 2018 and all follow-up visits were completed by 13th
120 February 2019. Within each group, participants were randomized to receive the ChAd63-MVA RH5
121 vaccine or rabies control vaccine in a ratio of 2:1. Participants and all investigatory staff (involved in
122 evaluation of safety and immunogenicity endpoints) were blinded to vaccine allocation. All participants
123 received their immunizations as scheduled and completed 6 months of follow-up post priming
124 vaccination, apart from three individuals who withdrew from the study after the prime. Similar numbers
125 of males and females were enrolled across the younger age groups (**Table S1**). Independent safety
126 reviews were conducted by the Safety Monitoring Committee (SMC) between every age de-escalation
127 and/or dose-escalation step in the protocol.

128
129 All participants were recruited from the district of Bagamoyo, Tanzania. Malaria prevalence in the district
130 ranges from 15.4% in the western part to at times as low as zero in Bagamoyo town; the population
131 average prevalence was 13% in 2013¹³. The mean age of adult participants in Group 1 was 25.8 years
132 (range 19.0 – 30.7 years), and they received the full dose of both vaccines via the intramuscular (IM)
133 route: 5×10^{10} viral particles (vp) ChAd63 RH5 to prime on day 0 and 2×10^8 plaque-forming units (pfu)
134 MVA RH5 to boost on day 56. The mean age of children across Groups 2A and 2B was 3.6 years (range
135 1.2 – 5.6 years), and infants across Groups 3A and 3B was 9.4 months (range 5.9 – 11.9 months). Those
136 in Groups 2A and 3A received lower “lead-in” doses of vaccine: 1×10^{10} vp ChAd63 RH5 and 1×10^8
137 pfu MVA RH5, and those in Groups 2B and 3B received the full vaccine dose (all via the IM route).

138
139 A total of sixty-three adults, children and infants were recruited and primed and sixty of these were
140 boosted (three withdrew post-prime). Thirty-nine participants were primed with ChAd63 RH5 and

141 boosted with MVA RH5, twenty-one participants received two doses of rabies vaccine 8 weeks apart. All
142 boosted participants completed six months of follow-up post-prime and analysis was by original assigned
143 groups. All participants (RH5 and rabies vaccinated) were negative for malaria by blood film throughout
144 the trial (tested at screening and monthly intervals post-vaccination). We also conducted retrospective
145 analysis for malaria parasitemia by highly sensitive qPCR on blood samples taken from every participant
146 at the day 0, day 63 and day 84 time-points. All samples tested negative, except for only one transient low
147 level asymptomatic parasitemia of 450 parasites/mL blood in a single RH5 vaccinated Group 2B child at
148 the day 63 time-point (7 days post-MVA RH5 boost).

149

150 **Outcomes and estimation**

151 **Reactogenicity and safety**

152 There were no serious adverse events (SAEs), AEs of special interest (AESIs) or unexpected reactions
153 and no safety concerns during the course of the trial. The local and systemic reactogenicity profile of the
154 full dose RH5 vaccines (**Figure 2**) was similar, if not reduced, as compared to that seen in healthy UK
155 adults immunized with the identical vaccines¹¹. Following the prime with ChAd63 RH5, the majority of
156 participants across all age groups experienced local pain at the injection site and systemic fever (>37.5°C)
157 (**Figure 2A-C**); all of these adverse events (AEs) were mild in severity, with similar profiles seen in those
158 that received the rabies control vaccine (**Figure S1A-C**). A very similar pattern was observed following
159 the boost with MVA RH5 (**Figure 2D-F**), with some of the older participants also experiencing Grade 1
160 (mild) erythema, in contrast to the boost with the rabies vaccine where hardly any local or systemic AEs
161 were observed (**Figure S1D-F**). Very similar AE profiles were observed in the Group 2A children and
162 Group 3A infants who received the lower “lead-in” doses of the ChAd63-MVA RH5 vaccine or the rabies
163 control vaccine (**Figure S2**). The majority of solicited AEs occurred within the first two days after
164 vaccination and the median duration of each local or systemic AE was between 1 and 2 days following
165 either vaccine. Of the unsolicited AEs recorded within 28 days of vaccination, none were considered

166 related to any of the vaccines (**Table S2**). None were recorded in the adult group or following the
167 ChAd63 RH5 prime. Those recorded following MVA RH5 or rabies vaccination were mostly mild in
168 nature, with only a few graded moderate and none graded severe, and all resolved spontaneously. There
169 were no Grade 3 unsolicited AEs. Viral upper respiratory tract infection (N=15) was the most common
170 unsolicited AE among the infants 6-11 months of age with similar numbers observed in the MVA RH5
171 vaccine and rabies vaccine groups.

172
173 The most frequent laboratory abnormality identified in the 28-day post vaccination period was increased
174 leukocytes, present in 9/42 vaccinees and 3/18 controls ($P=0.74$, 2-tailed Fisher's exact test). These
175 included four Grade 3 occurrences in the 6-11 month old infants (all in the ChAd63-MVA RH5 group).
176 These elevations were associated with concurrent infections and resolved with treatment of the infection
177 (**Table S3**). Other commonly observed laboratory abnormalities included mild and moderate anemia, mild
178 increases in alanine aminotransferase (ALT) levels, mild decreases in leukocytes, and mild decreases in
179 lymphocytes. The frequencies of these abnormalities did not differ between the vaccinees and controls
180 ($P=0.74$, $P=0.65$, $P=0.66$ and $P>0.99$, respectively; 2-tailed Fisher's exact test). Full blood count analyses
181 at the end of trial follow-up were all normal.

182

183 **Immunogenicity**

184 **IFN- γ T cell responses induced by ChAd63 and MVA RH5 are higher in adult than in** 185 **children and infant participants**

186 The kinetics and magnitude of the RH5-specific T cell response were assessed over time by *ex-vivo* IFN- γ
187 ELISPOT following re-stimulation of PBMC with 20mer peptides overlapping by 10 amino acids (aa)
188 spanning the entire RH5 insert present in the vaccines (**Figure 3A,B, S3A-D, Table S4**). Following
189 ChAd63 RH5 prime, comparable responses were detectable in the Tanzanian adults and children on day
190 14 (median 271 [range 7 – 2723] and median 243 [range 60 – 1147] spot forming units (SFU) / million

191 PBMC, respectively). These were approximately 3-fold lower than the responses previously observed in
192 UK adults who received the identical dose of vaccine ¹¹, but significantly higher than those seen in the
193 Tanzanian infants (median 19 [range 0 – 745] SFU / million PBMC) who also mirrored the rabies vaccine
194 control recipients (**Figure 3A**). Subsequently, administration of MVA RH5 boosted these responses in the
195 Tanzanian adults, as measured four weeks later on day 84 with a median of 1159 [range 740 – 1935] SFU
196 / million PBMC. These responses were now comparable to those previously seen post-boost in the UK
197 adults ¹¹, but significantly higher than those in the Tanzanian children (where no appreciable boost in the
198 T cell response was observed) and the Tanzanian infants where a modest boost reached a median level of
199 125 [range 0 – 793] SFU / million PBMC (**Figure 3B**). As expected for T cell responses, these then
200 contracted over time, but the same trend as day 84 was also observed 16 weeks post-boost at the end of
201 study period (day 168), with responses in all three age groups maintained above baseline and above those
202 observed in the rabies vaccine control recipients (**Figure S3D**). However, as expected, we also observed
203 that peripheral lymphocyte counts decreased with age as measured by routine hematology tests at each
204 time point (**Figure 3C**). Therefore, as reported in a previous pediatric study of a viral vectored malaria
205 vaccine ¹², we also analyzed the data by incorporating the lymphocyte count into the calculation of T cell
206 responses, to facilitate a more physiologically relevant comparison of cellular immunity across the
207 different age groups. Here, only the 6-11 month old infants still showed a significantly lower response
208 following the ChAd63 RH5 prime (**Figure S3E**), whilst there was no significant difference across the age
209 groups after the MVA RH5 boost (**Figure 3D**) and the end of follow up (**Figure S3F**). Finally, we also
210 assessed the responses at day 14 and day 84 in the Tanzanian children and infants receiving the lower
211 “lead in” doses of ChAd63 and MVA RH5. Here, these T cell responses were also comparable to those
212 induced by the full dose of each vaccine (**Figure S3G,H**).

213

214 **Anti-RH5 serum IgG and B cell responses induced by ChAd63 and MVA RH5 are higher**
215 **in children and infants than in adults.**

216 We next measured the kinetics and magnitude of the anti-RH5 serum IgG antibody response over time by
217 ELISA against full-length RH5 (RH5_FL) recombinant protein (**Figure 4A,B, S4A-D**). Following
218 ChAd63 RH5 prime, the magnitude of the antibody response showed a clear age-dependent hierarchy.
219 Here, on day 14, the lowest responses were detected in the Tanzanian adults, with higher responses
220 observed in the children 1-6 years of age, and the highest in the infants 6-11 months of age; median levels
221 of serum anti-RH5 IgG were 0.2, 1.8 and 4.9 $\mu\text{g/mL}$, respectively (in contrast to negligible median
222 responses in all groups at baseline). Median responses in UK adults who previously received the identical
223 dose of the ChAd63 RH5 vaccine ¹¹ or in the rabies vaccine control recipients were also negligible at this
224 time-point (**Figure 4A**). Subsequently, administration of MVA RH5 boosted these responses in all three
225 age groups, but the same hierarchy was maintained as measured by ELISA one week later on day 63
226 (**Figure 4B**). Median responses of serum anti-RH5 IgG in the Tanzanian adults were 14 $\mu\text{g/mL}$, now
227 highly comparable to those previously observed in UK adults who received the identical dose of the viral
228 vaccines ¹¹; whilst significantly higher responses were observed in the children (~6-fold, median of 93
229 $\mu\text{g/mL}$) and infants (~10-fold, median of 149 $\mu\text{g/mL}$). Antibody responses induced by the vaccines were
230 also dose dependent; here, responses at day 14 and day 63 were significantly lower in the Tanzanian
231 infants receiving the lower “lead in” doses of ChAd63 and MVA RH5 as compared to those receiving the
232 full dose of each vaccine, with a similar trend observed in the children (**Figure S4E,F**). As expected for
233 serum IgG responses, these contracted over time post-boost, but the same hierarchy of responses across
234 the age groups was maintained out to 16 weeks post-boost at the end of study period (day 168) (**Figure**
235 **S4A**).

236
237 Having observed such high anti-RH5 serum IgG responses in children and infants, we hypothesized this
238 could be due to anti-vector immunity that may increase with age and thereby negatively affect the priming

239 immunogenicity of adenovirus-vectored vaccines in adults. However, although the 6-11 month old
240 Tanzanian infants did show the lowest anti-ChAd63 antibody responses at baseline (consistent with
241 previous data on antibody responses to human adenovirus serotypes in this age group¹⁴), the children
242 aged 1-6 years and adults in our study were comparable (**Figure 4C**). Consequently, there was no
243 significant correlation between existing anti-ChAd63 antibody responses at baseline and the day14
244 ChAd63 RH5 humoral immunogenicity (Spearman's $r_s = -0.35$, $P > 0.05$, $N=30$). In line with this, we
245 also measured the anti-rabies glycoprotein serum antibody responses in the control participants. Here we
246 observed a similar age-dependent hierarchy, despite the small number of vaccinees, with responses in the
247 children and infants on average ~4-5-fold higher than in adults after the first and second immunizations
248 (**Figure 4D, S4G,H**).

249
250 We therefore next measured the underlying B cell response. Previous studies have shown that antibody-
251 secreting cells (ASC) can be detected in peripheral blood for a short time (around day 7) after MVA boost
252 when using the ChAd63-MVA regimen^{11,15,16}. RH5-specific ASC responses were assessed by *ex-vivo*
253 ELISPOT using fresh PBMC collected at the day 63 visit for participants. ASC responses were detectable
254 above baseline in all age groups vaccinated with ChAd63-MVA RH5, with the highest responses trending
255 to be measured in the Group 3B infants with a median of 60 RH5-specific ASC per million PBMC
256 (**Figure 5A**). We subsequently assessed RH5-specific B cell responses further by flow cytometry (**Figure**
257 **5B, S5A**). As noted during the analysis of the T cell responses, clinical hematology data showed a clear
258 age-dependent hierarchy in the lymphocyte counts per microliter of blood (**Figure S5B**). Within these
259 total lymphocyte populations, flow cytometry analysis showed the % live B cells to be approximately 2.5-
260 fold higher in the pediatric groups as compared to adults (**Figure S5C**). Combining these data showed
261 that 6-11 month old infants have significantly more B cells per microliter of blood (approximately 5-fold
262 on average) as compared to the adults 18-35 years of age, with children 1-6 years of age having
263 intermediate levels (**Figure 5C**). Following the MVA-RH5 vaccine boost, children and infants showed

264 comparable RH5-specific responses within the live CD19⁺ IgG⁺ B cell population. These frequencies
265 were significantly higher in the infants (approximately 10-fold on average) as compared to the vaccinated
266 adults (**Figure 5D**).

267

268 **Vaccine-induced antibodies show high-level functional GIA *in vitro*.**

269 Finally, serum samples were analyzed for functional anti-parasitic growth inhibition at the GIA Reference
270 Center at the NIH. Here, the standardized GIA assay of human samples typically tests purified total IgG
271 (normalized to a starting concentration of 10 mg/mL) against *P. falciparum* 3D7 clone parasites in the
272 absence of complement^{17,18}. However, we initially measured each serum sample's physiological total
273 serum IgG concentration by HPLC (**Figure S6A**). These data showed the median level in 6-11 month old
274 infants to be 10.1 mg/mL, highly comparable to that previously seen in healthy UK adults¹⁷. In contrast,
275 the levels in 1-6 year old children (median 13.7 mg/mL) and 18-35 year old adults (median 14.8 mg/mL)
276 were significantly higher. We therefore elected instead to initially screen all samples for GIA starting at
277 their physiological total IgG concentration as opposed to normalizing to 10 mg/mL. At these higher
278 starting levels of total IgG, some weak GIA was observed at baseline, especially in the adults (**Figure**
279 **6A**), in line with higher levels of prior malaria exposure in this age group as confirmed by an anti-parasite
280 lysate ELISA (**Figure 6B**). However, following ChAd63-MVA RH5 vaccination, very large increases in
281 GIA were observed in the Group 2B children and Group 3B infants (**Figure S6B**), reaching median levels
282 of 89.2% and 98.5%, respectively, at their physiological IgG concentration at the day 63 time-point
283 (**Figure 6A**). These increases were not observed in the RH5 vaccinated adults or any of the rabies vaccine
284 control groups.

285

286 We next titrated individual total IgG samples from the high responding pediatric groups using a 2-fold
287 dilution series in the GIA assay, starting at each individual sample's physiological total IgG concentration
288 (**Figure 6C**). Here, the observed GIA showed a very strong relationship to the concentration of RH5-

289 specific IgG present in the total IgG used in the assay as measured by ELISA (**Figure 6D**), as seen
290 previously for this antigen following viral vectored or protein-in-adjuvant immunization of healthy UK
291 adults ^{11,17}. The concentration of RH5_FL-specific polyclonal IgG required to give 50% GIA (EC₅₀) was
292 highly comparable in both age groups: 34 µg/mL (95% confidence interval [CI], 25–48) in children and
293 39 µg/mL (95% CI 31–53) in the infants. This functional “quality” readout of vaccine-induced RH5-
294 specific IgG was highly similar to that previously reported in RH5 vaccinated healthy UK adults ^{11,17},
295 which we also confirmed in an independent head-to-head repeat (**Figure S6C**).

296

297 Data from previous *Aotus* monkey *P. falciparum* challenge studies have suggested that levels of *in vitro*
298 GIA >60% at 2.5 mg/mL purified total IgG are associated with a protective outcome following blood-
299 stage vaccination ^{9,19}. We therefore next interpolated the levels of GIA at 10 mg/mL and 2.5 mg/mL total
300 IgG for both age groups, which showed these were significantly higher in the 6-11 month old infants
301 (**Figure 6E,F**). Notably, 6/11 vaccinated infants showed >60% GIA at 2.5 mg/mL total IgG (median
302 61%, range 41-78%). Finally, to relate the GIA assay results back to the original sera, and to account for
303 the significant differences in the physiological concentrations of total IgG across the age groups (**Figure**
304 **S6A**), we calculated the “GIA₅₀ titer”, defined in previous studies ^{9,11,17} as the dilution factor of each
305 serum sample required to reach the concentration of purified IgG that gives 50% GIA (i.e. the GIA EC₅₀).
306 Here, a GIA₅₀ titer >5 was previously associated with protection against a stringent *P. falciparum* blood-
307 stage challenge in *Aotus* monkeys vaccinated with RH5 ⁹. In this analysis of the GIA EC₅₀ (**Figure 6G**)
308 followed by GIA₅₀ titer (**Figure 6H**), the Group 3B vaccinated 6-11 month old infants again showed the
309 highest responses, with a median GIA₅₀ titer = 5.3 (range 3.2-10.6). In summary, these GIA responses in
310 the vaccinated 6-11 month old infants are the highest levels of GIA reported to-date following human
311 vaccination, and were higher than levels of GIA achieved in healthy UK adults using highly immunogenic
312 recombinant protein-in-adjuvant vaccine formulations targeting the RH5 or AMA1 antigens (**Figure 6F-**
313 **H**) ^{17,18}.

314 **Discussion**

315 This dose-escalation, age de-escalation, double-blind, randomized, controlled Phase 1b trial reports the
316 first data in a malaria-endemic population for a vaccine targeting the RH5 antigen from the blood-stage *P.*
317 *falciparum* merozoite. We show in healthy Tanzanian adults, children and infants that a recombinant
318 ChAd63-MVA heterologous prime-boost immunization regimen has a favorable safety profile, and can
319 induce robust functional RH5-specific serum antibody responses in addition to B cell responses in the
320 target infant age group. The local and systemic reactogenicity profile of the full dose RH5 vaccines in all
321 three age groups was similar, if not reduced, as compared to that seen in healthy UK adults immunized
322 with the identical vaccines ¹¹. These data are consistent with previous Phase 1a/b malaria vaccine trials
323 using the same viral vector delivery platforms at similar doses ^{16,20-30}.

324
325 The clinical safety of MVA as a recombinant vaccine vector for other infectious diseases and cancer is
326 well documented ³¹, whilst current efforts are also using this virus to vaccinate against mpox ³². Our data
327 with the ChAd63 vector also add support to existing data that suggest this simian adenovirus vector is
328 safe for clinical use. However, a very rare but serious adverse reaction to a similar adenovirus vector
329 (ChAdOx1) has been observed in the context of COVID-19 vaccines – vaccine-induced thrombosis with
330 thrombocytopenia ³³. The mechanism of this is not completely understood and so it is unclear whether this
331 risk is likely to apply to another serotype of adenovirus, delivering a non-coronavirus antigen to a
332 predominantly African target population. Reassuringly this phenomenon has not been observed in any
333 trials to-date of the same serotype of adenovirus (ChAdOx1) delivering a malarial or other non-
334 coronavirus antigen, nor has it been observed in trials of other serotypes ³⁴. However, all these trials have
335 been limited by their size (none reaching Phase 3 or more than hundreds of recipients) and so such a rare
336 adverse event, even if it were real, may not have been detected. Nevertheless, since the time of
337 undertaking this trial, a second Phase 1b trial has initiated in Tanzania using a soluble protein-in-adjuvant
338 formulation, RH5.1/Matrix-M™ (ClinicalTrials.gov NCT04318002) ^{17,35}. Future efforts will thus focus on

339 the clinical development of recombinant RH5 antigen-based vaccines formulated in adjuvant, rather than
340 ChAd63-MVA viral vectors, in order to align blood-stage vaccine delivery with the existing anti-
341 sporozoite vaccines RTS,S/AS01³⁶ and R21/Matrix-MTM² and to enable future multi-stage malaria
342 vaccine strategies.

343
344 The viral vectored vaccine platform was historically developed to induce T cell responses against the
345 encoded transgene³⁷, and in this trial RH5-specific IFN- γ T cell responses were induced by ChAd63
346 RH5 in the Tanzanian adults; these subsequently peaked at median levels of >1000 SFU/million PBMC
347 following the MVA RH5 boost. The magnitude and maintenance of these responses post-boost were
348 largely comparable to those seen previously in healthy UK adults immunized with the identical vaccine
349 regimen¹¹. In contrast, T cell priming by ChAd63 RH5 in the 6-11 month old infants, and boosting by
350 MVA RH5 in both the infants and the 1-6 year old children appeared much weaker, leading to
351 significantly lower responses (~5-10-fold) post-boost as compared to adults when using the standard
352 ELISPOT readout that reports per million PBMC. However, age-dependent variations are observed in the
353 numbers of lymphocytes circulating per mL of blood, with much higher lymphocyte frequencies
354 measured in young children and infants as compared to adults³⁸. Following incorporation of the
355 lymphocyte count to report ELISPOT responses per mL blood, only the 6-11 month old infants still
356 showed significantly lower responses post-prime, whilst far more comparable IFN- γ T cell responses
357 were observed across the age groups post-boost, as reported in previous West African Phase 1b clinical
358 trials using the same vectors recombinant for the pre-erythrocytic malaria antigen ME-TRAP¹².
359 Nonetheless, the possible contribution of IFN- γ -secreting T cells to vaccine-induced blood-stage malaria
360 immunity in humans remains unclear. The ChAd63-MVA viral vectors have routinely induced a mixed
361 antigen-specific CD4⁺/CD8⁺ T cell response in humans, and it is highly likely that RH5-specific CD4⁺ T
362 cells will provide key help to B cell responses^{39,40}. Indeed, previous work in UK adults has shown the
363 ChAd63-MVA RH5 vaccines can induce antigen-specific peripheral T follicular helper (Tfh) cell

364 responses⁴¹. Work therefore remains on-going to investigate the phenotypes of the Tfh cells induced in
365 the Tanzanian population and their contribution to vaccine-induced humoral immune responses.
366
367 We also optimized the design and delivery of these recombinant ChAd63 and MVA viruses to induce
368 strong antibody responses against blood-stage malaria antigens⁴²⁻⁴⁴. Here, the ChAd63-MVA RH5
369 vaccines induced RH5-specific IgG antibody responses in the Tanzanian adults that peaked at ~14
370 µg/mL; these levels of serum antibody were highly comparable to those seen previously in healthy UK
371 adults immunized with the identical vaccine regimen¹¹. More encouragingly, the levels of anti-RH5 IgG
372 induced by the same full-dose vaccines in the pediatric groups were significantly higher, reaching peak
373 levels of ~150 µg/mL in the 6-11 month old infants following the MVA RH5 boost. The ability to induce
374 such high levels of anti-RH5 IgG in the target age group for a blood-stage malaria vaccine (i.e. young
375 children and infants over 5 months of age) bodes well for future field efficacy testing of standalone blood-
376 stage malaria vaccines as well as multi-stage malaria vaccine strategies whereby a blood-stage RH5
377 component could be combined with existing pre-erythrocytic vaccines such as RTS,S/AS01³⁶ or
378 R21/Matrix-MTM². The data also support the ongoing and wider use of the chimpanzee adenovirus
379 vaccine platform to protect against outbreak or endemic viruses, such as Ebola, rabies and Rift Valley
380 fever⁴⁵⁻⁴⁷.
381
382 The induction of higher antibody levels in the pediatric groups, as compared to adults, is likely not
383 specific to the ChAd63-MVA RH5 vaccine. Indeed, although numbers were small, we observed the same
384 age-dependent hierarchy of antibody induction against the rabies virus glycoprotein in the rabies vaccine
385 controls. Elsewhere, younger children (aged 6-11 years) also induced higher antibodies than older
386 children (aged 12-17 years) vaccinated with the ChAdOx1-nCoV19 (AZD1222) Covid-19 vaccine⁴⁸, and
387 trials of the pre-erythrocytic vaccine ChAd63-MVA ME-TRAP also reported ten-fold higher anti-TRAP
388 antibody levels in West African infants as compared to UK adults and West African adults¹². Published

389 immunogenicity data spanning adults and infants 5-17 months of age vaccinated with the RTS,S/AS01
390 ^{36,49} and R21/Matrix-MTM 2,⁵⁰ malaria vaccines also suggest the same. However, responses in infants 6-12
391 weeks of age at first vaccination with RTS,S/AS01 showed lower levels of antibody that associated with
392 reduced efficacy against clinical malaria ³⁶. Consequently, given both anti-sporozoite and anti-merozoite
393 malaria vaccine strategies necessitate very high levels of antibody to protect against parasite infection,
394 current efforts remain focused on infants and young children over 5 months of age at the time of first
395 vaccination.

396
397 Why the infants and young children vaccinated with ChAd63-MVA RH5 induced such high levels of
398 antibody remains to be fully understood. Our results suggested that pre-existing anti-vector immunity is
399 unlikely to be the reason explaining the observed improvement in anti-RH5 humoral immunogenicity in
400 the younger age groups. Consistent with this, studies of the ChAdOx1-nCoV19 (AZD1222) Covid-19
401 vaccine given in a 2-dose homologous regimen reported similar findings across studies in adults and
402 children, with no evidence that anti-ChAd immune responses measured after the first vaccine dose
403 associated with the immunogenicity outcome measures of the second vaccine dose ^{48,51}. We therefore
404 sought to analyse the underlying B cell responses. Here, we observed higher peripheral ASC responses in
405 the infants, as well as higher absolute numbers of B cells per microliter of blood and a stronger RH5-
406 specific IgG⁺ B cell response within the CD19⁺ population in the younger age groups. The significantly
407 higher anti-RH5 serum IgG response induced by the ChAd63-MVA RH5 vaccine in children and infants,
408 as compared to adults, is therefore strongly associated with greater B cell immunogenicity. Given current
409 antibody-inducing vaccine strategies to protect against clinical malaria are focussed on infants 5-17
410 months of age, this observation warrants further investigation in the future.

411
412 Finally, we assessed functional GIA of the vaccine-induced anti-RH5 antibodies. Our previous work has
413 identified the *in vitro* assay of GIA as a highly significant predictor of *P. falciparum in vivo* growth

414 inhibition following blood-stage challenge of both vaccinated UK adults¹⁷ and *Aotus* monkeys^{9,19}. We
415 have also confirmed this association as a mechanistic correlate in *Aotus* monkeys, i.e. one that can cause
416 *in vivo* protection, via passive transfer of a GIA-positive RH5-specific IgG monoclonal antibody¹⁰, with
417 similar results observed in humanized mice⁵². Importantly, full protection of *Aotus* monkeys required a
418 serological threshold level of GIA, defined as i) a level of *in vitro* GIA >60% at 2.5 mg/mL purified total
419 IgG^{9,19}, or ii) a GIA₅₀ titer >5, with this latter measure also taking into account any differences in the
420 physiological concentrations of total IgG in the vaccinees⁹. Previous trials in healthy UK adults did not
421 exceed this threshold when using the ChAd63-MVA RH5 vaccine¹¹, or a more immunogenic soluble
422 RH5 protein-in-adjuvant, RH5.1/AS01_B¹⁷. In contrast, the 6-11 month old infants in this trial showed the
423 highest yet reported levels of GIA in vaccinated humans, with over half of the vaccinees in this target age
424 group exceeding the threshold. This related to the high quantity of anti-RH5 serum IgG induced by the
425 ChAd63-MVA RH5 vaccine, given the functional quality (i.e. GIA per unit anti-RH5 antibody) was
426 highly similar across the age groups tested. Moreover, the highly similar functional antibody quality data
427 across the UK and Tanzanian vaccine trials are encouraging because they indicate no obvious interference
428 from pre-existing and naturally-occurring anti-malarial antibody responses with the vaccine-induced anti-
429 RH5 IgG, as occurred for historical vaccines targeting AMA1⁵³.

430
431 The main limitations of the trial include relatively small numbers of participants and a limited follow-up
432 period of four months post-MVA RH5 booster vaccination, such that the longer-term kinetic of the
433 immune response has not been characterized. Improving the durability of protection against clinical
434 malaria will be critical for next-generation malaria vaccine strategies. We also only tested a single
435 heterologous prime-boost regimen in this trial, whereas a more extensive assessment of vaccine dose and
436 regimen, including the use of delayed boosting, in Phase 1a/b trials could be key to optimizing
437 immunogenicity and the longevity of vaccine-induced protection^{17,54}. In this regard, since undertaking
438 this trial, a second Phase 1b trial has initiated in Tanzania using a soluble protein-in-adjuvant formulation,

439 RH5.1/Matrix-M™ delivered in a variety of dosing regimens (ClinicalTrials.gov NCT04318002). This
440 trial will indicate whether even higher antibody and GIA levels can be achieved in the target 5-17 month
441 old infant population with a delivery platform that is more immunogenic than ChAd63-MVA.
442 Nonetheless, the data in the Phase 1b trial reported here confirm, for the first time, that substantial anti-
443 RH5 immune responses can be achieved safely by vaccination in infants from a malaria-endemic area, in
444 stark contrast to the poor immunogenicity seen to this antigen following natural *P. falciparum* infection
445 ^{6,11}. These data also justify onward progression to Phase 2b field efficacy trials to determine whether
446 growth inhibitory antibody levels of this magnitude can ultimately protect against clinical malaria.

447 **Methods**

448 **Study Design**

449 This was a randomized, controlled, age de-escalation, dose-escalation study called VAC070 that was
450 conducted according to the principles of the current revision of the Declaration of Helsinki 2008 and in
451 full conformity with the ICH guidelines for Good Clinical Practice (GCP). It was approved by the Oxford
452 Tropical Research Ethics Committee in the UK (OxTREC, reference 29-17), the Ifakara Health Institute
453 Institutional Review Board in Tanzania (reference: 20-2017), the National Institute for Medical Research
454 in Tanzania, the National Health Research Ethics Sub-Committee (NatHREC) and the then Tanzania
455 Food and Drugs Authority (now the Tanzania Medicines and Medical Devices Authority), reference:
456 TFDA0017/CTR/0015/3. The Consolidated Standards of Reporting Trials (CONSORT) guideline was
457 followed.

458
459 We report here the safety, reactogenicity and immunogenicity profile of heterologous prime-boost
460 ChAd63-MVA RH5 vaccination up until 168 days post-enrolment. The study was conducted (and
461 participants recruited and vaccinated) at Kingani Clinical Facility, Ifakara Health Institute, Bagamoyo
462 branch, Tanzania. The VAC070 trial was registered on ClinicalTrials.gov (NCT03435874), the Pan-
463 African Clinical Trials Registry (PACTR201710002722229) and ISRCTN (ISRCTN47448832).

464
465 **Participants**

466 Healthy adults (18-35 years), young children (1-6 years) and infants (6-11 months) residing in Bagamoyo,
467 Tanzania, with a negative malaria blood film at screening, were eligible for inclusion in the study and
468 enrolled into three groups according to age. A full list of inclusion and exclusion criteria are listed in the
469 study protocol which is included within the Supplementary Appendix. Each participant (or guardian)
470 signed or thumb-printed an informed consent form at the in-person screening visit and consent was
471 verified before each vaccination. A Safety Monitoring Committee (SMC) periodically reviewed the study

472 progress and safety data according to a safety review schedule, critically timed to age de-escalations and
473 dose escalations.

474

475 **Procedures**

476 **Randomization and Masking**

477 A randomization list was generated by an independent statistician. This contained sequential codes
478 (Treatment numbers) linking a study identification (ID) to a vaccine assignment. Study ID was assigned
479 to participants in the order in which they were enrolled in the trial. Access to the randomization list was
480 exclusively limited to the study pharmacist and the independent statistician(s). These individuals had no
481 role in the evaluation of the study participants.

482

483 Participants were assigned to groups based on their age, and groups (characterized by participant age and
484 dose of vaccine) were enrolled sequentially. Randomization into vaccine or control groups was performed
485 according to a 2:1 ratio.

486

487 Data pertaining to ChAd63 RH5, MVA RH5 or rabies vaccine were collected in a double-blinded
488 manner. Neither the vaccine recipient nor their parent(s)/guardian(s) or those members of the study team
489 responsible for administering the vaccines or evaluating safety and immunogenicity endpoints were aware
490 of individual vaccine allocation. Only those staff responsible for the storage and preparation of vaccines
491 were unblinded as both vaccines were distinguishable by their packaging and labelling; these staff played
492 no other role in the study and the vaccine preparation area was kept physically separate from the
493 immunization area. The Local Independent Safety Monitor was provided with sealed code-break
494 envelopes for each participant to facilitate unblinding for urgent clinical/ethical reasons. They also had
495 access to a copy of the master randomization list in a sealed envelope, in case emergency unblinding was
496 required.

497
498 Participants attended a two-part screening visit and those eligible returned for enrolment and were
499 randomized to either a dose of ChAd63 RH5 or rabies control vaccine. Eight weeks later all participants
500 were then randomized to either a dose of MVA RH5 or rabies vaccine. All vaccines were administered by
501 intramuscular injection in the upper arm.

502

503 **Safety Analysis**

504 Following each vaccination, each participant was visited at home on days 1, 3, 4, 5 and 6 by a community
505 health worker for assessment and recording of any solicited and unsolicited AEs. At days 2, 7, 14 and 28
506 post-vaccination participants were seen at the clinical research facility. Observations (heart rate,
507 temperature and blood pressure measurements) were taken at the clinic visits from the day of vaccination
508 until the 28 day follow-up visit. Blood tests for exploratory immunology were taken at all visits except
509 those occurring 2 days after each vaccination, 7 days after the first vaccination and 14 days after the
510 second vaccination. Blood samples for safety (full blood count, alanine aminotransferase (ALT) and
511 creatinine) were carried out at screening and on days 0, 7, 14, 28, 56, 63, 84 and 168 for all groups. Any
512 solicited AEs occurring during the 7 days post-vaccination were defined as being at least possibly related
513 to vaccination. The likely causality and grading of all other AEs were assessed as described in the
514 protocol. All unsolicited AEs are reported (**Table S2**) but none were considered possibly, probably or
515 definitely related to vaccination. The types of AEs were classified according to MedDRA (version 26.0).

516

517 **Outcomes**

518 Primary outcome measures for vaccine safety included numbers of solicited and unsolicited AEs after
519 each vaccination. The primary outcome analysis was conducted on the safety analysis population and
520 included participants who received at least the first dose of vaccine in the study. The maximum severity
521 for each solicited systemic AE across seven days after first and second vaccinations was derived for each

522 participant and summarized by group. Analyses were conducted similarly for local reactogenicity. Serious
523 adverse events (SAEs) were collected for the entire study period. The secondary outcome measures for
524 humoral immunogenicity were the concentration of anti-RH5 serum antibodies by ELISA and their
525 percentage GIA *in vitro* using purified IgG, and for cellular immunogenicity were T and B cell responses
526 to RH5 as measured by ELISPOT and/or flow cytometry.

527

528 **Vaccines**

529 The design, production and preclinical testing of the viral vector vaccines have been reported previously
530 in detail ^{6,9}. Briefly, both recombinant viruses express the same 1503 bp coding sequence of RH5 from the
531 3D7 clone of *P. falciparum*, aa E26–Q526 (NCBI Accession #XM_001351508.1). ChAd63 RH5 was
532 manufactured by Advent, Pomezia, Italy which is a daughter company of ReiThera. This production
533 facility meets current Good Manufacturing Practice (cGMP) requirements of the US Food and Drug
534 Administration (FDA) and the European Medicines Agency (EMA) to produce investigational vaccines to
535 be used in human clinical studies. MVA RH5 was manufactured under cGMP conditions by IDT
536 Biologika GmbH, Germany, as described in detail previously ¹¹. Control participants received
537 VERORAB, an inactivated rabies vaccine (Sanofi Pasteur).

538

539 **Peripheral Blood Mononuclear Cell (PBMC), Plasma and Serum Preparation**

540 Blood samples were collected into lithium heparin-treated vacutainer blood collection systems (Becton
541 Dickinson, UK). PBMC were isolated and used within 6 hours in fresh assays as previously described ²².
542 Excess cells were frozen in foetal calf serum (FCS) containing 10% dimethyl sulfoxide (DMSO) and
543 stored in liquid nitrogen. Plasma samples were stored at -80 °C. For serum preparation, untreated blood
544 samples were stored at room temperature (RT) and then the clotted blood was centrifuged for 5 min (1000
545 *xg*). Serum was stored at -80 °C.

546

547 **Peptides**

548 Peptides for *ex-vivo* IFN- γ ELISPOT were purchased from NEO Scientific (Cambridge, MA, USA).
549 Sequences are reported in **Table S4**. In brief, the peptides (20 amino acids (aa) in length and overlapping
550 by 10 aa) covered the entire RH5 sequence present in the RH5 vaccine. Peptides were reconstituted in
551 100% DMSO at 50-200 mg/mL and combined into various pools for the ELISPOT assay.

552

553 ***Ex-vivo* IFN- γ ELISPOT**

554 Fresh PBMC were used in all assays using a previously described protocol ¹¹. Spots were counted using
555 an ELISPOT counter (Autoimmun Diagnostika (AID), Germany). Results are expressed as IFN- γ spot-
556 forming units (SFU) per million PBMC. Background responses in unstimulated control wells were almost
557 always less than 20 spots, and were subtracted from those measured in peptide-stimulated wells.

558

559 **Recombinant RH5 Protein**

560 Recombinant full-length RH5 protein (also known as “RH5.1”) was used for all ELISA assays and B cell
561 ELISPOT assays. The protein was produced and purified from a stably transfected *Drosophila* S2 cell line
562 as previously described ³⁵.

563

564 **RH5 ELISA**

565 Anti-RH5 total IgG ELISAs were performed against full-length RH5 protein (RH5.1) using standardized
566 methodology as previously described in detail for other RH5 vaccine trials ^{11,17}. The reciprocal of the test
567 sample dilution giving an optical density at 405nm (OD₄₀₅) of 1.0 in the standardized assay was used to
568 assign an ELISA unit value of the standard. A standard curve and Gen5 ELISA software v3.04 (BioTek,
569 UK) were used to convert the OD₄₀₅ of individual test samples into arbitrary units (AU). These responses
570 in AU are reported in $\mu\text{g/mL}$ following generation of a conversion factor by calibration-free concentration
571 analysis (CFCA) as reported previously ¹¹.

572

573 **ChAd63 ELISA**

574 Antibody responses to the ChAd63 vaccine vector were determined by endpoint ELISA as previously
575 described^{42,55,56}. Briefly, a purified ChAd63 vector encoding an irrelevant antigen (ovalbumin)^{57,58} was
576 adsorbed overnight at 4 °C to 96 well NUNC-Immuno Maxisorp plates (Thermo Fisher Scientific) at 3 x
577 10⁸ vp/mL⁵⁹. Test sera were diluted 1:100, added in duplicate and serially diluted 3-fold. Bound
578 antibodies were detected using goat anti-human IgG conjugated to alkaline phosphatase (Sigma),
579 developed using 4-Nitrophenyl phosphate disodium salt hexahydrate (Sigma) and OD₄₀₅ determined on a
580 BioTek Elx808 reader with Gen5 ELISA software. Endpoint titers were calculated as the x-axis intercept
581 of the sample titration curve at OD₄₀₅ = 0.15 (equivalent to blank test samples). A positive control sample
582 from participants vaccinated with a different ChAd63 vectored vaccine²⁷ was included as an internal
583 reference.

584

585 **Parasite Lysate ELISA**

586 Serum antibody levels to parasite lysate were assessed by standardized ELISA methodology previously
587 described^{60,61}. Schizont extract from *P. falciparum* (3D7 clone) produced by the GIA Reference
588 Laboratory, NIAID, NIH, was adsorbed overnight at 4 °C to 96 well NUNC-Immuno Maxisorp plates
589 (Thermo Fisher Scientific) at equivalent to 5x10² parasites per µL. Test sera were diluted in 1% milk and
590 added in triplicate to plates following blocking with 5% milk in DPBS (Sigma). A reference standard and
591 internal control from a pool of N=34 high malaria pre-exposed serum samples, from the BCTF-IHI
592 Biobank, plus blank wells were included. Bound antibodies were detected using goat anti-human IgG
593 conjugated to alkaline phosphatase (Sigma), developed using 4-Nitrophenyl phosphate disodium salt
594 hexahydrate (Sigma) and absorbance (OD₄₀₅) was determined on a BioTek Elx808 reader with Gen5
595 software. Antibody units were assigned using the reciprocal dilution of the standard giving an optical

596 density of 1.0 at OD₄₀₅. The standard curve and Gen5 software v3.04 (Agilent) were then used to convert
597 the OD₄₀₅ of test samples to arbitrary units (AU).

598

599 **Rabies ELISA**

600 Binding antibody responses induced by the rabies control vaccine were determined by endpoint ELISA,
601 with the same methodology as for the anti-ChAd63 ELISA, but using recombinant rabies glycoprotein ⁴⁶
602 adsorbed to the plate at 2 µg/mL. Pooled sera from Group 3B participants vaccinated with the rabies
603 vaccine was included as a development control.

604

605 **Parasite qPCR**

606 Parasitemia was determined retrospectively by quantitative polymerase chain reaction (qPCR) performed
607 on blood samples taken at baseline, 7 days and 28 days post-boost as previously described ⁶². Briefly,
608 blood was collected in 2.0 mL tubes containing EDTA. DNA was extracted from whole blood using
609 Quick-DNA miniprep plus kit (Zymo research, USA), and 2 µL each extracted DNA was used per assay
610 well and run in triplicate. qPCR was conducted on a CFX96 real-time qPCR machine (Bio-Rad) and
611 analyzed with CFX Manager Software (v2.2) with the following cycling conditions: polymerase
612 activation at 95 °C for 1 min, 45 cycles of denaturation at 95 °C for 15 s and annealing and elongation at
613 57 °C for 45 s. Parasites per µL were calculated against a defined international standard for *P. falciparum*
614 DNA Nucleic Acid Amplification techniques (WHO reference from NIBSC #04/176) ⁶³ reconstituted in
615 0.5 mL sterile nuclease-free water to 5x10⁸ parasites per µL.

616

617 **Antibody-Secreting Cell (ASC) ELISPOT**

618 *Ex-vivo* ASC ELISPOT assays were performed against RH5.1 protein as described in detail elsewhere ¹¹
619 using fresh PBMC. Plates were counted using an AID ELISPOT plate reader. Results are reported as
620 RH5-specific ASC per million PBMC used in the assay.

621

622 **B cell Flow Cytometry**

623 Frequencies of live B cells within total lymphocytes and RH5-specific cells within IgG⁺ B cells (CD19⁺

624 IgD⁻ IgM⁻ IgA⁻) were measured by flow cytometry. In brief, cryopreserved PBMC were thawed and

625 washed in IMDM (12440053, Gibco) supplemented with 10% FCS (F9665, Sigma Aldrich), 0.2%

626 MycoZap (VZA-2031, Lonza), and 0.04% benzonase (71205-3, Merck). Thawed PBMC were stained

627 first with a viability stain (Live/Dead Aqua; L34966, Invitrogen), followed by a panel comprising anti-

628 human CD3-BV510 (317332, Biolegend); anti-human CD14-BV510 (301842, Biolegend); anti-human

629 CD56-BV510 (318430, Biolegend); anti-human CD27-A488 (393204, Biolegend); anti-human IgM-

630 PerCP-Cy5.5 (561285, BD Biosciences); anti-human CD19-ECD (IM2708U, Beckman Coulter); anti-

631 human IgD-PE-Cy7 (561315, BD Biosciences); anti-human IgA-A647 (109-475-011, Jackson); anti-

632 human CD38-APC-Cy7 (303534, Biolegend); and a RH5-PE probe (produced in-house as previously

633 described^{41,54}). Stained samples were washed and acquired on a FACS Aria Fusion Flow Cytometer (BD

634 Biosciences). See **Figure S5** for an example gating strategy of live B cells within single lymphocytes, and

635 RH5-specific (IgG⁺) cells within the CD19⁺ IgD⁻ IgM⁻ IgA⁻ B cell population.

636

637 **Haematology**

638 The lymphocyte count was obtained during Complete Blood Count (CBC) analysis using the automated

639 haematology analyzer, Sysmex XS 800i, for assessment of whole blood. The analyzer uses a fluorescence

640 flow cytometry method for analysis of white blood cells (WBC) and the five differentials by using the

641 semiconductor laser beam. A small volume of blood (20 µL) is aspirated by the analyzer to measure the

642 WBC differentials according to their size and structure.

643

644 **Serum IgG Concentration**

645 Total serum IgG concentrations were determined using a Bio-Monolith Protein G column on an Agilent
646 1260 HPLC system (Agilent, Cheshire, UK). Separation was performed at 1 mL/min using PBS and 0.2
647 M Glycine pH 2.0 as mobile phases with detection at UV 280 nm. A calibration curve was produced
648 using purified human IgG.

649

650 **Assay of Growth Inhibition Activity (GIA)**

651 Standardized assays were performed by the GIA Reference Center, NIH, USA, using previously
652 described methodology⁶⁴, with one modification. Here, each sample was tested in three independent
653 replication assays using three different batches of red blood cells (RBC), and the median of these three
654 results was used to generate the final dataset. Otherwise for each assay, in brief, protein G purified IgG
655 samples were incubated with RBC infected with synchronized *P. falciparum* 3D7 clone parasites in a
656 final volume of 40 μ L for 40 h at 37 °C, and the final parasitemia in each well was quantified by
657 biochemical determination of parasite lactate dehydrogenase. All purified IgG samples were tested at final
658 test well concentrations (reported in mg/mL) as described in Results. For certain samples a dilution series
659 was used to determine the concentration that gave 50% GIA (EC_{50}).

660

661 **Statistical Analysis**

662 Data were analyzed using GraphPad Prism version 9.5 for Windows (GraphPad Software Inc., California,
663 USA). All tests used were two-tailed and are described in the text. To analyze the relationship between
664 GIA and ELISA assay data, a Richard's five-parameter dose-response curve was fitted, constrained to 0
665 % GIA at the bottom and 100% GIA at the top. A value of $P < 0.05$ was considered significant.

666 **Author Contributions**

667 Conceived and performed the experiments: SES, WFK, IMM, NSL, MM, FM, SA, AD, FM, BS, TA,
668 MR, LM, OL, AMA, GN, BM, SM, TGM, JRB, LTW, YT, LDWK, SHH, ROP, CMN, CAL, KM, SJD,
669 AMM, AIO.
670 Analyzed the data: SES, WFK, IMM, NSL, TA, TGM, CMN, KM, SJD, AMM, AIO.
671 Project Management: SA, AML, FLN, J-SC.
672 Wrote the paper: SES, SJD, AMM.

673 **Acknowledgments**

674 The authors are grateful to Nick Edwards, Julie Furze, Mimi Hou, Jing Jin, Simon Kerridge, Penelope
675 Lane, Daniel Marshall-Searson, Ian Poulton and Alex Spencer for assistance, and to Sandy Douglas and
676 Adam Berg for provision of rabies glycoprotein (Jenner Institute, University of Oxford); Helena Parracho,
677 Tanja Brenner and Eleanor Berrie (Clinical Biomanufacturing Facility, University of Oxford); Amy
678 Duckett and Carly Banner for arranging contracts (University of Oxford); Robert Hedley and Vasiliki
679 Tsioligka for assistance with flow cytometry (Sir William Dunn School of Pathology, University of
680 Oxford); Loredana Siani, Stefania Di Marco (ReiThera SRL, formerly Okairòs SRL) and Annunziata Del
681 Gaudio (Advent SRL) for manufacture of ChAd63 RH5; Marc Lievens, Danielle Morelle and GSK for
682 review of the clinical trial design, supply of the ChAd63 vector and review of the draft manuscript, but
683 the authors are solely responsible for the final content and data interpretation; the Independent Safety
684 Monitoring Committee for their insightful safety oversight (Prof Brian Angus, Prof Karim Manji and Prof
685 Alison Elliot); and all the study participants.

686
687 This work was funded in part by an African Research Leader Award to AIO from the UK Medical
688 Research Council (MRC) [MR/P020593/1]. This award was jointly funded by the UK MRC and the UK
689 Department for International Development (DFID) under the MRC/DFID Concordat agreement and is

690 also part of the EDCTP2 programme supported by the European Union. This work was also supported in
691 part by the National Institute for Health Research (NIHR) Oxford Biomedical Research Centre (BRC).
692 The views expressed are those of the authors and not necessarily those of the NIHR or the Department of
693 Health and Social Care. The GIA work was supported by the United States Agency for International
694 Development (USAID) and the Intramural Program of the National Institute of Allergy and Infectious
695 Diseases, National Institutes of Health. SHH holds an NIHR Academic Clinical Lectureship. CMN held a
696 Sir Henry Wellcome Postdoctoral Fellowship [209200/Z/17/Z]. SJD is a Jenner Investigator and held a
697 Wellcome Trust Senior Fellowship [106917/Z/15/Z].

698

699 **Conflict of Interest Statement**

700 SJD is a named inventor on patent applications relating to RH5 malaria vaccines and adenovirus-based
701 vaccines, is an inventor on intellectual property licensed by Oxford University Innovation to AstraZeneca,
702 and has been a consultant to GSK on malaria vaccines. AMM has been a consultant to GSK on malaria
703 vaccines, and has an immediate family member who is an inventor on patent applications relating to RH5
704 malaria vaccines and adenovirus-based vaccines, and is an inventor on intellectual property licensed by
705 Oxford University Innovation to AstraZeneca. All other authors have declared that no conflict of interest
706 exists.

707

708 **Data and Materials Availability**

709 Requests for materials should be addressed to the corresponding authors.

710 **References**

- 711 1. World Health Organization. (2022). World Malaria Report.
- 712 2. Dattoo, M.S., Natama, M.H., Some, A., Traore, O., Rouamba, T., Bellamy, D., Yameogo, P.,
713 Valia, D., Tegneri, M., Ouedraogo, F., Soma, R., et al. (2021). Efficacy of a low-dose candidate
714 malaria vaccine, R21 in adjuvant Matrix-M, with seasonal administration to children in Burkina
715 Faso: a randomised controlled trial. *Lancet* 397, 1809-1818.
- 716 3. Rts, S.C.T.P. (2015). Efficacy and safety of RTS,S/AS01 malaria vaccine with or without a
717 booster dose in infants and children in Africa: final results of a phase 3, individually randomised,
718 controlled trial. *Lancet* 386, 31-45.
- 719 4. Draper, S.J., Sack, B.K., King, C.R., Nielsen, C.M., Rayner, J.C., Higgins, M.K., Long, C.A., and
720 Seder, R.A. (2018). Malaria Vaccines: Recent Advances and New Horizons. *Cell Host Microbe*
721 24, 43-56.
- 722 5. Ragotte, R.J., Higgins, M.K., and Draper, S.J. (2020). The RH5-CyRPA-Ripr Complex as a
723 Malaria Vaccine Target. *Trends Parasitol* 36, 545-559.
- 724 6. Douglas, A.D., Williams, A.R., Illingworth, J.J., Kamuyu, G., Biswas, S., Goodman, A.L.,
725 Wyllie, D.H., Crosnier, C., Miura, K., Wright, G.J., Long, C.A., et al. (2011). The blood-stage
726 malaria antigen PfRH5 is susceptible to vaccine-inducible cross-strain neutralizing antibody. *Nat*
727 *Commun* 2, 601.
- 728 7. Crosnier, C., Bustamante, L.Y., Bartholdson, S.J., Bei, A.K., Theron, M., Uchikawa, M., Mboup,
729 S., Ndir, O., Kwiatkowski, D.P., Duraisingh, M.T., Rayner, J.C., et al. (2011). Basigin is a
730 receptor essential for erythrocyte invasion by *Plasmodium falciparum*. *Nature* 480, 534-537.
- 731 8. Galaway, F., Yu, R., Constantinou, A., Prugnolle, F., and Wright, G.J. (2019). Resurrection of the
732 ancestral RH5 invasion ligand provides a molecular explanation for the origin of *P. falciparum*
733 malaria in humans. *PLoS Biol* 17, e3000490.
- 734 9. Douglas, A.D., Baldeviano, G.C., Lucas, C.M., Lugo-Roman, L.A., Crosnier, C., Bartholdson,
735 S.J., Diouf, A., Miura, K., Lambert, L.E., Ventocilla, J.A., Leiva, K.P., et al. (2015). A PfRH5-
736 Based Vaccine Is Efficacious against Heterologous Strain Blood-Stage *Plasmodium falciparum*
737 Infection in Aotus Monkeys. *Cell Host Microbe* 17, 130-139.
- 738 10. Douglas, A.D., Baldeviano, G.C., Jin, J., Miura, K., Diouf, A., Zenonos, Z.A., Ventocilla, J.A.,
739 Silk, S.E., Marshall, J.M., Alanine, D.G.W., Wang, C., et al. (2019). A defined mechanistic
740 correlate of protection against *Plasmodium falciparum* malaria in non-human primates. *Nat*
741 *Commun* 10, 1953.
- 742 11. Payne, R.O., Silk, S.E., Elias, S.C., Miura, K., Diouf, A., Galaway, F., de Graaf, H., Brendish,
743 N.J., Poulton, I.D., Griffiths, O.J., Edwards, N.J., et al. (2017). Human vaccination against RH5
744 induces neutralizing antimalarial antibodies that inhibit RH5 invasion complex interactions. *JCI*
745 *Insight* 2, 96381.
- 746 12. Bliss, C.M., Drammeh, A., Bowyer, G., Sanou, G.S., Jagne, Y.J., Ouedraogo, O., Edwards, N.J.,
747 Tarama, C., Ouedraogo, N., Ouedraogo, M., Njie-Jobe, J., et al. (2017). Viral Vector Malaria
748 Vaccines Induce High-Level T Cell and Antibody Responses in West African Children and
749 Infants. *Mol Ther* 25, 547-559.
- 750 13. Salim, N., Knopp, S., Lweno, O., Abdul, U., Mohamed, A., Schindler, T., Rothen, J., Masimba,
751 J., Kwaba, D., Mohammed, A.S., Althaus, F., et al. (2015). Distribution and risk factors for
752 *Plasmodium* and helminth co-infections: a cross-sectional survey among children in Bagamoyo
753 district, coastal region of Tanzania. *PLoS Negl Trop Dis* 9, e0003660.
- 754 14. Thorner, A.R., Vogels, R., Kaspers, J., Weverling, G.J., Holterman, L., Lemckert, A.A., Dilraj,
755 A., McNally, L.M., Jeena, P.M., Jepsen, S., Abbink, P., et al. (2006). Age dependence of
756 adenovirus-specific neutralizing antibody titers in individuals from sub-Saharan Africa. *Journal*
757 *of clinical microbiology* 44, 3781-3783.

- 758 15. Elias, S.C., Choudhary, P., de Cassan, S.C., Biswas, S., Collins, K.A., Halstead, F.D., Bliss,
759 C.M., Ewer, K.J., Hodgson, S.H., Duncan, C.J., Hill, A.V., et al. (2014). Analysis of human B-
760 cell responses following ChAd63-MVA MSP1 and AMA1 immunization and controlled malaria
761 infection. *Immunology* 141, 628-644.
- 762 16. Hodgson, S.H., Choudhary, P., Elias, S.C., Milne, K.H., Rampling, T.W., Biswas, S., Poulton,
763 I.D., Miura, K., Douglas, A.D., Alanine, D.G., Illingworth, J.J., et al. (2014). Combining Viral
764 Vected and Protein-in-adjuvant Vaccines Against the Blood-stage Malaria Antigen AMA1:
765 Report on a Phase 1a Clinical Trial. *Mol Ther* 22, 2142-2154.
- 766 17. Minassian, A.M., Silk, S.E., Barrett, J.R., Nielsen, C.M., Miura, K., Diouf, A., Loos, C., Fallon,
767 J.K., Michell, A.R., White, M.T., Edwards, N.J., et al. (2021). Reduced blood-stage malaria
768 growth and immune correlates in humans following RH5 vaccination. *Med* 2, 701-719.
- 769 18. Payne, R.O., Milne, K.H., Elias, S.C., Edwards, N.J., Douglas, A.D., Brown, R.E., Silk, S.E.,
770 Biswas, S., Miura, K., Roberts, R., Rampling, T.W., et al. (2016). Demonstration of the Blood-
771 Stage Controlled Human Malaria Infection Model to Assess Efficacy of the Plasmodium
772 falciparum AMA1 Vaccine FMP2.1/AS01. *J Infect Dis* 213, 1743-1751.
- 773 19. Singh, S., Miura, K., Zhou, H., Muratova, O., Keegan, B., Miles, A., Martin, L.B., Saul, A.J.,
774 Miller, L.H., and Long, C.A. (2006). Immunity to recombinant plasmodium falciparum merozoite
775 surface protein 1 (MSP1): protection in Aotus nancymai monkeys strongly correlates with anti-
776 MSP1 antibody titer and in vitro parasite-inhibitory activity. *Infect Immun* 74, 4573-4580.
- 777 20. de Barra, E., Hodgson, S.H., Ewer, K.J., Bliss, C.M., Hennigan, K., Collins, A., Berrie, E.,
778 Lawrie, A.M., Gilbert, S.C., Nicosia, A., McConkey, S.J., et al. (2014). A phase Ia study to assess
779 the safety and immunogenicity of new malaria vaccine candidates ChAd63 CS administered
780 alone and with MVA CS. *PLoS One* 9, e115161.
- 781 21. Sheehy, S.H., Duncan, C.J., Elias, S.C., Collins, K.A., Ewer, K.J., Spencer, A.J., Williams, A.R.,
782 Halstead, F.D., Moretz, S.E., Miura, K., Epp, C., et al. (2011). Phase Ia Clinical Evaluation of the
783 Plasmodium falciparum Blood-stage Antigen MSP1 in ChAd63 and MVA Vaccine Vectors. *Mol*
784 *Ther* 19, 2269-2276.
- 785 22. Sheehy, S.H., Duncan, C.J., Elias, S.C., Biswas, S., Collins, K.A., O'Hara, G.A., Halstead, F.D.,
786 Ewer, K.J., Mahungu, T., Spencer, A.J., Miura, K., et al. (2012). Phase Ia Clinical Evaluation of
787 the Safety and Immunogenicity of the Plasmodium falciparum Blood-Stage Antigen AMA1 in
788 ChAd63 and MVA Vaccine Vectors. *PLoS One* 7, e31208.
- 789 23. Sheehy, S.H., Duncan, C.J., Elias, S.C., Choudhary, P., Biswas, S., Halstead, F.D., Collins, K.A.,
790 Edwards, N.J., Douglas, A.D., Anagnostou, N.A., Ewer, K.J., et al. (2012). ChAd63-MVA-
791 vectored Blood-stage Malaria Vaccines Targeting MSP1 and AMA1: Assessment of Efficacy
792 Against Mosquito Bite Challenge in Humans. *Mol Ther* 20, 2355-2368.
- 793 24. Afolabi, M.O., Tiono, A.B., Adetifa, U.J., Yaro, J.B., Drammeh, A., Nebie, I., Bliss, C.,
794 Hodgson, S.H., Anagnostou, N.A., Sanou, G.S., Jagne, Y.J., et al. (2016). Safety and
795 Immunogenicity of ChAd63 and MVA ME-TRAP in West African Children and Infants. *Mol*
796 *Ther* 24, 1470-1477.
- 797 25. Kimani, D., Jagne, Y.J., Cox, M., Kimani, E., Bliss, C.M., Gitau, E., Ogwang, C., Afolabi, M.O.,
798 Bowyer, G., Collins, K.A., Edwards, N., et al. (2014). Translating the immunogenicity of prime-
799 boost immunization with ChAd63 and MVA ME-TRAP from malaria naive to malaria-endemic
800 populations. *Mol Ther* 22, 1992-2003.
- 801 26. Ogwang, C., Afolabi, M., Kimani, D., Jagne, Y.J., Sheehy, S.H., Bliss, C.M., Duncan, C.J.,
802 Collins, K.A., Garcia Knight, M.A., Kimani, E., Anagnostou, N.A., et al. (2013). Safety and
803 immunogenicity of heterologous prime-boost immunisation with Plasmodium falciparum malaria
804 candidate vaccines, ChAd63 ME-TRAP and MVA ME-TRAP, in healthy Gambian and Kenyan
805 adults. *PLoS One* 8, e57726.

- 806 27. Payne, R.O., Silk, S.E., Elias, S.C., Milne, K.H., Rawlinson, T.A., Llewellyn, D., Shakri, A.R.,
807 Jin, J., Labbe, G.M., Edwards, N.J., Poulton, I.D., et al. (2017). Human vaccination against
808 *Plasmodium vivax* Duffy-binding protein induces strain-transcending antibodies. *JCI Insight* 2,
809 93683.
- 810 28. de Graaf, H., Payne, R.O., Taylor, I., Miura, K., Long, C.A., Elias, S.C., Zaric, M., Minassian,
811 A.M., Silk, S.E., Li, L., Poulton, I.D., et al. (2021). Safety and Immunogenicity of ChAd63/MVA
812 Pfs25-IMX313 in a Phase I First-in-Human Trial. *Front Immunol* 12, 694759.
- 813 29. Mensah, V.A., Gueye, A., Ndiaye, M., Edwards, N.J., Wright, D., Anagnostou, N.A., Syll, M.,
814 Ndaw, A., Abiola, A., Bliss, C., Gomis, J.F., et al. (2016). Safety, Immunogenicity and Efficacy
815 of Prime-Boost Vaccination with ChAd63 and MVA Encoding ME-TRAP against *Plasmodium*
816 *falciparum* Infection in Adults in Senegal. *PLoS One* 11, e0167951.
- 817 30. Ogwang, C., Kimani, D., Edwards, N.J., Roberts, R., Mwacharo, J., Bowyer, G., Bliss, C.,
818 Hodgson, S.H., Njuguna, P., Viebig, N.K., Nicosia, A., et al. (2015). Prime-boost vaccination
819 with chimpanzee adenovirus and modified vaccinia Ankara encoding TRAP provides partial
820 protection against *Plasmodium falciparum* infection in Kenyan adults. *Sci Transl Med* 7,
821 286re285.
- 822 31. Gilbert, S.C. (2013). Clinical development of Modified Vaccinia virus Ankara vaccines. *Vaccine*
823 31, 4241-4246.
- 824 32. Mitja, O., Ogoina, D., Titanji, B.K., Galvan, C., Muyembe, J.J., Marks, M., and Orkin, C.M.
825 (2023). Monkeypox. *Lancet* 401, 60-74.
- 826 33. Greinacher, A., Thiele, T., Warkentin, T.E., Weisser, K., Kyrle, P.A., and Eichinger, S. (2021).
827 Thrombotic Thrombocytopenia after ChAdOx1 nCov-19 Vaccination. *N Engl J Med* 384, 2092-
828 2101.
- 829 34. Soboleva, K., Shankar, N.K., Yadavalli, M., Ferreira, C., Foskett, N., Putsepp, K., Ferstenberg,
830 L.B., Nord, M., da Silva, H.G., and Bhuyan, P. (2022). Geographical distribution of TTS cases
831 following AZD1222 (ChAdOx1 nCoV-19) vaccination. *Lancet Glob Health* 10, e33-e34.
- 832 35. Jin, J., Tarrant, R.D., Bolam, E.J., Angell-Manning, P., Soegaard, M., Pattinson, D.J., Dulal, P.,
833 Silk, S.E., Marshall, J.M., Dabbs, R.A., Nugent, F.L., et al. (2018). Production, quality control,
834 stability, and potency of cGMP-produced *Plasmodium falciparum* RH5.1 protein vaccine
835 expressed in *Drosophila* S2 cells. *NPJ Vaccines* 3, 32.
- 836 36. Rts, S.C.T.P. (2014). Efficacy and safety of the RTS,S/AS01 malaria vaccine during 18 months
837 after vaccination: a phase 3 randomized, controlled trial in children and young infants at 11
838 African sites. *PLoS Med* 11, e1001685.
- 839 37. Draper, S.J., and Heeney, J.L. (2010). Viruses as vaccine vectors for infectious diseases and
840 cancer. *Nat Rev Microbiol* 8, 62-73.
- 841 38. Erkeller-Yuksel, F.M., Deneys, V., Yuksel, B., Hannet, I., Hulstaert, F., Hamilton, C.,
842 Mackinnon, H., Stokes, L.T., Munhyeshuli, V., Vanlangendonck, F., and et al. (1992). Age-
843 related changes in human blood lymphocyte subpopulations. *J Pediatr* 120, 216-222.
- 844 39. Vargas-Inchaustegui, D.A., Demers, A., Shaw, J.M., Kang, G., Ball, D., Tuero, I., Musich, T.,
845 Mohanram, V., Demberg, T., Karpova, T.S., Li, Q., et al. (2016). Vaccine Induction of Lymph
846 Node-Resident Simian Immunodeficiency Virus Env-Specific T Follicular Helper Cells in Rhesus
847 Macaques. *J Immunol* 196, 1700-1710.
- 848 40. Wang, C., Hart, M., Chui, C., Ajuogu, A., Brian, I.J., de Cassan, S.C., Borrow, P., Draper, S.J.,
849 and Douglas, A.D. (2016). Germinal Center B Cell and T Follicular Helper Cell Responses to
850 Viral Vector and Protein-in-Adjuvant Vaccines. *J Immunol* 197, 1242-1251.
- 851 41. Nielsen, C.M., Ogbé, A., Pedroza-Pacheco, I., Doleman, S.E., Chen, Y., Silk, S.E., Barrett, J.R.,
852 Elias, S.C., Miura, K., Diouf, A., Bardelli, M., et al. (2021). Protein/AS01B vaccination elicits
853 stronger, more Th2-skewed antigen-specific human T follicular helper cell responses than
854 heterologous viral vectors. *Cell Rep Med* 2, 100207.

- 855 42. Draper, S.J., Moore, A.C., Goodman, A.L., Long, C.A., Holder, A.A., Gilbert, S.C., Hill, F., and
856 Hill, A.V. (2008). Effective induction of high-titer antibodies by viral vector vaccines. *Nat Med*
857 *14*, 819-821.
- 858 43. de Cassan, S.C., and Draper, S.J. (2013). Recent advances in antibody-inducing poxviral and
859 adenoviral vectored vaccine delivery platforms for difficult disease targets. *Expert Rev Vaccines*
860 *12*, 365-378.
- 861 44. Draper, S.J., Goodman, A.L., Biswas, S., Forbes, E.K., Moore, A.C., Gilbert, S.C., and Hill, A.V.
862 (2009). Recombinant viral vaccines expressing merozoite surface protein-1 induce antibody- and
863 T cell-mediated multistage protection against malaria. *Cell Host Microbe* *5*, 95-105.
- 864 45. Sebastian, S., Flaxman, A., Cha, K.M., Ulaszewska, M., Gilbride, C., Sharpe, H., Wright, E.,
865 Spencer, A.J., Dowall, S., Hewson, R., Gilbert, S., et al. (2020). A Multi-Filovirus Vaccine
866 Candidate: Co-Expression of Ebola, Sudan, and Marburg Antigens in a Single Vector. *Vaccines*
867 (Basel) *8*.
- 868 46. Jenkin, D., Ritchie, A.J., Aboagye, J., Fedosyuk, S., Thorley, L., Provstgaard-Morys, S., Sanders,
869 H., Bellamy, D., Makinson, R., Xiang, Z.Q., Bolam, E., et al. (2022). Safety and immunogenicity
870 of a simian-adenovirus-vectored rabies vaccine: an open-label, non-randomised, dose-escalation,
871 first-in-human, single-centre, phase 1 clinical trial. *Lancet Microbe* *3*, e663-e671.
- 872 47. Stedman, A., Wright, D., Wichgers Schreur, P.J., Clark, M.H.A., Hill, A.V.S., Gilbert, S.C.,
873 Francis, M.J., van Keulen, L., Kortekaas, J., Charleston, B., and Warimwe, G.M. (2019). Safety
874 and efficacy of ChAdOx1 RVF vaccine against Rift Valley fever in pregnant sheep and goats.
875 *NPJ Vaccines* *4*, 44.
- 876 48. Li, G., Cappuccini, F., Marchevsky, N.G., Aley, P.K., Aley, R., Anslow, R., Bibi, S., Cathie, K.,
877 Clutterbuck, E., Faust, S.N., Feng, S., et al. (2022). Safety and immunogenicity of the ChAdOx1
878 nCoV-19 (AZD1222) vaccine in children aged 6-17 years: a preliminary report of COV006, a
879 phase 2 single-blind, randomised, controlled trial. *Lancet* *399*, 2212-2225.
- 880 49. Leroux-Roels, G., Leroux-Roels, I., Clement, F., Ofori-Anyinam, O., Lievens, M., Jongert, E.,
881 Moris, P., Ballou, W.R., and Cohen, J. (2014). Evaluation of the immune response to
882 RTS,S/AS01 and RTS,S/AS02 adjuvanted vaccines: randomized, double-blind study in malaria-
883 naive adults. *Hum Vaccin Immunother* *10*, 2211-2219.
- 884 50. Venkatraman, N., Tiono, A.B., Bowyer, G., Powlson, J., Collins, K.A., Coulibaly, S., Dato, M.,
885 Silman, D., Ouedraogo, A., Nébié, I., Imoukhuede, E., et al. (2019). Phase I assessments of first-
886 in-human administration of a novel malaria anti-sporozoite vaccine candidate, R21 in matrix-M
887 adjuvant, in UK and Burkina Faso volunteers. *medRxiv*, 19009282.
- 888 51. Barrett, J.R., Belij-Rammerstorfer, S., Dold, C., Ewer, K.J., Folegatti, P.M., Gilbride, C.,
889 Halkerston, R., Hill, J., Jenkin, D., Stockdale, L., Verheul, M.K., et al. (2021). Phase 1/2 trial of
890 SARS-CoV-2 vaccine ChAdOx1 nCoV-19 with a booster dose induces multifunctional antibody
891 responses. *Nat Med* *27*, 279-288.
- 892 52. Foquet, L., Schafer, C., Minkah, N.K., Alanine, D.G.W., Flannery, E.L., Steel, R.W.J., Sack,
893 B.K., Camargo, N., Fishbaugher, M., Betz, W., Nguyen, T., et al. (2018). Plasmodium falciparum
894 Liver Stage Infection and Transition to Stable Blood Stage Infection in Liver-Humanized and
895 Blood-Humanized FRGN KO Mice Enables Testing of Blood Stage Inhibitory Antibodies
896 (Reticulocyte-Binding Protein Homolog 5) In Vivo. *Front Immunol* *9*, 524.
- 897 53. Miura, K., Perera, S., Brockley, S., Zhou, H., Aebig, J.A., Moretz, S.E., Miller, L.H., Doumbo,
898 O.K., Sagara, I., Dicko, A., Ellis, R.D., et al. (2011). Non-apical membrane antigen 1 (AMA1)
899 IgGs from Malian children interfere with functional activity of AMA1 IgGs as judged by growth
900 inhibition assay. *PLoS One* *6*, e20947.
- 901 54. Nielsen, C.M., Barrett, J.R., Davis, C., Fallon, J.K., Goh, C., Michell, A.R., Griffin, C., Kwok,
902 A., Loos, C., Darko, S., Laboune, F., et al. (2023). Delayed boosting improves human antigen-
903 specific Ig and B cell responses to the RH5.1/AS01B malaria vaccine. *JCI Insight* *8*.

- 904 55. Carey, J.B., Vrdoljak, A., O'Mahony, C., Hill, A.V., Draper, S.J., and Moore, A.C. (2014).
905 Microneedle-mediated immunization of an adenovirus-based malaria vaccine enhances antigen-
906 specific antibody immunity and reduces anti-vector responses compared to the intradermal route.
907 *Sci Rep* 4, 6154.
- 908 56. Vrdoljak, A., McGrath, M.G., Carey, J.B., Draper, S.J., Hill, A.V., O'Mahony, C., Crean, A.M.,
909 and Moore, A.C. (2012). Coated microneedle arrays for transcutaneous delivery of live virus
910 vaccines. *J Control Release* 159, 34-42.
- 911 57. Provine, N.M., Amini, A., Garner, L.C., Spencer, A.J., Dold, C., Hutchings, C., Silva Reyes, L.,
912 FitzPatrick, M.E.B., Chinnakannan, S., Oguti, B., Raymond, M., et al. (2021). MAIT cell
913 activation augments adenovirus vector vaccine immunogenicity. *Science* 371, 521-526.
- 914 58. de Cassan, S.C., Forbes, E.K., Douglas, A.D., Milicic, A., Singh, B., Gupta, P., Chauhan, V.S.,
915 Chitnis, C.E., Gilbert, S.C., Hill, A.V., and Draper, S.J. (2011). The requirement for potent
916 adjuvants to enhance the immunogenicity and protective efficacy of protein vaccines can be
917 overcome by prior immunization with a recombinant adenovirus. *J Immunol* 187, 2602-2616.
- 918 59. Flynn, O., Dillane, K., Lanza, J.S., Marshall, J.M., Jin, J., Silk, S.E., Draper, S.J., and Moore,
919 A.C. (2021). Low Adenovirus Vaccine Doses Administered to Skin Using Microneedle Patches
920 Induce Better Functional Antibody Immunogenicity as Compared to Systemic Injection. *Vaccines*
921 (Basel) 9.
- 922 60. Miura, K., Orcutt, A.C., Muratova, O.V., Miller, L.H., Saul, A., and Long, C.A. (2008).
923 Development and characterization of a standardized ELISA including a reference serum on each
924 plate to detect antibodies induced by experimental malaria vaccines. *Vaccine* 26, 193-200.
- 925 61. Mwamlima, T.G., Mwakasungula, S.M., Mkindi, C.G., Tambwe, M.M., Mswata, S.S.,
926 Mbwambo, S.G., Mboya, M.F., Draper, S.J., Silk, S.E., Mpina, M.G., Vianney, J.M., et al.
927 (2022). Understanding the role of serological and clinical data on assessing the dynamic of
928 malaria transmission: a case study of Bagamoyo district, Tanzania. *Pan Afr Med J* 43, 60.
- 929 62. Adegnika, A.A., Verweij, J.J., Agnandji, S.T., Chai, S.K., Breitling, L.P., Ramharter, M., Frolich,
930 M., Issifou, S., Kremsner, P.G., and Yazdanbakhsh, M. (2006). Microscopic and sub-microscopic
931 *Plasmodium falciparum* infection, but not inflammation caused by infection, is associated with
932 low birth weight. *Am J Trop Med Hyg* 75, 798-803.
- 933 63. Kamau, E., Alemayehu, S., Feghali, K.C., Saunders, D., and Ockenhouse, C.F. (2013). Multiplex
934 qPCR for detection and absolute quantification of malaria. *PLoS One* 8, e71539.
- 935 64. Malkin, E.M., Diemert, D.J., McArthur, J.H., Perreault, J.R., Miles, A.P., Giersing, B.K., Mullen,
936 G.E., Orcutt, A., Muratova, O., Awkal, M., Zhou, H., et al. (2005). Phase 1 clinical trial of apical
937 membrane antigen 1: an asexual blood-stage vaccine for *Plasmodium falciparum* malaria. *Infect*
938 *Immun* 73, 3677-3685.
- 939

940 **Figure Legends**

941 **Figure 1. VAC070 flow chart of study design and participant recruitment.**

942 Screening into the VAC070 study began in March 2018 and all follow-up visits were completed by
943 February 2019. All immunizations were administered IM into the left deltoid area.

944

945 **Figure 2. Solicited AEs following vaccination with full dose ChAd63 and MVA RH5.**

946 The solicited local and systemic AEs recorded for 7 days following ChAd63 RH5 prime and MVA RH5
947 boost are shown at the maximum severity reported by all participants within the VAC070 trial Groups 1,
948 2B and 3B. Participants aged (A) 18-35 years (n=6); (B) 1-6 years (n=12); and (C) 6-11 months (n=12)
949 received the full priming dose of 5×10^{10} vp ChAd63 RH5 on day 0. The same participants aged (D) 18-
950 35 years (n=5); (E) 1-6 years (n=12); and (F) 6-11 months (n=11) received the full boosting dose of $2 \times$
951 10^8 pfu MVA RH5 on day 56. Rabies vaccine control data are shown in **Figure S1** and data for the lower
952 lead-in dose of ChAd63-MVA RH5 (trial Groups 2A and 3A) are shown in **Figure S2**.

953

954 **Figure 3. *Ex-vivo* IFN- γ T cell response to ChAd63-MVA RH5 vaccination.**

955 Median and individual *ex-vivo* IFN- γ enzyme-linked immunospot (ELISPOT) responses to RH5 per
956 million peripheral blood mononuclear cells (PBMC) are shown for all groups receiving the full dose of
957 ChAd63-MVA RH5 vaccines and rabies vaccine control. Responses are shown at (A) day 14, two weeks
958 post-ChAd63 RH5 prime; and (B) day 84, four weeks post-MVA RH5 boost. Group 1 adults 18-35 years,
959 N=5 or 6; Group 2B children 1-6 years, N=12; Group 3B infants 6-11 months, N=11 or 12; and rabies
960 vaccine control, all groups and ages pooled, N=15. Historical data testing the identical dose and regimen
961 of the ChAd63-MVA RH5 vaccine in healthy UK adults 18-50 years are shown for comparison only: day
962 14, N=20; and day 84, N=8¹¹. (C) Lymphocyte counts per mL of blood for all vaccinated participants at
963 day 84. Group 1 adults 18-35 years, N=8; Group 2B children 1-6 years, N=18; Group 3B infants 6-11
964 months, N=17. (D) Median and individual *ex-vivo* IFN- γ ELISPOT responses to RH5 per mL peripheral

965 blood are shown for all groups receiving the full dose of ChAd63-MVA RH5 vaccines and rabies vaccine
966 control at day 84. Groups as per panel (B). Analyses using Kruskal-Wallis test with Dunn's multiple
967 comparison test across Groups 1, 2B and 3B only; * $P < 0.05$, ** $P < 0.01$, and **** $P < 0.0001$.

968

969 **Figure 4. Serum antibody response to vaccination.**

970 Median and individual anti-RH5_FL serum total IgG responses as measured by ELISA are shown for all
971 groups receiving the full dose of ChAd63-MVA RH5 vaccines and rabies vaccine control. Responses are
972 shown at (A) day 14, two weeks post-ChAd63 RH5 prime; and (B) day 63, one week post-MVA RH5
973 boost. Group 1 adults 18-35 years, N=5 or 6; Group 2B children 1-6 years, N=12; Group 3B infants 6-11
974 months, N=11 or 12; and rabies vaccine control, all groups and ages pooled, N=15. Baseline = day 0 pre-
975 vaccination time-point shown for Groups 1, 2B and 3B combined and color-coded. Historical data testing
976 the identical dose and regimen of the ChAd63-MVA RH5 vaccine in healthy UK adults 18-50 years are
977 shown for comparison only: day 14, N=20; and day 63, N=8¹¹. The horizontal dotted line indicates the
978 limit of detection of the assay. Responses to the "lead in" vaccine doses are shown in **Figure S4. (C)**
979 Median and individual anti-ChAd63 serum IgG endpoint titers as measured by ELISA in pre-vaccination
980 (day 0) samples. Solid symbols = participants who subsequently underwent ChAd63-MVA RH5
981 vaccination and open symbols = those who received rabies vaccine control. UK adults 18-50 years, N=8
982¹¹; Group 1 adults 18-35 years, N=9; Group 2B children 1-6 years, N=18; Group 3B infants 6-11 months,
983 N=18. (D) Median and individual anti-rabies glycoprotein serum total IgG responses as measured by
984 ELISA in day 63 samples (after two vaccine doses) are shown for all individuals receiving rabies vaccine
985 control: Group 1 adults 18-35 years, N=3; Group 2 children 1-6 years, N=9; Group 3 infants 6-11 months,
986 N=9; open symbols = Groups 2A and 3A, closed symbols = Groups 2B and 3B; as well as the ChAd63-
987 MVA RH5 vaccinees (N=28, Groups 1, 2B and 3B combined). Analyses shown used Kruskal-Wallis test
988 with Dunn's multiple comparison test across Groups 1, 2B and 3B only; * $P < 0.05$, ** $P < 0.01$.

989

990 **Figure 5. B cell response to ChAd63-RH5 vaccination.**

991 (A) RH5-specific antibody-secreting cell (ASC) responses were assessed by *ex-vivo* enzyme-linked
992 immunospot (ELISPOT) using RH5_FL protein and fresh peripheral blood mononuclear cells (PBMC)
993 from the day 63 time-point. Responses are reported as RH5-specific ASC per million PBMC used in the
994 assay. Group 1 adults 18-35 years, N=5; Group 2B children 1-6 years, N=12; Group 3B infants 6-11
995 months, N=11; Baseline = Groups 1, 2B and 3B combined, N=30; Rabies = Groups 1, 2B and 3B
996 combined, N=15. (B) Representative flow cytometry plots showing definition of RH5-specific cells
997 within the live CD19⁺ IgD⁻ IgM⁻ IgA⁻ B cell population (see **Figure S5A** for full gating strategy).
998 Representative plots are shown for matched day 0 (pre-vaccination) and day 63 (one week post-MVA
999 RH5 boost vaccination) samples from a single vaccinee. (C) Number of live CD19⁺ B cells per microliter
1000 of blood, as calculated by combining data from lymphocyte counts per microliter of blood (**Figure S5B**)
1001 and percentage of live CD19⁺ B cells within lymphocytes as assessed by flow cytometry (**Figure S5C**).
1002 (D) Percentage of RH5-specific cells within the CD19⁺ IgD⁻ IgM⁻ IgA⁻ B cell population as assessed by
1003 flow cytometry. Same individuals and color coding for age groups as shown in (C) measured at baseline
1004 day 0 (d0) or at 7 days post-MVA RH5 boost vaccination (V, closed symbols); N=3 individuals were
1005 measured at 28 days post-MVA RH5 boost due to sample availability (open symbols). Data from baseline
1006 samples for N=3 healthy UK adult participants 18-50 years of age enrolled in a previous trial of the
1007 ChAd63-MVA RH5 vaccine¹¹ are shown for comparison only (pink diamonds to left of dashed line).
1008 Individual and median results are shown in each panel. Analyses used Kruskal-Wallis test with Dunn's
1009 multiple comparison test across the three groups in (C) or the three vaccinated groups in (D); ** $P < 0.01$,
1010 *** $P < 0.001$.

1011

1012 **Figure 6. Functional GIA induced by ChAd63-MVA RH5 vaccination.**

1013 (A) *In vitro* growth inhibition activity (GIA) of purified IgG was assessed against 3D7 clone *P.*
1014 *falciparum* parasites. Total IgG purified from serum at baseline (day 0) and the day 63 time-point was

1015 initially tested at each sample's physiological total serum IgG concentration as measured by HPLC
1016 (**Figure S6A**). Data shown for Group 1 adults 18-35 years, N=5 vaccinees and N=3 control; Group 2B
1017 children 1-6 years, N=12 vaccinees and N=5 or 6 control (one sample was lost during analysis); and
1018 Group 3B infants 6-11 months, N=11 vaccinees and N=6 control. (**B**) Median and individual anti-parasite
1019 lysate serum IgG responses as measured by ELISA in arbitrary units (AU) in pre-vaccination (day 0)
1020 samples. Group 1 adults 18-35 years, N=9; Group 2A and 2B children 1-6 years, N=27; Group 3A and 3B
1021 infants 6-11 months, N=27. Analysis using Kruskal-Wallis test with Dunn's multiple comparison test; * P
1022 < 0.05 , *** $P < 0.001$, **** $P < 0.0001$. (**C**) Day 63 samples from Group 2B children 1-6 years (N=12)
1023 and Group 3B infants 6-11 months (N=11) vaccinated with ChAd63-MVA RH5 were subsequently
1024 titrated in the GIA assay using a 2-fold dilution series and starting at each sample's physiological total
1025 serum IgG concentration. GIA responses were later interpolated for 2.5 and 10 mg/mL total IgG (dotted
1026 lines), and the GIA assay EC_{50} of each purified IgG in mg/mL was also interpolated from the data, i.e. the
1027 concentration of total IgG that, following titration, showed 50% GIA (dashed line). (**D**) Relationship
1028 between GIA data from the dilution series shown in (**C**) and concentration of anti-RH5_FL purified IgG
1029 used in the assay as measured by ELISA. Non-linear regression curves are shown for all samples
1030 combined in Group 2B (blue line, $r^2 = 0.92$, N=108) and in Group 3B (green line, $r^2 = 0.95$, N=81). The
1031 EC_{50} (concentration of anti-RH5_FL polyclonal IgG that gives 50% GIA, dashed line) was calculated. (**E**)
1032 GIA data interpolated from (**C**) at 10 mg/mL total IgG and (**F**) at 2.5 mg/mL total IgG. Analysis using
1033 Mann-Whitney test comparing Groups 2B and 3B only; * $P < 0.05$, ** $P < 0.01$. Historical GIA assay data
1034 are also shown for comparison only in (**F**) for the RH5.1/AS01_B¹⁷ (N=15) and AMA1 FMP2.1/AS01¹⁸
1035 (N=12) protein-in-adjuvant vaccines previously tested in healthy UK adults. (**G**) Individual GIA assay
1036 EC_{50} of each purified IgG in mg/mL interpolated from the data in (**C**). Samples for which the GIA was
1037 $< 50\%$ at the top concentration of total IgG tested were plotted as 14 mg/mL (above the dashed black line).
1038 (**H**) To relate the GIA assay results (using purified total IgG) back to the original sera, the concentration
1039 of IgG in each original serum sample was also measured by HPLC (**Figure S6A**). This enabled

1040 calculation of the “GIA₅₀ Titer”, defined previously⁹ as the dilution factor of each serum sample required
1041 to reach the concentration of purified IgG that gives 50% GIA (i.e. the GIA EC₅₀). Samples for which the
1042 GIA₅₀ titer could not be calculated (because they did not achieve ≥50% GIA using purified IgG) are
1043 plotted arbitrarily at 0.8. In **(G)** and **(H)**, individual data and median results are shown for the same
1044 samples shown in **(F)**. Red dashed lines indicate threshold level of GIA at 2.5 mg/mL total IgG in **(F)** and
1045 a GIA₅₀ titer > 5 in **(H)** above which protection against a stringent *P. falciparum* blood-stage challenge
1046 has been reported in *Aotus* monkeys vaccinated with RH5⁹.

Figure 1





Figure 2

Prime

Boost


A

ChAd 63 RH5 Group 1 (18-35yrs)




D

MVA RH5 Group 1 (18-35yrs)




B

ChAd 63 RH5 Group 2B (1-6yrs)




E

MVA RH5 Group 2B (1-6yrs)




C

ChAd 63 RH5 Group 3B (6-11mos)



F

MVA RH5 Group 3B (6-11mos)



None Mild Moderate Severe

Figure 3






Figure 4


A



B



C



D




Figure 5






Figure 6


A




B




C




D




E



F



G



H

