Title: Patient consent for medical student pelvic exams under anesthesia: an exploratory retrospective chart review

Short title: Consent for pelvic exams under anesthesia

Authors: Jessica A. Jushchyshyn
Beth Israel Deaconess Medical Center
1309 Beacon, Office 214
Brookline, MA 02446
Email: jjushchy@bidmc.harvard.edu

Lakeisha Mulugeta-Gordon, MD
Resident, Department of Obstetrics & Gynecology
Hospital of the University of Pennsylvania
Philadelphia, PA 19104
Email: lakeisha.mulugeta-gordon@pennmedicine.upenn.edu

Cara Curley, DNP, CRNP
Department of Obstetrics & Gynecology
Perelman School of Medicine at the University of Pennsylvania
Email: cara.curley@pennmedicine.upenn.edu

Florencia Greer Polite, MD
Chief, Division of General Obstetrics & Gynecology
Perelman School of Medicine at the University of Pennsylvania
3701 Market Street
Philadelphia, PA 19104
Email: florencia.polite@pennmedicine.upenn.edu

Jon F. Merz, JD, PhD * (0000-0002-0612-7104)
Associate Professor of Medical Ethics & Health Policy
Perelman School of Medicine at the University of Pennsylvania
Blockley Hall Floor 14, 423 Guardian Drive
Philadelphia, Pennsylvania 19104-4884
Email: merz@pennmedicine.upenn.edu
Phone: 267-254-6470

* Communicating author.

Word count: Abstract: 357
Manuscript: 2223

Acknowledgements:

The authors thank Penn Data and Analytics and Ting-shan Chiu for their assistance in assembling the information used in this study, and Kavita Shah Arora, Andrew Fesnak, and several anonymous reviewers for comments. Responsibility for the study is solely that of the authors.

There was no funding for this study.

Author Contribution Statement

This study was conducted in partial fulfillment of the requirements for the Masters in Regulatory Affairs at the University of Pennsylvania (JJ). JFM and LMG independently conceived the idea for the study; JJ, JFM, and FGP laid out the approach; JJ, JFM, and CC worked iteratively with the data core to develop the EPIC search; JFM checked primary coding and performed secondary coding, JJ checked the coding, and JFM performed the data analysis; JJ wrote the first draft of the manuscript, JFM wrote a second draft and managed all subsequent editing, and the authors contributed equally to the interpretation of data and the editing of the manuscript.

A preprint of this paper has been posted on MedRxiv, DOI:

Competing Interests:

Jon Merz: Since 1997, JFM has been a paid expert witness in 3 civil cases involving the adequacy of informed consent and IRB review in research, twice for defense and once for plaintiff, as well as a 4th case involving the definition of human subjects research. In the last 3 years, JFM has received financial compensation for service on several Data and Safety Monitoring Boards for the NIH and the American College of Radiology Imaging Network, and

for service on a pharmacogenomics ethics advisory board for Merck. JFM received partial salary

support as moderator of the IRBForum (https://community.primr.org/home) by a grant from

Public Responsibility in Medicine & Research (PRIMR) from 2012 through 2020.

Title: Patient consent for medical student pelvic exams under anesthesia: an exploratory retrospective chart review

Precis: Racial, ethnic, and religious minority patients are more likely than others to refuse to allow medical students to perform pelvic examinations under anesthesia.

ABSTRACT

Background: Legal requirements and clinical practices of securing patient consent for medical student pelvic examinations under anesthesia (EUA) vary widely, while ethical arguments and patients' preferences for being asked for consent are well known.

Objective: This study was performed to examine patients' choices to permit or refuse medical student pelvic EUAs during planned gynecologic procedures.

Study Design: An exploratory retrospective chart review of electronic consent forms at a single academic medical center, using contingency table and logistic regression to explore relationships between patient and provider characteristics and consent.

Results: Electronic consent forms were downloaded for a census of 4000 patients undergoing gynecologic surgery from September 2020 through calendar year 2022 and linked to anonymized medical record information, including patient age, race, religion, and insurance carrier, along with physician name. Physicians were coded by gender, departmental affiliation, and status (attending, resident, or fellow). Of the 4000 patients, 142 (3.6%) were informed but not presented with a choice, and these patients were removed from further analysis. Of the remainder, 308 (8.0%) were asked for EUA consent more than once. Overall, of 3858 patients, 3308 (85.7%) consented every time asked and 550 (14.2%) refused or limited EUA consent at least once. Nine patients limited their consent to female students, and 2 patients refused medical student participation at all. Of the 308 asked more than once, 46 were not consistent. Exploratory

multiple logistic regression analysis showed that patients identifying as Black or African American (OR=0.492, p<0.001) or Asian (OR=0.292, p<0.001), or of Moslem/Muslim/Islamic faith (OR=0.579, p=0.006) were substantially less likely to grant EUA consent than other patients. Moreover, male physicians, most of whom were attendings, were much more likely to secure consent from patients than their female colleagues (OR=2.124, p<0.001).

Conclusions: The finding that some patients are more likely than others to refuse a pelvic EUA magnifies the dignitary harm from a nonconsensual invasion of intimate bodily integrity and perpetuates the historic wrongs visited upon vulnerable people of color and religious minorities. Patient's rights to control over their own bodies can only be respected if their physicians take seriously the ethical obligation to inform their patients and ask them for permission.

Introduction

Medical students typically perform examinations of patients for training purposes, normally securing permission from patients before doing so. It has not always been so, as ethical and legal standards for informed consent evolved only over the last 6 decades.[1-3] One continuing exception to this regards the performance of pelvic and prostate exams, which are often conducted on unconscious, sedated patients. Up until perhaps 30 years ago, it was common for such examinations under anesthesia (EUA) to be done without patient knowledge or consent, but the practice came under ethical and legal scrutiny in the 1980s and 1990s.[4-6] Patients and medical trainees almost uniformly view consent as necessary for such intimate exams, and professional guidelines recognized the need for prior consent.[7-10] Nonetheless, surveys of practices showed that consent was not uniformly sought from patients, nor is it legally mandated in all U.S. states, and this situation persists today.[11-18]

While patients express strong preferences to be informed and asked for consent, several small studies suggest that most patients will consent when asked. Bibby and colleagues reported that all 69 patients who were interviewed in their Birmingham (UK) hospital expressed the desire to be informed prior to a medical student EUA.[4] Wainberg and colleagues surveyed 102 patients at their Calgary clinic, asking about their preferences for being informed and willingness to permit a medical student EUA, reporting that a majority (72%) would want to be asked for consent. Of 93 patients who responded to a question about whether they would consent, 62% indicated they would give permission, 5% indicated they would only permit a female student to perform the EUA, 14% would refuse, and 18% were undecided.[19] Finally, one secondary source

refers to an unpublished 2021 retrospective chart review in a Chicago family planning clinic that reported 10.4% of patients refused medical student EUAs.[17]

The Department of Obstetrics and Gynecology (ObGyn) in the Perelman School of Medicine at the University of Pennsylvania instituted express consent for medical student EUAs 2 decades ago.[20] This was implemented as an electronic consent in 2016. For patients undergoing planned obstetric or gynecological procedures, the surgeon (who may be an attending, fellow, or resident) will ask patients if they are willing to allow a medical student to perform an EUA, and the patients' choice is recorded by capture of text and stored in EPIC back-end tables. While editable, two alternative forms of the last sentence of the text are provided for the physician from which to choose. See Figure 1. It is the purpose of this study to examine patients' actual choices and perform an exploratory analysis of demographic characteristics of patients and providers and consent to medical student EUAs.

[Insert Figure 1 about here]

Methods

We performed a single academic institution, multi-site, retrospective chart review, from a large urban teaching hospital. We drew a census from the electronic medical record of EUA consent forms for all patients from whom consent was sought, dating back to late 2018 through 2022. Consent forms were matched by date and patient IDs to patient demographics (age, race, religion, and insurance carrier) and visit details, which further enabled linking of physician name, department, specialty, and site location. Records were anonymized by the Penn Medicine Data and Analytics core, and provided to the researchers in anonymous form. Because some text

captured patient names, the study was approved by the Penn Institutional Review Board. All identifying information was irreversibly stripped from the records after coding.

Coding was initially conducted by the Data and Analytics analyst; one author manually checked 100% of the census by running text searches for key phrases unique to consent and refusal, as well as related key terms (e.g., male, refus!, decline, not, permit), reading the consent or refusal sentence for every form, and assigned numeric codes to categorical data. Coding from the manual step was checked by a second author.

We planned exploratory assessment using contingency tables and logistic regression to examine potential relationships between patient and provider characteristics and patients' choices to allow or restrict medical student EUA. All analyses were performed using Stata 12.1 (StataCorp, © 2014). Given the exploratory and hypothesis-generating nature of this analysis, we did not plan to adjust p-values for multiple comparisons.

Results

Data was captured and downloaded on February 2, 2023. While we attempted to search back to 2018 to enable inclusion of patients treated prior to the Covid-19 pandemic, our search method only provided access to consent forms on and after September 20, 2020; this was attributed to a change in system design. We thus captured every use of the EUA consent from September 20, 2020, through December 31, 2022. We have records for 4000 unique patients.

Patients were an average of 45.7 years of age (range: 16-88). There were 16 teenagers, and 39 patients in their 80s. Patients self-identified predominantly as white (1921, 48.0%), Black or African American (1613, 40.3%), Asian (165, 4.1%), unknown or patient declined (168, 4.2%), and other (including American Indian or Alaskan Native, East Indian, Hispanic/Latino Black, Hispanic/Latino White, and Native Hawaiian or other Pacific Island) (133, 3.3%).

Patients also self-identified with 2 dozen religions; grouping faiths in which fewer than 100 patients identified showed that just over half were Christian (2003, 50.1%), while religion is undocumented for many patients (1278, 32.0%). Other patients identified as Muslim/Moslem/Islamic (179, 4.5%), Jewish (151, 3.8%), and nondenominational and other (389, 9.7%). Information about insurance coverage or other payors was available for 3331 patients (83.3%); of whom almost two-thirds of patients had commercial health insurance (2170, 65.1%), while the remainder had Medicare, Medicaid, or other public coverage (1161, 34.8%).

Physicians were identified in 3440 of the cases in which EUA consent was sought. Attending physicians had the most patients (2704, 67.6%), while residents (643, 16.1%) and fellows (93, 2.3%) made up the remainder. Overall, 144 physicians were identified, 26 males and 118 females. Of the 26 males, 10 (38.5%) hold appointments in departments other than ObGyn, while only 5 of 118 female physicians (4.2%) are in other departments (χ^2=26.74 w/1 degree of freedom, p<0.001). As this implies, 113 of 129 ObGyn physicians (87.6%) are female. The 15 non-ObGyn physicians were involved in the care of only 26 patients, with 13 of the 15 involved in a single case. Because EUA consent was sought from these patients, we have left them in the census, but because of anonymization of records, we have no ready means for checking why EUA consent was sought in these cases.

Of the 4000 patients, 142 (3.6%) were informed but not presented with a choice, and these patients were removed from further analysis. For reasons unknown, these patients were only presented with a truncated consent form, which did not include the consent or refusal sentences (italicized sentences in Figure 1). These 142 cases have been dropped from further analysis. Of the remaining 3858 patients, 308 (8.0%) were asked for EUA consent more than once (on different days).

The unit of analysis for this study is the patient. We created 2 outcome variables: Consent and Flip. Consent is when a patient allows EUA every time asked. We assess nonconsent as any situation where a patient refuses EUA consent or limits consent to female medical students or otherwise expresses a wish that medical students not take part in their care. Flip is coded only for patients who were asked for EUA consent more than once, and reflects inconsistent choices for patients who sometimes said yes and at other times said no.

Overall, 550 of 3858 patients (14.2%) refused or limited EUA consent. Of the 550, 9 (1.6%) limited consent to only female medical students, including one who gave specific consent for one named female student, and 2 patients (0.4%) refused to have a medical student present at all during their surgery. Of the 308 patients asked more than once, 241 (78.2%) consistently said yes, 21 (6.8%) consistently refused, and 46 (14.9%) were inconsistent.

We performed exploratory univariate and multivariate contingency table and logistic regression analyses to examine whether any of the descriptive factors about patients or their physicians appear related to whether patients consented, and whether those who were asked for EUA consent more than once were consistent. Univariate analysis was consistent with multivariate analyses, so here we only present the results of a multiple logistic regression. This yields more robust estimates than univariate analysis, accounting for correlations amongst independent variables. Given the exploratory nature of this analysis, p-values have not been adjusted for multiple comparisons. Table 1 presents the results of this analysis of Consent. Note that there were no differences across physician specialties, so this variable was omitted from the model. A similar analysis of Flip found no relationship with any independent variables at lower than a 0.05 level of significance.

[Insert Table 1 about here]

Discussion

Our findings show that overall about 1 in 7 patients refuse consent for medical students to conduct an EUA. Consent rates vary substantially among identified racial groups, ranging from 9.5% of white patients to 18.9% of Black or African American patients and 25.3% of Asian patients. Overall, non-White patients were almost twice as likely as White patients to refuse (18.7%). Religious affiliation was also related, with 27.8% of patients identifying as Muslim/Moslem/Islamic refusing consent. A majority of patients identifying as Muslim are Black (147 of 169), and a test of the model with this interaction showed that the race and religion main effects remain significant while the interaction is not (p=0.09). Refusal rates amongst patients with other common religious affiliations varied from 0.0% of 58 Agnostic or Atheist patients, 28.6% of 21 Hindu patients, to 75% of 4 Christian Scientist patients. As shown in Table 1, patients on Medicare or Medicaid programs were also somewhat less likely to consent.

While patient age was not a significant predictor in our analysis, younger patients were the most likely to refuse medical student EUAs. The highest refusal rate was in patients under 30, with 17.1% of 422 patients saying "no". This included refusal by 6 of 16 teenagers (37.5%).

Physician characteristics also play a role. While the number of cases overseen by physicians who do not hold appointments in ObGyn was small, their patients were more likely to refuse the EUA (6 of 26 refused). Physician gender appears to play a role as well, with male physicians having a markedly lower refusal rate (9.2%) than their female colleagues (15.2%; χ^2=10.36 w/1 degree of freedom, p=0.001). As noted above, 87.6% of ObGyn physicians are female. Finally, we were unable to code physician role in 463 cases because of missing

information in the record, and these patients were substantially less likely to refuse consent than all others (5.2%).

We are unable to shed light on the reasons for the observed differences in rates of consent. Patients may distrust their providers, be concerned about their privacy, or simply not want unnamed medical students (of either gender) performing intimate examinations while they are unconscious. Patients may not fully grasp the role medical students may have during surgical procedures, which may involve genital exposure and contact, albeit for therapeutic and not training purposes. Moreover, the finding that male physicians are more likely to secure consent for a medical student EUA raises questions about the consent processes and about the perceived gender and power dynamics in the clinical encounter. It is possible that this finding reflects a selection bias, with patients who choose or accept a referral to a male physician perhaps more likely to agree to a pelvic examination by an unknown, and possibly male, medical student.

What is the moral significance of higher rates of refusal amongst patients in racial and religious groups? It is broadly (but not yet universally) viewed that consent is ethically necessary, and the fact that some patients would be more likely than others to say no heightens the dignitary harm from a nonconsensual invasion of intimate bodily integrity. The fact that this harm is more likely to fall on persons of color or religious minorities perpetuates the historic wrongs visited upon the vulnerable.[21] From our perspective, this bolsters the arguments for broader adoption of laws requiring informed consent from patients for nontherapeutic, educational interactions with medical trainees.[14-17]

One primary limitation on this analysis is that the physician identified in the medical record may not have been the physician seeking EUA consent, and there's no reliable way to determine from the medical record who secured consent in each case. Nonetheless, the difference

shown in rates of consent based on physician gender suggests that some aspect of the clinical encounter influences patients' choices. A second limitation is that physicians were not asked their preferred gender identities, with gender coded from each provider's institutional web page; we believe any imputed error is likely to be small, as a recent study showed that less than 2% of adults (including 5% of young adults under 30) identify as transgender or nonbinary.[22] Finally, our census is limited to patients treated during the Covid-19 pandemic and its aftermath, and whether this had an effect on patients' preferences is unknown.

In conclusion, we find that a majority of patients will permit medical students to conduct a pelvic EUA, when asked for their consent, but that some patients are substantially more likely to refuse than others. Patient's rights to control over their own bodies can only be respected if their physicians take seriously the ethical obligation to inform their patients and ask them for permission.

Figure 1 Alternative consent and refusal language of the electronic EUA consent form.

INFORMATION AND CONSENT FOR PELVIC EXAMINATION UNDER GENERAL ANESTHESIA After you are put to sleep with general anesthesia, A PELVIC EXAM may be performed as part of your planned gynecologic surgery. This examination is a standard part of many gynecologic procedures. Often, it provides the surgeon with valuable information for the safe conduct of the surgery. For example, findings on a pelvic examination might be used to determine the best surgical approach for your particular problem or the optimal location of the incision for your surgery. If a pelvic examination is planned as part of your surgical procedure, your physician will include this in the description of the surgical procedure for which you provide written consent. If a pelvic examination is performed after you are under general anesthesia, you will be draped in a manner similar to an examination in the office. Your surgeon may ask another physician member of the team to also perform a pelvic examination to confirm the findings or to provide an additional opinion. Performing a pelvic examination under anesthesia may also provide your surgeon with an opportunity to teach his/her surgical assistants regarding how to properly perform the procedure. If the surgical team will include a medical student, your physician also may ask you to read this information sheet and decide if you will permit a medical student, who will be assisting with the surgical procedure, to also perform a pelvic examination. Under no circumstances, will a medical student perform a pelvic examination under anesthesia without your permission. If you decide that you do not want a medical student to perform this examination, this decision will not affect your care in any way. I have read and understand this information sheet and my questions have been answered to my satisfaction. I certify that my physician has informed me that a pelvic examination is planned or may be necessary as part of my surgical procedure and I consented to this examination.*

Consent: *I understand that a medical student may assist my physician with the surgery. I hereby consent to permit a medical student who is part of the surgical team for my procedure to perform a pelvic examination under anesthesia as part of my procedure under the direct supervision of my physician.*

Refuse: *If a medical student is part of the surgical team, I request that the student not be permitted to perform a pelvic exam as part of my procedure.*

Notes: * The alternative italicized sentences appear without break in the paragraph, in normal (not italicized) font. This text is fully editable by the surgeon at the time consent is discussed with each patient.

Table 1 Multiple logistic regression results showing factors influencing the likelihood of pelvic EUA Consent

FACTOR	N	Odds Ratio	p-value
Patient Characteristics:			
Age (in years)	3858	1.0011	0.76
Race: (compared to White):			
Asian	158	0.292	<0.001
Black or African American	1541	0.492	<0.001
Other	115	0.726	0.26
Unknown	140	0.627	0.055
Religion: (compared to Christian)			
Jewish	144	0.608	0.064
Muslim/Moslem/Islamic	169	0.579	0.006
Other	374	1.104	0.58
Unknown	1236	0.847	0.15
Payors (compared to commercial health insurers)			
Public payors	1120	0.750	0.010
Unknown	625	1.154	0.34
Provider Characteristics:			
Department/Specialty (compared to non-ObGyn)			
ObGyn	3644	3.020	0.024
Gender (compared to female physician):			
Male physician	411	2.124	<0.001
Role (compared to attending physicians)			
Fellows	91	2.089	0.024
Residents	632	0.958	0.72
Unknown/other	464	6.517	<0.001

References

1. President's Commission for the Study of Ethical Problems in Medicine and Biomedical and Behavioral Research. Making health care decisions: the ethical and legal implications of informed consent in the practioner-patient relationship. Washington D.C., Vol. 1, October 1982, pp. 18-35.

2. Cohen DL, McCullough LB, Kessel RW, Apostolides AY, Heiderich KJ, Alden ER. A national survey concerning the ethical aspects of informed consent and role of medical students. *J Med Educ* 1988; 63:821-829.

3. Cohen DL, Kessel RWI, McCullough LB, Apostolides AY, Heiderich KJ, Alden ER. Pelvic examinations by medical students. *Am J Obstet Gynecol* 1989; 161:1013-1014.

4. Bibby J, Boyd N, Redman CWE, Luesley DM. Consent for vaginal examination by students on anaesthetised patients. *Lancet* 1988; 2(8620):1150.

5. York-Best, CM, Ecker JL. Pelvic examination under anesthesia: a teachable moment. *Obstet Gynecol* 2012; 120:741-742.

6. Bruce L. A pot ignored boils on: sustained calls for explicit consent of intimate medical exams. *HEC Forum* 2020; 32:125-145.

7. Adashi EY. Teaching pelvic examination under anesthesia without patient consent. *JAMA* 2019; 321:732-733.

8. Kaur K, Salwi S, McNew K, Kumar N, Millimet H, Ravichandran N, Tytus K, Zhang AY, Wood AG, Grimm B, Fairbrother EL. Medical student perspectives on the ethics of pelvic exams under anesthesia: a multi-institutional study. *J Surg Educ* 2022; 79:1413-1421.

9. Hammoud MM, Spector-Bagdady K, O'Reilly M, Major C, Baecher-Lind L. Consent for the pelvic examination under anesthesia by medical students: recommendations by the association of professors of gynecology and obstetrics. *Obstet Gynecol* 2019; 134:1303-1307.

10. Zuchelkowski BE, Eljamri S, McDonnell JE, Varma B, Stern NG, Rothenberger SD, Arora KS, Chang JC. Medical student attitudes on explicit informed consent for pelvic exams under anesthesia. *J Surg Educ* 2021; 79:676-685.

11. Coldicott Y, Pope C, Roberts C. The ethics of intimate examinations—teaching tomorrow's doctors. *BMJ* 2003; 326:97-101.

12. Ubel PA, Jepson C, Silver-Idenstadt A. Don't ask, don't tell: a chance in medical student attitudes after obstetrics/gynecology clerkships toward seeking consent for pelvic examinations on an anesthetized patient. *Am J Obstet Gynecol* 2003; 188:575-579.

13. Friesen P. Educational pelvic exams on anesthetized women: why consent matters. *Bioethics* 2018; 32:298-307.

14. Cundall HL, MacPhedran SE, Arora KS. Consent for pelvic examinations under anesthesia by medical students: historical arguments and steps forward. *Obstet Gynecol* 2019; 134:1298-1302.

15. Greene MF. Examining examinations conducted under anesthesia. *N Engl J Med* 2020; 383:1099-1101.

16. Friesen P, Wilson RF, Kim S, Goedken J. Consent for intimate exams on unconscious patients: sharpening legislative efforts. *Hast Cent Rep* 2022; 52(1):28-31.

17. Bruce L, Hannikainen IR, Earp BD. New findings on unconsented intimate exams suggest racial bias and gender parity. *Hast Cent Rep* 2022; 52(2):7-9.

18. Tillman S, Chor J. Educational pelvic examinations under anesthesia: recommendations for clinicians and learners. *J Clin Ethics* 2022; 33:347-351.

19. Wainberg A, Wrigley H, Fair J, Ross S. Teaching pelvic examinations under anaesthesia: what do women think? *J Obstet Gynaecol Can* 2010; 32:49-53.

20. University of Pennsylvania Health System, Department of Obstetrics & Gynecology Policy Statement. Pelvic Exams Under Anesthesia by Medical Students. Available at: https://www.med.upenn.edu/student/pelvic-exams-under-anesthesia-by-med-students.html (accessed March 25, 2023).

21. Cusanno BR. "What do you say if they say 'no'?": agency, materiality, and justice in pelvic exams. *Hlth Communication* 2023; doi: 10.1080/10410236.2022.2159134.

22. Brown A. About 5% of young adults in the U.S. say their gender is different from their sex assigned at birth. https://www.pewresearch.org/fact-tank/2022/06/07/about-5-of-young-adults-in-the-u-s-say-their-gender-is-different-from-their-sex-assigned-at-birth/ (accessed March 21, 2023).