

25

26 **Corresponding author:**

27 Lorena S. Pacheco, PhD

28 Harvard T.H. Chan School of Public Health

29 655 Huntington Ave., Bldg 2 room 302

30 Boston, MA 02115

31 Telephone: 617-981-4384

32 Email address: lpacheco@hsph.harvard.edu

33

34 The total word count of the manuscript (including Title Page, Abstract, Text, References, Tables and

35 Figures Legends): 7,947.

36

37

38

39

40

41

42

43

44

45

46

47

48

49 **Abstract**

50 **Background**

51 The extent to which physical activity attenuates the detrimental effects of sugar (SSBs)- or artificially-
52 sweetened beverages (ASBs) on the risk of cardiovascular disease is unknown.

53 **Methods**

54 We used Cox proportional-hazards models to calculate hazard ratios and 95% confidence interval [HR
55 (CI)] between SSB or ASB intake and physical activity with cardiovascular disease risk among 65,730
56 women in the Nurses' Health Study (1980-2016) and 39,418 men in the Health Professional's Follow-
57 up Study (1986-2016), who were free from chronic diseases at baseline. SSBs and ASBs were assessed
58 every 4-years and physical activity biannually.

59 **Results**

60 A total of 13,269 cardiovascular events were ascertained during 3,001,213 person-years of follow-up.
61 Compared with those that never/rarely consumed SSBs or ASBs, HR and 95% CI for cardiovascular
62 disease for participants consuming ≥ 2 servings/day were 1.21 (95% CI, 1.12 to 1.32; P -trend <0.001)
63 and 1.03 (95% CI, 0.97 to 1.09; P -trend=0.06), respectively. In the joint analyses, for participants
64 meeting and not meeting physical activity guidelines (<7.5 vs ≥ 7.5 MET-h/week) as well as consuming
65 ≥ 2 servings/day of SSBs or ASBs, the HRs for cardiovascular disease were 1.15 (95% CI, 1.08 to 1.23)
66 and 0.96 (95% CI, 0.91 to 1.02), and 1.47 (95% CI, 1.37 to 1.57) and 1.29 (95% CI, 1.22 to 1.37)
67 respectively, compared with participants who met physical activity guidelines and never/rarely
68 consumed these beverages. Similar patterns were observed when coronary heart disease and stroke
69 were analyzed.

70 **Conclusions**

71 Our findings suggest that among physically active participants, higher SSB intake, but not ASBs, is
72 associated with a higher cardiovascular risk. Our results support current recommendations to limit the
73 intake of SSB and maintain adequate physical activity levels.

74

75 **Keywords:** joint association, sucrose, aspartame, sugary beverages, diet drinks, coronary heart disease,
76 stroke

77

78 **Clinical Perspective**

79 What is new?

- 80 • Consumption of sugar- or artificially-sweetened beverages and physical activity are
81 independently associated with cardiovascular disease; nevertheless, the role that physical
82 activity plays in the associations between sugar- and artificially-sweetened beverages
83 consumption and cardiovascular disease warrants further analysis since physical activity could
84 offset the health risk that would have been induced by their consumption.
- 85 • In two US cohorts, not meeting physical activity guidelines jointly with a greater intake of
86 sugar- or artificially-sweetened beverages was associated with a higher risk of cardiovascular
87 disease; however, even when individuals were physically active, higher consumption of sugar-
88 sweetened beverages was associated with a higher cardiovascular risk.

89

90 What are the clinical implications?

- 91 • Our results provide further evidence of the intake of sugar-sweetened beverages and low
92 physical activity as factors negatively impacting cardiovascular health.
- 93 • Our results support public health recommendations and policies to limit the intake of sugar-
94 sweetened beverages and meet and maintain adequate physical activity levels.

95

96

97

98 **Introduction**

99 Sugar-sweetened beverages (SSBs) have a detrimental impact on health¹⁻⁵ and are associated
100 with a higher risk of cardiovascular disease (CVD),^{1,6-11} the leading cause of mortality worldwide.¹²
101 Although there has been a nationwide decline in SSB consumption since 2000,¹³ they remain the single
102 largest source of calories and added sugar in the diet.¹⁴ In contrast, the intake of artificially-sweetened
103 beverages (ASBs) has increased in the United States (US).¹⁵ They are used as ‘healthier’ alternatives to
104 SSBs and may be an effective strategy to help control caloric intake and support weight loss.¹⁶
105 However, evidence on the long-term health impact of ASBs on incident CVD is still inconclusive.¹⁷⁻¹⁹
106 Furthermore, the role that physical activity plays in the association between SSB or ASB consumption
107 and the risk of CVD warrants further analysis. Independently, physical activity is cardioprotective,^{20,21}
108 and the majority of SSB or ASB intake studies adjust for physical activity. A counterargument is the
109 probable role that physical activity could play to offset the potential health risk that would have been
110 induced by SSB or ASB intake. Thus, the implication is that given the same level of exposure to SSB
111 or ASB intake, those who meet physical activity recommendations would have a relatively lower risk
112 of CVD than those who are not physically active. Therefore, the aim of this study is to examine the
113 independent and joint associations between SSB or ASB consumption and physical activity in relation
114 to the risk of CVD in men and women from two large US prospective cohort studies.

115 **Methods**

116 *Study Population and Design*

117 The Nurses’ Health Study (NHS), a prospective study initiated in 1976, enrolled 121,700
118 registered nurses aged 30-55 years. The Health Professionals Follow-up Study (HPFS), a prospective
119 study established in 1986, enrolled 51,529 US male health professionals aged 40-75 years. Baseline
120 and follow-up questionnaires were sent to participants every 2-years to update medical, lifestyle, and
121 health information. Follow-up rates exceed 90% for each 2-year cycle. Diet was assessed using a self-
122 administered food frequency questionnaire every 4-years, first collected in 1980 in NHS and 1986 for

123 HPFS, used as this study's baseline cycles. A detailed description of the two cohorts has been
124 previously reported.^{22,23}

125 We excluded individuals with a history of diabetes mellitus, CVD, or cancer at or before
126 baseline; missing information regarding SSB and physical activity; or implausible caloric intake (<500
127 or >3500 kcal/day in women, and <800 kcal or >4200 kcal/day in men). The final analysis included
128 65,730 women and 39,418 men. The protocol was approved by the institutional review board of
129 Brigham and Women's Hospital and Harvard T.H. Chan School of Public Health. All participants gave
130 informed consent.

131 *Dietary Assessment and Sugar-Sweetened Beverage Intake*

132 Intakes of SSBs and ASBs were assessed using a validated and reproducible food frequency
133 questionnaire designed to assess the usual diet over the previous year. The reproducibility and validity
134 of these instruments have been described previously.^{24,25} Participants were asked how often, on
135 average, they consumed a standard portion of foods and beverages (one standard glass, bottle, or can),
136 with possible responses ranging from 'never or less than once/month to '≥6 times/day'. SSBs were
137 defined as carbonated sugar-sweetened beverages (caffeinated colas, caffeine-free colas, or other), and
138 noncarbonated sugar-sweetened beverages (fruit punches, lemonades, or other fruit drinks). Fruit juices
139 were not considered SSBs since sugar is naturally present. ASBs were defined as caffeinated, caffeine-
140 free, and noncarbonated low-calorie or diet beverages.

141 *Physical Activity Assessment*

142 Participants' physical activity was assessed every 4-years by reporting the average time per
143 week spent in various moderate- or vigorous-intensity leisure-time activities in the preceding year,
144 using a validated self-reported questionnaire.²⁶ Based on the intensity and duration of each activity,
145 weekly expenditure in metabolic equivalent tasks (MET-hr/week) was calculated and summed over all
146 activities for total physical activity.

147 *Ascertainment of Cardiovascular Disease Incidence*

148 The primary outcome was incident total CVD, defined as a composite of fatal and nonfatal
149 coronary heart disease (CHD, including myocardial infarction) and fatal and nonfatal stroke.
150 Secondary outcomes were CHD, defined as fatal CHD or nonfatal myocardial infarction, and stroke. In
151 both the NHS and HPFS cohorts, when a participant (or family members) reported an incident event,
152 permission was requested to examine their medical records by the study investigator-physicians who
153 were blinded to the participant's risk factor status. For each endpoint, the month and year of diagnosis
154 were recorded as the diagnosis date. Nonfatal events were confirmed through a review of medical
155 records. Myocardial infarction was defined according to the World Health Organization criteria.²⁷
156 Strokes were confirmed if data in the medical records fulfilled the National Survey of Stroke criteria.²⁸
157 Strokes were classified as ischemic stroke (thrombotic or embolic occlusion of a cerebral artery),
158 hemorrhagic stroke (subarachnoid and intraparenchymal hemorrhage), or stroke of probable and/or
159 unknown subtype (subtype data not available). Death ascertainment was performed by searching the
160 National Death Index,²⁹ by family members' responses to follow-up questionnaires, or by reports from
161 participants' professional organizations. We requested access to medical records, autopsy reports, and
162 death certificates to confirm all suspected deaths due to myocardial infarction. Fatal myocardial
163 infarction was confirmed by medical records or autopsy reports. We included all confirmed and
164 probable cases in our report because results were similar after probable cases were excluded. Follow-
165 up for deaths was >98% complete.

166 *Assessment of Covariates*

167 Information on lifestyle and CVD risk factors was assessed and updated every other year and
168 included: age; weight; smoking status; use of aspirin, multivitamins, postmenopausal hormone-
169 replacement therapy, and oral contraceptives; menopausal status; and hypertension or
170 hypercholesterolemia that had been recently diagnosed by a physician; and family history of chronic
171 diseases. Height was ascertained for women in 1976 and for men in 1986. Height and body weight
172 were used to calculate body mass index (BMI; weight in kilograms divided by height in meters

173 squared). Alcohol intake was assessed, and updated, from the food frequency questionnaires every 4
174 years. Ancestry information was collected in 1986 for men and in 1992 for women. The Alternate
175 Healthy Eating Index (AHEI, range 0-90; higher score indicating adherence to Dietary Guidelines for
176 Americans) was calculated from food frequency questionnaire data, without the exposure (SSBs) and
177 alcohol (we considered as a separate confounder) components, to indicate participant's overall diet
178 quality.³⁰

179 *Statistical Analysis*

180 Person-years of follow-up were calculated from the return of the baseline questionnaire to the date of
181 diagnosis of CVD, death, or end of follow-up whichever came first. We used Cox proportional-hazards
182 models to estimate hazard ratios and 95% confidence intervals (HRs [CIs]) of developing CVD
183 according to the independent and joint associations of SSB or ASB intakes and physical activity.

184 For the independent association analysis between SSB or ASB intakes and risk of CVD, we
185 calculated the cumulative averages of SSBs and ASBs, and subtypes, at each updated food-frequency
186 questionnaire to approximate long-term habitual intakes. Beverage intake was categorized by
187 frequency: <1/month (reference), 1–4/month, 2–6/week, 1–<2/day, and ≥ 2 /day, and linear trends were
188 evaluated using the Wald test on a continuous variable representing median intakes of each category.
189 The same categorization was applied to SSB and ASB sub-types. We also evaluated the association for
190 1 serving increase of SSB and ASB intakes per day in the multivariable models. In Model 1 we
191 adjusted for age; race; ancestry; alcohol intake; smoking status; physical activity (for independent SSB
192 and ASB models, in quintiles); family history of myocardial infarction; baseline hypertension or
193 antihypertensive medication; baseline hypercholesterolemia or cholesterol-lowering medication;
194 multivitamin and aspirin use; menopausal status and use of hormone-replacement therapy (in women);
195 total energy intake (continuous); and BMI (in quintiles). In Model 2 we additionally adjusted for the
196 Alternate Healthy Eating Index (in quintiles, without SSBs and alcohol; not applicable in the
197 independent physical activity models).

198 We conducted a series of stratified analyses for continuous servings/day of SSBs or ASBs with
199 incidence CVD by pre-specified subgroups and calculated p-values for heterogeneity with Wald tests
200 for the multiplication interaction of SSB or ASB exposures (continuous variable) with the effect
201 modifier variables.

202 For the joint analyses, SSB or ASB intakes were combined into three categories: (1) never or
203 less than 1 serving/month, (2) 1-4 servings/month, (3) ≥ 2 servings/week, with physical activity
204 dichotomized according to the current aerobic physical activity guidelines for adults (<7.5 MET-
205 hr/week is equivalent to 150 minutes/week of moderate-intensity physical activity)³¹: <7.5 MET-
206 hr/week vs. ≥ 7.5 MET-hr/week. We set participants who reported never/rarely consuming SSBs or
207 ASBs and meeting physical activity guidelines as the reference category. We tested the multiplicative
208 interaction with the cross-product interaction term between continuous servings/day of SSB or ASB
209 intakes*total physical activity added to the main effects models. To assess the additive interaction, we
210 derived the probability value of the relative excess risk due to interaction as an index of additive
211 interaction, also with SSB and ASB intakes and physical activity as continuous variables.³²⁻³⁴

212 Because participants may alter dietary patterns after the diagnosis of major illness, we stopped
213 updating dietary variables when participants reported a diagnosis of coronary artery bypass, angina, or
214 cancer, although follow-up continued until CVD endpoint occurrence, death, or the end of the study
215 period.³⁵ If food-frequency questionnaire exposure data were missing at a given updated time point, we
216 carried forward the exposure level from the previous cumulative cycles.

217 *Sensitivity Analyses*

218 We also ran a series of sensitivity analyses to assess the robustness of results in independent
219 and joint associations. First, the diet was continuously updated until the end of follow-up. Second, we
220 excluded BMI from the models. Third, models were adjusted for median household income and
221 education. Fourth, instead of adjusting for AHEI, we adjusted for a set of dietary variables. Lastly, we
222 further examined two sub-groups: 1) ‘heavy SSB or ASB consumers’ (top 25% SSB or ASB

223 consumers vs not) and the dichotomized physical activity variable, and 2) ‘heavy SSB or ASB
224 consumers’ and ‘high physical activity level’ (top 25% physical activity vs not).

225 All analyses were performed separately for each cohort and then pooled using inverse variance-
226 weighted fixed-effect meta-analysis, and variance-weighted meta-analysis. Statistical tests were two-
227 sided, and *P*-values of < 0.05 were considered to indicate statistical significance. Data were analyzed
228 with the SAS package, version 9.4 (SAS Institute, Cary, NC).

229 **Results**

230 A total of 13,269 CVD cases (8,438 CHD; 4,997 strokes) were documented, 6,156 in men and
231 7,113 in women, during a maximum follow-up of 30 (median 13.4) and 36 (median 17.9) years,
232 respectively. Men and women in the reference group, who jointly met physical activity guidelines and
233 consumed less than 1 SSB serving/month at baseline (21% women and 23% men), had on average
234 lower total energy intake and higher diet quality (**Table 1**). In contrast, men and women who did not
235 meet physical activity guidelines and consumed ≥ 2 servings/week of SSBs at baseline had a higher
236 total energy intake and lower Alternate Healthy Index diet quality score. **Table S1** shows baseline
237 characteristics by categories of SSBs and ASBs intakes separately, and quintiles of physical activity.
238 **Table S2** shows baseline characteristics jointly by categories of ASB and physical activity.

239 **Table 2** shows the pooled associations for intakes of SSBs, ASBs, and physical activity with
240 CVD, CHD, and stroke. In multivariable model 2, compared to participants who never/rarely
241 consumed SSBs, those consuming ≥ 2 SSB servings/day had a 21% higher risk of CVD (pooled HR
242 1.21, 95% CI, 1.12 to 1.32; *p*-trend<0.001). Per serving increment of SSB intake per day, the pooled
243 HR for CVD was 1.15 (95% CI, 1.09 to 1.20; *p*-trend<0.0001), for CHD 1.17 (95% CI, 1.10 to 1.24;
244 *p*-trend<0.0001), and for stroke 1.11 (95% CI, 1.02 to 1.20; *p*-trend=0.02). No evidence of significant
245 differences in ASB intake and risk of CVD or CHD was observed, with the exception of stroke where
246 per serving increment of ASB intake per day was 1.05 (95% CI, 1.01 to 1.10; *p*-trend=0.03).
247 Comparing the highest to the lowest quintile of physical activity, the pooled hazard ratios were 0.66

248 (95% CI, 0.63 to 0.70) for CVD, 0.67 (95% CI, 0.63 to 0.72) for CHD, and 0.64 (95% CI, 0.59 to 0.71)
249 for stroke. **Table S3** shows the results for stroke subtypes for SSB, ASB, and physical activity. **Table**
250 **S4** shows the results for the association between SSB or ASB subtypes and CVD. In the SSB subtype
251 pooled analysis, per serving increment of cola and non-cola carbonated drink intake per day was
252 associated with a 15% (pooled HR 1.15, 95% CI, 1.08 to 1.23; p-trend<0.0001) and a 12% (pooled HR
253 1.12, 95% CI, 1.04 to 1.21; p-trend=0.004) higher risk of CVD, respectively. The associations between
254 SSB, ASB, and physical activity with CVD risks were consistent across subgroups (**Table 3**). Cohort-
255 specific results are shown in **Table S5**.

256 Compared to the reference group (met physical activity guidelines and consumed SSBs
257 <1/month), participants who did not meet the physical activity guidelines and consumed ≥ 2
258 servings/week of SSBs, had a pooled HR of 1.47 (95% CI, 1.37 to 1.57) higher risk of CVD (**Figure 1,**
259 **panel A**). There was also a higher risk of CVD among those who met physical activity guidelines and
260 consumed ≥ 2 SSB servings/day (pooled HR 1.15, 95% CI, 1.08 to 1.23), with similar patterns for CHD
261 (**Figure 1 panel B**) and stroke (**Figure 1, panel C**). Participants who did not meet physical activity
262 guidelines and consumed ASBs <1/month had a 29% (pooled HR 1.29, 95% CI, 1.22 to 1.37) higher
263 risk of CVD (**Figure 1, panel A**), which was consistent with those who did not meet the physical
264 activity guidelines and consumed ≥ 2 servings/week of ASBs (pooled HR 1.29, 95% CI, 1.22 to 1.37),
265 compared to the reference group. Cohort-specific results for joint associations are shown in **Table S6**,
266 and stroke subtypes are included in **Table S7**. **Table S8** shows the results for the joint associations
267 between SSB or ASB subtypes and physical activity and risk of CVD. Although there was no evidence
268 of statistically significant multiplicative interactions between SSB or ASB and physical activity for any
269 outcome, we observed a significant additive interaction between SSB intake and physical activity and
270 risk of CVD (p<0.001) and CHD (p<0.001) (**Table S9**). In addition, we observed a significant additive
271 interaction between the intake of ASBs and physical activity and the risk of CVD (p=0.03) and stroke
272 (p=0.04).

273 The results of these analyses were consistent with those from primary analyses. When we
274 selectively stop updating diet, the independent associations between SSB or ASB consumption and risk
275 of CVD and CHD, were slightly attenuated, while the independent association between SSB intake and
276 risk of stroke remained consistent (**Table S10**). The results of the independent association between
277 physical activity and risk of CVD remained unchanged (**Table S10**). The SSB or ASB intakes and
278 physical activity joint associations and risk of CVD findings were consistent after we stopped updating
279 diet (**Table S11**). The independent and joint association results persisted after we excluded BMI from
280 the final model (**Tables S12 and S13**), except for the ASBs and physical activity joint association
281 results that were strengthened (**Table S13**). Results remained consistent when we additionally adjusted
282 for socioeconomic status (**Tables S14 and S15**) and when we adjusted for dietary factors in lieu of
283 AHEI (**Tables S16 and S17**). The pooled hazard ratios for risk of CVD in ‘heavy SSB consumers’
284 who met and did not meet the physical activity guidelines were 1.14 (95% CI, 1.08 to 1.21) and 1.44
285 (95% CI, 1.35 to 1.53), respectively (**Table S18**). The pooled hazard ratios for those ‘heavy SSB or
286 ASB consumers’ and that were in the top 25% of physical activity (‘high physical activity level’) were
287 slightly attenuated - vs using dichotomized physical activity - for all outcomes (**Table S19**).

288 **Discussion**

289 In two large prospective cohorts of US men and women followed for more than 30 years, we
290 found positive linear associations between SSB consumption and the risk of CVD and CHD, while
291 physical activity was inversely associated with CVD, CHD, and stroke risk, after adjusting for risk
292 factors and other dietary variables. Marginally significant associations were observed per serving
293 increment/day of ASB intake and the risk of CVD. Most importantly, our joint analysis showed that
294 individuals who met physical activity guidelines and consumed >2 servings/day of SSBs had a higher
295 risk of CVD, compared to those who met physical activity guidelines and consumed SSBs <1/month.
296 To our knowledge, the present study is the first large prospective study to examine the joint
297 associations of SSB or ASB consumption and physical activity and the incidence of CVD. Overall, our

298 study suggests that the protective effects of physical activity do not mitigate the detrimental impact of
299 SSB consumption on cardiovascular health.

300 Findings from several meta-analyses have consistently shown that SSB intake is associated
301 with CVD risk in adults, after adjustment for physical activity.^{3,5,36} The most recent meta-analysis
302 (seven prospective cohorts with 16,937,316 person-years of follow-up and 16,915 incident CVD cases),
303 found that SSB consumption was associated with a 9% higher risk of CVD (95% CI, 1.01 to 1.18)
304 when extreme categories of intake (none or less than one/month vs one or more/day) were compared.⁵

305 In regard to ASBs, a meta-analysis including six prospective studies (16,281,005 person-years
306 of follow-up with 18,077 incident CVD events) showed that per serving increment of ASBs per day
307 was associated with a 7% higher risk of incidence of CVD (95% CI, 1.05 to 1.10).⁵ Additionally, per
308 serving increase of ASBs per day was associated with a 6% higher risk of CHD and a 9% higher risk
309 of stroke. The dose-response results of our study were weaker because we observed a 3% higher risk of
310 CVD, with marginal significance, and a 5% higher risk of stroke.

311 Concerning physical activity, we observed a 34% lower risk in the incidence of CVD in those
312 in the highest quintile of physical activity, when compared to the lowest quintile and after accounting
313 for demographic and lifestyle factors. A consistent and strong protective association has been reported
314 with leisure time (relative risk 0.76, 95% CI, 0.70 to 0.82 in men 0.73, 95% CI, 0.68 to 0.78 in women)
315 and occupational physical activity (relative risk 0.89, 95% CI, 0.82 to 0.97 in men and 0.83, 95% CI,
316 0.67 to 1.03 in women).³⁷ Furthermore, a recent systematic review and meta-analysis (33 studies with
317 1,683,693 participants and 89,493 events, follow-up=12.8 years) reported that an increase from being
318 inactive to attaining recommended physical activity levels was associated with a lower risk of CVD
319 incidence by 17% (95% CI, 0.77 to 0.89), after adjusting for body weight.³⁸

320 The joint associations between SSB or ASB intakes and physical activity and risk of CVD
321 findings are novel. Findings provide evidence against the argument that physical activity could
322 counterbalance the potential health risk that would have been induced by SSB or ASB intake. Even

323 though participants who did not meet the physical activity recommendations had a higher risk of CVD
324 at all levels of SSB or ASB intakes, consuming ≥ 2 servings SSBs/week, compared to never/rarely
325 consumers of SSBs, while meeting physical activity guidelines was associated with a 15% higher risk
326 of CVD. Although we did not observe a multiplicative interaction between SSB or ASB and physical
327 activity and risk of CVD, we did detect significant additive interactions between SSB and physical
328 activity and risk of CVD and CHD; and ASB and physical activity and risk of CVD and stroke. Of
329 note, in a previous study in men, De Koning et al., reported no significant interaction between SSB and
330 physical activity.⁸ Hence, meeting physical activity recommendations does not modify the CVD risk
331 associated with higher SSB intake. Interestingly, the impact of SSBs on CVD risk is still observed
332 even in participants who met physical activity recommendations and were in the top 25% of physical
333 activity levels.

334 Strengths of this study include large sample size, prospective population-based design with
335 long and high rates of follow-up, well-defined outcomes, and the use of repeated and validated
336 measurements of diet and other lifestyle data. Limitations of this study include the difficulty in
337 inferring causality due to the observational design. Additionally, residual confounding remains a
338 possibility even though the analyses were extensively adjusted for potential confounders. As well,
339 reverse causation in the ASB analyses is a plausible limitation. Moreover, the dietary information was
340 self-reported and our assessment of SSBs and ASBs and physical activity inevitably had some degree
341 of measurement error. However, the use of repeated measurements lessens random measurement error
342 caused by within-person variation. Also, our study population consisted of primarily non-Hispanic
343 white health professionals, thus the generalizability of our results to other populations may be limited.

344 **Conclusion**

345 Consumption of SSBs or ASBs and physical activity are independently associated with the risk
346 of CVD. Not meeting physical activity guidelines jointly with a greater intake of SSBs or ASBs was
347 associated with a higher risk of CVD. Additionally, findings from this study suggest that even when

348 individuals are physically active, higher consumption of SSBs is associated with a higher risk of CVD.
349 Our results underscore the adverse health effects of both SSBs consumption and being physically
350 inactive and support public health recommendations and policies to limit the intake of SSBs and
351 maintain adequate physical activity levels.

352 **Acknowledgements**

353 The authors would like to thank the participants and staff of the NHS and HPFS cohorts for
354 their continuous involvement and valuable contributions.

355 **Sources of Funding**

356 The research reported in this manuscript were supported by National Institutes of Health grants
357 UM1 CA186107, U01 CA167552, R01 HL034594, R01 HL088521, and R01 HL35464. Additionally,
358 this work was supported by grant T32 DK007703-26 from the National Institute of Diabetes and
359 Digestive and Kidney Diseases and the Harvard Chan Yerby Fellowship at Harvard T.H. Chan School
360 of Public Health.

361 **Disclosures**

362 None.

363

364

365

366

367

368

369

370

371

372 **References:**

- 373 1. Malik VS, Hu FB. Sugar-Sweetened Beverages and Cardiometabolic Health: An Update of the
374 Evidence. *Nutrients*. 2019;11:1840. doi:10.3390/nu11081840
- 375 2. Malik VS, Hu FB. The role of sugar-sweetened beverages in the global epidemics of obesity and
376 chronic diseases. *Nat Rev Endocrinol*. 2022;18(4):205-218. doi:10.1038/S41574-021-00627-6
- 377 3. Narain A, Kwok CS, Mamas MA. Soft drinks and sweetened beverages and the risk of
378 cardiovascular disease and mortality: a systematic review and meta-analysis. *Int J Clin Pract*.
379 2016;70(10):791-805. doi:10.1111/ijcp.12841
- 380 4. Malik VS, Pan A, Willett WC, Hu FB. Sugar-sweetened beverages and weight gain in children
381 and adults: a systematic review and meta-analysis 1-3. *Am J Clin Nutr*. 2013;98(4):1084-1102.
382 doi:10.3945/ajcn.113.058362
- 383 5. Yin J, Zhu Y, Malik V, Li X, Peng X, Zhang FF, Shan Z, Liu L. Intake of Sugar-Sweetened and
384 Low-Calorie Sweetened Beverages and Risk of Cardiovascular Disease: A Meta-Analysis and
385 Systematic Review. *Adv Nutr*. 2021;12(1):89-101. doi:10.1093/ADVANCES/NMAA084
- 386 6. Malik VS, Popkin BM, Bray GA, Després J-P, Willett WC, Hu FB. Sugar-sweetened beverages
387 and risk of metabolic syndrome and type 2 diabetes: a meta-analysis. *Diabetes Care*.
388 2010;33(11):2477-2483. doi:10.2337/dc10-1079
- 389 7. Fung TT, Malik V, Rexrode KM, Manson JE, Willett WC, Hu FB. Sweetened beverage
390 consumption and risk of coronary heart disease in women. *Am J Clin Nutr*. 2009;89(4):1037-
391 1042. doi:10.3945/ajcn.2008.27140
- 392 8. de Koning L, Malik VS, Kellogg MD, Rimm EB, Willett WC, Hu FB. Sweetened beverage
393 consumption, incident coronary heart disease, and biomarkers of risk in men. *Circulation*.
394 2012;125(14):1735-1741, S1. doi:10.1161/CIRCULATIONAHA.111.067017
- 395 9. Pacheco LS, Lacey J V., Martinez ME, Lemus H, Araneta MRG, Sears DD, Talavera GA,
396 Anderson CAM. Sugar-Sweetened Beverage Intake and Cardiovascular Disease Risk in the
397 California Teachers Study. *J Am Heart Assoc*. 2020;9(10). doi:10.1161/JAHA.119.014883

- 398 10. Bernstein AM, de Koning L, Flint AJ, Rexrode KM, Willett WC. Soda consumption and the risk
399 of stroke in men and women. *Am J Clin Nutr.* 2012;95(5):1190-1199.
400 doi:10.3945/ajcn.111.030205
- 401 11. Larsson SC, Akesson A, Wolk A. Sweetened beverage consumption is associated with increased
402 risk of stroke in women and men. *J Nutr.* 2014;144(6):856-860. doi:10.3945/jn.114.190546
- 403 12. Roth GA, Mensah GA, Johnson CO, Addolorato G, Ammirati E, Baddour LM, Barengo NC,
404 Beaton A, ... Fuster V. Global Burden of Cardiovascular Diseases and Risk Factors, 1990–
405 2019: Update From the GBD 2019 Study. *J Am Coll Cardiol.* 2020;76(25):2982.
406 doi:10.1016/J.JACC.2020.11.010
- 407 13. Dai J, Soto MJ, Dunn CG, Bleich SN. Trends and patterns in sugar-sweetened beverage
408 consumption among children and adults by race and/or ethnicity, 2003–2018. *Public Health
409 Nutr.* 2021;24(9):2405-2410. doi:10.1017/S1368980021001580
- 410 14. Hu FB. Resolved: There is sufficient scientific evidence that decreasing sugar-sweetened
411 beverage consumption will reduce the prevalence of obesity and obesity-related diseases. *Obes
412 Rev.* 2013;14(8):606-619. doi:10.1111/obr.12040
- 413 15. Fakhouri T, Kit B, Ogden C. Consumption of diet drinks in the United States, 2009–2010.
414 *NCHS Data Brief.* 2012:1-8.
- 415 16. Johnson RK, Lichtenstein AH, Anderson CAM, Carson JA, Després JP, Hu FB, Kris-Etherton
416 PM, Otten JJ, Towfighi A, Wylie-Rosett J. Low-Calorie Sweetened Beverages and
417 Cardiometabolic Health: A Science Advisory From the American Heart Association.
418 *Circulation.* 2018;138(9):e126-e140. doi:10.1161/CIR.0000000000000569
- 419 17. Mossavar-Rahmani Y, Kamensky V, Manson Joann E., Silver B, Rapp SR, Haring B, Beresford
420 SAA, Snetselaar L, Wassertheil-Smoller S. Artificially Sweetened Beverages and Stroke,
421 Coronary Heart Disease, and All-Cause Mortality in the Women’s Health Initiative. *Stroke.*
422 2019;(50). doi:10.1161/STROKEAHA.118.023100

- 423 18. Chazelas E, Debras C, Srouf B, Fezeu LK, Julia C, Hercberg S, Deschasaux M, Touvier M.
424 Sugary Drinks, Artificially-Sweetened Beverages, and Cardiovascular Disease in the NutriNet-
425 Santé Cohort. *J Am Coll Cardiol.* 2020;76(18):2175-2177. doi:10.1016/J.JACC.2020.08.075
- 426 19. Toews I, Lohner S, Küllenberg De Gaudry D, Sommer H, Meerpohl JJ. Association between
427 intake of non-sugar sweeteners and health outcomes: Systematic review and meta-analyses of
428 randomised and non-randomised controlled trials and observational studies. *BMJ.*
429 2019;364:k4718. doi:10.1136/bmj.k4718
- 430 20. Franklin BA, Thompson CPD, Al-Zaiti SS, Albert CM, Hivert MF, Levine BD, Lobelo F,
431 Madan K, Sharrief AZ, Eijsvogels TMH. Exercise-Related Acute Cardiovascular Events and
432 Potential Deleterious Adaptations Following Long-Term Exercise Training: Placing the Risks
433 Into Perspective—An Update: A Scientific Statement From the American Heart Association.
434 *Circulation.* 2020;141:E705-E736. doi:10.1161/CIR.0000000000000749
- 435 21. Tian D, Meng J. Exercise for Prevention and Relief of Cardiovascular Disease: Prognoses,
436 Mechanisms, and Approaches. *Oxid Med Cell Longev.* 2019;2019:11.
437 doi:10.1155/2019/3756750
- 438 22. Colditz GA, Manson JE, Hankinson SE. The Nurses' Health Study: 20-year contribution to the
439 understanding of health among women. *J Women's Heal.* 1997;6(1):49-62.
440 doi:10.1089/jwh.1997.6.49
- 441 23. Rimm EB, Giovannucci EL, Willett WC, Colditz GA, Ascherio A, Rosner B, Stampfer MJ.
442 Prospective study of alcohol consumption and risk of coronary disease in men. *Lancet.*
443 1991;338(8765):464-468. doi:10.1016/0140-6736(91)90542-W
- 444 24. Yuan C, Spiegelman D, Rimm EB, Rosner BA, Stampfer MJ, Barnett JB, Chavarro JE, Rood
445 JC, Harnack LJ, Sampson LK, Willett WC. Practice of Epidemiology Relative Validity of
446 Nutrient Intakes Assessed by Questionnaire, 24-Hour Recalls, and Diet Records as Compared
447 With Urinary Recovery and Plasma Concentration Biomarkers: Findings for Women. 187(5).

448 doi:10.1093/aje/kwx328

- 449 25. Willett WC, Sampson L, Stampfer MJ, Rosner B, Bain C, Witschi J, Hennekens CH, Speizer
450 FE. Reproducibility and validity of a semiquantitative food frequency questionnaire. *Am J*
451 *Epidemiol.* 1985;122(1):51-65. doi:10.1093/oxfordjournals.aje.a114086
- 452 26. Wolf AM, Hunter DJ, Colditz GA, Manson JE, Stampfer MJ, Corsano KA, Rosner B, Kriska A,
453 Willett WC. Reproducibility and Validity of a Self-Administered Physical Activity
454 Questionnaire. *Int J Epidemiol.* 1994;23(5):991-999. doi:10.1093/IJE/23.5.991
- 455 27. Rose G. *Cardiovascular Survey Methods.* World Health Organization. Albany, N.Y.: WHO
456 Publications Centre distributor; 1982.
- 457 28. AE W, M R, FD W. The National Survey of Stroke. Clinical findings. *Stroke.* 1981;12(2 Pt 2
458 Suppl 1). <https://pubmed.ncbi.nlm.nih.gov/7222164/>. Accessed August 1, 2021.
- 459 29. Stampfer MJ, Willett WC, Speizer FE, Dysert DC, Lipnick R, Rosner B, Hennekens CH. Test of
460 the National Death Index. *Am J Epidemiol.* 1984;119(5):837-839.
461 <https://academic.oup.com/aje/article/119/5/837/125237>. Accessed May 11, 2021.
- 462 30. Chiuve SE, Fung TT, Rimm EB, Hu FB, Mccullough ML, Wang M, Stampfer MJ, Willett WC.
463 Alternative Dietary Indices Both Strongly Predict Risk of Chronic Disease. *J Nutr.*
464 2012;142:1009-1018. doi:10.3945/jn.111.157222
- 465 31. HHS. *Physical Activity Guidelines for Americans 2 Nd Edition.*
- 466 32. Vanderweele TJ, Knol MJ. A Tutorial on Interaction. *Epidemiol Methods.* 2014;3(1):33-72.
467 doi:10.1515/em-2013-0005
- 468 33. Vanderweele TJ, Tchetgen Tchetgen EJ. Attributing effects to interactions. *Epidemiology.*
469 2014;25(5):711. doi:10.1097/EDE.0000000000000096
- 470 34. Li R, Chambless L. Test for Additive Interaction in Proportional Hazards Models. *Ann*
471 *Epidemiol.* 2007;17(3):227-236. doi:10.1016/J.ANNEPIDEM.2006.10.009
- 472 35. Hu FB, Stampfer MJ, Rimm E, Ascherio A, Rosner BA, Spiegelman D, Willett WC. Dietary Fat

- 473 and Coronary Heart Disease: A Comparison of Approaches for Adjusting for Total Energy
474 Intake and Modeling Repeated Dietary Measurements. *Am J Epidemiol.* 1999;149(6):531-540.
475 doi:10.1093/oxfordjournals.aje.a009849
- 476 36. Huang C, Huang J, Tian Y, Yang X, Gu D. Sugar sweetened beverages consumption and risk of
477 coronary heart disease: a meta-analysis of prospective studies. *Atherosclerosis.* 2014;234(1):11-
478 16. doi:10.1016/J.ATHEROSCLEROSIS.2014.01.037
- 479 37. Li J, Siegrist J. Physical Activity and Risk of Cardiovascular Disease—A Meta-Analysis of
480 Prospective Cohort Studies. *Int J Environ Res Public Health.* 2012;9(2):391.
481 doi:10.3390/IJERPH9020391
- 482 38. Wahid A, Manek N, Nichols M, Kelly P, Foster C, Webster P, Kaur A, Friedemann Smith C,
483 Wilkins E, Rayner M, Roberts N, Scarborough P. Quantifying the Association Between Physical
484 Activity and Cardiovascular Disease and Diabetes: A Systematic Review and Meta-Analysis. *J*
485 *Am Heart Assoc.* 2016;5(9). doi:10.1161/JAHA.115.002495/FORMAT/EPUB

486

487

488

489

490

491

492


493

494

495

496

497 **Figures**



498

499 **Figure 1. Pooled Risk* of Cardiovascular Events Associated with the Joint Associations of Sugar-**
 500 **or Artificially-Sweetened Beverages and Physical Activity† in Two Large US Cohorts.**

501* Hazard ratios and 95% confidence intervals derived from the multivariable model adjusted for the following: age, ethnicity/race;

502 smoking status; alcohol intake; multivitamin use; in women, postmenopausal status and menopausal hormone use; family history of

503 myocardial infarction; baseline hypertension; baseline hypercholesterolemia; calorie intake; body mass index; and AHEI.

504† Physical activity was dichotomized to meet or do not meet the US Physical activity guidelines of ≥150 minutes/week of

505 moderate-vigorous physical activity recommendations (corresponds to ≥7.5 MET hr/week or ≥450 MET min/week).

506 AHEI = alternative healthy eating index, sugar-sweetened beverages and alcohol intake were removed from the index calculation; MET =

507 metabolic equivalent task.

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

529

530

1 Tables

Table 1. Baseline* Characteristics of Participants According to Sugar-Sweetened Beverage Intake and Physical Activity Joint Association Categories in Two Large US Cohorts.

	Meets the US Physical Activity Guidelines [†]			Does Not Meet the US Physical Activity Guidelines [†]		
	Never or <1 SSB serving per month	1-4 SSB servings per month	≥2 SSB servings per week	Never or <1 SSB serving per month	1-4 SSB servings per month	≥2 SSB servings per week
Nurses' Health Study						
Participants, n	13,895	7,636	12,014	11,331	7,367	13,487
Age, years	47 ± 7	47 ± 7	45 ± 7	48 ± 7	47 ± 7	45 ± 7
BMI, kg/m ²	23.7 ± 3.7	23.6 ± 3.8	24.0 ± 4.2	24.6 ± 4.5	24.5 ± 4.4	24.9 ± 4.9
Physical activity, MET-h/week	26.1 ± 24.5	24.3 ± 24.2	23.9 ± 22.3	2.8 ± 2.0	2.8 ± 2.0	2.8 ± 1.9
Family history of myocardial infarction (%)	39.6	38.5	37.5	39.5	37.7	37.1
Family history of diabetes (%)	28.2	27.9	27.8	30.3	29.5	30.5
Ethnicity, white (%)	98.7	97.9	97.4	98.7	98.0	97.0
Current smoker (%)	24.3	22.1	25.7	31.9	28.3	29.8
Hypertension (%)	5.2	4.6	5.6	6.3	5.8	6.2
Hypercholesterolemia (%)	4.1	3.6	3.6	4.2	3.6	3.6
Multivitamin supplement use (%)	37.8	36.0	34.5	33.1	31.9	31.0
Aspirin use (%)	44.2	46.5	49.1	46.2	45.9	49.6
Current menopausal hormone use (%)	7.3	6.8	6.3	7.1	6.8	6.3
Total energy intake, kcal/day	1,444 ± 461	1,534 ± 470	1,718 ± 500	1,445 ± 470	1,506 ± 473	1,705 ± 510
Artificially-sweetened beverages, servings/day	0.6 ± 1.0	0.3 ± 0.7	0.3 ± 0.6	0.6 ± 1.1	0.3 ± 0.7	0.3 ± 0.6
Sugar-sweetened beverages, servings/day	0.0 ± 0.0	0.1 ± 0.0	0.9 ± 0.9	0.0 ± 0.0	0.1 ± 0.0	1.0 ± 1.0
Total beverages, servings/day	5.4 ± 2.5	5.2 ± 2.4	5.7 ± 2.6	5.5 ± 2.6	5.3 ± 2.5	5.8 ± 2.6
Alcohol, g/day	8.0 ± 11.0	6.3 ± 9.6	5.5 ± 9.2	8.0 ± 12.3	5.5 ± 9.9	5.0 ± 9.7
Red and processed meat, servings/day	1.2 ± 0.8	1.4 ± 0.8	1.5 ± 0.8	1.3 ± 0.8	1.4 ± 0.8	1.6 ± 0.8
Nuts, servings/day	0.2 ± 0.3	0.2 ± 0.3	0.2 ± 0.3	0.1 ± 0.3	0.1 ± 0.3	0.1 ± 0.3
Whole grains, servings/day	2.3 ± 2.0	2.2 ± 1.9	2.0 ± 1.8	2.1 ± 2.0	2.1 ± 1.9	1.9 ± 1.8
Fruits and vegetables, servings/day	4.5 ± 2.2	4.2 ± 2.0	4.1 ± 2.0	4.0 ± 2.1	3.7 ± 1.9	3.6 ± 1.9
AHEI [‡]	44.6 ± 9.0	42.3 ± 8.5	40.3 ± 8.3	41.2 ± 8.7	39.4 ± 8.2	37.7 ± 7.8
Health Professionals Follow-up Study						
Participants, n	9,062	5,903	10,015	4,520	3,308	6,610
Age, years	54 ± 9	53 ± 9	51 ± 9	56 ± 9	54 ± 9	51 ± 9
BMI, kg/m ²	25.1 ± 3.1	25.2 ± 3.1	25.2 ± 3.1	26.2 ± 3.7	25.9 ± 3.5	25.9 ± 3.5
Physical activity, MET-h/week	32.6 ± 27.3	30.4 ± 25.5	30.7 ± 26.0	3.0 ± 2.2	3.1 ± 2.1	3.0 ± 2.1
Family history of myocardial infarction (%)	33.6	30.4	30.7	33.1	31.2	31.6

Family history of diabetes (%)	20.0	18.4	16.9	20.3	19.2	18.3
Ethnicity, white (%)	95.7	96.2	94.1	96.2	95.5	94.0
Current smoker (%)	6.7	7.4	9.1	12.1	11.6	13.6
Hypertension (%)	21.2	19.2	19.6	24.5	22.9	22.5
Hypercholesterolemia (%)	11.0	11.0	9.6	11.5	10.2	9.4
Multivitamin supplement use (%)	46.5	43.8	41.5	41.1	37.4	36.7
Aspirin use (%)	25.6	25.2	27.5	27.0	26.6	27.3
Total energy intake, kcal/day	1,821 ± 558	1,944 ± 575	2,219 ± 632	1,738 ± 545	1,872 ± 556	2,163 ± 639
Artificially-sweetened beverages, servings/day	0.7 ± 1.1	0.5 ± 0.8	0.3 ± 0.7	0.7 ± 1.2	0.5 ± 1.0	0.3 ± 0.7
Sugar-sweetened beverages, servings/day	0.0 ± 0.0	0.1 ± 0.0	0.7 ± 0.7	0.0 ± 0.0	0.1 ± 0.0	0.8 ± 0.8
Total beverages, servings/day	7.2 ± 3.0	7.2 ± 2.8	7.7 ± 2.9	7.2 ± 3.0	7.1 ± 2.9	7.4 ± 3.0
Alcohol, g/day	12.8 ± 16.2	11.6 ± 14.7	10.8 ± 14.3	12.4 ± 16.9	11.2 ± 16.0	10.0 ± 15.4
Red and processed meat, servings/day	0.9 ± 0.8	1.0 ± 0.8	1.3 ± 0.8	1.1 ± 0.8	1.2 ± 0.8	1.4 ± 0.9
Nuts, servings/day	0.3 ± 0.4	0.3 ± 0.4	0.3 ± 0.4	0.2 ± 0.4	0.2 ± 0.4	0.2 ± 0.4
Whole grains, servings/day	1.9 ± 1.6	1.8 ± 1.5	1.6 ± 1.4	1.5 ± 1.4	1.5 ± 1.3	1.4 ± 1.2
Fruits and vegetables, servings/day	5.3 ± 2.9	5.2 ± 2.9	5.1 ± 2.7	4.4 ± 2.6	4.3 ± 2.4	4.3 ± 2.4
AHEI [‡]	48.2 ± 10.2	45.9 ± 9.8	43.1 ± 9.4	43.9 ± 10.1	41.8 ± 9.6	39.6 ± 9.1

¹Values are mean ± SD or %, and are standardized to the age distribution of the study population. HPFS, n=39,418; NHS, n=65,730.

²* Baseline for NHS is 1980 and for HPFS is 1986.

³† US physical activity guidelines were met if participant reported ≥150 minutes/week of moderate-vigorous physical activity.

⁴‡ AHEI range 0-90, sugar-sweetened beverages and alcohol were removed from the index calculation.

⁵AHEI = alternative healthy eating index; BMI = body mass index; MET = metabolic equivalent task; MVPA = moderate-vigorous physical activity.

6

7

8

9

10

11

12

13

1 Table 2. Pooled Risk* of Cardiovascular Events According to Sugar- or Artificially Sweetened Beverage Intake Categories and Physical Activity†
2 Quintiles in Two Large US Cohorts.

Sugar-Sweetened Beverages							
	HR (95% CI)					P value for trend	HR (95% CI) per serving increment of SSB‡ intake per day
	Never or <1 serving per month	1-4 servings per month	2-6 servings per week	1 to <2 servings per day	≥2 servings per day		
Total Cardiovascular Disease							
HPFS, cases/person-years	1,354/187,404	1,906/266,714	2,195/338,803	276/41,556	425/74,663		
NHS, cases/person-years	991/392,997	2,669/727,712	2,789/735,164	238/85,332	426/150,868		
Pooled risk							
Age-adjusted model	1.00	1.00 (0.95, 1.05)	1.10 (1.05, 1.16)	1.26 (1.15, 1.39)	1.34 (1.24, 1.45)	<0.0001	1.26 (1.20, 1.32)
Multivariable model	1.00	1.04 (0.99, 1.10)	1.13 (1.07, 1.19)	1.23 (1.12, 1.36)	1.25 (1.16, 1.36)	<0.0001	1.17 (1.12, 1.23)
Multivariable model + AHEI	1.00	1.04 (0.98, 1.09)	1.11 (1.05, 1.17)	1.20 (1.08, 1.32)	1.21 (1.12, 1.32)	<0.0001	1.15 (1.09, 1.20)
Coronary Heart Disease							
HPFS, cases/person-years	1,001/187,692	1,435/267,136	1,649/339,278	220/41,622	336/74,765		
NHS, cases/person-years	548/393,295	1,389/728,575	1,477/736,011	138/85,404	245/151,000		
Pooled risk							
Age-adjusted model	1.00	1.01 (0.94, 1.07)	1.11 (1.04, 1.18)	1.34 (1.19, 1.51)	1.41 (1.28, 1.55)	<0.0001	1.30 (1.23, 1.38)
Multivariable model	1.00	1.06 (1.00, 1.13)	1.14 (1.07, 1.22)	1.30 (1.16, 1.46)	1.31 (1.19, 1.45)	<0.0001	1.21 (1.14, 1.28)
Multivariable model + AHEI	1.00	1.05 (0.99, 1.12)	1.12 (1.05, 1.19)	1.25 (1.11, 1.41)	1.25 (1.13, 1.39)	<0.0001	1.17 (1.10, 1.24)
Stroke							
HPFS, cases/person-years	353/187,871	471/267,471	546/339,696	56/41,678	89/74,850		
NHS, cases/person-years	476/393,327	1,326/728,488	1,381/735,934	107/85,423	192/151,020		
Pooled risk							
Age-adjusted model	1.00	0.97 (0.89, 1.06)	1.09 (1.00, 1.19)	1.14 (0.96, 1.34)	1.21 (1.05, 1.39)	<0.0001	1.18 (1.09, 1.28)
Multivariable model	1.00	0.99 (0.91, 1.08)	1.10 (1.01, 1.20)	1.11 (0.93, 1.31)	1.13 (0.98, 1.31)	0.01	1.11 (1.02, 1.21)
Multivariable model + AHEI	1.00	0.99 (0.91, 1.08)	1.09 (1.00, 1.19)	1.10 (0.92, 1.30)	1.12 (0.97, 1.29)	0.02	1.11 (1.02, 1.20)
Artificially-Sweetened Beverages							
	HR (95% CI)					P value for trend	HR (95% CI) per serving increment of ASB‡ intake per day
	Never or <1 serving per month	1-4 servings per month	2-6 servings per week	1 to <2 servings per day	≥2 servings per day		
Total Cardiovascular Disease							
HPFS, cases/person-years	1,906/266,941	1,378/188,107	1,774/266,083	296/50,244	802/137,764		

NHS, cases/person-years	2,692/774,993	1,199/344,166	1,585/487,216	632/184,325	1,005/301,373		
Pooled risk							
Age-adjusted model	1.00	0.91 (0.87, 0.96)	0.93 (0.89, 0.97)	1.04 (0.97, 1.12)	1.09 (1.04, 1.16)	<0.0001	1.07 (1.04, 1.10)
Multivariable model	1.00	0.94 (0.90, 0.99)	0.94 (0.90, 0.98)	1.02 (0.95, 1.10)	1.02 (0.96, 1.08)	0.11	1.02 (0.99, 1.05)
Multivariable model + AHEI	1.00	0.95 (0.90, 1.00)	0.95 (0.91, 0.99)	1.03 (0.96, 1.10)	1.03 (0.97, 1.09)	0.06	1.03 (1.00, 1.06)
Coronary Heart Disease							
HPFS, cases/person-years	1,448/267,352	1,059/188,387	1,308/266,499	219/50,313	607/137,939		
NHS, cases/person-years	1,452/775,826	629/344,545	849/487,721	329/184,504	538/301,689		
Pooled risk							
Age-adjusted model	1.00	0.91 (0.86, 0.97)	0.92 (0.87, 0.97)	1.01 (0.92, 1.11)	1.10 (1.03, 1.18)	0.0003	1.07 (1.03, 1.11)
Multivariable model	1.00	0.95 (0.89, 1.01)	0.92 (0.87, 0.98)	0.97 (0.88, 1.07)	1.00 (0.93, 1.08)	0.61	1.01 (0.97, 1.05)
Multivariable model + AHEI	1.00	0.95 (0.90, 1.02)	0.94 (0.88, 0.99)	0.98 (0.89, 1.08)	1.01 (0.94, 1.09)	0.42	1.02 (0.98, 1.05)
Stroke							
HPFS, cases/person-years	458/267,729	319/188,627	466/266,738	77/50,370	195/138,102		
NHS, cases/person-years	1,292/775,823	596/344,501	781/487,702	321/184,495	492/301,670		
Pooled risk							
Age-adjusted model	1.00	0.92 (0.85, 1.00)	0.96 (0.89, 1.03)	1.11 (1.00, 1.24)	1.09 (1.00, 1.20)	0.002	1.07 (1.03, 1.12)
Multivariable model	1.00	0.95 (0.87, 1.03)	0.98 (0.91, 1.06)	1.11 (1.00, 1.24)	1.06 (0.97, 1.17)	0.03	1.05 (1.00, 1.10)
Multivariable model + AHEI	1.00	0.95 (0.88, 1.03)	0.98 (0.91, 1.06)	1.12 (1.00, 1.25)	1.07 (0.97, 1.17)	0.03	1.05 (1.01, 1.10)

Physical Activity

	HR (95% CI)					P value for trend
	Quintile 1 (0 to ≤3.0)	Quintile 2 (>3.0 to ≤8.4)	Quintile 3 (>8.4 to ≤17.8)	Quintile 4 (>17.8 to ≤35.1)	Quintile 5 (>35.1)	
Total Cardiovascular Disease						
HPFS, cases/person-years	1,924/178,321	1,239/181,529	1,105/182,492	990/183,382	898/183,415	
NHS, cases/person-years	2,205/413,395	1,520/418,327	1,262/417,294	1,163/421,176	963/421,881	
Pooled risk						
Age-adjusted model	1.00	0.72 (0.69, 0.76)	0.66 (0.62, 0.69)	0.61 (0.58, 0.64)	0.54 (0.51, 0.57)	<0.0001
Multivariable model	1.00	0.78 (0.74, 0.82)	0.74 (0.70, 0.78)	0.71 (0.67, 0.75)	0.65 (0.61, 0.69)	<0.0001
Multivariable model + AHEI	1.00	0.78 (0.74, 0.82)	0.74 (0.71, 0.78)	0.72 (0.68, 0.76)	0.66 (0.63, 0.70)	<0.0001
Coronary Heart Disease						
HPFS, cases/person-years	1,524/178,645	916/181,811	847/182,731	697/183,661	657/183,644	
NHS, cases/person-years	1,221/414,038	806/418,789	657/417,706	589/421,559	524/422,193	
Pooled risk						
Age-adjusted model	1.00	0.68 (0.64, 0.73)	0.63 (0.59, 0.67)	0.55 (0.52, 0.59)	0.52 (0.49, 0.56)	<0.0001
Multivariable model	1.00	0.75 (0.71, 0.80)	0.73 (0.68, 0.78)	0.67 (0.62, 0.71)	0.65 (0.61, 0.70)	<0.0001
Multivariable model + AHEI	1.00	0.76 (0.71, 0.80)	0.74 (0.70, 0.79)	0.68 (0.64, 0.73)	0.67 (0.63, 0.72)	<0.0001
Stroke						
HPFS, cases/person-years	400/178,905	323/182,010	258/182,987	293/183,828	241/183,837	

NHS, cases/person-years	1,033/413,964	746/418,790	634/417,681	608/421,538	461/422,220	
Pooled risk						
Age-adjusted model	1.00	0.79 (0.73, 0.86)	0.70 (0.64, 0.76)	0.72 (0.66, 0.78)	0.58 (0.53, 0.63)	<0.0001
Multivariable model	1.00	0.83 (0.76, 0.90)	0.75 (0.69, 0.82)	0.79 (0.73, 0.86)	0.64 (0.59, 0.71)	<0.0001
Multivariable model + AHEI	1.00	0.83 (0.77, 0.90)	0.75 (0.69, 0.82)	0.80 (0.73, 0.87)	0.64 (0.59, 0.71)	<0.0001

1 * Hazard ratios and 95% confidence intervals derived from the age-adjusted and multivariable model adjusted for age; race; ethnicity; physical activity (excluded in the physical activity as main exposure models); smoking status; alcohol intake; multivitamin use; in women, postmenopausal status and menopausal hormone use; family history of myocardial infarction; baseline hypertension; baseline hypercholesterolemia; calorie intake; and body mass index.

2 † Physical activity was measured in MET-hr/week.

3 ‡ 1 serving increment of sugar- or artificially-sweetened beverage intake per day, where appropriate.

4 AHEI = alternative healthy eating index, sugar-sweetened beverages and alcohol intake were removed from the index calculation; HPFS = Health Professionals Follow-up Study; MET = metabolic equivalent task; NHS = Nurses' Health Study.

5

6

7

8

9

10

11

12

13

14

15

16

17

1
2

Table 3. Pooled Subgroup Analyses for Risk* of Cardiovascular Events According to 1 Serving Increment in Sugar- or Artificially-Sweetened Beverage Intake Per Day† in Two Large US Cohorts.

	Cardiovascular Disease		Coronary Heart Disease		Stroke	
	Pooled Adjusted HR (95% CI)	p-value for interaction	Pooled Adjusted HR (95% CI)	p-value for interaction	Pooled Adjusted HR (95% CI)	p-value for interaction
Sugar-Sweetened Beverages						
Age						
<65 years	1.15 (1.06, 1.25)	0.05	1.21 (1.10, 1.34)	0.04	1.01 (0.87, 1.18)	0.94
≥65 years	1.12 (1.05, 1.20)		1.12 (1.03, 1.21)		1.14 (1.03, 1.26)	
Body mass index, kg/m²						
<25	1.20 (1.12, 1.28)	0.76	1.24 (1.14, 1.35)	0.45	1.12 (0.99, 1.26)	0.69
≥25	1.09 (1.02, 1.17)		1.09 (1.00, 1.19)		1.10 (0.97, 1.24)	
Physical activity, meet US guidelines						
No (<150 min/week MVPA)	1.12 (1.04, 1.20)	0.14	1.12 (1.03, 1.23)	0.07	1.09 (0.97, 1.24)	0.63
Yes (≥150 min/week MVPA)	1.18 (1.10, 1.26)		1.22 (1.12, 1.32)		1.12 (1.00, 1.26)	
Television watching, hour/week						
<8	1.18 (1.08, 1.29)	0.45	1.22 (1.10, 1.36)	0.59	1.12 (0.96, 1.30)	0.75
8 to 15.5	1.12 (1.00, 1.26)		1.13 (0.98, 1.30)		1.13 (0.93, 1.37)	
≥15.5	1.15 (1.07, 1.24)		1.18 (1.07, 1.29)		1.12 (0.98, 1.26)	
Smoking status						
Never	1.13 (1.05, 1.21)	0.01	1.16 (1.06, 1.28)	0.08	1.07 (0.94, 1.21)	0.10
Former/current	1.20 (1.13, 1.29)		1.21 (1.12, 1.31)		1.19 (1.06, 1.33)	
Family history of myocardial infarction						
No	1.17 (1.09, 1.25)	0.89	1.19 (1.10, 1.29)	0.98	1.13 (1.02, 1.26)	0.69
Yes	1.11 (1.03, 1.20)		1.15 (1.05, 1.26)		1.06 (0.93, 1.22)	
Calories per day, kcal						
<median	1.19 (1.10, 1.29)	0.17	1.17 (1.06, 1.29)	0.67	1.24 (1.09, 1.41)	0.06
≥median	1.09 (1.02, 1.16)		1.12 (1.03, 1.21)		1.03 (0.92, 1.15)	
AHEI						
<median	1.13 (1.06, 1.20)	0.26	1.12 (1.04, 1.21)	0.02	1.15 (1.03, 1.28)	0.34
≥median	1.20 (1.11, 1.30)		1.30 (1.18, 1.43)		1.05 (0.92, 1.21)	
Artificially-Sweetened Beverages						
Age						
<65 years	0.99 (0.93, 1.04)	0.52	0.95 (0.89, 1.01)	0.16	1.08 (0.99, 1.19)	0.07
≥65 years	1.04 (1.01, 1.08)		1.05 (1.00, 1.10)		1.04 (0.99, 1.10)	
Body mass index, kg/m²						
<25	1.05 (1.01, 1.10)	0.72	1.02 (0.96, 1.08)	0.46	1.10 (1.03, 1.18)	0.25
≥25	1.03 (0.99, 1.07)		1.04 (0.99, 1.09)		1.02 (0.96, 1.09)	

Physical activity, meet US guidelines						
No (<150 min/week MVPA)	1.04 (1.00, 1.09)	0.70	1.04 (0.98, 1.10)	0.60	1.05 (0.98, 1.13)	0.88
Yes (≥150 min/week MVPA)	1.02 (0.98, 1.06)		1.00 (0.95, 1.05)		1.06 (0.99, 1.13)	
Television watching, hour/week						
<8	1.09 (1.04, 1.15)	0.01	1.10 (1.03, 1.18)	0.004	1.08 (1.00, 1.18)	0.62
8 to 15.5	1.03 (0.96, 1.10)		1.03 (0.95, 1.12)		1.03 (0.93, 1.14)	
≥15.5	0.98 (0.93, 1.03)		0.96 (0.90, 1.02)		1.01 (0.94, 1.09)	
Smoking status						
Never	1.06 (1.02, 1.11)	0.0004	1.05 (0.99, 1.11)	0.002	1.08 (1.01, 1.16)	0.10
Former/current	1.02 (0.98, 1.07)		1.03 (0.97, 1.09)		1.03 (0.96, 1.11)	
Family history of myocardial infarction						
No	1.03 (0.99, 1.07)	0.96	1.01 (0.96, 1.06)	1.00	1.06 (1.00, 1.13)	0.98
Yes	1.03 (0.98, 1.07)		1.02 (0.96, 1.08)		1.04 (0.97, 1.12)	
Calories per day, kcal						
<median	1.05 (1.01, 1.09)	0.21	1.05 (0.99, 1.10)	0.10	1.06 (1.00, 1.13)	0.75
≥median	1.01 (0.96, 1.05)		0.98 (0.93, 1.04)		1.04 (0.97, 1.11)	
AHEI						
<median	1.03 (0.98, 1.07)	0.97	1.01 (0.96, 1.07)	0.99	1.05 (0.98, 1.12)	0.84
≥median	1.03 (0.99, 1.07)		1.02 (0.96, 1.07)		1.06 (0.99, 1.13)	

1 * Hazard ratios and 95% confidence intervals derived from the multivariable model adjusted for age; race; ethnicity; physical activity (excluded in the physical activity as main exposure
2 models); smoking status; alcohol intake; multivitamin use; in women, postmenopausal status and menopausal hormone use; family history of myocardial infarction; baseline
3 hypertension; baseline hypercholesterolemia; calorie intake; body mass index; and AHEI.

4 † 1 serving of sugar- or artificially-sweetened beverage. HRs for 1 serving increment of sugar- or artificially-sweetened beverage intake per day in each subgroup category.

5 AHEI = alternative healthy eating index, sugar-sweetened beverages and alcohol intake were removed from the index calculation; MVPA = moderate-vigorous physical activity.

6