Title:

Histological analysis of breast implant capsules: A validated scoring system based on 480 patients

Author list:

 Andreas Larsen, MD[1]; Adam M Timmermann, BMSc[1]; Mikela Kring, BMSc[1]; Tim K Weltz, MD[1]; Mathias Ørholt, MD[1]; Peter Vester-Glowinski, MD, PhD[1]; Jens Jørgen Elberg, MD[2]; Jesper Trillingsgaard, MD[3]; Louise V Mielke, MD[4]; Lisbet R Hölmich, MD, DMSc[5,6]; Tine E Damsgaard, MD, PhD, MRBS[1,6]; Anne Roslind, MD, PhD[7]; Mikkel Herly, MD, PhD[1,8]

Affiliations:

[1]Copenhagen University Hospital, Department of Plastic Surgery and Burns Treatment, Rigshospitalet, Copenhagen, Denmark

[2]Amalieklinikken, Copenhagen, Denmark

[3]AK Nygart, Copenhagen, Denmark

[4]Aleris-Hamlet, Copenhagen, Denmark

[5]Copenhagen University Hospital, Department of Plastic and Reconstructive Surgery, Herlev and Gentofte, Copenhagen, Denmark

[6]University of Copenhagen, Department of Clinical Medicine, Copenhagen, Denmark

[7]Copenhagen University Hospital, Department of Pathology, Herlev and Gentofte Hospital, Copenhagen, Denmark

[8]University of Copenhagen, Department of Immunology and Microbiology, Copenhagen, Denmark

Abstract

Capsular contracture is a severe complication to breast implants, but the pathophysiology remains unclear, and consensus is lacking on how to analyze implant capsules histologically. In this study, we developed and validated a histological scoring system for analyzing breast implant capsules. Biopsies of breast implant capsules from 480 patients who had undergone breast augmentation or reconstruction were collected and stained with hematoxylin and eosin. First, biopsies from 100 patients were used to select parameters for the scoring system. Then, biopsies from the remaining 380 patients were used to determine intra- and interobserver agreement of two blinded non-pathologist-observers and agreement with a pathologist. Finally, we tested the association between the parameters and capsular contracture. The histological scoring system consisted of ten parameters assessed on a two-, three-, or four-point scale. Intra- and interobserver agreement was almost perfect (0.83 and 0.80), and agreement with the pathologist was substantial (0.67). Inflammatory infiltration ($p<0.01$), thickness of the collagen layer ($p<0.0001$), fiber organization ($p<0.01$) and calcification ($p<0.001$) were significantly correlated with capsular contracture. This is the first validated histological scoring system for breast implant capsules that provides an objective and precise assessment of breast implant capsules which can be used in research and clinical practice.

Main text

Introduction

Capsular contracture is one of the most common severe complications of breast implants. Within 10 years after surgery, capsular contracture affects 3.6-19%(1–5) of women after breast augmentation and 3.3-25%(1,3–6) of women after implant-based breast reconstruction. The severity of capsular contracture is most often clinically evaluated using the Baker Classification System, which consists of a four-point scale based on the firmness and distortion of the breast.(7,8) Despite the importance of capsular contracture, its pathogenesis is largely unknown. Previous studies suggest that capsular contracture is caused by a low-grade inflammatory response in the tissue around the breast implant, which results in excessive fibrosis.(9–11) In rare cases, the immunological reaction may potentially be the cause of breast implant-associated anaplastic large cell lymphoma (BIA-ALCL); however, our knowledge of these immunological processes is still very limited.(12,13)

Several studies have investigated the histological characteristics of breast implant capsules, but they are limited by descriptive histological approaches in small, heterogeneous patient populations.(14) The lack of a standardized and reproducible tool for analyzing breast implant capsules prevents an effective comparison of results between studies and thus impairs our ability to build on the knowledge of previous studies. An objective and validated tool for systematically characterizing the histological appearance of the breast implant capsule is needed to create a common language between studies. Such a tool may help us understand why some but not all women develop capsular contracture, and it may be of use in developing new preventive strategies. In this study, we aimed to develop and validate a histological scoring system that covers the histological characteristics that are seen across a large sample of hematoxylin and eosin (H&E)-stained breast implant capsules. Furthermore, we aimed to show the applicability of the scoring system by analyzing differences between patients with and without capsular contracture.

Materials and methods

Patients and sample material

The histological samples were obtained from a biobank of tissue biopsies excised from the breast implant capsule from patients undergoing exchange or removal of their breast implants after breast augmentation or implant-based breast reconstruction. Patients were included from October 2019 to August 2022 from two plastic surgery departments (Rigshospitalet, Department of Plastic Surgery and Burns Treatment and Herlev and Gentofte Hospital, Department of Plastic Surgery) and three private hospitals (Amalieklinikken, AK Nygart and Aleris-Hamlet). The biobank consisted of 400 patients who had their implants removed after breast augmentation and 80 patients who had their implants removed after an implant-based breast reconstruction. Patient and implant characteristics are described in Table 1. Biopsies of approximately 1 cm^2 were excised from the anterior-inferior part of the fibrous capsule adjacent to the inframammary fold, fixated in a 4% formaldehyde solution and embedded in paraffine. Sections of 3-4 µm were cut and stained with hematoxylin and eosin (H&E). All sections were digitalized using a whole slide scanner and analyzed using NDP.viewer2 (Hamamatsu Photonics).

Design of the scoring system

The scoring system was designed and validated according to principles for valid histologic scoring in research.[15] Potential parameters for the histological scoring system were chosen by the authors based on a systematic review of histological studies of breast implant capsules.[14] Each selected parameter was transferred to a binary scale or an ordinal scale from absent to severe to assess the prevalence and abundance of each parameter in the biopsies. The reproducibility and applicability of each potential parameter were tested by two observers on breast implant biopsies from 100 randomly selected women from the biobank. Disagreement between observers was discussed with a breast pathologist (AR) until a consensus was reached. Parameters with low applicability or rare incidence in the sample were discarded. The ordinal scales were modified to ensure that each scale point was used in the sample. To avoid

diagnostic drift, a reference library of histological images illustrating each parameter was developed. The reference library includes detailed instructions on how to score each parameter as well. It can be found in Supplemental Digital Content 1. In the next phase, the histological scoring system was validated in terms of observer reproducibility in the samples from the remaining 380 women in the biobank. Two observers (AT and MK) scored all 380 samples. In addition, 30 of these samples were scored twice to calculate intraobserver agreement. The biopsies from 100 randomly selected patients were scored by the breast pathologist (AR) to assess the agreement between the pathologist and the nonpathologist observers. The results from the histological scorings were used to assess the association between each parameter and the patient's clinical severity of capsular contracture using the Baker grade. A flowchart of the study design is presented in Figure 1.

Ethical considerations

Informed consent was obtained from all patients prior to inclusion in the study. The study was approved by the Regional Ethics Committee (H-20015276) and the Danish Data Protection Agency (P-2020-176 and P-2020-177).

Statistics

Agreement between and within observers was calculated with linear weighted Kappa scores. Interobserver agreement compared the scores of each histological parameter between two observers, and intraobserver agreement compared the duplicated scores of each histological parameter. Kappa scores were evaluated as no (0.01-0.20), fair (0.21-0.40), moderate (0.41-0.60), substantial (0.61-0.80) and almost perfect (0.81-1.00) agreement.(16) The association between the Baker grade (Baker I/II vs. Baker III/IV) and the histological parameter was estimated with a multivariable logistic regression with ridge penalization and robust covariance estimation to adjust for clustering of observers. All associations are reported as odds ratios with 95% CIs, and p values were evaluated at an alpha level of 5%. All statistics and plots were performed in R, version 4.2.0.

Results

Selection of parameters

We identified 16 potential parameters for the scoring system. Four parameters were related to fibrosis, including fibroblast and myofibroblast quantification, thickness of the dense collagen layer and organization of the collagen fibers. Four parameters were related to inflammation, including acute inflammation (neutrophil granulocytes), chronic inflammation (lymphocytes), vascularization and calcification. Five potential parameters were related to a foreign-body reaction, including synovial-like metaplasia, free silicone vacuoles, intracellular located silicone in macrophages (foam cells), multinucleated giant cells and granulomatous inflammation (silicone granulomas). Furthermore, the total number of layers in the capsule, activity in the stromal layer and neuromas were identified as potential parameters for the scoring system.

Based on the initial scoring of the 100 randomly selected biopsies, five parameters were excluded, including acute inflammation, granulomatous inflammation, number of layers, neuromas and histomorphometric grading of synovial-like metaplasia(17) due to low prevalence, low applicability and difficulty to standardize. Ten parameters were included in the final scoring system, each with its own two-, three- or four-point scale. An overview of the parameters in the histological scoring system can be seen in Table 2. The thickness of the collagen layer was divided into four intervals based on the median and interquartile range (IQR). Reliable morphological distinction between fibroblasts, myofibroblasts and resident macrophages is challenging in H&E-stained sections. Consequently, we chose to combine these into one parameter that evaluates the number of resident cells in the collagen layer. Furthermore, we combined free silicone vacuoles and foam cells into one parameter for silicone infiltration.

Reliability and reproducibility of the histological scoring system

The intra- and interobserver agreements were 0.85 (95% CI 0.85-0.85) and 0.80 (95% CI 0.80-0.80), respectively, which indicate almost perfect agreement. Six biopsies (1.5%) from patients with breast augmentation and six biopsies (7.5%) from patients with breast reconstruction were excluded due to poor quality of the histological slides (for instance, insufficient representation of the full capsule, tissue folding or sectioning artifacts). Examples of poor-quality biopsies have been included in the reference library. The agreement between the nonpathologist observers and the breast pathologist was 0.67 (95% CI 0.67-0.67), indicating substantial agreement.

<u>Histological features of the breast implant capsule</u>

When investigating the fibrotic changes, we found that the collagen layer generally constituted the majority of the implant capsule biopsies and varied greatly in thickness and fiber organization. The thickness of the collagen layer ranged from 72 μm to over 3000 μm. We found that 19% of the samples were highly organized with fibers running parallel to the implant surface and fibroblast nuclei elongated in the direction of traction and parallel to the fibers (Fig. 2A). In contrast, 14% of the samples were highly disorganized with multidirectional bundles of fibers and crosscut, rounder fibroblast nuclei. The cellularity of the resident cells in the collagen layer also varied greatly, with 24% of the samples containing many fibrocytes, fibroblasts, myofibroblasts and macrophages, whereas 12% of the samples had a low cellularity with an almost acellular appearance (Fig. 2A/2B).

When investigating the inflammatory response, we found chronic inflammatory infiltration to be present in 12% of the samples (Fig. 2C). Tissue calcification was generally rare and only seen in 6% of the samples. (Fig. 2B). Furthermore, we found a low degree of vascularization in most of the biopsies (85%); however, as expected, a high degree of vascularization was often seen in biopsies with severe inflammatory infiltration (Fig. 2C).

Silicone was observed extracellularly as small round to irregular translucent droplets of amorphous refractile material as well as intracellularly when phagocytized by histocytes, giving them a foam cell

appearance (Fig. 2D). Silicone was identified in over half of the included samples (54%). Multinucleated

giant cells, which are known to be part of a foreign body reaction, were seen in 22% of the samples and

were closely related to the presence of silicone vacuoles and foam cells (Fig. 2D). A layer of synovial-like

metaplasia was found adjacent to the implant surface in 70% of the samples with a varying thickness from

1 to >5 cell layers (Fig. 2E) but was absent in 30% of the samples (Fig. 2B). Moreover, we found a layer of

stromal tissue to be present between the synovial-like metaplasia and the layer of dense collagen fibers or

between bundles of densely packed collagen fibers in 42% of the samples (Fig. 2E). See Table 3 for the

distributions of histological scores between biopsies from patients with breast augmentation and breast

reconstruction.

Associations between the histological parameters and capsular contracture

To demonstrate the applicability of the scoring system, the association between the histological scores and

Baker grade was analyzed in 468 patients. We found that four fibrotic and inflammatory parameters were

significantly associated with capsular contracture. The thickness of the collagen layer had the strongest

association with capsular contracture: 40% of the capsules from breasts with capsular contracture (Baker

III/IV) had the highest score with a thickness above 600 μm compared with only 13% of the capsules from

breasts without capsular contracture (Baker I/II) (p<0.0001). Moreover, the collagen fibers were

significantly more organized with fibers running parallel to the implant surface in capsules from breasts

with capsular contracture than in capsules from breasts without capsular contracture (28% vs. 14%,

p<0.01). Infiltration of lymphocytes in the fibrous capsule was also significantly more present in capsules

from breasts with capsular contracture than in capsules from breasts without capsular contracture (17% vs.

6%, p<0.01). Furthermore, the presence of calcification was also significantly associated with capsular

contracture (14% vs. 0.7%, p<0.001). The remaining parameters were not significantly associated with

capsular contracture (Fig. 3).

Discussion

To the best of our knowledge, this is the first validated histological scoring system that can be used to evaluate the immunological and fibrotic response in breast implant capsules. The scoring system comprises 10 semiquantitative parameters that are highly reliable in terms of intra- and interobserver agreement and with substantial agreement between nonpathologist observers and a breast pathologist. To improve the applicability and usability of the scoring system, we created a large histological reference index with histological examples of each parameter and thorough instructions on how to score each parameter. We found no acute inflammation or neuromas and only one true silicone granuloma, and consequently, these parameters were discarded from the scoring system. Our histological scoring system provides an objective and standardized measure for characterizing the histology of breast implant capsules. It was validated in 480 patients after breast augmentation or breast reconstruction. The provided distributions of the individual parameters may serve as a strong reference point for future studies. The scoring system was developed for H&E-stained sections and does not require special immunohistochemical staining, which makes it cost-efficient to include in future studies. The scoring system can be applied to other important research questions, such as the impact of implant texture, radiotherapy and biological or artificial meshes. Therefore, we recommend that future studies include all 10 parameters and not just those significantly associated with capsular contracture, as we might find other important patterns when including all the visible histological characteristics of breast implant capsules.

As an example of the applicability of the scoring system, we analyzed the differences between patients with and without capsular contracture and found two fibrotic and two inflammatory parameters significantly associated with capsular contracture. The correlation between capsule thickness and capsular contracture is well established in the literature.(10,18–21) In a study by Siggelkow et al.(10) analyzing 53 biopsies from 43 patients, a significant association of capsule thickness and Baker grade was found, but they did not find an association with inflammation, silicone deposition or calcification. Bui et al.(19) also found capsules from

breasts with capsular contracture to be significantly thicker than those from breasts without capsular contracture based on the analysis of 49 samples from 41 patients. In line with our findings, they found capsules from breasts with capsular contracture to have a significantly higher degree of fiber organization than capsules from breasts without capsular contracture based on vector analysis. Most previous studies investigating the inflammatory reaction to breast implants have either been based on qualitative analysis or have not included statistical tests.(22–24) In a study by Kamel et al.(25) analyzing 93 biopsies, they did, however, find a significant association between the Baker grade and CD3-positive lymphocytes, but they did not find a correlation with calcification. In histological studies of capsular contracture, calcification is described as the end stage of chronic inflammation and the fibrotic reaction.(26,27) In a study by Peters et al.,(28) calcification was clinically present in 16% of breast implant capsules and was significantly more frequent in first-generation implants(29) than in second- and third-generation implants. In our study, we primarily included biopsies from breasts with textured fourth- or fifth-generation implants and found that calcification was present histologically in 14% of capsules from breasts with capsule contracture and 0.7% of capsules from breasts without capsular contracture. This may be the explanation why previous studies with smaller sample sizes have not been able to find a significant correlation between calcification and capsular contracture.(10,22,23,25) Previous studies have described a loss of cellularity in the fibrous capsule when the time of implantation increases.(27) Myofibroblasts are hypothesized to undergo apoptosis in the later stages of capsular contracture development, similar to what has been observed in wound healing and scar formation.(19) The role of fibroblasts and myofibroblasts in capsular contracture and how the cellularity change over time may be elucidated using specific immunohistochemical stains that improve the distinction between the two cell types, which is not possible in H&E-stained sections.(30,31) Previous studies have found an inverse relationship between the presence of synovial-like metaplasia and capsular contracture and have hypothesized that it plays an important protective role against capsular contracture.(18,32) Raso et al.(33) hypothesized that the protection is caused by the production of lubricating factors such as proteoglycans, whereas De Bakker(18) et al. hypothesized that synovial-like

metaplasia is lost due to the increasing amount of tissue and, thereby, loss of nutrition. Other potential

reasons could be that the layer is torn off during implant removal due to a strong adhesion between the

implant surface and the capsule, which has been described as a "Velcro effect"(14), or that the layer is lost

during sample preparation. The results from this study showed no significant association between the

absence of synovial-like metaplasia and capsular contracture. Silicone leakage is believed to trigger capsular

contracture by inducing an inflammatory foreign body response.(34) However, we did not find the

semiquantitative amount of silicone deposition or the presence of multinucleated giant cells in the capsule

to be independently associated with capsular contracture. This could be due to a potential

underestimation, as silicone is known to be lost during tissue preparation. Future studies performing a

quantitative analysis of silicone leakage adjusted for rupture status may have the sensitivity to find a

possible correlation between the amount of silicone in the capsule and capsular contracture.

Limitations

The study is based on a population of patients among whom the vast majority had textured implants

(>90%). This limits the generalizability of the scoring system in regard to smooth implants. The parameters

of the scoring system are scaled after a population with an even distribution of patients with and without

capsular contracture. Therefore, a study that, for example, is concerned with patients with very short

implantation times and no capsular contracture will probably find that their patients cluster in one extreme

of many of the scoring system's parameters. Standardization of the sample site has improved the

comparability between samples; however, this might impose a sampling bias, as potential variability of the

histological parameters between different locations of the capsule has not been investigated.

Conclusion

To our knowledge, this is the largest histological study of the breast implant capsule and the morphological

changes from capsular contracture. We have developed and validated a histological scoring system that is

highly reliable in terms of agreement and can be used with a standard H&E stain. Thus, it can provide an

accurate and objective assessment of breast implant capsules for future research studies and clinical

practice. We strongly encourage others to use and perhaps refine this scoring system.

Funding information: The work was funded by the Research Fund of Rigshospitalet and the Novo Nordisk

Foundation (grant number 052198)

Conflict of interest statement: The authors have no conflicts of interest to declare.

Data availability statement: The data that support the findings of this study are available from the

corresponding author upon reasonable request

References

1.	Hammond DC, Canady JW, Love TR, Wixtrom RN, Caplin DA. Mentor Contour Profile Gel Implants:

Clinical Outcomes at 10 Years. Plast Reconstr Surg. 2017;140(6):1142–50.

2.	Calobrace MB, Stevens WG, Capizzi PJ, Cohen R, Godinez T, Beckstrand M. Risk Factor Analysis for

Capsular Contracture: A 10-Year Sientra Study Using Round, Smooth, and Textured Implants for

Breast Augmentation. Plast Reconstr Surg. 2018 Apr;141(4S Sientra Shaped and Round Cohesive Gel

Implants):20S-28S.

3.	Spear SL, Murphy DK. Natrelle round silicone breast implants: Core study results at 10 years. Plast

Reconstr Surg. 2014;133(6):1354–61.

4.	Cunningham B. The Mentor Core Study on Silicone MemoryGel Breast Implants. Plast Reconstr Surg.

2007;120(7 Suppl 1):19S-32S.

5.	Grant Stevens W, Bradley Calobrace M, Alizadeh K, Zeidler KR, Harrington JL, d'Incelli RC. Ten-year

core study data for sientra's food and drug Administration–Approved round and shaped breast

implants with cohesive silicone gel. Plast Reconstr Surg. 2018;141(4S):7S-19S.

6.	Hvilsom GB, Hölmich LR, Steding-Jessen M, Frederiksen K, Henriksen TF, Lipworth L, et al. Delayed

breast implant reconstruction: A 10-year prospective study. J Plast Reconstr Aesthetic Surg. 2011;64(11):1466–74.

7.   Spear SL, Baker JL, Coffee HH. Classification of Capsular Contracture after Prosthetic Breast Reconstruction. Plast Reconstr Surg. 1995;96(5):1124.

8.   Hu H, Jacombs A, Vickery K, Merten SL, Pennington DG, Deva AK. Reply: Chronic Biofilm Infection in Breast Implants Is Associated with an Increased T-Cell Lymphocytic Infiltrate: Implications for Breast Implant-Associated Lymphoma. Plast Reconstr Surg. 2015 Jun;135(6):319–29.

9.   Doloff JC, Veiseh O, de Mezerville R, Sforza M, Perry TA, Haupt J, et al. The surface topography of silicone breast implants mediates the foreign body response in mice, rabbits and humans. Nat Biomed Eng. 2021;

10.   Siggelkow W, Faridi A, Spiritus K, Klinge U, Rath W, Klosterhalfen B. Histological analysis of silicone breast implant capsules and correlation with capsular contracture. Biomaterials. 2003;24(6):1101–9.

11.   Basu CB, Leong M, Hicks MJ. Acellular cadaveric dermis decreases the inflammatory response in capsule formation in reconstructive breast surgery. Plast Reconstr Surg. 2010;126(6):1842–7.

12.   Collett DJ, Rakhorst H, Lennox P, Magnusson M, Cooter R, Deva AK. Current Risk Estimate of Breast Implant-Associated Anaplastic Large Cell Lymphoma in Textured Breast Implants. Plast Reconstr Surg. 2019 Mar;143(3S A Review of Breast Implant-Associated Anaplastic Large Cell Lymphoma):30S-40S.

13.   Blombery P, Thompson ER, Prince HM. Molecular Drivers of Breast Implant-Associated Anaplastic Large Cell Lymphoma. Plast Reconstr Surg. 2019;143(3):59S-64S.

14.   Larsen A, Rasmussen LE, Rasmussen LF, Weltz TK, Hemmingsen MN, Poulsen SS, et al. Histological Analyses of Capsular Contracture and Associated Risk Factors: A Systematic Review. Aesthetic Plast Surg. 2021;45(6):2714–28.

15. Gibson-Corley KN, Olivier AK, Meyerholz DK. Principles for Valid Histopathologic Scoring in Research. Vet Pathol. 2013;50(6):1007–15.

16. McHugh M L. Interrater reliability: the kappa statistic. Biochem Medica [Internet]. 2012;22(3):276–82. Available from: https://hrcak.srce.hr/file/132393

17. Cheriyan T, Guo L, Orgill DP, Padera RF, Schmid TM, Spector M. Lubricin in human breast tissue expander capsules. J Biomed Mater Res B Appl Biomater. 2012/08/04. 2012;100(7):1961–9.

18. de Bakker E, van den Broek LJ, Ritt MJPF, Gibbs S, Niessen FB. The Histological Composition of Capsular Contracture Focussed on the Inner Layer of the Capsule: An Intra-Donor Baker-I Versus Baker-IV Comparison. Aesthetic Plast Surg. 2018/09/05. 2018;42(6):1485–91.

19. Bui JM, Perry T, Ren CD, Nofrey B, Teitelbaum S, Van Epps DE. Histological characterization of human breast implant capsules. Aesthetic Plast Surg. 2015/03/06. 2015;39(3):306–15.

20. Ersek RA, Burroughs JR, Ersek CL, Navarro A. Interrelationship of capsule thickness and breast hardness confirmed by a new measurement method. 1991;1069–73.

21. Prantl L, Pöppl N, Horvat N, Heine N, Eisenmann-Klein M. Serologic and histologic findings in patients with capsular contracture after breast augmentation with smooth silicone gel implants: Is serum hyaluronan a potential predictor? Aesthetic Plast Surg. 2005;29(6):510–8.

22. Prantl L, Schreml S, Fichtner-Feigl S, Pöppl N, Eisenmann-Klein M, Schwarze H, et al. Clinical and morphological conditions in capsular contracture formed around silicone breast implants. Plast Reconstr Surg. 2007;120(1):275–84.

23. Luke JL, Kalasinsky VF, Turnicky RP, Centeno JA, Johnson FB, Mullick FG. Pathological and biophysical findings associated with silicone breast implants: A study of capsular tissues from 86 cases. Vol. 100, Plastic and Reconstructive Surgery. 1997. p. 1558–65.

24. Carpaneda CA. Inflammatory reaction and capsular contracture around smooth silicone implants.

Aesthetic Plast Surg. 1997;21(2):110–4.

25.   Kamel M, Protzner K, Fornasier V, Peters W, Smith D, Ibanez D. The peri-implant breast capsule: an

        immunophenotypic study of capsules taken at explantation surgery. J Biomed Mater Res.

        2001;58(1):88–96.

26.   Yeoh G, Russell P, Jenkins E. Spectrum of histological changes reactive to prosthetic breast implants:

        a clinopathological study of 84 patients. Pathology. 1996;28(3):232–5.

27.   Legrand AP, Marinov G, Pavlov S, Guidoin M-F, Famery R, Bresson B, et al. Degenerative

        mineralization in the fibrous capsule of silicone breast implants. J Mater Sci Mater Med.

        2005;16(5):477–85.

28.   Peters W, Pritzker K, Smith D, Fornasier V, Holmyard D, Lugowski S, et al. Capsular calcification

        associated with silicone breast implants: Incidence, determinants, and characterization. Ann Plast

        Surg. 1998;41(4):348–60.

29.   Bradley Calobrace M, Caprizi PJ. The biology and evolution of cohesive gel and shaped implants.

        Plast Reconstr Surg. 2014;134(1s):6–11.

30.   Tevlin R, Borrelli MR, Irizarry D, Nguyen D, Wan DC, Momeni A. Acellular Dermal Matrix Reduces

        Myofibroblast Presence in the Breast Capsule. Plast Reconstr surgery Glob open. 2019;7(5):e2213–

        e2213.

31.   Moyer HR, Pinell-White X, Losken A. The effect of radiation on acellular dermal matrix and capsule

        formation in breast reconstruction: clinical outcomes and histologic analysis. Plast Reconstr Surg.

        2014;133(2):214–21.

32.   Bassetto F, Scarpa C, Caccialanza E, Montesco MC, Magnani P. Histological features of periprosthetic

        mammary capsules: silicone vs. polyurethane. Aesthetic Plast Surg. 2010/03/13. 2010;34(4):481–5.

33.   Raso DS, Schulte BA. Immunolocalization of keratan sulfate, chondroitin-4-sulfate, and chondroitin-

6-sulfate in periprosthetic breast capsules exhibiting synovial metaplasia. Plast Reconstr Surg. 1996;98(1):78–82.

34.   Katzin WE, Centeno JA, Feng LJ, Kiley M, Mullick FG. Pathology of lymph nodes from patients with breast implants: A histologic and spectroscopic evaluation. Am J Surg Pathol. 2005;29(4):506–11.

Table 1. Patient- and implant characteristics

	Breast augmentation	Breast reconstruction
Patients, n	400	80
Breasts, n	400	80
Median age, yr (IQR)	45.9 (38.1–54.2)	56.9 (49.7–67.2)
Median BMI, kg/m^2 (IQR)	22.3 (20.5–24.6)	24.5 (22.1–26.9)
Smoking, pt (%)		
Yes	98 (24)	4 (5)
No	302 (76)	76 (95)
ASA score, pt (%)		
I	258 (65)	23 (29)
II	123 (31)	47 (59)
III	8 (2)	8 (10)
Unknown	10 (2.5)	2 (2.5)
Indication for revisional surgery, pt (%)		
Capsular contracture	200 (50)	33 (41)
Obs. Rupture	64 (16)	10 (13)
Obs. BIA-ALCL	10 (2.5)	2 (2.5)
Implant age	7 (1.8)	0 (0.0)
Cosmetic reason	119 (30)	35 (44)
Baker grade, breast (%)		
I	147 (37)	23 (29)
II	53 (13)	24 (30)
III	110 (28)	23 (29)

IV	90 (23)	10 (13)
Debut of capsular contracture, breast (%)		
Under 1 year	47 (24)	7 (30)
Over 1 year	145 (73)	16 (70)
Unknown	8 (4.0)	0 (0.0)
Median time of implantation, yr, breast (IQR)	14.0 (8.0–18.0)	5.0 (2.2–11.8)
Implant plane, breast (%)		
Subpectoral	336 (84)	67 (84)
Prepectoral	63 (16)	2 (2.5)
Unknown	1 (0.3)	11 (14)
Implant brand, breast (%)		
Mentor	157 (39)	51 (64)
EuroSilicone	92 (23)	1 (1.3)
Nagor	6 (1.5)	0 (0.0)
Allergan/McGhan	40 (10)	14 (18)
CUI	19 (4.8)	0 (0.0)
Motiva	11 (2.8)	1 (1.3)
Silimed	4 (1.0)	0 (0.0)
Polytech	2 (0.5)	2 (2.5)
Other	12 (3.0)	0 (0.0)
Unknown	56 (14)	11 (14)
Implant rupture, breast (%)		
Yes	105 (26)	10 (13)
No	295 (74)	63 (79)
Implant texture, breast (%)		

Smooth	36 (9.0)	1 (1.3)
Textured	364 (91)	79 (99)
Previous implant exchange, breast (%)		
Yes	67 (17)	17 (21)
No	333 (83)	63 (79)
Radiotherapy, breast (%)		
Yes	-	11 (14)
No	-	69 (86)
Acellular dermal matrix, breast (%)		
Yes	-	22 (28)
No	-	58 (72)

Yr = year, IQR = interquartile range, BMI = body mass index, pt = patients, ASA score = American Society of

Anesthesiology Physical Status Classification Score

Table 2. The histological scoring system for breast implant capsules

	1	2	3	4
Fibrosis				
Thickness of collagen layer	<400 µm	400-600 µm	600-800 µm	>800 µm
Fiber organization	Highly disorganized	Moderately disorganized	Highly organized	-
Resident cells in the collagen layer	Few fibroblasts and macrophages	Several fibroblasts and macrophages	Many fibroblasts and macrophages	-
Inflammation				
Inflammatory infiltration	Few single inflammatory cells	Several single inflammatory cells and few aggregates	Severe infiltration with several follicular aggregates	-
Calcification	Absent	Present	-	-
Vascularization	Low	High	-	-
Foreign-body-reaction				
Synovial-like metaplasia	Absent	1-2 cell layers	3-4 cell layers	>5 cell layers
Silicone deposition	Absent or nearly absent	Few silicone vacuoles and/or foam cells	Several silicone vacuoles and/or foam cells	Abundant silicone vacuoles and/or foam cells
Giant cells	Absent	Few giant cells	Several giant cells	-
Activity in stromal layer	Absent stromal layer	Present with low cellularity	Present with high cellularity	-

Table 3. Distribution of histological scores between biopsies from patients with breast augmentation and

breast reconstruction

	Breast augmentation	Breast reconstruction	Total
Patients, n	394	74	468
Thickness of collagen layer (%)			
<400 µm	170 (43)	35 (47)	205 (44)
400–600 µm	87 (22)	14 (19)	101 (22)
600–800 µm	51 (13)	12 (16)	63 (14)
>800 µm	86 (22)	13 (18)	99 (21)
Fiber organization (%)			
Highly disorganized	57 (15)	7 (9.5)	64 (14)
Moderately disorganized	255 (65)	59 (80)	314 (67)
Highly organized	82 (21)	8 (11)	90 (19)
Resident cells in the collagen layer (%)			
Few fibroblasts, myofibroblasts and macrophages	51 (13)	6 (8.1)	57 (12)
Several fibroblasts and macrophages	244 (62)	56 (76)	300 (64)
Many fibroblasts and macrophages	99 (25)	12 (16)	111 (24)
Inflammatory infiltration (%)			
Few single inflammatory cells	348 (88)	65 (88)	413 (88)
Several single inflammatory cells and few aggregates	33 (8.4)	8 (11)	41 (8.8)

Severe infiltration with several follicular aggregates	13 (3.3)	1 (1.4)	14 (3.0)
Calcification (%)			
Absent	365 (93)	74 (100)	439 (94)
Present	29 (7.4)	0 (0.0)	29 (6.2)
Vascularization (%)			
Low	334 (85)	63 (85)	397 (85)
High	60 (15)	11 (15)	71 (15)
Synovial-like metaplasia (%)			
Absent	127 (32)	12 (16)	139 (30)
1–2 cell layers	129 (33)	31 (42)	160 (34)
3–4 cell layers	97 (25)	26 (35)	123 (26)
>5 cell layers	41 (10)	5 (6.8)	46 (9.8)
Silicone deposition (%)			
Absent or nearly absent	172 (44)	41 (55)	213 (46)
Few silicone vacuoles and/or foam cells	105 (27)	21 (28)	126 (27)
Several silicone vacuoles and/or foam cells	55 (14)	8 (11)	63 (14)
Abundant silicone vacuoles and/or foam cells	62 (16)	4 (5.4)	66 (14)
Giant cells (%)			
Absent	305 (77)	61 (82)	366 (78)
Few giant cells	35 (8.9)	5 (6.8)	40 (8.5)
Several giant cells	54 (14)	8 (11)	62 (13)

Activity in stromal layer (%)

Absent stromal layer	238 (60)	42 (57)	280 (60)
Present with low cellularity	42 (11)	7 (9.5)	49 (11)
Present with high cellularity	114 (29)	25 (34)	139 (30)

Figure 1. Flow diagram of the study design

Figure 2. Examples of the histological appearance of breast implant capsules

Footnote: *(A)* capsule with highly organized fibers running parallel to the implant surface and fibroblast nuclei elongated in the direction of traction, *(B)* capsule with a low cellularity of resident cells and the presence of calcification, *(C)* capsule with severe inflammatory infiltration and high vascularization, *(D)* capsule with abundant silicone vacuoles, foam cells and several multinucleated giant cells, and *(E)* capsule with synovial-like metaplasia of 3-4 cell layers and the presence of a stromal layer with low cellularity.

Figure 3. The association between each histological parameter and the clinical degree of capsular contracture estimated with a multivariable logistic regression

	OR	95% CI	p value
Collagen thickness	9.10	5.18-15.97	<0.0001*
Fiber organization	2.61	1.38-4.94	<0.01*
Resident cells	1.19	0.55-2.55	0.66
Inflammatory infiltration	3.12	1.37-7.15	<0.01*
Calcification	5.81	2.16-15.65	<0.001*
Vascularization	0.94	0.60-1.47	0.79
Synovial-like metaplasia	0.72	0.46-1.15	0.17
Silicone deposition	1.36	0.86-2.14	0.18
Giant cells	0.69	0.37-1.29	0.24
Stromal layer	1.01	0.62-1.66	0.95

Risk of capsular contracture (OR)