ORIGINAL ARTICLE

An exploratory machine learning study on paediatric abdominal pain

phenotyping and prediction

Kazuya Takahashi[1,2], Michalina Lubiatowska[1], Huma Shehwana[3], James K. Ruffle[1,4],

John A Williams[5,6,7], Animesh Acharjee[5,6,7], Shuji Terai[2], Georgios V Gkoutos[5,6,7],

Humayoon Satti[3], and Qasim Aziz[1]

1. Centre for Neuroscience and Trauma, Wingate Institute of Neurogastroenterology,

Blizard Institute, Barts and The London School of Medicine and Dentistry, Queen

Mary University of London, London, UK.

2. Division of Gastroenterology and Hepatology, Graduate School of Medical and

Dental Sciences, Niigata University, Niigata, Japan.

3. Department of Biological Sciences, National University of Medical Sciences,

Rawalpindi, Pakistan.

4. Queen Square Institute of Neurology, University College London, UK

5. College of Medical and Dental Sciences, Institute of Cancer and Genomic Sciences,

University of Birmingham, Birmingham, UK.

6. Health Data Research UK, Midlands Site, Birmingham, UK.

7. Centre for Health Data Science, Birmingham, UK

H.Satti and Q.Aziz are joint senior authors.

Corresponding Author

Qasim Aziz

Centre for Neuroscience and Trauma, Wingate Institute of Neurogastroenterology,

Blizard Institute, Barts and The London School of Medicine and Dentistry, Queen

Mary University of London, London, UK

Tel: +44 2078822655

E-mail: q.aziz@qmul.ac.uk

Conflicts of interest

None declared.

Funding information

This study was supported by JA Niigata Kouseiren Grant (Niigata University School of Medicine) and JSPS Grants-in-Aid for Scientific Research (grant number: 22K16013).

Author contributions

K Takahashi planned the study, performed the analysis, and drafted the manuscript; M Lubiatowska revised the manuscript; H Shehwana performed the analysis; JK Ruffle, JA Williams, A Acharjee, S Terai, and GV Gkoutos revised the manuscript; H Satti and Q Aziz planned the study, interpreted the data, and revised the manuscript.

ABSTRACT

Background: The exact mechanisms underlying paediatric abdominal pain (AP) remain unclear due to patient heterogeneity. This study aimed to identify AP phenotypes and develop predictive models to explore associated factors.

Methods: In 13,790 children from a large birth cohort, data on paediatric and maternal demographics and comorbidities were extracted from general practitioner records. Machine learning (ML) clustering was used to identify distinct AP phenotypes, and an ML-based predictive model was developed using demographics and clinical features.

Results: 1,274 children experienced AP (9.2 %) (average age: 8.4 ± 1.1 years, male/female: 615/659), who clustered into three distinct phenotypes: Phenotype 1 with an allergic predisposition (n = 137), Phenotype 2 with maternal comorbidities (n = 676), and Phenotype 3 with minimal other comorbidities (n = 340). As the number of allergic diseases or maternal comorbidities increased, so did the frequency of AP, with 17.6% of children with ≥ 3 allergic diseases and 25.6% of children with ≥ 3 maternal comorbidities. The predictive model demonstrated moderate performance in predicting paediatric AP (AUC 0.67), showing that a child's ethnicity, paediatric allergic diseases, and maternal comorbidities were key predictive factors. When stratified by

ML-predicted probability, observed AP rates were 18.9% in the <40% group, 44.8% in

the 40–50% group, 60.6% in the 50–60% group, and 100.0% in the >60% group.

Conclusions: Our findings reveal distinct phenotypes and associated factors of

paediatric AP by an ML approach. These insights suggest potential targets for future

research to clarify the underlying mechanisms of paediatric AP.

Keywords: paediatric abdominal pain, machine learning, artificial intelligence,

Uniform Manifold Approximation and Projection, Shapley Additive exPlanations

1 | Introduction

Abdominal pain (AP) is one of the most common symptoms among children and adolescents, with prevalence rates across the USA and Europe ranging from 0.3% to 19.0% [1–3]. In primary care, children presenting with AP are diagnosed with functional or medically un-explained AP in 80% of the cases, while organic causes are considerably less frequent [4,5]. Although rarely life-threatening, AP is often refractory to treatment and associated with psychiatric comorbidities, such as anxiety and depressive disorders [6], significantly affecting health-related quality of life [7]. Therefore, early recognition and intervention are warranted.

Organic diseases, as well as early life events ranging from allergy status, psychological comorbidity, and parental factors, are proposed risk factors for paediatric AP [5,8–10]. Moreover, these diverse risk factors can interact, contributing to its complexity and heterogeneity. A more robust stratification with large patient cohorts is essential for understanding the aetiology of paediatric AP, plausibly disclosing novel insights into its underlying mechanisms.

A robust approach for identifying subgroups of patients with shared characteristics is data-driven clustering by unsupervised machine learning (ML) [11,12]. Any yielded subgroups may share an underlying mechanism associated with AP. Furthermore,

supervised ML is considered a powerful tool for clinical outcome prediction [13], and

it could aid clinicians in assessing the risk of AP development in early childhood [14].

Based on this background, we hypothesised that an ML algorithm could classify

children with AP into distinct phenotypes based on common characteristics, thereby

helping to unravel the complex underlying AP mechanisms. Additionally, we

hypothesised that ML algorithms could predict the development of paediatric AP and

help identify the key factors associated with it.

In this exploratory study, we comprehensively evaluated the risk factors of paediatric

patients with AP in a large birth cohort, covering a broad spectrum of cases. Using

unsupervised ML, we delineated phenotypes of paediatric AP using paediatric and

maternal clinical data. Moreover, ML models tasked to predict the development of

paediatric AP uncovered underlying factors linked to its frequency, catalytic for future

hypothesis-driven research.

2 | Materials and Methods

2.1 | Participants

Between 2007 and 2011, 12,453 pregnant women (recruited at 26–28 weeks) and 13,858 children were registered in the Born in Bradford (BiB) cohort. The detailed demographics of the whole cohort are described elsewhere [15]. Data for this study were obtained from the BiB cohort under agreement, with informed consent collected at the time of data gathering. For this study, we included children who had linked general practitioner (GP) records and whose comorbidities could be identified using the systematized nomenclature of medicine clinical terms (SNOMED-CT), a standardized terminology system used in medicine and healthcare. The details of the study population are illustrated in Fig 1. Ethical approval was obtained from the Bradford Research Ethics Committee (Ref 07/H1302/112). This study was conducted following the principles of the Declaration of Helsinki.

2.2 | Definition of AP and frequency survey of paediatric and maternal comorbidities

AP was defined as the presence of one of the following diagnoses in SNOMED-CT, regardless of its cause or whether it was acute or chronic: AP, abdominal wall pain, or generalized AP. The frequencies of comorbidities, such as gastrointestinal (GI), psychological, and allergic diseases (asthma, eczema, urticaria, or hay fever), as well

as diseases causing somatic pain, were also retrieved using SNOMED-CT, since these are associated with both organic and functional AP [8,10,16–18]. Additionally, the comorbidities of the children's genetic mothers were extracted to identify any associations between the maternal comorbidities and paediatric AP. The diseases of the children and their mothers investigated in this study and their SNOMED-CT codes are listed in S1 Table. Fathers were not included in our analysis pipeline due to a significant proportion of missing values.

2.3 | Study 1. Identifying factors associated with paediatric AP

Study 1 aimed to determine the frequency of AP, elucidate the clinical characteristics of children with AP, and identify its associated factors. First, to adjust for background differences at the time of extraction between children with and without AP, we employed propensity score matching. After matching, we compared the two matched groups to assess differences in demographics and the frequency of comorbidities.

Next, we identified factors associated with paediatric AP. For this analysis, all comorbidities listed in S1 Table were included, irrespective of whether they occurred before or after the onset of AP, as the aim was to examine their overall association with paediatric AP.

2.4 | Study 2. Phenotyping paediatric AP using unsupervised ML clustering

Study 2 focused on children experiencing AP, using unsupervised ML clustering to delineate phenotypes. In Python (version 3.7.12) [19], Uniform Manifold Approximation and Projection (UMAP), a non-linear dimensionality reduction technique, was applied to embed all variables into a three-dimensional latent space for subsequent clustering [20]. A scatterplot was then generated from the latent features, and phenotypic AP clusters were identified using Hierarchical Density-based Spatial Clustering of Applications with Noise (HDBSCAN), a density-based hierarchical method [21]. Demographics and comorbidities of both children and their mothers were subsequently compared across the identified phenotypes to elucidate their distinct characteristics. Furthermore, we investigated whether the number of significant comorbidities identified in the phenotype analysis was associated with the frequency of paediatric AP.

2.5 | Study 3. Using supervised ML to predict the development of paediatric AP

To construct a balanced dataset, we randomly selected an equal number of children without AP from the control group to match those with AP (Fig 1), thereby mitigating potential model bias due to class imbalance. Predictive variables included demographic

information and pre-existing comorbidities diagnosed prior to the onset of AP.

Comorbidities with a prevalence of less than 1% were excluded to reduce the risk of

model overfitting. The variables used in the predictive model, along with their

frequencies in children with and without AP, are summarised in S2 Table. The overall

workflow for model development is illustrated in S1 Fig.

We initially developed predictive models for paediatric AP using several ML

algorithms, including eXtreme Gradient Boosting, Categorical Boosting (CatBoost),

Light Gradient Boosting Machine, and Random Forest. Model performance was

assessed using receiver operating characteristic (ROC) curves and the corresponding

area under the curve (AUC). In addition, accuracy, sensitivity, specificity, positive

predictive value (PPV), and negative predictive value (NPV) were calculated. The

model with the highest AUC was selected for subsequent feature importance analysis

and risk stratification.

To interpret model predictions, we employed Shapley Additive exPlanations (SHAP)

values. SHAP quantifies the contribution of each feature to individual predictions,

allowing us to identify which variables had the greatest influence on the model's

outputs and to enhance interpretability [22,23].

Finally, because the ML models produce continuous probability estimates for AP, we stratified children in the test dataset into four probability groups: <40%, 40–50%, 50–60%, and >60%. The observed risk of paediatric AP was then compared across these strata.

2.6 | Statistical analysis

Continuous data were expressed as mean ± standard deviation (SD). Categorical data were expressed as numbers and percentages. The Student's t-test and Chi-squared test were used for numerical and categorical data, respectively.

To compare children with AP and those without, the propensity score was calculated using logistic regression analysis, with age at extraction and gender as covariates. Matched children with and without AP were then extracted. A 1:2 nearest-neighbour matching without replacement was performed, with a calliper width set to 0.2 times the standard deviation of the logit of the propensity score. Model discrimination was assessed using the c-statistic, and covariate balance was evaluated using standardized mean differences (SMDs), with SMD < 0.1 indicating good balance.

Univariate and multivariate logistic regression analyses were utilised to calculate odds ratios (ORs) and 95% confidence intervals (CI) for each variable, determining

variables associated with paediatric AP. Variables significant in the univariate analysis were included in the multivariate analysis. As the comparisons between phenotypes were descriptive and intended to characterize group differences, Bonferroni correction was not applied. The Wald test was applied to compare ORs among the predicted probability groups. A p-value < 0.05 was considered statistically significant. EZR (Saitama Medical Center, Jichi Medical University, Saitama, Japan) was used for statistical analysis [24].

3 | Results

3.1 | Study 1. Identifying factors associated with paediatric AP

A total of 13,790 children were included in the analysis, of whom 1,274 (9.2%) experienced AP (male/female = 615/659). The average age at diagnosis of AP was 5.6 $\pm$ 2.7 years. The characteristics of all children are summarised in S3 Table.

As a result of propensity score matching, 1,239 children with AP and 2,478 children without AP were selected (Table 1). Compared to children without AP, more children of Pakistani origin were in the AP group, while fewer were of White origin (all $p <$

0.01). Children with AP had higher rates of comorbidities, including allergic diseases, appendicitis, Celiac disease, constipation, and gastro-oesophageal reflux disease (GORD), and migraine (all $p < 0.01$). Moreover, mothers of children with AP reported higher incidences of AP, allergic diseases, arthritis, chronic muscle pain, functional dyspepsia, GORD, irritable bowel syndrome (IBS), and migraine (all $p < 0.01$).

Multivariate logistic regression analysis revealed that Pakistani ethnicity, allergic diseases, GI diseases, migraine, maternal AP, maternal allergic diseases, maternal GORD, and maternal migraine were significantly associated with paediatric AP (Table 2). Non–significant results are summarized in S4 Table.

3.2 | Study 2. Phenotyping children using unsupervised ML clustering

The unsupervised model classified children with AP into three distinct phenotypes: 137 children in Phenotype 1 (10.8%), 677 children in Phenotype 2 (53.1%), and 340 children in Phenotype 3 (26.7%) (Figure 2). The remaining 120 children (9.4%) exhibited miscellaneous characteristics, making them unclassifiable by the model; these children were excluded from downstream analyses.

Subsequently, we compared the clinical characteristics among the three phenotypes (Table 3). Age at diagnosis of AP, gender, route of birth, and ethnicity showed almost

no significant differences among the three phenotypes. The frequency of GI disorders was low in all phenotypes and showed few significant differences between the phenotypes. The clinical characteristics of each phenotype are summarised below.

Phenotype 1

Most children in Phenotype 1 had allergic diseases (99.3%), and all mothers had allergic diseases (100%), which were significantly higher than in Phenotypes 2 and 3 ($p < 0.01$). In short, this phenotype was characterized as 'AP with allergic predisposition', suggesting relevance of allergic mechanisms in AP development in children.

Phenotype 2

In this phenotype, the frequency of allergic diseases is relatively high at 68.7%, but other comorbidities in the children were uncommon. In contrast, maternal comorbidities showed the highest frequencies in this phenotype: AP (70.0%), allergic diseases (40.0%), depressive and/or bipolar disorder (37.2%), GORD (18.9%), and migraine (34.0%). We termed Phenotype 2 as 'AP with mothers' comorbidities'.

Phenotype 3

The frequency of mother's AP (46.2%) was the second highest in Phenotype 3. However, the frequencies of other comorbidities, including children's allergic diseases and other maternal comorbidities were uncommon. This phenotype was termed 'AP with the least comorbidities'.

Impact of allergic diseases and maternal comorbidities on paediatric AP

We further investigated the effect of allergic diseases and maternal comorbidity burdens on the frequency of paediatric AP. As the number of paediatric allergic diseases increased, the frequency of AP in children increased commensurately (Fig 3A). Specifically, 17.6% of children with ≥ 3 allergic diseases experienced AP, which was significantly more frequent than those with 0–2 allergic diseases ($p < 0.01$). Similarly, as the number of maternal comorbidities increased, the frequency of AP in children also increased (Fig 3B). In cases where mothers had ≥ 3 comorbidities, the frequency of paediatric AP was 25.6%, which was significantly higher than in cases with 0–2 comorbidities ($p < 0.01$). Furthermore, when comparing the high-risk children (those who met the criteria of having both ≥ 3 allergic diseases and ≥ 3 maternal comorbidities) to the others (control), the high-risk group had a significantly higher frequency of AP ($p < 0.01$) (Fig 3C).

3.3 | Study 3. Development of ML predictive models for paediatric AP

Finally, we developed predictive models for paediatric AP. After evaluating several

algorithms, CatBoost yielded the highest AUC of 0.67 (95% CI: 0.63–0.71) and was

therefore selected for subsequent analyses (Fig 4A and S5 Table). The accuracy,

sensitivity, specificity, PPV, and NPV of the CatBoost-based model were 0.62 (95%

CI: 0.58–0.65), 0.68 (95% CI: 0.63–0.73), 0.55 (95% CI: 0.50–0.61), 0.62 (95% CI:

0.57–0.66), and 0.62 (95% CI: 0.57–0.68), respectively (Table 4).

SHAP analysis indicated that ethnicity had a greater influence on the prediction than

comorbidities, with White children being less likely and Pakistani children more likely

to develop AP (Figs 4C and 4D). Maternal AP emerged as the most influential

maternal comorbidity, while allergic diseases were the strongest predictor among

paediatric comorbidities.

When stratified by the probability estimated by the CatBoost-based model, the

observed rates of AP were 18.9%, 44.8%, 60.6%, and 100.0% in the <40%, 40–50%,

50–60%, and >60% probability groups, respectively (Fig 4B and Table 5). Compared

to the lowest-probability group (<40%), the odds of AP were significantly higher in the

40–50% group (OR 3.49, 95% CI: 1.94–6.26, p < 0.01), 50–60% group (OR 6.61, 95%

CI: 3.76–11.6, $p < 0.01$), and >60% group (OR could not be calculated, but $p < 0.01$).

4 | Discussion

The prevalence of paediatric AP in our study cohort was 9.2 %. Several child and

maternal factors, including paediatric ethnicity, GI and allergic diseases, as well as

maternal AP, were associated with paediatric AP. ML-based clustering successfully

identified three distinct phenotypes of paediatric AP, suggesting the presence of

subgroups among children with AP. The characteristics of these phenotypes indicate

that allergic diseases and maternal comorbidities are relevant to the development of AP

in two of the three phenotypes. The predictive performance of the ML model was

moderate when using the data from GP records. The most influential determinants of

paediatric AP in the predictive model were child ethnicity, allergic predisposition, and

maternal comorbidities, particularly maternal AP.

In this study, maternal GI comorbidities, such as AP and GORD, were significantly

associated with paediatric AP. This finding is consistent with previous research

showing that parental factors are related to functional AP in children [8,10,18]. For

example, children of mothers with IBS or chronic pain tended to experience more GI symptoms or functional AP [18,25]. This association may be explained by parental influence on children's pain experiences, whereby children may model their parents' pain behaviours, and parents may reinforce their children's pain complaints through solicitous responses [18,25–27]. Moreover, functional GI disorders, which are common causes of AP, are influenced by genetic factors [28], suggesting that the observed association may be partly due to shared genetic predisposition between mothers and children. Our findings underscore the important link between maternal factors and paediatric AP.

The ML clustering revealed three distinct phenotypes of paediatric AP. In all phenotypes, paediatric AP could not be simply explained by GI diseases. Phenotype 1 was characterized by the presence of allergic diseases. Some studies have reported that allergic diseases are significant risk factors for functional GI disorders such as IBS in children [9,17,29]. A recent study reported that IBS may be a food-induced disorder mediated by mast cell activation localized to the intestine [30]. Mast cells are known to be associated with visceral hypersensitivity [31]. These previous reports, along with our findings, imply that GI neuro-immune reactions could be one of the possible mechanisms underlying AP in children with allergic predispositions.

In Phenotype 2, where maternal comorbidities were associated, aforementioned parental mechanisms may play a role in the development of paediatric AP [18,25–28]. In contrast, in Phenotype 3, maternal AP was relatively common, occurring in 46.2% of cases. Therefore, some of the AP in children within Phenotype 3 may be explained by maternal influence. However, since the frequency of maternal comorbidities was much lower than in Phenotype 2, it is unlikely that all cases can be attributed to this factor. It is possible that the information available from GP records was insufficient to deeply phenotype children in Phenotype 3. Furthermore, 9.4% of children in this study were categorized as unclassified. With more detailed AP-related data, such as socioeconomic status, lifestyle, autonomic nervous system function, and gut microbiota composition, it may be possible to perform a deeper phenotypic analysis of children in Phenotype 3 and identify unclassified children as a distinct phenotype with specific characteristics [32–34].

The CatBoost-based model achieved a moderate AUC of 0.67 and identified the child's ethnicity as a key factor, consistent with its significance in the logistic regression model. The BiB cohort is unique in that it primarily includes children of White (29.3%) and Pakistani ethnicities (39.5%) [15]. Socioeconomic status varies by ethnicity within the cohort [35], and lower socioeconomic status is a known risk factor

for pain conditions in children, including AP [36]. These disparities may partly explain why ethnicity emerged as an important predictor.

Notably, approximately 60% of Pakistani couples represented in this cohort are reported to have consanguineous marriages, with 37% being first-cousin unions [39]. Children born from consanguineous marriages have been shown to have higher mortality rates and an increased number of primary care appointments compared to those born to non-consanguineous couples, indicating a higher burden of health issues [40]. Such consanguinity may also help explain why being of Pakistani ethnicity was a significant risk factor for paediatric AP in our model.

While the ML predictive model enabled a certain degree of risk stratification according to predicted probabilities, the substantial number of AP cases observed even in lower (40-50%) and intermediate (50-60%) probability groups suggests that the model's discriminative power remains limited. This may be attributable to the fact that, although important associated factors such as allergic diseases and maternal AP were included as predictors in the ML model, variables directly reflecting the underlying pathophysiology of AP were not incorporated. Future models may benefit from incorporating variables that capture the underlying pathophysiology of AP, such as

visceral hypersensitivity, altered gut-brain signalling, dysbiosis, and subclinical

intestinal inflammation [41–43].

Our study is not without limitations. Firs, we were unable to distinguish between

chronic and acute AP due to the lack of temporal detail in the available GP records.

While this limits the precision of phenotype classification, it reflects real-world clinical

practice where such distinctions are not always made or documented clearly. Moreover,

the identified risk factors–such as allergic predisposition– are likely relevant to both

types of AP, as supported by previous literature [17,44]. Most of our findings are also

consistent with prior studies, lending support to the credibility and reliability of the

data [40]. Therefore, despite this limitation, our findings provide valuable insights into

the broader mechanisms of paediatric AP and remain relevant for understanding its

development. Second, at the GP level, the exact cause of AP might not always be

identifiable. Therefore, the criteria used for each diagnosis and the exact causes of AP

in each case are unknown and was beyond our control. However, functional AP is

likely the predominant cause, as previous community-based studies of AP in children

suggest that approximately 80% of cases are classified as functional or non-medically

explained, whereas organic diseases account for only 5–10% of cases [42]. Despite its

complexity, real-world data of this nature holds significant potential for generating

valuable insights, particularly when analysed using advanced methods such as ML

[45]. Finally, although we evaluated the performance of our predictive models in an

out-of-sample test partition, we did not fully validate them using a different cohort of

children, which would further maximize generalizability. Moreover, a different cohort

would also be useful for verifying the reproducibility of clustering. However, our study

was an exploratory one and further research is now warranted to investigate the factors

we have identified in detail.

In conclusion, our exploratory study identified paediatric and maternal comorbidities

as significant associated factors for paediatric AP. Using data-driven clustering

techniques, we uncovered three distinct phenotypes of paediatric AP. These findings

provide a foundation for future research aimed at elucidating the precise mechanisms

underlying paediatric AP. The predictive models developed in this study highlight the

potential for early identification and intervention. Further studies incorporating more

detailed data could refine phenotyping and predictive models, leading to more accurate

predictions and paving the way for optimised, patient-personalised treatment strategies.

References

1. Chitkara DK, Rawat DJ, Talley NJ. The epidemiology of childhood recurrent abdominal pain in western countries: A systematic review. Am J Gastroenterol. 2005;100:1868–75.

2. Korterink JJ, Diederen K, Benninga MA, Tabbers MM. Epidemiology of pediatric functional abdominal pain disorders: A meta-analysis. PLoS One. 2015;10:1–17.

3. Devanarayana NM, de Silva DGH, de Silva HJ. Recurrent abdominal pain syndrome in a cohort of Sri Lankan children and adolescents. J Trop Pediatr. 2008;54:178–83.

4. Vila M, Kramer T, Obiols JE, Garralda ME. Abdominal pain in British young people: Associations, impairment and health care use [Internet]. Journal of Psychosomatic Research. 2012. p. 437–42. Available from: http://dx.doi.org/10.1016/j.jpsychores.2012.09.009

5. Spee LA, Van Den Hurk AP, Van Leeuwen Y, Benninga MA, Bierma-Zeinstra SM, Passchier J, et al. Childhood abdominal pain in primary care: Design and patient selection of the HONEUR abdominal pain cohort. BMC Fam Pract. 2010;11:6–8.

6. Dufton LM, Dunn MJ, Compas BE. Anxiety and somatic complaints in children with recurrent abdominal pain and anxiety disorders. J Pediatr Psychol. 2009;34:176–86.

7. Warschburger P, Hänig J, Friedt M, Posovszky C, Schier M, Calvano C. Health-related quality of life in children with abdominal pain due to functional or organic gastrointestinal disorders. J Pediatr Psychol. 2014;39:45–54.

8. Korterink J, Devanarayana NM, Rajindrajith S, Vlieger A, Benninga MA. Childhood functional abdominal pain: Mechanisms and management. Nat Rev Gastroenterol Hepatol. 2015;12:159–71.

9. Sjölund J, Kull I, Bergström A, Järås J, Ludvigsson JF, Törnblom H, et al. Allergy-related diseases in childhood and risk for abdominal pain-related functional gastrointestinal disorders at 16 years—a birth cohort study. BMC Med. 2021;19:1–11.

10. Zeevenhooven J, Rutten JMTM, van Dijk M, Peeters B, Benninga MA. Parental Factors in Pediatric Functional Abdominal Pain Disorders: A Cross-sectional Cohort Study. J Pediatr Gastroenterol Nutr. 2019;68:e20–6.

11. Ruffle JK, Farmer AD, Aziz Q. Artificial Intelligence-Assisted Gastroenterology — Promises and Pitfalls. Am J Gastroenterol. 2019;114:422–8.

12. Ko S, Pareek A, Jo C, Han H-S, Lee MC, Krych AJ, et al. Automated Risk Stratification of Hip Osteoarthritis Development in Patients With Femoroacetabular Impingement Using an Unsupervised Clustering Algorithm: A Study From the Rochester Epidemiology Project. Orthop J Sports Med. 2021;9:23259671211050612.

13. Takahashi K, Sato H, Shimamura Y, Abe H, Shiwaku H, Shiota J, et al. Achalasia phenotypes and prediction of per‐oral endoscopic myotomy outcomes using machine learning. Dig Endosc [Internet]. 2023; Available from: http://dx.doi.org/10.1111/den.14714

14. Bravo-Merodio L, Acharjee A, Russ D, Bisht V, Williams JA, Tsaprouni LG, et al. Translational biomarkers in the era of precision medicine. Adv Clin Chem. 2021;102:191–232.

15. Wright J, Small N, Raynor P, Tuffnell D, Bhopal R, Cameron N, et al. Cohort profile: The born in bradford multi-ethnic family cohort study. Int J Epidemiol. 2013;42:978–91.

16. Spee LAA, Lisman-Van Leeuwen Y, Benninga MA, Bierma-Zeinstra SMA, Berger MY. Prevalence, characteristics, and management of childhood functional abdominal pain in general practice. Scand J Prim Health Care. 2013;31:197–202.

17. Olén O, Neuman A, Koopmann B, Ludvigsson JF, Ballardini N, Westman M, et al. Allergy-related diseases and recurrent abdominal pain during childhood - A birth cohort study. Aliment Pharmacol Ther. 2014;40:1349–58.

18. Levy RL, Whitehead WE, Walker LS, Von Korff M, Feld AD, Garner M, et al. Increased somatic complaints and health-care utilization in children: Effects of parent IBS status and parent response to gastrointestinal symptoms. Am J Gastroenterol. 2004;99:2442–51.

19. Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. Scikit-learn: Machine learning in Python. the Journal of machine Learning research. 2011;12:2825–30.

20. McInnes L, Healy J, Melville J. UMAP: Uniform Manifold Approximation and Projection for Dimension Reduction [Internet]. arXiv [stat.ML]. 2018. Available from: http://arxiv.org/abs/1802.03426

21. Melvin RL, Xiao J, Godwin RC, Berenhaut KS, Salsbury FR. Visualizing correlated motion with HDBSCAN clustering. Protein Sci. 2018;27:62–75.

22. Lundberg S, Lee S-I. A unified approach to interpreting model predictions [Internet]. arXiv [cs.AI]. 2017. Available from: http://arxiv.org/abs/1705.07874

23. Lundberg SM, Erion GG, Lee S-I. Consistent individualized feature attribution for tree ensembles [Internet]. arXiv [cs.LG]. 2018. Available from: http://arxiv.org/abs/1802.03888

24. Kanda Y. Investigation of the freely available easy-to-use software "EZR" for medical statistics. Bone Marrow Transplant. 2013;48:452–8.

25. Sherman AL, Bruehl S, Smith CA, Walker LS. Individual and additive effects of mothers' and fathers' chronic pain on health outcomes in young adults with a childhood history of functional abdominal pain. J Pediatr Psychol. 2013;38:365–75.

26. Van Der Veek SMC, Derkx HHF, De Haan E, Benninga MA, Plak RD, Boer F. Do parents maintain or exacerbate pediatric functional abdominal pain? A systematic review and meta-analysis. J Health Psychol. 2012;17:258–72.

27. Walker LS, Smith CA, Garber J, Van Slyke DA. Development and validation of the pain response inventory for children. Psychol Assess. 1997;9:392–405.

28. D'Amato M. Genes and functional GI disorders: From casual to causal relationship. Neurogastroenterol Motil. 2013;25:638–49.

29. Jones MP, Walker MM, Ford AC, Talley NJ. The overlap of atopy and functional gastrointestinal disorders among 23 471 patients in primary care. Aliment Pharmacol Ther. 2014;40:382–91.

30. Aguilera-Lizarraga J, Florens MV, Viola MF, Jain P, Decraecker L, Appeltans I, et al. Local immune response to food antigens drives meal-induced abdominal pain. Nature. 2021;590:151–6.

31. Takahashi K, Khwaja IG, Schreyer JR, Bulmer D, Peiris M, Terai S, et al. Post-inflammatory abdominal pain in Patients with inflammatory bowel disease during remission: A comprehensive review. Crohn's & Colitis 360 [Internet]. 2021;3. Available from: https://academic.oup.com/crohnscolitis360/article/doi/10.1093/crocol/otab073/6423267

32. Zhou Q, Verne GN. Epigenetic modulation of visceral nociception. Neurogastroenterol Motil. 2022;34:e14443.

33. Gibler RC, Jastrowski Mano KE. Systematic Review of Autonomic Nervous System Functioning in Pediatric Chronic Pain. Clin J Pain. 2021;37:281–94.

34. Rajindrajith S, Zeevenhooven J, Devanarayana NM, Perera BJC, Benninga MA. Functional abdominal pain disorders in children. Expert Rev Gastroenterol Hepatol. 2018;12:369–90.

35. Petherick ES, Pearce N, Sunyer J, Wright J. Ethnic and socio-economic differences in the prevalence of wheeze, severe wheeze, asthma, eczema and medication usage at 4 years of age: Findings from the Born in Bradford birth cohort. Respir Med. 2016;119:122–9.

36. Grøholt E-K, Stigum H, Nordhagen R, Köhler L. Recurrent pain in children, socio-economic factors and accumulation in families. Eur J Epidemiol. 2003;18:965–75.

37. Chuah K-H, Mahadeva S. Cultural factors influencing functional gastrointestinal disorders in the East. J Neurogastroenterol Motil. 2018;24:536–43.

38. Borrello K, Lim U, Park S-Y, Monroe KR, Maskarinec G, Boushey CJ, et al. Dietary intake mediates ethnic differences in gut microbial composition. Nutrients. 2022;14:660.

39. Bradford Institute for Health Research. Born in Bradford. *Evidence Briefing: Genes and Health – Inheritance and Risk* [Internet]. 2022. Available from: https://borninbradford.nhs.uk/wp-content/uploads/2023/04/HG2954-BIHR-BiB-Evidence-Briefing-Genes-and-Health-4.pdf#:~:text=Evidence%20from%20the%20Born%20in,UK%20this%20fell%20to%2030

40. Small N, Kelly B, Malawsky DS, Lodh R, Oddie S, Wright J. Mortality, morbidity and educational outcomes in children of consanguineous parents in the Born in Bradford cohort. Wellcome Open Res. 2024;9:319.

41. Goudman L, Demuyser T, Pilitsis JG, Billot M, Roulaud M, Rigoard P, et al. Gut dysbiosis in patients with chronic pain: a systematic review and meta-analysis. Front Immunol. 2024;15:1342833.

42. Shulman RJ, Devaraj S, Heitkemper M. Activation of the innate immune system in children with irritable bowel syndrome evidenced by increased fecal human β-defensin-2. Clin Gastroenterol Hepatol. 2021;19:2121–7.

43. Castillo DF, Denson LA, Haslam DB, Hommel KA, Ollberding NJ, Sahay R, et al. The microbiome in adolescents with irritable bowel syndrome and changes with percutaneous electrical nerve field stimulation. Neurogastroenterol Motil. 2023;35:e14573.

44. Oishi K, Nakano N, Ota M, Inage E, Izawa K, Kaitani A, et al. MHC class II-expressing mucosal mast cells promote intestinal mast cell hyperplasia in a mouse model of food allergy. Allergy [Internet]. 2025; Available from: http://dx.doi.org/10.1111/all.16477

45. Liu F, Panagiotakos D. Real-world data: a brief review of the methods, applications, challenges and opportunities. BMC Med Res Methodol. 2022;22:287.

Table 1. Comparison of children with and without abdominal pain

	Before matching (n = 13,790)			After matching (n = 3,717)		
	Without AP (n=12,516)	With AP (n=1,274)	*p* value	Without AP (n=2,478)	With AP (n=1,239)	*p* value
Average age at the extraction ± SD (years)	8.3 ± 1.1	8.4 ± 1.1	<0.01	8.6 ± 1.1	8.6 ± 1.1	0.83
Average age at the diagnosis of abdominal pain ± SD (years)	N/A	5.6 ± 2.7	N/A	N/A	7.5 ± 2.3	N/A
Gender, female, n (%)	6,015 (48.1 %)	659 (51.7)	0.01	1,304 (52.6)	634 (51.2)	0.40
Route of birth, (vaginal), n (%)	9,415 (77.0 %)	943 (75.3)	0.17	1,866 (75.3)	916 (73.9)	0.36
Ethnicity*, n (%)						
Pakistani	4,750 (45.8)	703 (58.6)	<0.01	970 (46.9)	681 (58.4)	<0.01
White	3,771 (36.4)	267 (22.3)	<0.01	724 (35.0)	259 (22.2)	<0.01
Other ethnicities	1,847 (17.8)	229 (19.1)	0.27	374 (18.1)	227 (19.5)	0.34
Frequency of diseases, n (%)						
Allergic diseases	4,665 (37.3)	601 (47.2)	<0.01	918 (37.0)	584 (47.1)	<0.01
Appendicitis	24 (0.2)	27 (2.1)	<0.01	4 (0.2)	27 (2.2)	<0.01
Arthritis	6 (0.0)	1 (0.1)	0.49	0 (0.0)	1 (0.1)	0.16
Celiac disease	24 (0.2)	9 (0.7)	<0.01	3 (0.1)	9 (0.7)	<0.01
Constipation	142 (1.1)	57 (4.5)	<0.01	30 (1.2)	56 (4.5)	<0.01

FD	3 (0.0)	4 (0.3)	<0.01	2 (0.1)	4 (0.3)	0.08
GORD	342 (2.7)	66 (5.2)	<0.01	54 (2.2)	63 (5.1)	<0.01
IBD, colitis	4 (0.0)	2 (0.2)	0.1	1 (0.0)	2 (0.2)	0.22
Migraine	53 (0.4)	16 (1.3)	<0.01	10 (0.4)	16 (1.3)	<0.01
EDS, JHS	9 (0.1)	4 (0.3)	<0.01	1 (0.0)	4 (0.3)	0.03
Autism	36 (0.3)	4 (0.3)	0.87	9 (0.4)	4 (0.3)	0.84
Intellectual disability	16 (0.1)	2 (0.2)	0.78	3 (0.1)	2 (0.2)	0.75
Mother's abdominal pain	3,985 (31.8)	631 (49.5)	<0.01	782 (31.6)	616 (49.7)	<0.01
Mother's allergic disease	4,931 (39.4)	609 (47.8)	<0.01	965 (38.9)	597 (48.2)	<0.01
Mother's appendicitis	94 (0.8)	13 (1.0)	0.31	14 (0.6)	13 (1.0)	0.10
Mother's arthritis	495 (4.0)	109 (8.6)	<0.01	24 (1.0)	28 (2.3)	<0.01
Mother's Celiac disease	73 (0.6)	14 (1.1)	0.04	17 (0.7)	13 (1.0)	0.24
Mother's chronic fatigue syndrome	25 (0.2)	3 (0.2)	0.74	6 (0.2)	3 (0.2)	1.00
Mother's chronic muscle pain	167 (1.3)	30 (2.4)	0.01	30 (1.2)	29 (2.3)	<0.01
Mother's constipation	128 (1.0)	10 (0.8)	0.55	36 (1.5)	10 (0.8)	0.09
Mother's depressive disorder, bipolar disorder	2,632 (21.0)	309 (24.3)	0.01	555 (22.4)	300 (24.2)	0.22
Mother's FD	162 (1.3)	33 (2.6)	<0.01	36 (1.5)	33 (2.7)	<0.01
Mother's GORD	876 (7.0)	171 (13.4)	<0.01	172 (6.9)	170 (13.7)	<0.01
Mother's IBD	84 (0.7)	10 (0.8)	0.59	12 (0.5)	9 (0.7)	0.35

Mother's IBS	877 (7.0)	126 (9.9)	<0.01	171 (6.9)	125 (10.1)	<0.01
Mother's migraine	1,845 (14.7)	285 (22.4)	<0.01	378 (15.3)	283 (22.8)	<0.01
Mother's EDS, JHS	47 (0.3)	5 (0.4)	0.93	10 (0.4)	5 (0.4)	1.00
Mother's obsessive-compulsive disorder	61 (0.5)	10 (0.8)	0.15	13 (0.5)	10 (0.8)	0.30
Mother's schizophrenia	6 (0.0)	2 (0.2)	0.17	1 (0.0)	1 (0.1)	0.62
Mother's intellectual disability	44 (0.4)	1 (0.1)	0.1	8 (0.3)	1 (0.1)	0.16

* Gender and age at extraction were used to calculate the propensity score. Before matching, the standardized mean differences (SMD) for gender and age were 0.073 and 0.337, respectively. After matching, the SMD values for gender and age were reduced to 0.029 and 0.007, respectively, indicating that the data were well balanced. The c-statistic was 0.60 (95% CI: 0.58–0.61).

AP, abdominal pain; CI, confidence interval; SD, standard deviation; FD, functional dyspepsia; GORD, gastro-oesophageal reflux disease; IBD, inflammatory bowel disease; IBS, irritable bowel syndrome; EDS, Ehlers-Danlos syndrome; JHS, joint hypermobility syndrome

Table 2. Significant results of logistic regression analysis for the diagnosis of paediatric abdominal pain

	Univariate		Multivariate	
	OR (95% CI)	*p* value	OR (95% CI)	*p* value
Pakistani (vs. other ethnicities)	1.59 (1.37–1.83)	<0.01	1.58 (1.39–1.79)	<0.01
Allergic diseases	1.52 (1.32–1.74)	<0.01	1.19 (1.05–1.34)	0.01
Appendicitis	13.8 (4.81–39.5)	<0.01	8.92 (4.94–16.1)	<0.01
Celiac disease	6.04 (1.63–22.3)	0.01	3.35 (1.53–7.34)	<0.01
Constipation	3.86 (2.47–6.05)	<0.01	3.11 (2.23–4.33)	<0.01
GORD	2.40 (1.66–3.48)	<0.01	1.64 (1.23–2.18)	<0.01
Migraine	3.23 (1.46–7.14)	<0.01	2.95 (1.64–5.33)	<0.01
Mother's abdominal pain	2.14 (1.86–2.47)	<0.01	1.68 (1.48–1.90)	<0.01
Mother's allergic diseases	1.46 (1.27–1.67)	<0.01	1.17 (1.03–1.33)	0.01
Mother's arthritis	2.36 (1.36–4.10)	<0.01	1.39 (0.89–2.16)	0.14
Mother's chronic fatigue syndrome	1.96 (1.17–3.27)	0.01	0.98 (0.28–3.41)	0.98
Mother's chronic muscle pain	1.96 (1.17–3.27)	0.01	1.12 (0.73–1.72)	0.61
Mother's FD	1.86 (1.15–2.99)	0.01	1.15 (0.76–1.74)	0.51

Mother's GORD	2.13 (1.70–2.67)	<0.01	1.46 (1.21–1.76)	<0.01
Mother's IBS	1.51 (1.19–1.93)	<0.01	1.07 (0.86–1.33)	0.55
Mother's migraine	1.64 (1.38–1.95)	<0.01	1.24 (1.06–1.44)	0.01

FD, functional dyspepsia; GORD, gastro-oesophageal reflux disease; IBD, inflammatory bowel disease; IBS, irritable bowel syndrome; EDS, Ehlers-Danlos syndrome; JHS, joint hypermobility syndrome

Table 3. Comparison of clinical characteristics among 3 pain phenotypes

	Phenotype 1 (n = 137)	Phenotype 2 (n = 677)	Phenotype 3 (n = 340)	P value (Phenotype 1 vs. 2)	P value (Phenotype 1 vs. 3)	P value (Phenotype 2 vs. 3)
Average age at the extraction ± SD (years)	8.5±1.1	8.7±1.1	8.6 ±1.2	0.15	0.52	0.34
Average age at the diagnosis of abdominal pain ± SD (years)	5.6±2.8	5.6±2.7	5.6±2.7	0.95	0.99	0.91
Gender, female, n (%)	67 (48.9)	334 (49.3)	172 (50.6)	0.93	0.74	0.71
Route of birth, (vaginal), n (%)	111 (81.0)	477 (70.5)	260 (76.5)	0.01	0.28	0.04
Ethnicity, n (%)						
White	31 (22.6)	146 (22.8)	62 (20.1)	0.97	0.54	0.34
Pakistani	74 (54.0)	382 (59.6)	183 (59.2)	0.23	0.30	0.91
Others	32 (23.4)	149 (22.0)	95 (27.9)	0.73	0.31	0.04

Frequency of comorbidities, n (%)

	Phenotype 1 (n = 137)	Phenotype 2 (n = 677)	Phenotype 3 (n = 340)	P value (Phenotype 1 vs. 2)	P value (Phenotype 1 vs. 3)	P value (Phenotype 2 vs. 3)
Allergic disease	136 (99.3)	465 (68.7)	0 (0.0)	<0.01	<0.01	<0.01
Appendicitis	3 (2.2)	17 (2.5)	5 (1.5)	0.83	0.58	0.28
Arthritis	0 (0.0)	1 (0.1)	0 (0.0)	0.65	N/A	0.48
Celiac disease	3 (2.2)	1 (0.1)	2 (0.6)	0.05	0.44	0.78
Constipation	6 (4.4)	38 (5.6)	10 (2.9)	0.56	0.43	0.06
FD	0 (0.0)	3 (0.4)	0 (0.0)	0.44	N/A	0.22
GORD	11 (8.0)	40 (5.9)	13 (3.8)	0.35	0.06	0.16
IBD	1 (0.7)	0 (0.0)	1 (0.3)	0.03	0.51	0.16
Migraine	1 (0.7)	12 (1.8)	3 (0.9)	0.38	0.87	0.27
EDS, JHS	1 (0.7)	3 (0.4)	0 (0.0)	0.66	0.12	0.22
Autism	0 (0.0)	0 (0.0)	2 (0.6)	N/A	0.37	0.05

Intellectual disability	0 (0.0)	0 (0.0)	0 (0.0)	N/A	N/A	N/A
Mother's abdominal pain	0 (0.0)	474 (70.0)	157 (46.2)	<0.01	<0.01	<0.01
Mother's allergic diseases	137 (100.0)	271 (40.0)	81 (23.8)	<0.01	<0.01	<0.01
Mother's appendicitis	2 (1.5)	7 (1.0)	4 (1.2)	0.66	0.80	0.84
Mother's arthritis	10 (7.3)	77 (11.4)	14 (4.1)	0.53	0.50	<0.01
Mother's Celiac disease	0 (0.0)	9 (1.3)	4 (1.2)	0.18	0.20	0.84
Mother's chronic fatigue syndrome	1 (0.7)	2 (0.3)	0 (0.0)	0.44	0.12	0.32
Mother's chronic muscle pain	1 (0.7)	24 (3.5)	4 (1.2)	0.08	0.67	0.03
Mother's constipation	0 (0.0)	8 (1.2)	2 (0.6)	0.56	0.37	0.37
Mother's depressive disorder, bipolar disorder	28 (20.4)	252 (37.2)	0 (0.0)	<0.01	<0.01	<0.01
Mother's FD	1 (0.7)	27 (4.0)	4 (1.2)	0.21	1.00	0.04
Mother's GORD	14 (10.2)	128 (18.9)	14 (4.1)	0.04	0.05	<0.01
Mother's IBD	0 (0.0)	5 (0.7)	5 (1.5)	0.31	0.15	0.26
Mother's IBS	12 (8.8)	86 (12.7)	24 (7.1)	0.75	1.00	0.02
Mother's migraine	15 (10.9)	230 (34.0)	20 (5.9)	<0.01	0.24	<0.01
Mother's EDS, JHS	0 (0.0)	3 (0.4)	1 (0.3)	0.44	0.53	0.72
Mother's obsessive-compulsive disorder	2 (1.5)	5 (0.7)	3 (0.9)	0.40	0.58	0.81
Mother's intellectual disability	0 (0.0)	0 (0.0)	0 (0.0)	N/A	N/A	N/A
Mother's schizophrenia	1 (0.7)	0 (0.0)	1 (0.3)	0.03	0.51	0.16

FD, functional dyspepsia; GORD, gastro-oesophageal reflux disease; IBD, inflammatory bowel disease; IBS, irritable bowel syndrome; EDS, Ehlers-Danlos syndrome; JHS, joint hypermobility syndrome

Table 4. Results of predictive models for paediatric abdominal pain

Performance metrics	Outcomes
AUC (95% CI)	0.67 (0.63–0.71)
Accuracy (95% CI)	0.62 (0.58–0.65)
Sensitivity (95% CI)	0.68 (0.63–0.73)
Specificity (95% CI)	0.55 (0.50–0.61)
PPV (95% CI)	0.62 (0.57–0.66)
NPV (95% CI)	0.62 (0.57–0.68)

Abbreviations: CI, confidence interval; PPV, positive predictive value; NPV, negative predictive value;

AUC, area under the curve

Table 5. Risk stratification of paediatric abdominal pain based on the probability predicted by the CatBoost-based ML model

AP probability by ML	Observed AP rate, % (n/N)	OR (95% CI)	p value
<40%	18.9 (17/90)	ref	
40–50%	44.8 (108/241)	3.49 (1.94–6.26)	<0.01
50–60%	60.6 (257/424)	6.61 (3.76–11.6)	<0.01
>60%	100.0 (10/10)	Inf	<0.01
Abbreviations: AP, abdominal pain; CI, confidence interval; ML, machine learning; OR, odds ratio			

Figure legends

FIGURE 1. Study population

Among 13,858 children in the Born in Bradford cohort, 13,790 with linked general practitioner (GP) records were included. Study 1 identified factors associated with paediatric abdominal pain (AP) using propensity score matching (1:2 ratio). Study 2 performed phenotyping of AP using unsupervised machine learning (ML) clustering. Study 3 developed predictive models using supervised ML, including 1,274 children with AP and 1,274 randomly selected children without AP (1:1 ratio).

FIGURE 2. Phenotyping of children with abdominal pain

Scattering plots of based on three new variables created by unsupervised Uniform Manifold Approximation and Projection (UMAP). Hierarchical Density-Based Spatial Clustering of Applications with Noise (HDBSCAN) found 3 distinct phenotypes.

FIGURE 3. The frequency of paediatric abdominal pain depending on the number of allergic diseases and the number of maternal comorbidities

A–B. The bar graphs illustrate the relationship between the frequency of paediatric abdominal pain (AP) and the number of allergic diseases in children, as well as the number of comorbidities in their mothers. Error bars in the figure represent the 95% confidence intervals (CI) for the frequency of paediatric AP. **C.** The high-risk group is defined as children with both ≥ 3 allergic diseases and ≥ 3 maternal comorbidities while the remaining children are regarded as the control group. The frequency of AP is compared between the high-risk and control groups.

FIGURE 4. Results of supervised machine learning predictive models for paediatric abdominal pain

A. Receiver operating characteristic (ROC) curve of the CatBoost-based model. **B.** The y-axis represents the observed abdominal pain (AP) rate, while the x-axis indicates the predicted probability groups. Error bars represent the 95% confidence intervals of the observed AP rates. **C.** The mean Shapley Additive exPlanations (SHAP) value of each variable. Variables written in red represent maternal comorbidities, while those written in black represent variables related to the children themselves. **D.** The beeswarm plots show the SHAP feature importance and the direction of the effect of each variable on

the model. The horizontal axis represents the SHAP value. Since all the variables used in the predictive models were binary, red and blue plots indicate positive or negative for each variable, respectively. For example, in the case of 'Pakistani', red plots tend to be distributed in the positive SHAP value and blue plots in the negative SHAP value. This indicates that Pakistani children are more likely to have AP.

S1 FIGURE. Workflow for developing predictive models

The entire dataset was randomly partitioned into 70% for model training and 30% for out-of-sample testing. Using the training data, we trained predictive models. Grid-search with 10-fold cross-validation was employed to tune the hyperparameters of each model using the training dataset. The training dataset was divided into 10 folds. Nine folds were used as the training dataset, and the remaining one-fold was used as the validation dataset. This process was repeated 10 times to optimize the hyperparameters of each predictive model. The performance of each model was evaluated using the test dataset.

A. ROC curve

B. Abdominal pain rate

C. SHAP feature importance

D. Effect direction of each variable