Prevalence and Factors Associated with Overweight and Obesity among Women of Reproductive Age in Cambodia: Analysis of Cambodia Demographic and Health Survey 2021-2022

Samnang Um[1,2]*, Yom An[1]

2 The National Institute of Public Health, Phnom Penh, Cambodia
2 Doctor of Philosophy in Social Science, Faculty of Social Science and Humanities, Royal University of Phnom Penh, Cambodia
umsamnang56@gmail.com, ORCID: https://orcid.org/0000-0002-1521-1260

Abstract

Introduction. Overweight and obesity are associated with increased rates of chronic disease and death globally. In Cambodia, the prevalence of overweight and obesity among women is high and may be growing. This study aimed to determine the prevalence and factors associated with overweight and obesity among women of reproductive age (WRA) in Cambodia. **Methods.** We analyzed data from the 2021-2022 Cambodia Demographic and Health Survey (CDHS) that used a two-stage stratified cluster sampling design. Data analysis was restricted to non-pregnant women, resulting in an analytic sample of 9,417 WRA. Multivariable logistic regressions were performed using STATA V17 to examine factors associated with overweight and obesity. **Results**. Prevalence of overweight and obesity among non-pregnant women of reproductive age were 22.56% and 5.61% were overweight and obese respectively. Factors independently associated with increased odds of overweight and/or obesity included women aged 20-29 years [AOR=1.85; 95% CI: 1.22 - 2.80], 30-39 years [AOR=3.34; 95% CI: 2.21 - 5.04] and 40-49 years [AOR=5.57; 95% CI: 3.76 - 8.25], married women [AOR=2.49; 95% CI: 1.71 - 3.62], women from middle wealth quintile [AOR=1.21; 95% CI : 1.02 - 1.44], and rich wealth quintile [AOR=1.44 ; 95% CI: 1.19 - 1.73], having at least three children or more [AOR=1.40; 95% CI: 1.00 - 1.95], ever drink alcohol [AOR=1.24 ; 95% CI: 1.04 - 1.47], and current drink alcohol [AOR=1.2; 95% CI: 1.01 - 1.45]. On the contrary, following factors were independently associated with decreased odds of having overweight and obese: women with at least secondary education

[AOR=0.73; 95% CI: 0.58-0.91], and working in in manual labor jobs [AOR=0.76; 95% CI: (0.64 - 0.90]. **Conclusion.** Increased age, married women, having at least three children and alcohol consumption were the main risk factors associated with overweight and/or obesity. Conversely, higher education, and working in manual labor were negatively associated with overweight and/or obesity. Cambodia's non-communicable disease (NCD) public health programs should consider these characteristic for targeting interventions to further reduce overweight and/or obesity in the coming years.

Keywords: Overweight, obesity, non-communicable diseases, NCDs, women at reproductive age, CDHS, Cambodia

Introduction

Overweight and obesity are major global public health challenges that have been rapidly on the rise in the last four decades and are regarded as an epidemic [1]. In 2016, approximately 39% (1.9 billion) of adults aged 18 years and older were overweight and 13% (650 million) were obese worldwide [1]. In general, being overweight and obesity are major risk factors for several non-communicable diseases (NCDs) including cardiovascular and kidney diseases, type 2 diabetes, some cancers, musculoskeletal disorders, and other chronic diseases [2,3]. Moreover, among women of reproductive age (WRA), overweight and obesity have been associated with increased risk of pregnancy complications, caesarean section births, adverse birth outcomes, and infant mortality [4]. Overweight and obesity are the fourth leading cause of risk-attributable mortality [5], with a reduced life expectancy of 5-20 years, depending on the severity of the condition and the presence of comorbidities [1]. According to the WHO, being overweight and obese are the leading risks for global deaths with at least 2.8 million adults dying annually due to these conditions [1]. Overweight and obesity were higher proportion among women as compared to men in both developed and developing countries; for instance, overweight was 40% in women vs 39% in men and obesity 15% in women vs. 11% in men [1]. The Global Nutrition Report 2019 showed that 26.1% of women and 20.4% of men were overweight, while obesity was 6.3% in women

compared to 3.5% in men [6]. Along with the rapidly increasing population in Cambodia,

from 15.42 million in 2015 to nearly 16.21 million in 2019 [7], the prevalence of

overweight and obesity are also steadily on the rise. The 2021-22 Cambodia

Demographic and Health Survey (CDHS) report indicated that overweight and obesity

among non-pregnant women at reproductive age increased from 18% in 2014 to 39% in

2021-22 [8]. It was estimated that overweight and obesity contribute to rising healthcare

costs in Cambodia (approximately 1.7% of its annual gross domestic product (GDP) per

capita) and is a major contributor to mortality and decreased general health and

productivity [9]. Regarding predictors of overweight and obesity among women include

higher socioeconomic status, older age, marriage, living in an urban residence, and lack

of education [3,10-12]. Women with formal employment had higher odds of being

overweight or obese than informally employed women [3,13]. Also, women who used

hormonal contraceptives such as oral contraceptive pills, implants, patches, and rings

were at higher risk of being overweight or obese [14,15]. Globally, 30% of daily smokers

are overweight or obese [16], with women smokers at greater risk for overweight or

obesity than men smokers [17,18]. Women with frequent television watching [19],

alcohol drinking [20,21], frequent consumption of sweets foods and unhealthy foods [22]

were found to have higher odds of being overweight or obese. To our knowledge,

factors associated with overweight and obesity specifically among WRA in Cambodia

using an updated data have not been explored. A previous study on prevalence of

overweight and obesity among WRA and its associated factors utilized data since 2014

[3]. In the context of Cambodia where the prevalence of overweight and obesity in WRA

have increased rapidly, a thorough and comprehensive investigation of a wide range of

socio-demographic and behavioral factors is warranted to identify the factors

independently associated with having overweight and/or obesity. Identifying the key

modifiable socio-demographic and behavioral factors, as well as women who are at a

high-risk of having overweight and obesity, may help guide the timely development of

promising and feasible public health intervention strategies to address the growing

overweight and obesity pandemic. Therefore, we aimed to determine the prevalence

and examined sociodemographic and behavioral factors associated with overweight and

obesity among WRA in Cambodia.

Methods

Data sources

We followed the methods of Um et al., 2023 [3]. To analyze prevalence and factors associated with overweight and obesity among WRA in Cambodia, we used existing women's data from the 2021-2022 CDHS. The CDHS is a nationally representative population-based household survey implemented by the National Institute of Statistics (NIS) in collaboration with the Ministry of Health (MoH) with financial support from the Government of the Cambodia, and technical assistance from ICF International (USA) thorough the Demographic and Health Survey program. from September 15, 2021 to February 15, 2022 [8]. The sampling frame used for the 2021-2022 CDHS was taken from the Cambodia General Population Census 2019 [7]. Thereafter, the participants in CDHS 2021-2022 were selected using probability proportion based on two-stage stratified cluster sampling from the chosen sampling frame. In the initial stage, 709 enumeration area (EAs) (241 urban areas and 468 rural areas) were selected. In the second stage, an equal systematic sample of 30 households was selected from each cluster, for a total sample size of 21,270 households. In total, 19,496 women aged 15-49 years were interviewed with a response rate of 98.2%. Survey data were obtained through face-to-face survey interviews using a standardized survey instrument by trained interviewers. Anthropometric of weight and height measurements for adult women age 15-49 were taken by trained female field staff using standardized instruments and procedures [7]. Weight measurements were taken using scales with a digital display (UNICEF model S0141025). Height and length were measured using a portable adult height measurement system (UNICEF model S0114540). The detailed protocol and methods were published previously [8]. Eligible participants for this study were women of reproductive age 15-49 years, with exclusion of pregnant women and those who had a birth within 2 months before the survey, with available body mass index (BMI) data. As a result, we excluded a total of 828 pregnant women and 9,249 women with missing BMI data. A final sample including in this analysis was 9,417 women of reproductive age 15-49 years.

Measurements

Outcome variable

The primary outcome variable of this study was overweight and obese for women of
reproductive age. BMI - defined by dividing a person's weight in kilograms by the square
of their height in metres (kg/m^2) – was used to measure the outcome [8,23]. For adults
over 15 years old, the following BMI ranges were used: underweight (<18.0 kg/m^2),
normal weight (18.5-24.9 kg/m^2), overweight (25.0-29.9 kg/m^2) and obesity (≥
30.0 kg/m^2) [8,23]. **Overweight/Obese** was defined as a binary outcome for which
women with a BMI ≥ 25.0 kg/m^2 were classified as overweight and obese (coded = 1),
while women with a BMI < 25.0 kg/m^2 were coded as other (coded = 0).

Covariate variables

Covariates included socio-demographic characteristics of women of reproductive age:
Women's age in years (15-19, 20-29, 30-39, and 40-49), marital status (not married,
married or living together, and divorced or widowed or separated), educational level (no
formal education, primary, secondary or higher education), occupation (not working,
agriculture, manual labor or unskilled, professional or sealer or services), number of
children ever born (no children, one-two child, and three and more children).
Households' wealth status was represented by a wealth index calculated via principal
component analysis (PCA) and using variables for household assets and dwelling
characteristics. Weighted scores divided into five wealth quintiles (poorest, poorer,
medium, richer, and richest) each comprising 20% of the population [8], place of
residence (rural vs urban). Cambodia's provinces were regrouped for analytic purposes
into a categorical variable with 4 geographical regions: plains, Tonle Sap, costal/sea,
and mountains [7]. Behavioral factors including smoking (non-smoker vs smoker),
alcohol consumption in the past month (never drink, ever drink, and current drink)
current drink of alcohol corresponds to one can or bottle of beer, one glass of wine, or
one shot of spirits in past month, watching television at least once a week (yes vs no),

contraceptive usage (not used, hormonal methods (using a pill, emergency
contraceptive pill, Norplant, injection, and vaginal rings), non-hormonal methods
(condoms, the diaphragm, the IUD, spermicides, lactational amenorrhea, and
sterilization), and traditional methods (periodic abstinence and withdrawal) [8].
160

161 **Statistical analysis**
162

All statistical analyses were performed using STATA version 17 (Stata Corp 2021,
College Station, TX) [24] and accounted for the CDHS sampling weights and complex
survey design. Socio-demographic characteristics and behavioral factors were
described in weighted frequency and percentage. The provincial variation in the
prevalence of overweight and/or obesity was done using ArcGIS software version 10.8
[24]. Shapefiles for administrative boundaries in Cambodia are publicly accessible
through the DHS website (URL: https://spatialdata.dhsprogram.com/covariates).
Descriptive analyses that coupled cross-tabular frequency distributions with chi-square
tests were used to assess associations between explanatory variables and the outcome
variable Overweight/Obese. Variables associated with the outcome variable with a
significance level of p-value ≤ 0.10 or that had theoretical justification (for example,
women's age, wealth index, alcohol consumption, residence, and geographic of region)
were included in the unadjusted and adjusted logistic regression analyses [3].
Unadjusted logistic regression was used to determine the magnitude effect of
associations between overweight and obesity with socio-demographic characteristics
and behavioral factors reported as odds ratios (OR) with 95% confidence intervals (CI).
Then, adjusted logistics regression was used to assess independent associations,
reported as adjusted odds ratios (AOR), with overweight and obesity after adjusting for
other independent variables included in the model. Multi-collinearity between
independent variables was checked before fitting the final regression model
(supplementary table S1).
184

185 **Results**
186

Characteristics of the study samples

The mean age of women was 31 years old (SD = 9.6 years) in which age group 30-49 years old accounted for 57.11%. Almost 67.82% were married; 49.47% had completed at least secondary education and 11.90% did not receive formal schooling. A total of 30.30% of women did professional work, 25.86% were unemployed. Of the total sample, 35.97% women were from the poor households. Over half (57.27%) of women resided in rural areas. About 27.79% had three and more children while 29.49% had no children. Only 1.40% of women reported cigarette smoking, 16.60% reported currently drink alcohols and 18.60 reported ever drink, 12.35% reported using hormonal contraceptives. Mean women's BMI was 22.9 kg/m^2 (SD = 3.9 kg/m^2) and 22,56% and 5.61% were overweight and obese respectively (**Table 1**).

Table 1. Socio-demographic and behaviors characteristics of the weighted samples of women aged 15-49 years old in Cambodia, 2021-2022 (n□=□9,417, weighted count)

Variables	Freq.	Percent (95% CI)
Women's mean age in years (SD)		**30.9 (9.6)**
15-19	1,488	15.80
20-29	2,551	27.09
30-39	3,220	34.19
40-49	2,158	22.92
Marital status		
Not married	2,429	25.79
Married or living together	6,387	67.82
Widowed/divorced/separated	601	6.38
Education		
No education	1,121	11.90
Primary	3,637	38.62
Secondary and higher	4,659	49.47
Current work status		
Not working	2,435	25.86
Agricultural	1,607	17.06
Professional	2,853	30.30
Manual labor and unskilled	2,331	24.75
Number of children born		

No birth	2,777	29.49
1-2 child	4,023	42.72
Three and above	2,617	27.79
Household wealth index		
Poor	3,387	35.97
Middle	1,822	19.35
Rich	4,207	44.67
Smoking		
Non smoker	9,285	98.60
Smoker	132	1.40
Current drinking alcohol		
Never drink	6103	64.80
Ever drink	1751	18.60
Current drink	1562	16.60
Frequency of watching television		
Not at all	5,808	61.68
Less than once a week	1,493	15.85
At least once a week	2,116	22.47
Ever report of contraceptive use		
No method	5,152	54.71
Traditional method	2,398	25.46
Non-hormonal method	704	7.48
Hormonal method	1,163	12.35
Place of residence		
Urban	4,024	42.73
Rural	5,393	57.27
Region		
Plain	4,708	49.99
Tonle Sap	2,859	30.36
Coastal	595	6.32
Plateau/Mountain	1,255	13.33
BMI mean in kg/m2 (SD)		22.9(3.9)
Underweight	1013	10.76 (9.9-11.7)
Normal weight	5751	61.08 (59.7-62.5]
Overweight	2124	22.56 (21.5-23.7)
Obese	528.2	5.61 (5.0-6.3]

202 **Notes: Survey weights applied to obtain weighted percentages. Plains:** Phnom Penh, Kampong

203 Cham, Tbong Khmum, Kandal, Prey Veng, Svay Rieng, and Takeo; **Tonle Sap:** Banteay Meanchey,

204 Kampong Chhnang, Kampong Thom, Pursat, Siem Reap, Battambang, Pailin, and Otdar Meanchey;

Coastal/sea: Kampot, Kep, Preah Sihanouk, and Koh Kong; **Mountains:** Kampong Speu, Kratie, Preah Vihear, Stung Treng, Mondul Kiri, and Ratanak Kiri.

Prevalence of overweight and obesity among WRA by provinces

The prevalence of overweight and obesity is highest among WRA in Kampong Cham (34.1%), Kandal (32.3%), Svay Rieng (32%), Preah Sihanouk (31.2%), Phnom Penh, Kompong Thom, Pailin, and Tboung Khmum (30% each) and lowest among WRA in Ratanak Kiri (9.4%) and Kampong Chhnang (14.6%) (**Fig 1,** supplementary table S2).

Fig 1. Prevalence of overweight and obesity in Women Reproductive Age by province. Map created using ArcGIS software version 10.8 [24]. Shapefiles for administrative boundaries in Cambodia are publicly accessible through DHS website (URL: https://spatialdata.dhsprogram.com/covariates).

Factors associated with overweight and obesity in bivariate analysis

Socio-demographic characteristics and behavioral factors were significantly associated with overweight and obesity (**Table 2**). Women had higher prevalence of overweight/obesity if they were aged 40-49 years (47.31%) compared to younger age groups, (p value <0.001), married (35.87%) compared to other relationship status (p value <0.001), had no education (37.91%) compared to higher education (p value <0.001), were employed in a professional job (34.03%) compared to other jobs, or were from the rich households (30.90%) compared to poor households (p value <0.001). In addition, Overweight/obesity increased with parity. Women reported at least three children had a significantly highest overweight and obesity (42.53%), (p value < 0.001). Women who current drink alcohol had a significant higher proportion of overweight and obesity (33.87%) than women did not (p value <0.001). Women used non-hormonal

contraceptives methods had a significant higher proportion of overweight and obesity (37.92%) than women using other methods (p value <0.019). Geographic regions of residence were likewise associated with a woman having overweight and obesity. Women who lived in urban areas had a higher prevalence of overweight and obesity than those living in rural areas (29.90% vs 26.87%, p value = 0.028). Plain regions were positively associated with overweight and obesity (29.91%) compared other regions (p value = 0.006).

Table 2. Factors associated with overweight and obesity among women aged 15-49 years in Chi^2 analysis (n □=□ 9,417, weighted count)

Characteristics	Overweight/Obese (n=2,652)		Normal/Underweight (n= 6,764)		P value
	Freq.	**%**	**Freq.**	**%**	
Women's age group in years					
15-19	80	5.38	1,408	94.62	**<0.001**
20-29	444	17.40	2,107	82.60	
30-39	1,108	34.41	2,112	65.59	
40-49	1,021	**47.31**	1,137	52.69	
Marital status					
Not married	181	7.45	2,248	92.55	**<0.001**
Married or living together	2,291	**35.87**	4,096	64.13	
Widowed/divorced/separated	181	30.12	420	69.88	
Education					
No education	425	**37.91**	695	62.00	**<0.001**
Primary	1,254	34.48	2,383	65.52	
Secondary and higher	973	20.88	3,686	79.12	
Current work status					
Not working	586	24.07	1,849	75.93	**<0.001**
Agricultural	485	30.18	1,121	69.76	
Professional	971	**34.03**	1,881	65.93	
Manual labor and unskilled	563	24.15	1,768	75.85	
Number of children born					
No birth	264	9.51	2,512	90.46	**<0.001**
1-2 child	1,275	31.69	2,748	68.31	
Three and above	1,113	**42.53**	1,504	57.47	
Household wealth index					
Poor	841	24.83	2,546	75.17	**<0.001**
Middle	512	28.10	1,311	71.95	
Rich	1,300	**30.90**	2,907	69.10	
Smoking					
Non smoker	2,621	28.23	6,664	71.77	0.325

Smoker	32	24.24	100	75.76	
Current drinking alcohol					
Never drink	1,573	25.77	4,530	74.23	**<0.001**
Ever drink	550	31.41	1,201	68.59	
Current drink	529	**33.87**	1,033	66.13	
Frequency of watching television					
Not at all	1,623	27.94	4,186	72.07	0.901
Less than once a week	423	28.33	1,069	71.60	
At least once a week	606	**28.64**	1,509	71.31	
Ever report of contraceptive use					
No method	1,132	21.97	4,020	78.03	**<0.001**
Traditional method	821	34.24	1,577	65.76	
Non-hormonal method	259	36.79	446	63.35	
Hormonal method	441	**37.92**	722	62.08	
Place of residence					
urban	1,203	**29.90**	2,821	70.10	**0.028**
rural	1,449	26.87	3,944	73.13	
Region					
Plain	1,408	**29.91**	3,299	70.07	**0.006**
Tonle Sap	768	26.86	2,092	73.17	
Coastal	171	28.74	425	71.43	
Plateau/Mountain	306	24.38	949	75.62	

Factors associated with overweight and obesity in adjusted logistic regression

As shown in **Table 3**, several factors were independently associated with increased odds of having overweight and obesity among women. These factors included age group 20-29 years [AOR=1.85; 95% CI: 1.22 - 2.80], 30-39 years [AOR=3.34; 95% CI: 2.21 - 5.04] and 40-49 years [AOR=5.57; 95% CI: 3.76 - 8.25] married [AOR=2.49; 95% CI: 1.71 - 3.62] and widowed/divorced/separated [AOR=1.14; 95% CI: 1.14 - 2.63], middle wealth quintile [AOR=1.21; 95% CI : 1.02 - 1.44], and rich wealth quintile [AOR=1.44 ; 95% CI: 1.19 - 1.73], having at least three children or more [AOR=1.40 ; 95% CI: 1.00 - 1.95], ever drink alcohol [AOR=1.24 ; 95% CI: 1.04 - 1.47], and current drink alcohol [AOR=1.21 ; 95% CI: 1.01 - 1.45]. On the contrary, following factors were independently associated with decreased odds of having overweight and obese: women

with at least secondary education [AOR=0.73; 95% CI: 0.58-0.91], working in in manual

labor jobs [AOR=0.76; 95% CI: (0.64 - 0.90].

Table 3. Risk factors associated with overweight and obesity in unadjusted and adjusted logistic regression analysis

Characteristics	Total (n=9,417)		Total (n= 9,225)	
	OR	95%CI	AOR	95%CI
Women's age group in years				
15-19	Ref.		Ref.	
20-29	3.71***	(2.61 - 5.27)	**1.85***	**(1.22 - 2.80)**
30-39	9.25***	(6.70 - 12.77)	**3.34***	**(2.21 - 5.04)**
40-49	15.84***	(11.68 - 21.47)	**5.57***	**(3.76 - 8.25)**
Marital status				
Not married	Ref.		Ref.	
Married or living together	6.96***	(5.67 - 8.54)	**2.49***	**(1.71 - 3.62)**
Widowed/divorced/separated	5.34***	(3.81 - 7.50)	**1.73**	**(1.14 - 2.63)**
Education				
No education	Ref.		Ref.	
Primary	0.86	(0.72 - 1.03)	0.97	(0.80 - 1.17)
Secondary and higher	0.43***	(0.35 - 0.53)	**0.73***	**(0.58 - 0.91)**
Current work status				
Not working	Ref.		Ref.	
Agricultural	1.37***	(1.14 - 1.64)	0.85	(0.69 - 1.03)
Professional	1.63***	(1.40 - 1.89)	1.10	(0.93 - 1.31)
Manual labor and unskilled	1.00	(0.85 - 1.19)	**0.76***	**(0.64 - 0.90)**
Number of children born				
No birth	Ref.		Ref.	
1-2 child	4.41***	(3.61 - 5.38)	1.23	(0.89 - 1.68)
Three and above	7.03***	(5.79 - 8.55)	**1.40**	**(1.00 - 1.95)**
Household wealth index				
Poor	Ref.		Ref.	
Middle	1.18**	(1.02 - 1.37)	**1.21**	**(1.02 - 1.44)**
Rich	1.35***	(1.18 - 1.56)	**1.44***	**(1.19 - 1.73)**
Smoking				

	OR	95% CI	AOR	95% CI
Non smoker	Ref.		-	
Smoker	0.81	(0.52 - 1.24)	-	
Current drinking alcohol				
Never drink	Ref.		Ref.	
Ever drink	1.32***	(1.13 - 1.54)	**1.24****	**(1.04 - 1.47)**
Current drink	1.47***	(1.23 - 1.76)	**1.21****	**(1.01 - 1.45)**
Watching television				
Not at all	Ref.			
Less than once a week	1.02	(0.85 - 1.23)	-	-
At least once a week	1.04	(0.89 - 1.21)	-	-
Ever report of contraceptive use				
No method	Ref.		Ref.	
Traditional method	1.85***	(1.61 - 2.12)	1.02	(0.86 - 1.21)
Non-hormonal method	2.06***	(1.64 - 2.60)	0.84	(0.65 - 1.10)
Hormonal method	2.17***	(1.75 - 2.69)	1.01	(0.81 - 1.26)
Place of residence				
Rural	Ref.		Ref.	
Urban	1.16**	(1.02 - 1.33)	1.11	(0.94 - 1.31)
Region				
Plain	Ref.		Ref.	
Tonle Sap	1.33***	(1.11 - 1.58)	1.17	(0.95 - 1.43)
Coastal	1.14	(0.95 - 1.36)	1.11	(0.91 - 1.35)
Plateau/Mountain	1.25**	(1.00 - 1.55)	1.10	(0.86 - 1.41)

*** p<0.01, ** p<0.05, * p<0.10, OR= unadjusted odds ratio, AOR= adjusted odds ratio

Discussion

The overall prevalence of overweight and obesity among WRA were 22,56% (95% CI: 21.5-23.7) and 5.61% (95% CI: 5.0-6.3) were overweight and obese respectively. Factors associated with being overweight or obese for non-pregnant women include older age, increased parity, marriage and ever married, middle and rich household

wealth index, and alcohol consumption. Women with higher education as well as those
working in manual labor were less likely to be overweight or obese.

Being older aged women were more likely overweight or obese as compared to younger
women aged 15-19 years. This is consistent with other studies that showed obesity was
more prevalent in older WRA [3,25]. The risk of women becoming overweight or obese
rises with age, possibly due to unhealthy food consumption and/or a lack of physical
activity [26]. Women with better socioeconomic status and who live in urban areas had
higher odds of being overweight and obese, which is consistent with prior research
[27,28]. Women with higher socioeconomic status tend to use more improved
technologies for a more comfortable lifestyle [19,29]. Women have at least three and
more children at risk of being overweight and obese, similar other studies [3,30]. During
pregnancy, factors such as stress, depression, and/or anxiety may play a role in
hypothalamic-pituitary-adrenal hyperactivity [31,32]. Women with several children may
also have gained weight as a result of their reduced physical activity and have less time
to focus on health behaviors including weight management. In addition, women ever
and current consuming alcohol were significantly associated with higher risks of
overweight and obesity. The association between alcohol consumption status and
overweight and obesity has been studied thoroughly in various populations such as in
Urban Cambodia and in Hawassa city, southern Ethiopia [20,21]. Commonly, alcohol
consumption is considered to increase appetite, resulting in excessive energy intake
and leading to overweight and obesity [24]. In contrast, women with higher education
are less likely to be overweight or obesity than women with no school due to increased
knowledge and awareness; for example, studies in Cambodia, Nigeria and South Korea
reported that educated women had a lower risk of being overweight or obese and that
this may be linked to education influencing healthy behavior [3,33-36]. Education is a
critical predictor of women's healthy behaviors and health outcomes, including diet,
physical activity [37]. Although overweight and obesity was not significantly associated
with geographical regions of residence, those from Plain and Coastal region had a high
rate of overweight and obesity, 29.91% and 28.74%, respectively. Also, women who
lived in urban areas had a higher prevalence of overweight and obesity 29.90%. Prior

research on overweight and obesity among women reproductive age in Cambodia
found women resided in urban had significant associated with overweight and obesity
[3]; our findings suggest a need for more regionally representative studies.
310

311 **Conclusion**
312

The overall prevalence of overweight and obesity was very high among non-pregnant in
Cambodia. Women being older age, higher education levels, higher income and wealth,
and behavioral factors such as alcohol consumption, type of employment as risk factors
associated with overweight and obesity among women of reproductive age in Cambodia.
It is crucial to design intervention programs that target these socio-demographic factors
and to raise awareness on the importance of consuming healthy food as well as the
benefits of regular physical activity, especially among older women and those with
professional jobs.
321

323 **limitations and strengths of this study**
324

Our study has several limitations. First, because CDHS 2021-2022 collects cross-
sectional data, our analysis could not explore changes over time. Second, data on
several important variables, such as women's food-consumption behaviors, were not
collected by CDHS, and CDHS did not objectively measure physical activity; therefore,
our ability to examine the association of these variables with overweight and obesity
was restricted. Third, numerous psychological factors (e.g., depressive and anxiety
disorders) and physiological factors may also be associated with obesity, but these
factors were not included in this study because they were not available in the data set.
Finally, because the data were available only for women aged 15 to 49, our results may
not be generalizable to girls younger than 15 and women older than 49. Despite these
limitations, our results contribute to the literature on the prevalence and association
between sociodemographic and behavioral variables and the status of overweight and
obesity among women reproductive age in Cambodia. The major strength of this study

is that it is the first to use nationally representative data with a high response rate of 97%

to examine the prevalence and predictors of overweight and obesity in this country.

Data were collected using validated survey methods including calibrated measurement

tools and highly trained data collectors, which contribute to improved data quality [38].

Formally incorporating the complex survey design and sampling weights into the

analysis bolsters the rigor of the analysis and enables generalizing our findings to the

population of non-pregnant women in Cambodia.

Abbreviations

BMI, Body mass index; NCDs, Non-communicable diseases; WHO, World Health

Organization; CDHS, Cambodia Demographic Health Survey; WRA, Women

reproductive age; AOR, Adjusted odds ratio; PPS, Probability proportional to size. EA,

Enumeration areas

Data Availability

Our study used 2021-2022 Cambodia Demographic and Health Survey (CDHS)

datasets. The DHS data are publicly available from the website at

(URL:https://www.dhsprogram.com/data/available-datasets.cfm).

The Shapefiles for administrative boundaries in Cambodia are publicly accessible

through DHS website at (URL: https://spatialdata.dhsprogram.com/covariates).

Ethical Approval

The data used in this study were extracted from CDHS 2021-22 data [8], which are

publicly available with all personal identifiers of study participants removed. Permission

to analyze the data was granted through registering with the DHS program website and

submitting an application outlining the intended use of the datasets [8]. Informed

consent was obtained from the study participants before data collection. The data

collection tools and procedures for CHDS 2021-22 was approved by the Cambodia

National Ethics Committee for Health Research and the Institutional Review Board (IRB)

of ICF in Rockville, Maryland, USA.

Funding Statement

The authors received no specific funding for this work.

Consent

Consent for publication is not necessary because this manuscript did not contain any

personal details such as photos, images, videos, or quote data.

Competing interests

The authors have declared that no competing interests exist.

Authors' Contributions

Samnang Um. contributed to conceptualization, methods, performed the data analysis, interpreted the data, writing the original draft, and reviewing/editing the manuscript. **Yom An.** contributed to technical methods evaluation, support data analysis, and reviewing/editing the manuscript. Finally, all authors read and approved the final manuscript.

Acknowledgments

The authors would like to thank DHS-ICF, who approved the data used for this paper. We would like to acknowledge management team of National Institute of Public Health, Phnom Penh, Cambodia namely Professor. **Chhea Chhorvann**. Professor. **Heng Sopheab** for their continued support and encouragement.

References

1. **World Health Organization, Fact sheets, Obesity and overweight, Available from [https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight], Accessed on April 10, 2023**.
2. **Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19·2 million participants**. *Lancet (London, England)* 2016, **387**(10026):1377-1396.
3. Um S, Yom A, J AM, Sopheab H: **Overweight and obesity among women at reproductive age in Cambodia: Data analysis of Cambodia Demographic and Health Survey 2014**. *PLOS global public health* 2023, **3**(3):e0001768.
4. Yu Z, Han S, Zhu J, Sun X, Ji C, Guo X: **Pre-pregnancy body mass index in relation to infant birth weight and offspring overweight/obesity: a systematic review and meta-analysis**. *PLoS One* 2013, **8**(4):e61627.
5. Stanaway JD, Afshin A, Gakidou E, Lim SS, Abate D, Abate KH, Abbafati C, Abbasi N, Abbastabar H, Abd-Allah F: **Global, regional, and national**

comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *The Lancet* 2018, **392**(10159):1923-1994.

6. Report GN. Country Nutrition Profiles, Available from; [https://globalnutritionreport.org/resources/nutrition-profiles/asia/south-eastern-asia/cambodia/]; Accessed on April 10, 2023.

7. National Institute of Statistics (NIS). (2020). General population census of the Kingdom of Cambodia 2019; accessed on November 4th 2021, availbale from: [https://www.nis.gov.kh/nis/Census2019/Final%20General%20Population%20Census%202019-English.pdf].

8. National Institute of Statistics (NIS) [Cambodia], Ministry of Health (MoH) [Cambodia], and ICF. 2023. Cambodia Demographic and Health Survey 2021–22 Final Report. Phnom Penh, Cambodia, and Rockville, Maryland, USA: NIS, MoH, and ICF.

9. World Food Programme in Cambodia; [https://www.wfp.org/countries/cambodia]: Accessed on April 10, 2023.

10. Balarajan Y, Villamor E: **Nationally representative surveys show recent increases in the prevalence of overweight and obesity among women of reproductive age in Bangladesh, Nepal, and India**. *The journal of nutrition* 2009, **139**(11):2139-2144.

11. Abrha S, Shiferaw S, Ahmed KY: **Overweight and obesity and its socio-demographic correlates among urban Ethiopian women: evidence from the 2011 EDHS**. *BMC Public Health* 2016, **16**:1-7.

12. Yeshaw Y, Kebede SA, Liyew AM, Tesema GA, Agegnehu CD, Teshale AB, Alem AZ: **Determinants of overweight/obesity among reproductive age group women in Ethiopia: multilevel analysis of Ethiopian demographic and health survey**. *BMJ open* 2020, **10**(3):e034963.

13. Oddo VM, Bleich SN, Pollack KM, Surkan PJ, Mueller NT, Jones-Smith JC: **The weight of work: the association between maternal employment and overweight in low-and middle-income countries**. *International Journal of Behavioral Nutrition and Physical Activity* 2017, **14**(1):1-10.

14. Lopez LM, Bernholc A, Chen M, Grey TW, Otterness C, Westhoff C, Edelman A, Helmerhorst FM: **Hormonal contraceptives for contraception in overweight or obese women**. *Cochrane Database of Systematic Reviews* 2016(8).

15. Aryeetey R, Lartey A, Marquis GS, Nti H, Colecraft E, Brown P: **Prevalence and predictors of overweight and obesity among school-aged children in urban Ghana**. *BMC obesity* 2017, **4**:1-8.

16. de Munter Js, Tynelius P, Magnusson C, Rasmussen F: **Longitudinal analysis of lifestyle habits in relation to body mass index, onset of overweight and obesity: results from a large population-based cohort in Sweden**. *Scandinavian journal of public health* 2015, **43**(3):236-245.

17. Piirtola M, Jelenkovic A, Latvala A, Sund R, Honda C, Inui F, Watanabe M, Tomizawa R, Iwatani Y, Ordonana JR: **Association of current and former**

smoking with body mass index: A study of smoking discordant twin pairs from 21 twin cohorts. *PloS one* 2018, **13**(7):e0200140.

18. Chiolero A, Jacot-Sadowski I, Faeh D, Paccaud F, Cornuz J: **Association of cigarettes smoked daily with obesity in a general adult population**. *Obesity* 2007, **15**(5):1311-1318.

19. Ghose B: **Frequency of TV viewing and prevalence of overweight and obesity among adult women in Bangladesh: a cross-sectional study**. *BMJ open* 2017, **7**(1):e014399.

20. Dagne S, Gelaw YA, Abebe Z, Wassie MM: **Factors associated with overweight and obesity among adults in northeast Ethiopia: a cross-sectional study**. *Diabetes, metabolic syndrome and obesity: targets and therapy* 2019:391-399.

21. Tamaoki M, Honda I, Nakanishi K, Nakajima M, Cheam S, Okawada M, Sakakibara H: **Lifestyle Factors Associated with Metabolic Syndrome in Urban Cambodia**. *International Journal of Environmental Research and Public Health* 2022, **19**(17):10481.

22. Darebo T, Mesfin A, Gebremedhin S: **Prevalence and factors associated with overweight and obesity among adults in Hawassa city, southern Ethiopia: a community based cross-sectional study**. *BMC obesity* 2019, **6**:1-10.

23. **World Health Organization, Fact sheets, A healthy lifestyle - WHO recommendations. Available from [https://www.who.int/europe/news-room/fact-sheets/item/a-healthy-lifestyle---who-recommendations], Accessed on April 10, 2023**.

24. Yeomans MR: **Alcohol, appetite and energy balance: is alcohol intake a risk factor for obesity?** *Physiology & behavior* 2010, **100**(1):82-89.

25. Biswas T, Uddin MJ, Mamun AA, Pervin S, P Garnett S: **Increasing prevalence of overweight and obesity in Bangladeshi women of reproductive age: Findings from 2004 to 2014**. *PloS one* 2017, **12**(7):e0181080.

26. Alemu E, Atnafu A, Yitayal M, Yimam K: **Prevalence of overweight and/or obesity and associated factors among high school adolescents in Arada Sub city, Addis Ababa, Ethiopia**. *Journal of nutrition & food sciences* 2014, **4**(2):1.

27. Mangemba NT, San Sebastian M: **Societal risk factors for overweight and obesity in women in Zimbabwe: a cross-sectional study**. *BMC public health* 2020, **20**(1):1-8.

28. Kamal SM, Hassan CH, Alam GM: **Dual burden of underweight and overweight among women in Bangladesh: patterns, prevalence, and sociodemographic correlates**. *Journal of health, population, and nutrition* 2015, **33**(1):92.

29. Islam M, Sathi NJ, Abdullah HM, Naime J, Butt ZA: **Factors Affecting the Utilization of Antenatal Care Services During Pregnancy in Bangladesh and 28 Other Low-and Middle-income Countries: A Meta-analysis of Demographic and Health Survey Data**. *Dr Sulaiman Al Habib Medical Journal* 2022, **4**(1):19-31.

30.   Mbochi RW, Kuria E, Kimiywe J, Ochola S, Steyn NP: **Predictors of overweight**
and obesity in adult women in Nairobi Province, Kenya. *BMC public health*
2012, **12**(1):1-9.
31.   Li W, Wang Y, Shen L, Song L, Li H, Liu B, Yuan J, Wang Y: **Association**
between parity and obesity patterns in a middle-aged and older Chinese
population: a cross-sectional analysis in the Tongji-Dongfeng cohort study.
Nutrition & metabolism 2016, **13**(1):1-8.
32.   Pasquali R, Vicennati V, Cacciari M, Pagotto U: **The**
hypothalamic-pituitary-adrenal axis activity in obesity and the metabolic
syndrome. *Annals of the New York Academy of Sciences* 2006, **1083**(1):111-128.
33.   Hamad R, Elser H, Tran DC, Rehkopf DH, Goodman SN: **How and why studies**
disagree about the effects of education on health: A systematic review and
meta-analysis of studies of compulsory schooling laws. *Social Science &*
Medicine 2018, **212**:168-178.
34.   Chung A, Backholer K, Wong E, Palermo C, Keating C, Peeters A: **Trends in**
child and adolescent obesity prevalence in economically advanced
countries according to socioeconomic position: a systematic review.
Obesity reviews 2016, **17**(3):276-295.
35.   Barlow P: **The effect of schooling on women's overweight and obesity: A**
natural experiment in Nigeria. *Demography* 2021, **58**(2):685-710.
36.   McLaren L: **Socioeconomic status and obesity**. *Epidemiologic reviews* 2007,
29(1):29-48.
37.   Liu H, Guo G: **Lifetime socioeconomic status, historical context, and genetic**
inheritance in shaping body mass in middle and late adulthood. *American*
sociological review 2015, **80**(4):705-737.
38.   BOWLER P: **CUNNINGHAM, A. and JARDINE, N.(editors). Romanticism and**
the sciences. Cambridge University Press, Cambridge: 1990. Pp xxii, 345;
illustrated. Price:£ 15, US $19.95. ISBN: 0-521-35685-7 (paperback). In.; 1992.

Supplemental materials

Table S1. Results of checking multicollinearity using Variance Inflation Factor (VIF)

Table S2. Prevalence of overweight and/or obesity among women reproductive age, CDHS

2021-2022 **(n=9,417)**

Percent	
	9 - 15
	16 - 25
	26 - 30
	31 - 35