Combined assessment of *KRAS* mutational status and tumor size has no impact on prognosis in early-stage non-small cell lung cancer

Ella A. Eklund[1,2,3], Ali Mourad[1,2], Clotilde Wiel[1], Sama I. Sayin[3], Henrik Fagman[4,5], Andreas Hallqvist [3,7], Volkan I. Sayin[1,2]

[1]Department of Surgery, Institute of Clinical Sciences, Sahlgrenska Center for Cancer Research, University of Gothenburg, Gothenburg, Sweden
[2]Wallenberg Centre for Molecular and Translational Medicine, University of Gothenburg, Gothenburg, Sweden
[3]Department of Oncology, Sahlgrenska University Hospital, Gothenburg, Sweden;
[4]Department of Laboratory Medicine, Institute of Biomedicine, University of Gothenburg, Gothenburg, Sweden
[5]Department of Clinical Pathology, Sahlgrenska University Hospital, Gothenburg, Sweden
[6]Department of Microbiology and Immunology, Institute for Biomedicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden
[7]Department of Oncology, Institute of Clinical Sciences, University of Gothenburg, Gothenburg, Sweden

Correspondence to:
volkan.sayin@gu.se

Keywords: lung cancer, *KRAS*, tumor size, stage I and II, clinical outcome
Abbreviations:[1]
Word count: 3024
Number of main figures: 4
Number of tables: 2

---

[1] CT: Computed Tomography; ECOG: Eastern Cooperative Oncology Group; HR: Hazard Ratio; NSCLC: Non-Small Cell Lung Cancer; NGS: Next Generation Sequencing; PS: Performance Status; OS: Overall Survival;

Abstract

Background: *KRAS* mutation status, stage and tumor size at the time of diagnosis are well-established independent prognostic factors in non-small cell lung cancer (NSCLC). Here, we investigate the prognostic value of combining survival data on *KRAS* mutation status and tumor size in early-stage NSCLC.

Methods: We studied the combined impact of *KRAS* mutational status and tumor size on overall survival (OS) and risk of death in patients with stage I-II NSCLC. We performed a retrospective study including 310 consecutively diagnosed patients with early (stage I-II) NSCLCs. All consecutive patients molecularly assessed and diagnosed between 2016-2018 with stage I-II NSCLC in the Västra Götaland region of western Sweden were included in this multi-center retrospective study. The primary study outcome was OS and risk of death (hazard ratio).

Results: Out of 310 patients with stage I-II NSCLC, 37% harbored an activating mutation in the *KRAS* gene. Our study confirmed staging and tumor size as prognostic factors. However, *KRAS* mutational status was not found to impact OS and there was no difference in the risk of death when combining *KRAS* mutational status and primary tumor size.

Conclusions: In our patient cohort, *KRAS* mutations in combination with primary tumor size are not associated with a worse prognosis in stage I-II NSCLC.

Introduction

Non-small cell lung cancer (NSCLC) is the second most common cancer worldwide with 2.1 million new cases annually and the highest mortality rate with 1.8 million deaths. [1] Staging is a crucial aspect of NSCLC management, as it is one of the most important predictors of survival. The TNM staging system describes key tumor characteristics such as size, location, and whether the disease has spread to lymph nodes and/or distant organs [2-5]. There are four main stages in NSCLC (stage I-IV), with stage IV having the worst prognosis. Pathological stage is considered the most important prognostic factor for resected patients, with 5-year survival rates, gradually decreasing across stages, of 83% for stage IA, 71% for IB, 57% for IIA, 49% for IIB, 36% for IIIA, and 23% for IIIB [4].

The most frequent oncogenic driver in NSCLC is the Kirsten rat sarcoma viral oncogene (*KRAS*), which is present in up to 40% of all cases, with the most common mutations being *G12C, G12V,* and *G12D* [6]. *KRAS* mutations are associated with worse outcomes after chemotherapy and radiotherapy, with shorter OS in stage III and IV patients [7-14]. In early-stage NSCLC, however, while several studies have shown that *KRAS* mutations negatively influence the prognosis [15-17], others have shown no significant effect [18]. Most recently, it was reported that *KRAS G12C* mutation (but not other *KRAS* mutations or with no mutation in *KRAS*) significantly increased risk of disease recurrence in stage I surgically resected lung adenocarcinomas [19]. However, while the study found this in two distinct local cohorts of IRE-LUAD (Rome, Italy) and MSK-LUAD (New York, USA), data extracted from The Cancer Genome Atlas (TCGA) showed no significant difference. Hence, the debate about the prognostic value of *KRAS* mutational status in early NSCLC is ongoing [20, 21]. In fact, given the lack of consensus regarding its effects on prognosis, testing for *KRAS* mutations for resectable stage I and II tumors is currently not recommended in clinical guidelines [22]. In addition, several inhibitors that specifically bind *KRAS-G12C* have been investigated in clinical trials, with sotorasib becoming the first treatment to gain approval for adults with stage IV NSCLC harboring a *KRAS-G12C* mutation as second-line therapy [23-27]. However, treatment with sotorasib is not currently recommended for patients with early-stage NSCLC due to lack of evidence showing positive outcomes of treatment in this group.

Therefore, further investigations are warranted to identify potential subgroups in Stage I-III disease who may still have to gain from effective and well-established treatments, and to add to the pool of clinical data required to study this further. One strategy is to stratify patients according to *KRAS* mutational status together with other key prognostic factors, such as tumor size. Primary tumor size is an established prognostic factor in NSCLC, with larger tumors being associated with poorer survival [21, 28-31]. The reason for this association is not yet fully understood but larger tumors may be more resistant to therapy due to having poorer blood supply, differential metabolism, and potentially a higher likelihood of micrometastatic disease compared to smaller tumors [32-35]. Further research is needed to elucidate the underlying mechanisms. However, when considering primary tumor size, the grouping as early (I-II), advanced (III), and metastasized (IV) NSCLC can be argued to be more clinically relevant due to that stage I-II is primarily based on tumor size whereas a spread to the lymph nodes, a negative prognostic factor, is more common in stage III [3, 21].

To our knowledge, no one has investigated the combined impact of primary tumor size and *KRAS* mutational status on OS and risk of death in stage I-II NSCLC. However, in Sweden, reflex testing for targetable alterations in NSCLC, including *KRAS* mutational status, has been widely implemented since 2015 for all stages. By including all consecutive patients diagnosed with stage I-II NSCLC and molecularly assessed between 2016-2018 in Västra Götaland, the second largest county in Sweden with a population of 1.7 million, the current retrospective cohort study provides a unique real-world dataset for assessing the impact of combining *KRAS* mutations with primary tumor size.

To summarize, primary tumor size is a key determinant of prognosis especially in the early stages of NSCLC. At the same time, the prognostic value of *KRAS* mutational status in early disease stages remains unclear. Hence patients diagnosed at an early stage are not

113   automatically tested for *KRAS* mutations and recommended treatment with *KRAS*-targeted
114   therapy. Here, we investigate whether there is prognostic value in combining *KRAS* mutational
115   status with tumor size to aid in clinical stratification of potentially treatment-responsive
116   subgroups in early-stage NSCLC.
117
118
119   **Materials and Methods**
120
Patient population
We conducted a multi-center retrospective study including all consecutive NSCLC patients
diagnosed with stage I-II NSCLC and molecular assessment performed between 2016-2018
in Västra Götaland, Sweden (*n* = 310). Further inclusion criteria included the availability of
tumor size from CT scanning or a pathology report as well as follow-up data. Patients were
excluded if diagnosed before 2016, had no digitally accessible patient charts, no tumor
measurements noted in the patient charts, or had recurrent disease.
128
Patient demographics (age, sex, Eastern Cooperative Oncology Group [ECOG] performance
status [PS], and smoking history), cancer stage, pathological details (histology, mutational
status including *KRAS* mutational status and subtype), first-line treatment and outcome data
were retrospectively collected from patient charts and the Swedish Lung Cancer Registry.
Clinical staging was based on TNM staging guidelines 7th edition [4]. TNM staging 8th edition
released in 2017 was introduced in Swedish guidelines in 2018, and full implementation was
reached in 2019. Ethical approval was obtained from the Swedish Ethical Review Authority
prior to study commencement (Dnr 2019-04771 and 2021-04987). No informed consent was
required due to all data presented in a de-identified form according to the Swedish Ethical
Review Authority.
139
Mutational status
Patients were assessed with next-generation sequencing (NGS) for mutational status on DNA
from formalin-fixed paraffin-embedded (FFPE) blocks or cytological smears using the Ion
AmpliSeq™ Colon and Lung Cancer Panel v2 from Thermo Fisher Scientific as part of the
diagnostic workup process at the Department of Clinical Pathology at Sahlgrenska University
Hospital, assessing hotspot mutations in *EGFR, BRAF, KRAS,* and *NRAS*. Until June 2017,
ALK-fusions were assessed with immunohistochemistry (IHC), and with fluorescence in situ
hybridization (FISH) if positive or inconclusive IHC. *ROS1* was analyzed upon request with
FISH. Thereafter, *ALK, ROS1,* and *RET* fusions were assessed on RNA using the Oncomine
Solid Tumor Fusion Panel from Thermo Fisher Scientific.
150
Tumor size
To obtain the most recent and accurate untreated primary tumor size, measurements were
collected from the radiology report of computed tomography (CT) performed before a final
diagnosis of NSCLC was established; this is referred as clinical staging. In patients who
underwent surgical resection, the actual primary tumor size was also collected from the
pathology report, also referred as pathological staging (PAD). The largest tumor diameter was
collected and reported in millimeters.
158
Study objectives
The primary outcome of this study was OS and risk of death, defined as the interval between
the date of first treatment and the date of death from any cause. Patients alive or lost to follow-
up at the cut-off date were censored at last contact. Median follow-up time was estimated
using the reverse Kaplan-Meier method. We compared OS and risk of death stratified by
KRAS mutational status, i.e., with no mutation in *KRAS* (wildtype, *KRAS*^{WT}), all *KRAS*
mutations (*KRAS*^{MUT}), *KRAS G12C* mutations (*KRAS*$^{MUT\ G12C}$) and all *KRAS* mutations other
than *G12C* (*KRAS*$^{MUT\ not\ G12C}$).
167
168
169

Statistical analysis
Clinical characteristics were summarized using descriptive statistics and evaluated with univariate analysis in table form. Survival was estimated using the Kaplan-Meier method. The log-rank test was used to assess significant differences in OS between $KRAS^{WT}$ and $KRAS^{MUT}$ groups. To evaluate if there was a significant difference in primary tumor size between $KRAS^{MUT}$ and $KRAS^{WT}$, the Mann-Whitney U test was used. Cox proportional hazard regression was conducted to measure the influence of tumor size on the risk of death (hazard ratio [HR]) stratified by *KRAS* mutational status. We defined an interaction term between tumor size (largest diameter in mm) and *KRAS* mutational status to assess the combined impact on the risk of death (HR). Statistical significance was set at $p < 0.05$ and no adjustments were made for multiple comparisons. Data analysis was conducted using IBM SPSS Statistics version 27 and GraphPad Prism version 9.

Results

Patients and tumor characteristics
A total of 310 consecutive patients, who were diagnosed with stage I-II NSCLC during 2016-2018 in Västra Götaland, Sweden and for whom genetic data was available, were included in this retrospective cohort study (Fig. 1). In the total population, majority of patients were female (187, 60.3%), with a median age of 70 years, and most were current or former smokers (267, 86%) (Table 1). Most patients had good PS with ECOG 0-1 at diagnosis (285, 92%) and the proportion of N1 was low (18, 5.8%). NSCLC was predominantly adenocarcinoma of the lung (281, 90.6%), while squamous cell carcinoma incidence was relatively low (11, 3.5%), which was expected due to the selection of histological type for NGS assessment. Of included patients, over a third (115, 37%) had a *KRAS* mutation (Table 1). This percentage matches what has been previously reported [9], showing good representativeness of the patient group studied here. When comparing the baseline characteristics of $KRAS^{WT}$ with $KRAS^{MUT}$ patients, a greater proportion of those with $KRAS^{MUT}$ were female and current or former smokers. There were no cases of squamous cell carcinoma in the $KRAS^{MUT}$ group. The most common KRAS mutation was G12C (47%). In the total population, majority of patients underwent surgical resection (273, 88%; Table 2). Three patients did not receive any treatment and were excluded from further survival analyses. Median follow-up time was 63 months (95% CI, 59.7-68.3) and the data cut-off date was 31 October 2022.

No significant difference in survival for all patients stratified by *KRAS* mutations
When comparing OS for all (stage I-II) patients stratified by *KRAS* mutational status, no significant difference was detected with a mean OS (median not reached) of 74 months for $KRAS^{WT}$ vs 63 months for $KRAS^{MUT}$ ($p = 0.847$; Fig 2A). Further stratification of the *KRAS* mutated group by the G12C mutation also did not significantly change survival: 74 months for $KRAS^{WT}$, 61 months for $KRAS^{MUT\ not\ G12C}$ and 63 months for $KRAS^{MUT\ G12C}$ ($p = 0.834$; Fig 2D).

No significant difference in survival for patients in Stage I or Stage II disease combined with *KRAS* mutations
There were also no significant differences according to *KRAS* mutational status in stage I (Fig 2B, 2E) or Stage II (Fig 2C, 2F). Similarly in resected patients, no significant difference was observed with a mean OS (median not reached) of 78 months for $KRAS^{WT}$ vs 65 months for $KRAS^{MUT}$ ($p = 0.856$; Supplemental Fig. 1A), or between the subgroups of $KRAS^{MUT}$ ($p = 0.471$; Supplemental Fig.1B).

Next, we stratified by stage and found mean OS (median not reached) of 79 months for stage I vs 50 months for stage II (Fig. S2A). We then conducted the analysis separately according to *KRAS* mutational status. For $KRAS^{WT}$, the mean OS (median not reached) was 78 months for stage I vs 46 months for stage II (Fig. S2B), and for $KRAS^{MUT}$, 65 months for stage I vs 53 months for stage II (Fig. S2C).

227
No significant difference in survival for patients with TNM-stage T1, T2 or T3 disease combined with *KRAS* mutations

Next, we stratified patients using T-staging and studied OS according to *KRAS* mutational status. Among those with T1 disease, $KRAS^{WT}$ had 83 months while $KRAS^{MUT}$ had 66 months OS (p = 0.751; Fig 3A). Further, $KRAS^{MUT\ not\ G12C}$ patients had survival of 70 months and $KRAS^{MUT\ G12C}$ had 61 months (p = 0.344; Fig 3D). In the T2 group, $KRAS^{WT}$ had 53 months while $KRAS^{MUT}$ had 59 months OS (p = 0.495; Fig 3B). $KRAS^{MUT\ not\ G12C}$ patients had survival of 51 months and $KRAS^{MUT\ G12C}$ had 66 months (p = 0.389; Fig 3E). Similarly, in the T3 group, $KRAS^{WT}$ had 47 months while $KRAS^{MUT}$ had 50 months OS (p = 0.966; Fig 3C). $KRAS^{MUT\ not\ G12C}$ patients had survival of 50 months and $KRAS^{MUT\ G12C}$ had 53 months (p = 0.984; Fig 3F).

We further analyzed the impact of T stage on survival and found that it correlated as expected with mean OS of 82 months for T1, 55 months for T2, and 46 months for T3 (p < 0.001; Fig. S3A). The same trend was observed when separately analyzing $KRAS^{WT}$ with a mean OS (median not reached) of 83 months for T1, 53 months for T2, and 45 months for T2 (p < 0.001; Fig. S3B), and $KRAS^{MUT}$ with a mean OS of 65 months for T1, 58 months for T2, and 48 months for T3 (p < 0.023; Fig. S3C).

***KRAS* mutations are associated with smaller tumor size measured from CT scans, but not resection specimens/PAD**

To evaluate differences between primary tumor size from CT scans at diagnosis stratified by *KRAS* mutational status, we used the Mann-Whitney U test. The test revealed that $KRAS^{MUT}$ primary tumors were significantly smaller at diagnosis, with a median size of 20 mm (n = 115) vs $KRAS^{WT}$ primary tumors with a median size of 25 mm (n = 190) (p = 0.043; Fig. 4A). However, when looking at tumor size as assessed in resected specimens, there were no differences; $KRAS^{WT}$ median size 22 mm (n = 171) vs $KRAS^{MUT}$ median size 21 mm (n = 102) (p = 0.16; Fig. 4B).

Larger tumor size measured from resection specimens/PAD, but not CT scans, is associated with a higher risk of death

We found that increase in primary tumor size determined from CT scans did not have a significant effect on risk of death (HR, 1.006; 95% CI, 0.922-1.021; p > 0.5) (Fig 4C). However, when testing the correlation between primary tumor size as assessed in resection specimens, we found a significantly increased risk of death (HR, 1.029; 95% CI, 1.012-1.046; p < 0.001) (Fig 4D). The risk of death increases with 2.9% for every mm increase of size.

The combination of *KRAS* mutational status and tumor size does not impact the risk of death

To test if the combination of tumor size and *KRAS* mutational status impacts the risk of death, we defined an interaction term including both variables. For primary tumor size from CT scans and *KRAS* mutational status, no significant difference in the risk of death was detected (HR, 1.008; 95% CI, 0.988-1.030; p > 0.5) (Fig 4C). Similarly, there were no significant differences for primary tumor size and *KRAS* mutational status when measured in resection specimens (HR, 1.002; 95% CI, 0.978-1.027; p = 0.807) (Fig 4D).

Discussion

In this study, we assessed the prognostic value of combining *KRAS* mutational status with tumor size in early-stage NSCLC. We found that combining these variables had no significant effect on overall survival or the risk of death.

In alignment with previous findings, we found in our patient cohort that later disease stage and larger primary tumor size is associated with worse survival. Interestingly, we found that these

correlations are sustained independent of *KRAS* mutational status. Importantly, the established literature on how KRAS mutations affect outcomes in early-stage NSCLC is varying between worse survival and no significant difference. Here, we find as the latter, that *KRAS* mutational status alone does not significantly impact OS or risk of death in patients with stage I-II NSCLC. Patients harboring a *KRAS* G12C mutation did not have a worse OS. Taken together, these findings show good representativeness of this well-defined patient cohort.

Our study included only patients with stage I-II disease due to the focus on primary tumor size and to limit the prognostic impact of local invasion and regional lymph node involvement. Only 5.8% of the patients had N1 disease that could affect the prognosis. The major portion of the patients had tumor resection and more than 90% of tumors were adenocarcinoma. During this period, patients diagnosed with squamous cell carcinoma were molecularly assessed to a lesser extent, thus our study is more representative of adenocarcinoma. Even though most tumors were classified according the TNM staging guidelines 7[th] edition, changes included in the 8[th] edition, mainly covering substages that were not analyzed in this study, do not alter our findings [4].

When combining tumor size and *KRAS* mutational status there was no increased risk of death. No significant differences were observed when comparing OS for all stage I-II patients stratified by *KRAS* mutational status. However, the mean OS was 11 months shorter for $KRAS^{MUT}$ patients. The same trend was observed when looking at resected patients with a 13-month shorter mean survival for $KRAS^{MUT}$ patients. In this case, our results could differ from former studies due to earlier studies not being corrected for tumor size [15, 16]. Our cohort could be used as a historical reference if, in the future, sotorasib would be approved for adjuvant treatment for patients harboring *KRAS* G12C mutation.

Outcome variables other than survival such as recurrence rates and progression-free survival were not examined here. In addition, there remain confounders that were not include in the analyses such as the effect of different treatment methods on survival. Further, we use the T descriptor of the TNM staging system for tumor size even though the descriptor also includes invasion status and or intrapulmonary metastasis. We found that larger tumor size measured from resection specimens, but not CT scans, is associated with a higher risk of death. However, one confounder here is that non-resected patients are included in the CT group but not in the PAD group, which biases towards worse prognosis.

Going forward, much remains to be explored on the role of *KRAS* mutation in early NSCLC. In the age of precision medicine, our study contributes towards the detailed level clinical data that is required for future pooled analysis of prognosis assessments that can help guide clinical decisions.

In conclusion, we confirm the importance of primary tumor size and stage as a prognostic factor for survival in stage I-II NSCLC. *KRAS* mutations were not found to impact OS and no difference in the risk of death was observed when combining *KRAS* mutations and primary tumor size.

Declarations
Acknowledgments
We thank Sayin lab members and Nesrin Vurgun, Scientific editor at the Institute of Clinical Sciences, University of Gothenburg for a critical review of the manuscript. In addition, we thank members of the Swedish Lung Cancer Registry and the continuous reporting by Swedish healthcare employees.

Authors' contributions
Conceptualization, E.A.E., H.F., A.H. and V.I.S.; Data curation, E.A.E., A.M., H.F.; Formal analysis, E.A.E., A.M. and C.W.; Funding acquisition, E.A.E., C.W., H.F., A.H. and V.I.S.;

341 Methodology, E.A.E.; Resources, H.F.; Supervision, A.H. and V.I.S.; Visualization, E.A.E.,
342 S.I.S., C.W. and V.I.S.; Writing—original draft, E.A.E., S.I.S and V.I.S.; Writing—review &
343 editing, E.A.E., S.I.S, C.W., H.F., A.H. and V.I.S., Project coordination, V.I.S. All authors have
344 read and agreed to the published version of the manuscript.
345

Funding
This work was supported by the Swedish Research Council (2018-02318 and 2022-00971 to
VIS, 2021-03138 to CW), the Swedish Cancer Society (23-3062 to VIS, 22-0612FE to CW),
the Gothenburg Society of Medicine (2019; 19/889991 to EAE), Assar Gabrielsson Research
Foundation (to EAE, CW, and VIS), the Swedish state under the agreement between the
Swedish government and the county councils, the ALF-agreement (to HF), Department of
Oncology, Sahlgrenska University Hospital (to EAE and AH), the Swedish Society for Medical
Research (2018; S18-034 to VIS), the Knut and Alice Wallenberg Foundation, and the
Wallenberg Centre for Molecular and Translational Medicine (to VIS).
355

356 **Declaration of potential conflict of interest**
357 The authors declare no conflicts of interest.
358

359 **Institutional Review Board Statement**
360 Approval from the Swedish Ethical Review Authority (Dnr 2019-04771 and 2021-04987) was
361 obtained prior to the commencement of the study. No informed consent was required due to
362 all data presented in a de-identified form according to the Swedish Ethical Review Authority.
363

365 **Consent for publication**
366 Not applicable. Patient consent statements were not required due to the retrospective nature
367 of this study. No informed consent was required due to all data presented in a de-identified
368 form according to the Swedish Ethical Review Authority.
369

371 **Data Availability Statement**
372 The datasets used and/or analysed during the current study available from the corresponding
373 author on reasonable request.
374

References

1. *World Health Organization, International Agency for Research on Cancer. Globocan 2020: Lung Cancer. International Agency for Research on Cancer. Available at* http://gco.iarc.fr/today/data/factsheets/cancers/15-Lung-fact-sheet.pdf*. Accessed: March 2, 2021.*

2. Amin, M.B., et al., *The Eighth Edition AJCC Cancer Staging Manual: Continuing to build a bridge from a population-based to a more "personalized" approach to cancer staging.* CA Cancer J Clin, 2017. **67**(2): p. 93-99.

3. Detterbeck, F.C., *The eighth edition TNM stage classification for lung cancer: What does it mean on main street?* J Thorac Cardiovasc Surg, 2018. **155**(1): p. 356-359.

4. Goldstraw, P., et al., *The IASLC Lung Cancer Staging Project: Proposals for Revision of the TNM Stage Groupings in the Forthcoming (Eighth) Edition of the TNM Classification for Lung Cancer.* J Thorac Oncol, 2016. **11**(1): p. 39-51.

5. Wankhede, D., *Evaluation of Eighth AJCC TNM Sage for Lung Cancer NSCLC: A Meta-analysis.* Ann Surg Oncol, 2021. **28**(1): p. 142-147.

6. Tartarone, A., et al., *Beyond Conventional: The New Horizon of Targeted Therapy for the Treatment of Advanced Non Small Cell Lung Cancer.* Front Oncol, 2021. **11**: p. 632256.

7. Brady, A.K., et al., *Survival outcome according to KRAS mutation status in newly diagnosed patients with stage IV non-small cell lung cancer treated with platinum doublet chemotherapy.* Oncotarget, 2015. **6**(30): p. 30287-94.

8. Eklund, E.A., et al., *KRAS Mutations Impact Clinical Outcome in Metastatic Non-Small Cell Lung Cancer.* Cancers (Basel), 2022. **14**(9).

9. Goulding, R.E., et al., *KRAS mutation as a prognostic factor and predictive factor in advanced/metastatic non-small cell lung cancer: A systematic literature review and meta-analysis.* Cancer Treat Res Commun, 2020. **24**: p. 100200.

10. Hallqvist, A., et al., *Mutated KRAS Is an Independent Negative Prognostic Factor for Survival in NSCLC Stage III Disease Treated with High-Dose Radiotherapy.* Lung Cancer Int, 2012. **2012**: p. 587424.

11. Hames, M.L., et al., *Correlation between KRAS mutation status and response to chemotherapy in patients with advanced non-small cell lung cancer ☆.* Lung Cancer, 2016. **92**: p. 29-34.

12. Marabese, M., et al., *KRAS mutations affect prognosis of non-small-cell lung cancer patients treated with first-line platinum containing chemotherapy.* Oncotarget, 2015. **6**(32): p. 34014-22.

13. Mellema, W.W., et al., *KRAS mutations in advanced nonsquamous non-small-cell lung cancer patients treated with first-line platinum-based chemotherapy have no predictive value.* J Thorac Oncol, 2013. **8**(9): p. 1190-5.

14. Rodenhuis, S., et al., *Mutational activation of the K-ras oncogene and the effect of chemotherapy in advanced adenocarcinoma of the lung: a prospective study.* J Clin Oncol, 1997. **15**(1): p. 285-91.

15. Meng, D., et al., *Prognostic value of K-RAS mutations in patients with non-small cell lung cancer: a systematic review with meta-analysis.* Lung Cancer, 2013. **81**(1): p. 1-10.

16. Kadota, K., et al., *KRAS Mutation Is a Significant Prognostic Factor in Early-stage Lung Adenocarcinoma.* Am J Surg Pathol, 2016. **40**(12): p. 1579-1590.

17. Izar, B., et al., *The prognostic impact of KRAS, its codon and amino acid specific mutations, on survival in resected stage I lung adenocarcinoma.* J Thorac Oncol, 2014. **9**(9): p. 1363-9.

18. D'Angelo, S.P., et al., *Distinct clinical course of EGFR-mutant resected lung cancers: results of testing of 1118 surgical specimens and effects of adjuvant gefitinib and erlotinib.* Journal of Thoracic Oncology, 2012. **7**(12): p. 1815-1822 %@ 1556-0864.

19. Gallina, F.T., et al., *KRAS G12C mutation and risk of disease recurrence in stage I surgically resected lung adenocarcinoma.* Lung Cancer, 2023. **181**: p. 107254.

20. Ihle, N.T., et al., *Effect of KRAS oncogene substitutions on protein behavior: implications for signaling and clinical outcome.* J Natl Cancer Inst, 2012. **104**(3): p. 228-39.

21.   Garinet, S., et al., *Updated Prognostic Factors in Localized NSCLC.* Cancers (Basel), 2022.
14(6).
22.   Remon, J., J.C. Soria, and S. Peters, *Early and locally advanced non-small-cell lung cancer: An*
update of the ESMO Clinical Practice Guidelines focusing on diagnosis, staging, systemic and
local therapy. Annals of Oncology, 2021. **32**(12): p. 1637-1642 %@ 0923-7534.
23.   Addeo, A., G.L. Banna, and A. Friedlaender, *KRAS G12C Mutations in NSCLC: From Target to*
Resistance. Cancers (Basel), 2021. **13**(11).
24.   Burns, T.F., et al., *Targeting KRAS-Mutant Non-Small-Cell Lung Cancer: One Mutation at a*
Time, With a Focus on KRAS G12C Mutations. J Clin Oncol, 2020. **38**(35): p. 4208-4218.
25.   Indini, A., et al., *Targeting KRAS in Solid Tumors: Current Challenges and Future*
Opportunities of Novel KRAS Inhibitors. Pharmaceutics, 2021. **13**(5).
26.   Mathieu, M., et al., *KRAS G12C fragment screening renders new binding pockets.* Small
GTPases, 2021: p. 1-14.
27.   Skoulidis, F., et al., *Sotorasib for Lung Cancers with KRAS p.G12C Mutation.* N Engl J Med,
2021.
28.   Chen, Y., et al., *Prognostic factors of survival in patients with non-small cell lung cancer: a*
competing risk model using the SEER database. Transl Cancer Res, 2022. **11**(11): p. 3974-
3985.
29.   Gerber, D.E., et al., *Baseline tumour measurements predict survival in advanced non-small*
cell lung cancer. Br J Cancer, 2013. **109**(6): p. 1476-81.
30.   Okada, M., et al., *Effect of tumor size on prognosis in patients with non-small cell lung*
cancer: the role of segmentectomy as a type of lesser resection. J Thorac Cardiovasc Surg,
2005. **129**(1): p. 87-93.
31.   Zhang, J., et al., *Relationship between tumor size and survival in non-small-cell lung cancer*
(NSCLC): an analysis of the surveillance, epidemiology, and end results (SEER) registry. J
Thorac Oncol, 2015. **10**(4): p. 682-90.
32.   Goldie, J.H. and A.J. Coldman, *A mathematic model for relating the drug sensitivity of tumors*
to their spontaneous mutation rate. Cancer treatment reports, 1979. **63**(11-12): p. 1727-
1733.
33.   Skipper, H.E., *Thoughts on Cancer Chemotherapy and Combination Modality Therapy*
(1974). JAMA, 1974. **230**(7): p. 1033-1035.
34.   Jain, R.K., *Physiological Barriers to Delivery of Monoclonal Antibodies and Other*
Macromolecules in Tumors1. Cancer Research, 1990. **50**(3_Supplement): p. 814s-819s.
35.   Trédan, O., et al., *Drug Resistance and the Solid Tumor Microenvironment.* JNCI: Journal of
the National Cancer Institute, 2007. **99**(19): p. 1441-1454.
463
464

Figure legends

Table 1 Characteristics of the total cohort as well as stratified by $KRAS^{WT}$ and $KRAS^{MUT}$. Data are presented as n (%).
ECOG PS, Eastern Cooperative Oncology Group Performance Status. T, Tumor. N, Nodulus.

Table 2 Summary of first-line treatments in the total cohort as well as stratified by $KRAS^{WT}$ and $KRAS^{MUT}$. Data are presented as n (%).

Figure 1. Patient selection
Flow chart showing patient selection for the study.

Figure 2. Impact of *KRAS* mutational status on overall survival in Stage I and II NSCLC.
Kaplan-Meier estimates comparing overall survival between (**A, D**) all patients, (**B, E**) Stage I and (**C, F**) Stage II patients with no mutation in *KRAS* (wildtype, $KRAS^{WT}$), with all *KRAS* mutations ($KRAS^{MUT}$), only *KRAS-G12C* mutations ($KRAS^{MUT\ G12C}$) and *KRAS* mutations other than *G12C* ($KRAS^{MUT\ not\ G12C}$).

Figure 3. Impact of *KRAS* mutational status on overall survival across TNM-stages in NSCLC
Kaplan-Meier estimates comparing overall survival between (**A, D**) T1, (**B, E**) T2 and (**C, F**) T3 patients with no mutation in *KRAS* (wildtype, $KRAS^{WT}$), with all *KRAS* mutations ($KRAS^{MUT}$), only *KRAS-G12C* mutations ($KRAS^{MUT\ G12C}$) and *KRAS* mutations other than *G12C* ($KRAS^{MUT\ not\ G12C}$).

Figure 4. KRAS mutations are associated with smaller tumor size at diagnosis
Primary tumor size from (**A**) CT scans and (**B**) resection specimens (PAD) in patients with no mutation in *KRAS* (wildtype, $KRAS^{WT}$) or with *KRAS* mutations ($KRAS^{MUT}$). Forest plot of multivariate COX regression analysis for patients with tumor size from (**C**) CT scan and (**D**) resection specimens (PAD).

Supplemental Figure 1
Kaplan-Meier estimates of overall survival for resected Stage I-II NSCLC patients stratified by KRAS mutational status. (**A**) No mutation in *KRAS* (wildtype, $KRAS^{WT}$), with all *KRAS* mutations ($KRAS^{MUT}$). (**B**) Only *KRAS-G12C* mutations ($KRAS^{MUT\ G12C}$), *KRAS* mutations other than *G12C* ($KRAS^{MUT\ not\ G12C}$).

Supplemental Figure 2
Kaplan-Meier estimates comparing overall survival for (**A**) all patients, (**B**) patients with no mutation in *KRAS* (wildtype, $KRAS^{WT}$), and (**C**) with all *KRAS* mutations ($KRAS^{MUT}$), stratified by Stages I and II.

Supplemental Figure 3
Kaplan-Meier estimates comparing overall survival for (**A**) all patients, (**B**) patients with no mutation in *KRAS* (wildtype, $KRAS^{WT}$), and (**C**) with all *KRAS* mutations ($KRAS^{MUT}$), stratified by TNM-stages T1, T2 and T3.

Table 1. Characteristics all patients stage I-II NSCLC

	Total	KRAS WT	KRAS MUT
	n (%)	n (%)	n (%)
Total	310 (100)	195 (63.0)	115 (37.0)
Age in years, median (range)	70 (35-85)	70 (35-85)	70 (48-84)
Sex			
Male	123 (39.7)	88 (45.1)	35 (30.4)
Female	187 (60.3)	107 (54.9)	80 (69.6)
Smoking history			
Current smoker	99 (31.9)	51 (26.2)	48 (41.7)
Former smoker	168 (54.2)	106 (54.4)	62 (53.9)
Never smoker	43 (13.9)	38 (19.5)	5 (4.3)
Perfomance status			
ECOG 0	144 (46.5)	82 (42.1)	62 (53.9)
ECOG 1	141 (45.5)	96 (49.2)	45 (39.1)
ECOG 2	24 (7.7)	16 (8.2)	8 (7.0)
ECOG 3	1 (0.3)	1 (0.5)	0
ECOG 4	0	0	0
Histology			
Adenocarcinoma	281 (90.6)	168 (86.2)	113 (98.3)
Squamous cell carcinoma	11 (3.5)	11(5.6)	0
NCSLC NOS	18 (5.9)	16 (8.2)	2 (1.7)
Mutation status			
None known	124 (40.2)	124 (63.6)	
KRAS	115 (37.0)	0	
EGFR	54 (17.4)	54 (27.7)	
BRAF	6 (1.9)	6 (3.1)	
ALK	4 (1.3)	4 (2.1)	
ROS1	3 (1.0)	3 (1.5)	
RET	1 (0.3)	1 (0.5)	
Other	3 (1.0)	3 (1.5)	
KRAS submutation			
G12A			9 (7.8)
G12C			54 (47.0)
G12D			15 (13.0)
G12V			26 (22.6)
Other			11 (9.6)
TNM			
T1a	99 (31.9)	55 (28.2)	44 (38.3)
T1b	76 (24.5)	54 (27.7)	22 (19.1)
T1c	12 (3.9)	7 (3.6)	5 (4.3)
T2a	67 (21.6)	46 (23.6)	21 (18.3)
T2b	28 (9.0)	15 (7.8)	13 (11.3)
T3	28 (9.0)	18 (9.2)	10 (8.7)
N0	292 (94.2)	185 (94.9)	107 (93.0)
N1	18 (5.8)	10 (5.1)	8 (7.0)
Measurement modality			
CT-scan (mm)	305 (98.4)	190 (97.4)	115 (100)
PAD	273 (88.0)	171 (88.7)	102 (88.7)
At last follow up 31/10-2022			
Alive	206 (66.5)	128 (65.6)	78 (67.8)
Deceased	104 (33.5)	67 (34.4)	37 (32.2)
Survival			
Mean survival (months)	63	62	64

515
516

Table 2. Treatment table all patients stage I-II NSCLC

	Total	*KRAS* WT	*KRAS* MUT
	n (%)	n (%)	n (%)
Total	310 (100)	195 (63.0)	115 (37.0)
Surgery	273 (88.0)	171 (87.7)	102 (88.7)
Curative chemoradiotherapy	7 (2.3)	6 (3.1)	1 (0.9)
Medical treatment	2 (0.6)	1 (0.5)	1 (0.9)
Stereotactic radiotherapy	11 (3.5)	11 (5.6)	0 (0)
Radiotherapy	14 (4,5)	3 (1,5)	11 (9.6)
No treatment	3 (1.0)	3 (1.5)	0 (0)

Eklund et al. Figure 1

Eklund et al. Figure 2

Eklund et al. Figure 3

Eklund et al. Figure 4

A CT

p = 0.046

Tumor size CT (mm)

KRAS: WT MUT

KRAS^WT *n* = 190 Median: 25 mm
KRAS^MUT *n* = 115 Median: 20 mm

B PAD

p = 0.161

Tumor size PAD (mm)

KRAS: WT MUT

KRAS^WT *n* = 171 Median: 22 mm
KRAS^MUT *n* = 102 Median: 21 mm

C CT

	HR	[95% CI]	*p* value
Tumor Size (CT - mm) + *KRAS* mut	1.008	[0.986, 1.021]	0.493
Stage II	1.947	[1.201, 3.155]	0.007
Tumor size (CT - mm)	1.006	[0.922, 1.021]	0.398
KRAS mut (Yes)	0.993	[0.651, 1.513]	0.973
Age at diagnosis	1.060	[1.031, 1.091]	< 0.001
Female	0.628	[0.421, 0.938]	0.023

Hazard ratio (Log scale)

D PAD

	HR	[95% CI]	*p* value
Tumor Size (PAD - mm) + *KRAS* mut	1.002	[0.978, 1.027]	0.843
Stage II	1.015	[0.530, 1.947]	0.963
Tumor size (PAD - mm)	1.029	[1.012, 1.046]	< 0.001
KRAS mut (Yes)	1.042	[0.631, 1.722]	0.872
Age at diagnosis	1.043	[1.020, 1.087]	0.002
Female	0.637	[0.397, 1.023]	0.062

Hazard ratio (Log scale)