**Serendipitous detection of *Anopheles stephensi* in Kisumu, Kenya in June 2022**

Bryson Alberto Ndenga, Sammy Wambua, Kevin Omondi Owuor, Rodney Omukuti, Salome

Chemutai, Daniel Arabu, Irene Miringu, Carren Bosire, Kavinya Mwendwa, Christabel Achieng

Winter, Francis Maluki Mutuku, Donal Bisanzio, Angelle Desiree LaBeaud, Keli Nicole Gerken

Author affiliations: Kenya Medical Research Institute, Kisumu, Kenya (B.A. Ndenga, K.O.

Owuor, C.A. Winter); Pwani University, Kilifi, Kenya; Research and Conservation Support

Sciences, Kilifi, Kenya; and University of Glasgow, Glasgow, UK (S. Wambua); Pwani

University, Kilifi, Kenya, and Research and Conservation Support Sciences, Kilifi, Kenya (R.

Omukuti, S. Chemutai, D. Arabu, I. Miringu); Technical University of Mombasa, Mombasa,

Kenya (C. Bosire, K. Mwendwa, F.M. Mutuku); RTI International, Washington, DC, USA (D.

Bisanzio); Stanford University School of Medicine, Stanford, California, USA (A.D. LaBeaud,

K.N. Gerken).

**Abstract**

In June 2022, a pool of five mosquitoes that were morphologically classified as *Anopheles*

*gambiae* and caught in Kisumu (Kenya) were tested for blood-meal analysis. Of the 19.6%

(11/56) amplicon sequence variants assigned to mosquito species using basic local alignment

search tool (BLAST), one had 15 hits matching *Anopheles stephensi*.

**Research letter**

*Anopheles stephensi* is a native urban malaria vector in south Asia and the Arabian

Peninsula. It is capable of transmitting both *Plasmodium falciparum* and *P. vivax* and is

projected to expand further as urbanization increases, especially in Africa (*1*). It was first

reported on the African continent in Djibouti in 2012 (*2*). Since then, it has been reported in

other sub-Saharan countries including Ethiopia in 2019 (*3*), Sudan in 2018 (*4*), Somali-land in

2020 (*5*), Nigeria in 2020 (*6*) and for the first time in Marsabit County in northern Kenya in

December 2022 (*7*). This demonstrates a clear and rapid expansion of its geographical range

within the continent.

During a One Health study on livestock and vector dynamics at urban slaughterhouses,

we carried out mosquito collections using ovitraps, larval/pupae sampling, Prokopack aspirator

and Biogents (BG)-trap baited with $CO_2$. This study was titled "Urban Rift Valley Fever Virus as

a New Ecological Niche: Continuous Introduction from Animal Products" and performed at the

Mamboleo Slaughterhouse (0° 3' 25.4" N, 34° 47' 10.9" E, Figure) in Kisumu City, Kenya, and

was approved by Stanford University (61386), Kenya Medical Research Institute Scientific and

Ethics Review Unit (KEMRI/SERU/CGHR/03-07-390/4293) and obtained research license from

the National Commission for Science, Technology & Innovation (NACOSTI/P/21/13557).

Adult mosquitoes resting indoors and outdoors were sampled using a Prokopack aspirator

from May 16[th] 2022 to July 1[st] 2022. Sampling lasted one hour and was conducted between

08:45 HRS and 01:30 HRS. Subsequent samplings were completed biweekly. Collected

mosquitoes were killed using a pyrethrum aerosol spray before morphological identification in a

field station and recorded on a data form. Female mosquitoes were further sorted according to

their blood-feeding stages as unfed, blood-fed, half-gravid or gravid. Mosquitoes were preserved

in silica gel self-indicating 6-20mesh (Blue) - 500gm (Loba Chemie Pvt. Ltd, Jehangir Villa,

107, Wodehouse Road, Colaba, Mumbai 400 005, India) with the intention for further blood

meal testing.

A total of 2,891 mosquitoes were collected: 66.2% (1,782) indoors and 33.8% (765) outdoors; 50.7% (1,465) males and 49.3% (1,426) females. Species identified morphologically were: 0.3% (9) *Aedes aegypti*; 6.3% (182) *Anopheles coustani*; 0.3% (8) *An. funestus*; 10.9% (315) *An. gambiae*, and 82.2% (2,377) *Culex* spp. Over a half, 61.6% (194/315), of the *An. gambiae* were collected outdoors and 38.4% (121) indoors; 46.3% (146) were males; 19.0% (60) un-blood-fed; 25.1% (79) blood-fed; 6.0% (19) half-gravid and 3.5% (11) gravid.  A sample of the blood-fed mosquitoes were transported on dry ice to Pwani University Biosciences Research Centre (PUBReC) in Kilifi, Kenya, for determination of blood meal sources.

Genomic DNA was extracted from the mosquito abdomens using the TIANamp Genomic DNA Kit (Tiangen, Beijing, China). Quality and quantity of DNA was analyzed by NanoDrop 2000C spectrophotometer (Thermo Scientific Inc, USA) and verified on 1% agarose gel electrophoresis. To identify blood meal sources, we PCR-amplified ~300 base pair (bp) of the cytochrome b barcode (*8*) which was high-throughput-sequenced, at Macrogen Inc., Seoul Korea, using the Illumina 300 × 2 bp platform (Illumina, USA). The amplicon sequence data were analyzed with the DADA2 (version 1.21.0) bioinformatics pipeline (Callahan et al., 2016) implemented in the R programming language (version 4.1.1). After quality preprocessing, high quality reads were denoised and clustered into unique sequences i.e., amplicon sequence variants (ASVs). We assigned taxonomy to the ASVs using two approaches. First, assignment was by exact ASV matching against DADA2-trained MIDORI2 reference database (version GB254) which includes eukaryotic mitochondrial sequences (*9*). ASVs that were not assigned by the first approach were compared against the NCBI database using BLAST (*10*).

In all, 248 ASVs were inferred, 192 of which were classified to 14 vertebrate species (Mammalia and Aves) by exact matching against MIDORI2 reference database. Other than

vertebrates, 19.6% (11/56) of the ASVs assigned using BLAST matched to mosquito species one

of which was *An. stephensi* with a total of 15 hits (89-93% identity, 67-85% coverage, E-value <

1e-46 (Table). The reads matching *An. stephensi* were tracked to a pool of five (5) mosquitoes

that had been morphologically classified as *An. gambiae* in the field. These mosquitoes were

collected from Mamboleo Slaughterhouse in Kisumu City, Kenya in June 2022.

Our findings report the first detection of *An. stephensi* south of the Equator. Furthermore,

this detection represents the presence of an invasive vector in an urban area that is malaria

endemic. Our study suggests that the detections of *An. stephensi* within Africa (*1-5, 7*) could be

dramatically underreported. The initial false identification of this vector also indicates that to

fully capture this vector's boundary, molecular methods and confirmatory testing may be

required. The presence of a malaria vector adapted to breeding in open water containers

prevalent in urban areas is particularly concerning. *Anopheles stephensi* can effectively transmit

*P. falciparum* and *P. vivax* and entry into malaria endemic zones could shift malaria transmission

dynamics considerably.

**Acknowledgements**

We thank Charles Otieno Adipo for sampling the vectors, the management of Mamboleo

Slaughterhouse for allowing us to sample mosquitoes in the premises and our colleagues in

KEMRI and Stanford University for their support to approve this research study. This study was

funded by the Stanford University Center for Innovations in Global Health Seed Grant 2021 and

the Robert E Shope ASTMH (American Society of Tropical Medicine and Hygiene) to Dr. Keli

Nicole Gerken.

**About the Author**

Bryson Alberto Ndenga, PhD is a Senior Research Scientist in Kenya Medical Research

Institute, Centre for Global Health Research. His primary interest is in the biology, ecology and

control of mosquitoes that transmit malaria, dengue, chikungunya and Rift Valley fever.

**References**

1.      Sinka M, Pironon S, Massey N, Longbottom J, Hemingway J, Moyes C, et al. A new

99              malaria vector in Africa: Predicting the expansion range of *Anopheles stephensi* and

100             identifying the urban populations at risk. PNAS. 2020;117.

101     2.      Faulde M, Rueda L, Khaireh B. First record of the Asian malaria vector *Anopheles*

102             *stephensi* and its possible role in the resurgence of malaria in Djibouti, Horn of Africa.

103             Acta Trop. 2014;139:39-43.

104     3.      Tadesse F, Ashine T, Teka H, Esayas E, Messenger L, Chali W, et al. *Anopheles*

105             *stephensi* mosquitoes as vectors of *Plasmodium vivax* and *P. falciparum*, horn of Africa,

106             2019. Emerg Infect Dis. 2021;27.

107     4.      Ahmed A, Khogali R, Elnour M, Nakao R, Salim B. Emergence of the invasive malaria

108             vector *Anopheles stephensi* in Khartoum State, Central Sudan. Parasit Vectors. 2021;14.

109     5.      Ali S, Samake J, Spear J, Carter T. Morphological identification and genetic

110             characterization of *Anopheles stephensi* in Somaliland. Parasit Vectors. 2022;15.

111     6.      World Health Organization initiative to stop the spread of *Anopheles stephensi* in Africa.

112             Geneva: WHO; 2022.

7.    Ochomo E, Milanoi S, Abong'o B, Onyango B, Muchoki M, Omoke D, et al. Molecular

surveillance leads to the first detection of *Anopheles stephensi* in Kenya. Res Square. In

press 2023.

8.    Hadj-Henni L, De-Meulemeester T, Depaquit J, Noël P, Germain A, Helder R, et al.

Comparison of vertebrate cytochrome b and prepronociceptin for blood meal analyses in

Culicoides. Front Vet Sci. 2015;2.

9.    Leray M, Knowlton N, Machida R. MIDORI2: A collection of quality controlled,

preformatted, and regularly updated reference databases for taxonomic assignment of

eukaryotic mitochondrial sequences. Environ DNA. 2022;4:894–907.

10.    Altschul S, Gish W, Miller W, Myers E, Lipman D. Basic local alignment search tool. J

Mol Bio. 1990;215:403–410.

Address for correspondence: Dr. Bryson Alberto Ndenga, Kenya Medical Research Institute,

Centre for Global Health Research, P.O. Box 1578 Code 40100 Kisumu City, Kenya, email:

bndenga@yahoo.com

**Tables and Figures**

**Figure 1.** Map showing the location in Kisumu, Kenya where *Anopheles stephensi* was collected

in June 2022 and other sites in Africa where it has been reported (Source: *Anopheles stephensi*

reports (red points): https://apps.who.int/malaria/maps/threats/).

**Table 1.** Summary of coverage and identity values for each of the top ten (10) hits in the BLAST

search matching *Anopheles stephensi*.

| Description | Maximum score | Query coverage | E value | Identity | Accession |
|---|---|---|---|---|---|
| *Anopheles stephensi* strain Indian chromosome 2L | 326 | 84% | 2.00E-84 | 92.51% | CP032300.1 |
| *Anopheles stephensi* strain Indian chromosome 3L | 320 | 85% | 8.00E-83 | 92.07% | CP032302.1 |
| *Anopheles stephensi* strain Indian chromosome 3R | 320 | 82% | 8.00E-83 | 92.07% | CP032301.1 |
| *Anopheles stephensi* strain Indian chromosome 2R | 315 | 81% | 4.00E-81 | 91.63% | CP032299.1 |
| *Anopheles stephensi* strain SDA-500 chromosome 3R | 315 | 83% | 4.00E-81 | 91.63% | CP032234.1 |
| *Anopheles stephensi* strain SDA-500 chromosome 2R | 309 | 81% | 2.00E-79 | 91.19% | CP032232.1 |
| *Anopheles stephensi* strain SDA-500 chromosome 2L | 303 | 84% | 9.00E-78 | 90.75% | CP032233.1 |

| | | | | | |
|---|---|---|---|---|---|
| *Anopheles stephensi* strain SDA-500 chromosome 3L | 298 | 88% | 4.00E-76 | 90.31% | CP032235.1 |
| *Anopheles ziemanni* genome assembly, chromosome: 2 | 291 | 83% | 7.00E-74 | 89.52% | OX030919.2 |