# Prevalence and determinants of not testing for HIV among young adult women in Papua New Guinea: findings from the 2016–2018 Demographic and Health Survey

McKenzie K. Maviso[1], Fatch Welcome Kalembo[2]

[1]School of Medicine and Health Sciences, University of Papua New Guinea, Port Moresby, Papua New Guinea

[2]School of Nursing, Curtin University, Perth, Western Australia, Australia

Correspondence to McKenzie K. Maviso; mckenzie.maviso@upng.ac.pg

## Abstract

### Objective

The study investigated the factors associated with not ever testing for HIV among women aged 15–29 years in Papua New Guinea (PNG).

### Design and setting

The study used secondary data from the 2016–2018 Demographic and Health Survey (DHS) of PNG, a nationally representative cross-sectional survey that used a two-stage stratified sampling.

### Participants

A total weighed sample of 5,164 young adult women aged 15–29 years were included in the study.

### Primary outcome measure

Ever been tested for HIV was the primary outcome of the study. All analyses were adjusted using survey weights to account for unequal sampling probabilities.

### Results

The prevalence of not testing for HIV was 58.8% (95% CI: 57.4%, 60.1%). The mean age was 21.65 (SD ± 4.23) years. The majority (62.9%) of young adult women in rural areas were not tested for HIV. In the multivariable analysis, those who were never married (AOR: 4.9, 95% CI: 3.6–6.6), had poor wealth index (AOR: 1.8, 95% CI: 1.3–2.5), from rural areas (AOR: 2.0, 95% CI: 1.5–2.6), from the Momase region (AOR: 1.3, 95% CI: 1.0–1.7, did not read newspaper or magazine (AOR: 1.7, 95% CI: 1.3–2.1), did not listen to radio (AOR: 1.5, 95% CI: 1.1–2.0), experienced early sexual debut (AOR: 1.5, 95% CI: 1.1–1.9), had one sexual partner (AOR: 1.5, 95% CI: 1.2–2.0), and reported no STI in the past 12 months (AOR: 1.8, 95% CI: 1.1–3.1) had higher odds of not testing for HIV.

### Conclusions

Our study found a very high unmet need for HIV testing among young adult women in PNG. Health promotion programmes should be designed to increase HIV knowledge and access to HIV testing services. Furthermore, efforts to optimise HIV testing services should target women who are disadvantaged and from rural areas.

## Strengths and limitations of this study

- This study used a nationally representative large sample of young adult women aged 15–29 years from the 2016–2018 PNGDHS; thus, the findings are generalisable to the entire population of this age group.

- The study provides much-needed data for strategic planning and programming to improve the health outcomes of young adult women in the country.

- This is a cross-sectional study, and as such, it is not possible to make causal inferences.

- The level of knowledge or awareness of where to get tested for HIV was not included in the analysis, and it remains unknown whether this variable contributes to the high proportion of respondents who remain unaware of their HIV status.

- Responses to the questions from respondents were self-reported, and this could have resulted in behavioural desirability bias.

## Introduction

Globally, an estimated 39 million people were living with HIV in 2022, and approximately 17% (6.5 million) of these were from the Asia-Pacific region. [1] Among people living with HIV, young women continue to endure a disproportionate burden of the virus. [2] Young women are twice as likely as young men of the same age to have HIV; [3] are frequently unaware of their HIV status and lack access to antiretroviral therapy (ART) and care. [4] They are more likely to experience physical or sexual violence from intimate partners compared to their male counterparts. [5,6] The lifetime prevalence of intimate partner violence among young women is 16%. [6,7] The violence experienced by young women is a significant factor in causing unequal infection rates in settings with high HIV prevalence. [8] Such disproportionately high rates of HIV infection in young women compared to young men are illuminated by an array of influences. For instance, social and institutional norms (e.g., male superiority) and gender-based power inequalities (e.g., sexual entitlement within relationships) prevent women from negotiating safe sex practices, which frequently impact their access to sexual and reproductive health care and HIV prevention services. [9–12]

Voluntary counselling and testing (VCT) for HIV is a gateway to HIV treatment and care for those who test positive and a prevention strategy for those who test negative. [13] Awareness of one's HIV

serostatus is associated with risk reduction and behaviour change. [14,15] Newly diagnosed HIV individuals can be promptly connected to ART, care, and support services. Timely HIV diagnosis and treatment access are crucial for preventing transmission, as newly diagnosed individuals can immediately be linked to ART, care, and support services. [16]

While access to HIV testing, treatment, and support services has been expanding globally, the response to HIV testing has remained inadequate, particularly among young women who are recognised for having poor health-seeking behaviours in highly prevalent settings. [17,18] Factors associated with low coverage or not testing for HIV are multifactorial and include the fear of HIV test results and stigma, the perceived low risk of contracting HIV, and the psychological burden of living with HIV. [17,19] In addition, knowledge and attitude towards HIV testing services, age, education, engagement in sexual relationships, and distance to access testing facilities are associated with low uptake of HIV testing among young people. [20,21]

PNG is one of the most HIV-affected countries in the Western Pacific Region, with a high HIV prevalence projected at 0.9% in the general adult population. [22,23] In 2022, an estimated 72,000 people were living with HIV in PNG. [24] In addition, women of childbearing age (15–49 years) in PNG represent a disproportionate population that requires adequate HIV prevention services, including sexual and reproductive health care. [25–27] In 2022, women accounted for about 67% (41,000) of HIV, with 3,600 new HIV infections in the country. [24] It has been reported that HIV prevalence among PNG women is attributed to multiple factors such as gender inequality, sociocultural norms, household poverty, and unsupported social and structural factors, including unequal access to sexual and reproductive health services, increasing women's vulnerability to HIV infection. [28–31]

To address the burden of HIV, the government of PNG has integrated the WHO-recommended VCT for HIV into the country's critical HIV prevention and control strategies. [32] With more ART programmes available, efforts to strengthen HIV testing have become increasingly important for early diagnosis and linking access to treatment, adherence to treatment, and viral load suppression. [32,33] Despite the country's efforts to reduce HIV transmission, VCT uptake, linkage to ART, and support services remain problematic. [33–35] In particular, young women's awareness of HIV prevention strategies, such as consistent condom use and reducing the number of sexual partners, remains low (24%). [2] For this study, young adult women aged 15–29 years were included in the analysis. Previous

studies have investigated sexual, social, and structural determinants of risk behaviours and adverse

health outcomes, including HIV acquisition, among this group (aged 15–29). [36–38] However,

research on factors associated with not testing is limited, especially in young women, who face a high

risk of HIV. Furthermore, there have been no studies investigating how these factors influence HIV

testing, an integral component of an HIV prevention package. Therefore, this study aimed to examine

factors associated with not testing for HIV among young adult women in PNG. Recognising and

addressing the challenges that young women face in accessing HIV testing services is critical to

achieving the ambitious UNAIDS 95-95-95 targets to end the HIV/AIDS epidemic. [39]

## Methods

### Study design and sampling

This study is a result of the analysis of the 2016–2018 PNGDHS data, which is a nationally

representative cross-sectional survey that provides data to monitor and evaluate population, health, and

nutrition programmes. The PNGDHS researchers used a two-stage stratified sampling technique to

recruit participants for the study. The first stage involved selecting 800 census units (CUs), which was

achieved through a probability proportional to CU size. The second stage involved the systematic

selection of 24 households from each cluster (CU) through equal probability sampling, with the

resulting sample consisting of 19,200 households. The survey used the CUs from the 2011 PNG

National Population and Housing Census, which used structured sampling. Additionally, PNG has 22

administrative provinces, and each province is subdivided into urban and rural areas. Each province

also has districts, and each district is divided into local-level government wards, where each ward is

comprised of several CUs. The PNG National Statistical Office (NSO) and ICF International collected

the data. In the interviewed households, 18,175 women aged 15–49 years were identified for individual

interviews. Of these, 15,198 women participated in the survey successfully, yielding a response rate of

84%. Among those who were interviewed successfully, 5,164 (the weighted sub-sample) were young

adult women aged 15–29 years who had already initiated sex and were included in this study.

Information about the methodology, pretesting, and training of field officers, the sampling design, and

sample selection is available in the PNGDHS final report [40] and online at

https://dhsprogram.com/methodology/survey/survey-display-499.cfm.

143
144
145
146

## Definition of variables

*Outcome variable*

The outcome variable was HIV testing among young adult women, a dichotomous variable coded as "1" when a woman reported having "ever been tested for HIV" and "0" when she reported otherwise (never tested for HIV). The information about the dependent variable was generated from this question.

*Explanatory variables*

The explanatory variables used in this study were selected based on the available data in the 2016–2018 PNGDHS datasets and previously published literature. [17,18,41] The independent variables were categorised into socio-demographic factors (age, marital status, education level, occupation, wealth index, place of residence, and region). Age was categorised into "15–19," 20–24," and "25–29." Marital status was recoded into "never married," "married/living with a partner," and "previously married." The occupation was recoded into "not working" and "working." The wealth indices were determined by household assets owned (e.g., televisions and radio), housing construction materials, and access to water and sanitation facilities [40], and were reported as being "poorer," "poor," "middle," "rich," and "richer." A new variable for the wealth index was created and recoded into "poor," "middle," and "rich." Behaviour-related variables comprised the frequency of exposure to mass media (reading a newspaper or magazine, listening to the radio, and watching television), mobile phone ownership, and internet use. Exposure to media was determined by the frequency of reading a newspaper or magazine, listening to the radio, and watching television. Each of these variables had three responses: "not at all," "less than once a week," and "at least once a week." These variables were recoded as "yes" and "no" responses to exposure to media. Use of the internet was identified as "never," "yes, last 12 months," "yes, before last 12 months," and "yes, cannot establish." The internet use variable was recoded into "no" and "yes." Furthermore, health risk behaviours include age at first sex, the total number of sexual partners (in the last 12 months), ever being paid for sex (in exchange for cash, gifts, or others), always using a condom during sex, and history of any STI (in the last 12 months). Age at sexual debut was recoded as "less than 20 years" and "20 years or older." The total number of sexual partners (in the last 12 months) was recoded into "1" and "2 or more." Incomplete and missing data in the "ever tested for HIV" group remained low (1.2%); therefore, they were excluded from the final analysis.

## Statistical analysis

The final sample data was weighted to restore representativeness and produce a reliable estimate and standard error. Descriptive statistics were used to describe the sociodemographic and HIV-related characteristics of the participants. Bivariate analysis using the Pearson Chi-square ($\chi^2$) test of independence was performed to examine the relationship between the explanatory variables and the outcome variable. Variables significant at $p < 0.25$ in univariate analysis were then used in the final multivariable logistic regression model. [42] A Complex Sample Analysis Procedure that accounted for the cluster sampling design and sample weight to provide generalisable and accurate estimates of proportion, probability values, and odds ratios was employed in this analysis. [43,44] The Hosmer-Lemeshow goodness-of-fit was used to test for model fitness. Measures of association were presented as adjusted odds ratios (AORs) with their corresponding 95% confidence intervals (CIs). A $p$-value $\leq$ 0.05 was used to determine the statistical significance of all analyses. All analyses were conducted using the IBM Statistical Package for the Social Sciences (SPSS), Version 26 (Armonk, NY: IBM Corp.).

## Patient and public involvement

Patients and the public were not involved in the study. We analysed the secondary data available in the public repository.

## Results

### Characteristics of participants

Participants' demographic characteristics between those who were tested for HIV and those who were not are presented in **Table 1**. Of the total weighted sample of 5,164 young adult women, 3,010 [58.8% (95% CI 57.4%, 60.1%)] were never tested for HIV. The mean age was 21.65 (SD ± 4.23) years. Over half of the women who were not tested for HIV were young (aged < 20 years) (55.1%), never married (79.4%), lacked formal education (63%), not working (60.2%), poor (71.2%), and from rural areas (62.9%). Nearly two-thirds were not exposed to the media (i.e., newspaper or magazine, radio, television). In terms of risky sexual behaviours and related factors, most women who had not tested for HIV reported a sexual debut at age < 20 (61.3%), having one sexual partner (61.7%), having ever been paid for sex (64.9%), always using a condom during sex (51.1%), and not having had an STI in the last 12 months (59.3%). There were statistically significant differences in the proportions of participants who had never been tested for HIV compared to those who had in all the demographic variables, age at

210 sexual debut, the total number of sexual partners, ever being for sex, and having had an STI for the past

211 12 months in the HIV-related variables (p < 0.05).

**Table 1. Comparison of participants' demographic characteristics between those who were ever tested for HIV and those who were not in PNG.**

| Characteristics | Participants 5,164 (%) | Ever tested for HIV | | p-value* |
| | | No 3,010 (%) | Yes 2,154 (%) | |
|---|---|---|---|---|
| **Age group (years)** | | | | **0.001** |
| 15–19 | 1,880 | 1,035 (55.1) | 845 (44.9) | |
| 20–24 | 1,725 | 1,028 (59.6) | 697 (40.4) | |
| 25–29 | 1,559 | 947 (60.7) | 612 (39.3) | |
| Mean age = 21.65 (SD ± 4.23) | | | | |
| **Marital status** | | | | **< 0.001** |
| Never married | 832 | 661 (79.4) | 171 (20.6) | |
| Married | 4,005 | 2,182 (54.5) | 1,823 (45.5) | |
| Divorced/separated | 327 | 167 (51.1) | 160 (48.9) | |
| **Education[#]** | | | | **0.001** |
| No formal education | 894 | 563 (63.0) | 331 (37.0) | |
| Primary | 2,652 | 1,535 (57.9) | 1,117 (42.1) | |
| High/Secondary | 1,422 | 785 (55.2) | 637 (44.8) | |
| College/Tertiary | 195 | 127 (65.1) | 68 (34.9) | |
| **Occupation*** | | | | **< 0.001** |
| Not working | 3,475 | 2,091 (60.2) | 1,384 (39.8) | |
| Working | 1,620 | 883 (54.5) | 737 (45.5) | |
| **Wealth Index[#]** | | | | **< 0.001** |
| Poor | 1,977 | 1,407 (71.2) | 570 (28.8) | |
| Middle | 943 | 540 (57.3) | 403 (42.7) | |
| Rich | 2,243 | 1,063 (47.4) | 1,180 (52.6) | |
| **Place of residence** | | | | **< 0.001** |
| Urban | 1,054 | 424 (40.2) | 630 (59.8) | |
| Rural | 4,110 | 2,586 (62.9) | 1,524 (37.1) | |
| **Region** | | | | **< 0.001** |
| Southern | 999 | 617 (61.8) | 382 (38.2) | |
| Highlands | 2,259 | 1,168 (51.7) | 1,091 (48.3) | |
| Momase | 1,272 | 870 (68.4) | 402 (31.6) | |
| Islands | 633 | 355 (56.1) | 278 (43.9) | |
| **Read newspaper/magazine[#]** | | | | **< 0.001** |
| No | 3,311 | 2,160 (65.2) | 1,151 (34.8) | |

| | | | |
|---|---|---|---|
| Yes | 1,834 | 835 (45.5) | 999 (54.5) |
| **Listen to the radio**[#] | | | < 0.001 |
| No | 3,253 | 2,107 (64.8) | 1,146 (35.2) |
| Yes | 1,893 | 896 (47.3) | 997 (52.7) |
| **Watch television**[#] | | | < 0.001 |
| No | 3,814 | 2,366 (62.0) | 1,448 (38.0) |
| Yes | 1,298 | 602 (46.4) | 696 (53.6) |
| **Owns a mobile phone**[#] | | | <0.001 |
| No | 3,352 | 2,135 (63.7) | 1,217 (36.3) |
| Yes | 1,800 | 867 (48.2) | 933 (51.8) |
| **Use the internet**[#] | | | < 0.001 |
| No | 4,362 | 2,676 (61.3) | 1,686 (38.7) |
| Yes | 783 | 325 (41.5) | 458 (58.5) |
| **Age of sexual debut (years)** | | | < 0.001 |
| < 20 | 3,710 | 2,275 (61.3) | 1,435 (38.7) |
| 20 or more | 1,454 | 735 (50.6) | 719 (49.4) |
| Mean age = 14.86 (SD ± 8.80) | | | |
| **Total number of sexual partners**[#] | | | < 0.001 |
| 1 | 3,564 | 2,198 (61.7) | 1,366 (38.3) |
| 2 or more | 1,486 | 755 (50.8) | 731 (49.2) |
| **Ever being paid for sex**[#] | | | 0.050 |
| No | 476 | 375 (78.8) | 101 (21.2) |
| Yes | 37 | 24 (64.9) | 13 (35.1) |
| **Always condom use**[#] | | | 0.261 |
| No | 1,542 | 761 (49.4) | 781 (50.6) |
| Yes | 2,771 | 1,417 (51.1) | 1,354 (48.9) |
| **Had any STI (last 12 months)**[#] | | | < 0.001 |
| No or don't know | 4,859 | 2,882 (59.3) | 1,977 (40.7) |
| Yes | 221 | 85 (38.5) | 136 (61.5) |

*$\chi$ test, $p = 0.05$. [#]Variables with missing values.

**Factors associated with not testing for HIV**

Factors associated with not testing for HIV among young adult women are presented in **Table 2**. In the multivariable logistic regression analysis, those who were never married (AOR: 4.9, 95% CI: 3.6–6.6), had poor wealth index (AOR: 1.8, 95% CI: 1.3–2.5), from rural areas (AOR: 2.0, 95% CI: 1.5–2.6), from the Momase region (AOR: 1.3, 95% CI: 1.0–1.7), did not read newspapers or magazines (AOR:

219 1.7, 95% CI: 1.3–2.1), did not listen to radio (AOR: 1.5, 95% CI: 1.1–2.0), experienced early sexual

220 debut (< 20 years) (AOR: 1.5, 95% CI: 1.1–1.9), had one sexual partner (AOR: 1.5, 95% CI: 1.2–2.0),

221 and reported no STI in the past 12 months (AOR: 1.8, 95% CI: 1.1–3.1) had higher odds of not testing

222 for HIV. Conversely, young women from the Highlands region had lower odds of not testing for HIV

223 compared with those in the Southern region (AOR: 0.4, 95% CI: 0.4–0.6).

**Table 2. Factors associated with not ever being tested for HIV in the multivariable logistic regression analysis.**

| Characteristics | COR (95% CI) | AOR (95% CI) | *p*-value |
|---|---|---|---|
| **Age group (years)** | | | 0.133 |
| 15–19 | Ref | Ref | |
| 20–24 | 1.2 (0.9, 1.6) | 0.9 (0.8–1.2) | |
| 25–29 | 1.3 (1.0, 1.6) | 1.2 (0.9–1.5) | |
| **Marital status** | | | **< 0.001** |
| Married | Ref | Ref | |
| Never married | 3.2 (2.4, 4.3) | 4.9 (3.6–6.6) | |
| Divorced/separated | 0.9 (0.6, 1.2) | 0.9 (0.7–1.4) | |
| **Education** | | | 0.210 |
| No formal education | 0.9 (0.3, 2.6) | 0.9 (0.3–2.7) | |
| Primary | 0.7 (0.3, 2.1) | 0.7 (0.3–2.0) | |
| High/Secondary | 0.7 (0.2, 2.0) | 0.6 (0.2–2.0) | |
| College/Tertiary | Ref | Ref | |
| **Occupation** | | | 0.702 |
| Not working | 1.3 (1.0, 1.5) | 1.0 (0.9–1.3) | |
| Working | Ref | Ref | |
| **Wealth Index** | | | **< 0.001** |
| Poor | 2.7 (2.2, 3.5) | 1.8 (1.3–2.5) | |
| Middle | 1.5 (1.1, 1.9) | 1.2 (0.9–1.7) | |
| Rich | Ref | Ref | |
| **Place of residence** | | | **< 0.001** |
| Urban | Ref | Ref | |
| Rural | 2.5 (2.0, 3.1) | 2.0 (1.5–2.6) | |
| **Region** | | | **< 0.001** |
| Southern | Ref | Ref | |
| Highlands | 0.7 (0.5, 0.9) | 0.4 (0.4–0.6) | |

| | COR (95% CI) | AOR (95% CI) | p-value |
|---|---|---|---|
| Momase | 1.3 (0.9, 1.9) | 1.3 (1.0–1.7) | |
| Islands | 0.8 (0.6, 1.0) | 1.0 (0.7–1.3) | |
| **Read newspaper/magazine** | | | **< 0.001** |
| No | 2.2 (1.9, 2.7) | 1.7 (1.3–2.1) | |
| Yes | Ref | Ref | |
| **Listen to radio** | | | **0.004** |
| No | 2.1 (1.7, 2.4) | 1.5 (1.1–2.0) | |
| Yes | Ref | Ref | |
| **Watch television** | | | 0.120 |
| No | 1.9 (1.5, 2.3) | 0.8 (0.6–1.1) | |
| Yes | Ref | Ref | |
| **Owns a mobile phone** | | | 0.118 |
| No | 1.9 (1.5, 2.3) | 1.2 (0.9–1.6) | |
| Yes | Ref | Ref | |
| **Use the internet** | | | 0.328 |
| No | 2.2 (1.8, 2.8) | 1.1 (0.9–1.5) | |
| Yes | Ref | Ref | |
| **Age of sexual debut (years)** | | | **0.006** |
| < 20 | 1.6 (1.2, 2.0) | 1.5 (1.1–1.9) | |
| 20 or older | Ref | Ref | |
| **Total number of sexual partners** | | | **0.001** |
| 1 | 1.6 (1.2, 1.9) | 1.5 (1.2–2.0) | |
| 2 or more | Ref | Ref | |
| **Had any STI (last 12 months)** | | | **0.024** |
| No or don't know | 2.3 (1.5, 3.6) | 1.8 (1.1–3.1) | |
| Yes | Ref | Ref | |

Ref = Reference Category; COR = Crude Odds Ratio; AOR = Adjusted Odds Ratio; CI = Confidence Interval; $p$-value = 0.05

## Discussion

This study investigated factors associated with not testing for HIV among young adult women aged

15–29, based on a nationally representative sample. In our analysis of 5,164 young women from the

2016–2018 DHS in PNG, we found that 71% had never tested for HIV. Not testing for HIV was

associated with marital status, wealth index, place of residence, region, reading of newspapers and

magazines, listening to the radio, age of sexual debut, the total number of sexual partners, and having

any STI in the last 12 months preceding the survey, which were significantly associated with not

testing for HIV among young adult women. Given the nature of the study, a causal association cannot

be established. Also, behavioural desirability bias might have some effect on the underreporting of

sexual behaviours since all the responses were self-reported.

Seven out of every 10 young adult women were never tested for HIV, suggesting a significantly low

proportion of HIV testing. This finding corresponds with previous studies from sub-Saharan Africa

(63.5%) [41], including Ethiopia (66.5%) and Nigeria (76.3%), [17,18] which showed significantly low

HIV testing rates among young women. The low HIV testing uptake in the country may be due to

differences in HIV testing modalities implemented for young women. Strategies to optimise access to

HIV testing need to shift from conventional facility-based testing to community- and home-based

testing services. Community and home-based HIV testing have been proven to enhance HIV testing

coverage among young women through self-testing and linking them to ART treatment. [45–47] Self-

testing for HIV could increase the proportion of individuals who have ever tested, increase testing

frequency, and promote early detection and treatment. [48]

In this study, the majority (80%) of women were from poor households and reported to have never

been tested for HIV. This finding supports the recent studies showing that young women, particularly

in low- and middle-income countries, encounter a heterogeneous risk for HIV based on their household

socioeconomic conditions. [49,50] Furthermore, various aspects of socio-cultural structure can

potentiate the HIV risk burden for poor women, such as lack of access to information, health care,

formal education, economic opportunities, and exposure to violence. [49]

Adolescent girls and young women from economically disadvantaged and poor households are at

higher risk for HIV due to their lack of awareness about HIV transmission and their less frequent use of

condoms. [49] However, the authors of studies conducted in Malawi and Rwanda found an inverse

association between rich households, higher education, and the uptake of HIV testing among young

women. [51,52] While earlier studies in PNG have suggested strategies for reducing the gap between

socioeconomic quintiles and health among young women, [29] there is still evidence of inequalities.

Studies have shown links between HIV testing and better adherence to safe sex practices and women's

empowerment in terms of education, financial independence, control over household financial

decisions, and the power of negotiating sex. [53–55] Our findings suggest that there is an urgent need

to tailor prevention strategies for HIV for this group that are sensitive to their culture and

socioeconomic context. The focus on improving access to HIV testing services should first be on

conceptualising gender in a way that is inclusive. Furthermore, there is a need to address barriers

related to poor care-seeking behaviours through the provision of youth-friendly services.

Additionally, the findings of this study showed that rural women are twice as likely not to be tested for

HIV compared to their urban counterparts, which corresponds with recent studies conducted in

Ethiopia [18] and eastern Africa. [56] Low uptake of HIV testing could be due to the difficulty in

accessing sexual and reproductive health and HIV services in primary healthcare facilities, including

gender issues, societal norms, and a lack of HIV knowledge, as reported in other studies. [27,57,58]

Contrary to the findings of our study, the authors of the Ethiopian study have reported an increased

uptake of HIV testing among women in rural areas who have higher socio-economic status, non-

farming occupations, female-headed households, live within proximity of health facilities, and

participate in community dialogues. [59] The utilisation of HIV testing services in rural areas is largely

influenced by socio-economic, behavioural, and health service factors.

Our study further found a negative association between the region of residence and not testing for HIV,

with women in the Momase region being more likely to not be tested for HIV compared to those in the

Southern region. The regional disparities in not testing for HIV in PNG could be a result of the unequal

allocation and distribution of health resources, including access to healthcare services [26,31]. This

finding highlights the need for interventions aimed at improving HIV testing, especially targeting

women from disadvantaged regions of the country.

Women who had no access to at least one media platform (newspapers or radio) were less likely to

have been tested for HIV. This confirms the findings from recent Ethiopian studies that a lack of media

access and comprehensive HIV knowledge is associated with lower odds of HIV testing. [60,61] It is

most likely that young women do not read newspapers or listen to the radio. However, a recent

Rwandan study revealed that young women who have access to at least one media platform at least

once a week (newspapers, TV, or radio) are more likely to have been tested for HIV. [52] Media

exposure increases awareness about the importance of HIV testing and improves understanding of the

prevention approach, which might encourage women to be tested for HIV. [60] The results of the study

underscore the need to utilise various media platforms in disseminating HIV information to sensitise

young women to constantly learn about the benefits of HIV prevention.

Findings from this study showed that the majority (61.3%) of women who debuted sex were aged < 20

and were not tested for HIV compared to their counterparts. The study supports previous findings

suggesting a low HIV testing prevalence among this age group in high-prevalence settings. [17] Early

sexual debut with risky behaviours has been posited as an important risk factor for HIV infection.

Studies have shown that factors such as peer pressure, lower education, coercion, substance use

(tobacco and alcohol use), poor parental or guardian attachment, and exposure to pornographic

materials were associated with an early sexual debut. [62–64] Furthermore, a lack of HIV knowledge

and low-risk perception have been considered a potential barrier to HIV testing among adolescent

women. [65] This could be a likely reason why young women were not tested for HIV in this study.

The precocious sexual debut predisposes young women to adverse psychological and health outcomes.

Thus, the need for timely, age-appropriate, continuous, and enhanced awareness and reproductive

health services and interventions is imperative for this priority group.

Regarding the number of sexual partners, women who had one sexual partner had lower odds of being

tested for HIV compared with their counterparts. This finding is in contrast with other studies [56],

where women with multiple sex partners had a lower chance of being tested for HIV. The low uptake

of HIV testing among young women in this study could be due to the perception that they have a lower

risk of contracting HIV because they have one sexual partner. Another reason could be that women

refuse or forgo testing out of fear of the consequences they may encounter, such as HIV-related stigma

and discrimination, including violence from their partners and family for disclosing their HIV-positive

status, as reported in previous studies. [27,66,67] There are common notions that female biological

vulnerability makes women more likely to contract HIV than men and that women's socio-economic

marginalisation increases their HIV risk. [57] HIV-related stigma can manifest in a variety of ways,

including isolation, mockery, condemnation, physical and verbal abuse, and denial of healthcare

access, including HIV services. [68] Interventions to address stigma should not be confined to health

facilities; instead, they need to be tailored and focus on creating changes in individual knowledge,

attitudes, and behaviours rather than broader social and environmental changes.

In this study, participants who had no STIs in the last 12 months were likely not to have ever been tested for HIV, indicating a lack of awareness and knowledge about STIs. Studies have shown that understanding of STIs in terms of acquisition, protection, prevention, and clinical signs and symptoms is limited among young people. [69] Poor knowledge about STI screening and services could be one possible explanation for young women not knowing their STI statuses. Lack of STI information and screening services, acceptability of services, fear of positive test results, fear of invasive procedures, self-consciousness of genital examination, anticipated stigma, and confidentiality concerns have been found to impede STI screening uptake among young people. [70–72] Also, young women may not have access to the required STI information and services to avoid infections or may be hesitant to access the facilities where information is available. This could be another reason for being unsure or not having any STIs. This study highlights the unmet need for STI screening among young adult women in PNG. Urgent efforts are needed to improve STI knowledge through tailored sexual and reproductive health services while taking into account social and cultural norms surrounding gender and sexuality.

**Implications for practice and research**

Increasing uptake of HIV testing and counselling among young women can lead to earlier diagnosis, linkage to ART, and retention in care. Interventions that are affordable, feasible, and acceptable for this population to increase access to and the likelihood of being tested remain imperative. To optimise testing coverage, it is necessary to determine young women's preferences about HIV testing modalities and where they choose to access the service. HIV testing options for young women that include self-testing are promising strategies that may overcome an array of access barriers in this context. In addition, sexual and reproductive health for young women remains crucial to increasing the uptake of HIV testing. There is a need for more sensitisation programmes for healthcare providers, for instance, education and training to sensitise them to reduce HIV-related fear and stigma among young women through user-friendly services. Given that the majority of young women do not seek HIV testing services in PNG, there is a critical need for implementation research on strategies for increasing access to testing, prevention, and treatment. While the findings of this study have provided information about the proportion of young women who have never tested for HIV, it is also important to have an in-depth understanding of the barriers to HIV testing in this age group. Further qualitative research is required to explore barriers and facilitators to HIV testing among this population, which could inform the development of novel HIV prevention strategies.

## Conclusions

HIV testing among young adult women remained low in PNG, with more than half having not been tested for HIV. Factors associated with not testing for HIV included never being married, poor households, rural areas, not having access to media (newspaper, magazine, or radio), early sexual debut, having one sex partner, and an unknown STI status. These factors could allow the epidemic to intensify as young adult women with undiagnosed infections are not likely to be tested for HIV, adopt behaviour change and risk reduction strategies, or access treatment. The use of various media platforms is crucial for disseminating information on HIV/STI prevention, risk reduction, and increasing HIV testing uptake among young women. Efforts to optimise access to HIV testing should also shift from conventional facility-based testing to community- and home-based testing services to target those who are disadvantaged and from rural areas. Furthermore, programmatic approaches from the government, civil society, and communities to overcome barriers associated with the uptake of HIV testing are thus warranted.

## Acknowledgements

We are grateful to the ICF International and DHS Program for permitting us to use the 2016–2018 PNGDHS dataset to conduct this study.

## References

[dataset] 1    UNAIDS. Global HIV & AIDS statistics - Fact sheet. Global HIV & AIDS statistics — Fact sheet. 2023. https://www.unaids.org/en/resources/fact-sheet

[dataset] 2    UNAIDS. UNAIDS Data 2021. Geneva, Switzerland: UNAIDS 2021. https://www.unaids.org/en/resources/documents/2021/2021_unaids_data

[dataset] 3    Karim SSA, Baxter C. HIV incidence rates in adolescent girls and young women in sub-Saharan Africa. *The Lancet Global Health*. 2019;7:e1470–1. doi: 10.1016/S2214-109X(19)30404-8

[dataset] 4    Stoner MCD, Nguyen N, Kilburn K, *et al.* Age-disparate partnerships and incident HIV infection in adolescent girls and young women in rural South Africa. *AIDS*. 2019;33:83–91. doi: 10.1097/QAD.0000000000002037

[dataset] 5    De Veauuse Brown NF, Annor FB, Swahn MH, *et al.* Sexual Violence Experience Among Nigerian Girls and Young Women: What Are the Roles of Early Sexual Debut, Multiple Sex Partnerships, and Traditional Gender Role Beliefs? *J Interpers Violence*. 2022;37:NP2747–67.

[dataset] 6    Sardinha L, Maheu-Giroux M, Stöckl H, *et al.* Global, regional, and national prevalence estimates of physical or sexual, or both, intimate partner violence against women in 2018. *The Lancet*. 2022;399:803–13.

[dataset] 7　　World Health Organization. *Violence against women prevalence estimates, 2018:*
*global, regional and national prevalence estimates for intimate partner violence against women*
*and global and regional prevalence estimates for non-partner sexual violence against women.*
World Health Organization 2021.

[dataset] 8　　Annor FB, Chiang LF, Oluoch PR, *et al.* Changes in prevalence of violence and risk
factors for violence and HIV among children and young people in Kenya: a comparison of the 2010
and 2019 Kenya Violence Against Children and Youth Surveys. *The Lancet Global Health*.
2022;10:e124–33.

[dataset] 9　　Brown R, Kismödi E, Khosla R, *et al.* A sexual and reproductive health and rights
journey: from Cairo to the present. *Sexual and Reproductive Health Matters*. 2019;27:326–8. doi:
10.1080/26410397.2019.1676543

[dataset] 10　　Imo CK, Odimegwu CO, De Wet-Billings N. Women's attitudes towards negotiating
safe sexual practices in Nigeria: Do family structure and decision-making autonomy play a role?
*BMC Women's Health*. 2022;22:16.

[dataset] 11　　Mabaso M, Sokhela Z, Mohlabane N, *et al.* Determinants of HIV infection among
adolescent girls and young women aged 15-24 years in South Africa: a 2012 population-based
national household survey. *BMC Public Health*. 2018;18:183–183. doi: 10.1186/s12889-018-5051-
3

[dataset] 12　　Seidu A-A, Aboagye RG, Okyere J, *et al.* Women's autonomy in household decision-
making and safer sex negotiation in sub-Saharan Africa: An analysis of data from 27 Demographic
and Health Surveys. *SSM - Population Health*. 2021;14:100773.

[dataset] 13　　World Health Organization. *Consolidated guidelines on HIV testing services, 2019*.
Geneva, Switzerland: World Health Organization 2020.
https://apps.who.int/iris/bitstream/handle/10665/336323/9789241550581-eng.pdf

[dataset] 14　　Ryan S, Hahn E, Rao A, *et al.* The impact of HIV knowledge and attitudes on HIV
testing acceptance among patients in an emergency department in the Eastern Cape, South Africa.
*BMC Public Health*. 2020;20:1066.

[dataset] 15　　Scott-Sheldon LA, Carey MP, Carey KB, *et al.* HIV testing is associated with increased
knowledge and reductions in sexual risk behaviours among men in Cape Town, South Africa. *null*.
2013;12:195–201.

[dataset] 16　　Sanga ES, Lerebo W, Mushi AK, *et al.* Linkage into care among newly diagnosed HIV-
positive individuals tested through outreach and facility-based HIV testing models in Mbeya,
Tanzania: a prospective mixed-method cohort study. *BMJ Open*. 2017;7:e013733.

[dataset] 17　　Ajayi AI, Awopegba OE, Adeagbo OA, *et al.* Low coverage of HIV testing among
adolescents and young adults in Nigeria: Implication for achieving the UNAIDS first 95. *PLOS*
*ONE*. 2020;15:e0233368. doi: https://doi.org/10.1371/journal.pone.0233368

[dataset] 18　　Bekele YA, Fekadu GA. Factors associated with HIV testing among young females;
further analysis of the 2016 Ethiopian demographic and health survey data. *PLOS ONE*.
2020;15:e0228783. doi: 10.1371/journal.pone.0228783

[dataset] 19　　MacCarthy S, Saya U, Samba C, *et al.* "How am I going to live?": exploring barriers to
ART adherence among adolescents and young adults living with HIV in Uganda. *BMC Public*
*Health*. 2018;18:1158. doi: 10.1186/s12889-018-6048-7

[dataset] 20    MacKellar DA, Hou S-I, Whalen CC, *et al.* Reasons for Not HIV Testing, Testing
Intentions, and Potential Use of an Over-the-Counter Rapid HIV Test in an Internet Sample of Men
Who Have Sex With Men Who Have Never Tested for HIV. *Sexually Transmitted Diseases*.
2011;38.

[dataset] 21    Sanga Z, Kapanda G, Msuya S, *et al.* Factors influencing the uptake of Voluntary HIV
Counseling and Testing among secondary school students in Arusha City, Tanzania: a cross
sectional study. *BMC Public Health*. 2015;15:452.

[dataset] 22    Grundy J, Dakulala P, Wai K, *et al.* Independent State of Papua New Guinea Health
System Review. World Health Organization. Regional Office for South-East Asia 2019.

[dataset] 23    World Health Organization. HIV data and statistics in the Western Pacific: HIV in the
Region. 2022. https://www.who.int/westernpacific/health-topics/hiv-aids/regional-data

[dataset] 24    UNAIDS. Country Factsheet: HIV and AIDS Estimates 2022. Country: Papua New
Guinea. 2023. https://www.unaids.org/en/regionscountries/countries/papuanewguinea (accessed 26
May 2023)

[dataset] 25    Gray N, Azzopardi P, Kennedy E, *et al.* Improving Adolescent Reproductive Health in
Asia and the Pacific: Do We Have the Data? A Review of DHS and MICS Surveys in Nine
Countries. *Asia Pac J Public Health*. 2013;25:134–44. doi: 10.1177/1010539511417423

[dataset] 26    Robin Klaver, Jennifer Ruth Coe. Barriers to healthcare for female patients in Papua
New Guinea. *BMJ Case Reports*. 2018;2018:bcr-2017-224005.

[dataset] 27    Redman-MacLaren ML, Mills J, Tommbe R, *et al.* Women and HIV in a moderate
prevalence setting: an integrative review. *BMC Public Health*. 2013;13:552. doi: 10.1186/1471-
2458-13-552

[dataset] 28    Eves R. Marital sexual violence and conjugality in highlands Papua New Guinea.
*Culture, Health & Sexuality*. 2020;1–15.

[dataset] 29    Hinton R, Earnest J. The right to health: Overcoming inequalities and barriers to
women's health in Papua New Guinea. *Women's Studies International Forum*. 2010;33:180–7.

[dataset] 30    Shih P, Worth H, Travaglia J, *et al.* 'Good culture, bad culture': polygyny, cultural
change and structural drivers of HIV in Papua New Guinea. *Culture, Health & Sexuality*.
2017;19:1024–37. doi: 10.1080/13691058.2017.1287957

[dataset] 31    Kassens AL, van der Meulen Rodgers Y. Health and Distance to Healthcare in Papua
New Guinea. In: Batabyal AA, Higano Y, Nijkamp P, eds. *Disease, Human Health, and Regional
Growth and Development in Asia*. Singapore: Springer Singapore 2019:203–29.
https://doi.org/10.1007/978-981-13-6268-2_10

[dataset] 32    National AIDS Council Secretariat (NACS). PNG National STI and HIV Strategy 2018-
2022. 2018.

[dataset] 33    Gare J, Kelly-Hanku A, Ryan CE, *et al.* Factors Influencing Antiretroviral Adherence
and Virological Outcomes in People Living with HIV in the Highlands of Papua New Guinea.
*PLoS One*. 2015;10:e0134918–e0134918. doi: 10.1371/journal.pone.0134918

[dataset] 34    Kelly A, Worth H, Man N, *et al.* Barriers and Facilitators for Adherence to
Antiretroviral Therapy in Papua New Guinea. *Current HIV Research*. 2010;8:630–7. doi:
10.2174/157016210794088191

[dataset] 35    Mitchell E, Hakim A, Nosi S, *et al.* A socio-ecological analysis of factors influencing
HIV treatment initiation and adherence among key populations in Papua New Guinea. *BMC Public*
*Health*. 2021;21:2003.

[dataset] 36    Romero-Estudillo E, González-Jiménez E, Mesa-Franco MC, *et al.* Gender-based
differences in the high-risk sexual behaviours of young people aged 15-29 in Melilla (Spain): a
cross-sectional study. *BMC Public Health*. 2014;14:745.

[dataset] 37    Marianne Johansson Jørgensen, Helle Terkildsen Maindal, Kaj Sparle Christensen, *et al.*
Sexual behaviour among young Danes aged 15–29 years: a cross-sectional study of core indicators.
*Sex Transm Infect*. 2015;91:171.

[dataset] 38    Raiford JL, Herbst JH, Carry M, *et al.* Low Prospects and High Risk: Structural
Determinants of Health Associated with Sexual Risk Among Young African American Women
Residing in Resource-Poor Communities in the South. *American Journal of Community*
*Psychology*. 2014;54:243–50.

[dataset] 39    Frescura L, Godfrey-Faussett P, Feizzadeh A. A, *et al.* Achieving the 95 95 95 targets
for all: A pathway to ending AIDS. *PLOS ONE*. 2022;17:e0272405.

[dataset] 40    National Statistics Office (NSO) [PNG], ICF. Papua New Guinea Demographic and
Health Survey 2016-18. Port Moresby, Papua New Guinea & Rockville, Maryland, USA: NSO and
ICF 2019. https://dhsprogram.com/methodology/survey/survey-display-499.cfm

[dataset] 41    Asaolu IO, Gunn JK, Center KE, *et al.* Predictors of HIV Testing among Youth in Sub-
Saharan Africa: A Cross-Sectional Study. *PLOS ONE*. 2016;11:e0164052. doi:
https://doi.org/10.1371/journal.pone.0164052

[dataset] 42    Zhang Z. Model building strategy for logistic regression: purposeful selection. *Annals of*
*Translational Medicine; Vol 4, No 6 (March 25, 2016): Annals of Translational Medicine (Toward*
*Precision Medicine in Neurological Diseases)*. Published Online First: 2016. doi:
http://dx.doi.org/10.21037/atm.2016.02.15

[dataset] 43    Archer KJ, Lemeshow S, Hosmer DW. Goodness-of-fit tests for logistic regression
models when data are collected using a complex sampling design. *Computational Statistics & Data*
*Analysis*. 2007;51:4450–64.

[dataset] 44    West BT. Statistical and methodological issues in the analysis of complex sample
survey data: Practical guidance for trauma researchers. *Journal of Traumatic Stress*. 2008;21:440–
7.

[dataset] 45    Sibanda EL, Neuman M, Tumushime M, *et al.* Community-based HIV self-testing: a
cluster-randomised trial of supply-side financial incentives and time-trend analysis of linkage to
antiretroviral therapy in Zimbabwe. *BMJ Glob Health*. 2021;6:e003866.

[dataset] 46    Kyaddondo D, Wanyenze RK, Kinsman J, *et al.* Home-based HIV counseling and
testing: Client experiences and perceptions in Eastern Uganda. *BMC Public Health*. 2012;12:966.

[dataset] 47    Sekandi JN, Sempeera H, List J, *et al.* High acceptance of home-based HIV counseling
and testing in an urban community setting in Uganda. *BMC Public Health*. 2011;11:730.

[dataset] 48    Johnson CC, Kennedy C, Fonner V, *et al.* Examining the effects of HIV self-testing
compared to standard HIV testing services: a systematic review and meta-analysis. *Journal of the*
*International AIDS Society*. 2017;20:21594.

[dataset] 49    Gutiérrez JP, Trossero A. Socioeconomic inequalities in HIV knowledge, HIV testing,
and condom use among adolescent and young women in Latin America and the Caribbean. *Rev*
*Panam Salud Publica*. 2021;45:e47.

[dataset] 50    Mabaso M, Sokhela Z, Mohlabane N, *et al.* Determinants of HIV infection among
adolescent girls and young women aged 15-24 years in South Africa: a 2012 population-based
national household survey. *BMC Public Health*. 2018;18:183–183. doi: 10.1186/s12889-018-5051-
3

[dataset] 51    Mandiwa C, Namondwe B, Munthali M. Prevalence and correlates of comprehensive
HIV/AIDS knowledge among adolescent girls and young women aged 15–24 years in Malawi:
evidence from the 2015–16 Malawi demographic and health survey. *BMC Public Health*.
2021;21:1508.

[dataset] 52    Musekiwa A, Silinda P, Bamogo A, *et al.* Prevalence and factors associated with self-
reported HIV testing among adolescent girls and young women in Rwanda: evidence from 2019/20
Rwanda Demographic and Health Survey. *BMC Public Health*. 2022;22:1281.

[dataset] 53    Sherafat-Kazemzadeh R, Gaumer G, Hariharan D, *et al.* Between a Rock and a Hard
Place: How poverty and lack of agency affect HIV risk behaviors among married women in 25
African countries: A cross-sectional study. *J Glob Health*. 2021;11:04059.

[dataset] 54    Zegeye B, Adjei NK, Ahinkorah BO, *et al.* HIV testing among women of reproductive
age in 28 sub-Saharan African countries: a multilevel modelling. *International Health*.
2023;15:573–84.

[dataset] 55    Schierl T, Tanaka LF, Klug SJ, *et al.* The Association of Women's Empowerment with
HIV-Related Indicators: A Pooled Analysis of Demographic and Health Surveys in Sub-Saharan
Africa. *Journal of Epidemiology and Global Health*. 2023;13:816–24.

[dataset] 56    Worku MG, Teshale AB, Tesema GA. Prevalence and associated factors of HIV testing
among young (15–24) women in eastern Africa: a multilevel analysis of demographic health survey
data (2008-2018). *Archives of Public Health*. 2022;80:117.

[dataset] 57    Eves R. Masculinity matters: Men, gender-based violence and the AIDS epidemic in
Papua New Guinea. *Civic Insecurity: Law, Order and HIV in Papua New Guinea*. 2010;1:47–79.

[dataset] 58    McCall C. In Papua New Guinea, HIV/AIDS-related stigma is still rife. *The Lancet*.
2016;387:215–6.

[dataset] 59    Teklehaimanot HD, Teklehaimanot A, Yohannes M, *et al.* Factors influencing the
uptake of voluntary HIV counseling and testing in rural Ethiopia: a cross sectional study. *BMC*
*Public Health*. 2016;16:239.

[dataset] 60    Diress G, Ahmed M, Adane S, *et al.* Barriers and Facilitators for HIV Testing Practice
Among Ethiopian Women Aged 15-24 years: Analysis of the 2016 Ethiopian Demographic and
Health Survey. *HIV/AIDS - Research and Palliative Care*. 2020;12:963–70.

[dataset] 61    Alem AZ, Liyew AM, Guadie HA. Spatial pattern and associated factors of HIV testing
and counselling among youths (15–24 years) in Ethiopia. *BMC Public Health*. 2021;21:644.

[dataset] 62    Omona K, Ssuka JK. Early sexual debut and associated factors among adolescents in
Kasawo Sub-county, Mukono district, Uganda. *Cogent Public Health*. 2023;10:2183561.

[dataset] 63    Ferede TA, Muluneh AG, Wagnew A, *et al.* Prevalence and associated factors of early
sexual initiation among youth female in sub-Saharan Africa: a multilevel analysis of recent
demographic and health surveys. *BMC Women's Health*. 2023;23:147.

[dataset] 64    Millanzi WC, Osaki KM, Kibusi SM. Attitude and prevalence of early sexual debut and
associated risk sexual behavior among adolescents in Tanzania; Evidence from baseline data in a
Randomized Controlled Trial. *BMC Public Health*. 2023;23:1758.

[dataset] 65    Kamire V, Magut F, Khagayi S, *et al.* HIV Risk Factors and Risk Perception Among
Adolescent Girls and Young Women: Results From a Population-Based Survey in Western Kenya,
2018. *J Acquir Immune Defic Syndr*. 2022;91:17–25.

[dataset] 66    Colombini M, James C, Ndwiga C, *et al.* The risks of partner violence following HIV
status disclosure, and health service responses: narratives of women attending reproductive health
services in Kenya. *Journal of the International AIDS Society*. 2016;19:20766.

[dataset] 67    Mengwai K, Madiba S, Modjadji P. Low Disclosure Rates to Sexual Partners and
Unsafe Sexual Practices of Youth Recently Diagnosed with HIV; Implications for HIV Prevention
Interventions in South Africa. *Healthcare*. 2020;8. doi: 10.3390/healthcare8030253

[dataset] 68    Pulerwitz J, Michaelis A, Weiss E, *et al.* Reducing HIV-Related Stigma: Lessons
Learned from Horizons Research and Programs. *Public Health Rep*. 2010;125:272–81.

[dataset] 69    Malli IA, Kabli BA, Alhakami LA. Sexually Transmitted Diseases among Saudi
Women: Knowledge and Misconceptions. *International Journal of Environmental Research and*
*Public Health*. 2023;20. doi: 10.3390/ijerph20064858

[dataset] 70    Avuvika E, Masese LN, Wanje G, *et al.* Barriers and Facilitators of Screening for
Sexually Transmitted Infections in Adolescent Girls and Young Women in Mombasa, Kenya: A
Qualitative Study. *PLoS One*. 2017;12:e0169388.

[dataset] 71    Denison HJ, Bromhead C, Grainger R, *et al.* Barriers to sexually transmitted infection
testing in New Zealand: a qualitative study. *Australian and New Zealand Journal of Public Health*.
2017;41:432–7.

[dataset] 72    Newton-Levinson A, Leichliter JS, Chandra-Mouli V. Sexually Transmitted Infection
Services for Adolescents and Youth in Low- and Middle-Income Countries: Perceived and
Experienced Barriers to Accessing Care. *Journal of Adolescent Health*. 2016;59:7–16.

**Supplementary materials**

STROBE Checklist

**Footnotes**

**Contributors:** MKM conceptualized and designed the study. MKM and FWK conducted statistical

analyses. MKM drafted the original manuscript. FWK contributed to the writing and editing of the

manuscript. MKM and FWK reviewed and edited the manuscript. All the authors reviewed and

approved the final version of the manuscript. MKM is the guarantor and accepts full responsibility for

the work.

**Funding:** The authors have not declared a specific grant for this research from any funding agency in
the public, commercial, or not-for-profit sectors.
598
599    **Competing interests:** None declared.
600
601    **Patient consent for publication:** Not required.
602
603    **Ethics approval:** The 2016–2018 PNGDHS protocol was reviewed and approved by the Institutional
604    Review Board of ICF and Medical Research Advisory Committee of PNG. Informed consent was
605    obtained from all participants before the interviews were conducted. For this study, a formal request to
606    use the raw data was obtained from the DHS program. Permission was granted by the DHS program to
607    access the data, which was anonymized for analysis.
608
609    **Provenance and peer review:** Not commissioned; externally peer reviewed.
610
611    **Data availability statement:** The 2016–2018 PNGDHS data are publicly available in a public, open-
612    access repository. Data can be obtained upon a registration-access request from the DHS Program
613    website at https://dhsprogram.com/data/available-datasets.cfm.