Inferring Alzheimer's disease pathologic traits from clinical measures in living adults

Author(s):

Jingjing Yang, PhD[1]; Xizhu Liu[2]; Shahram Oveisgharan, MD[3]; Andrea R. Zammit, PhD[3]; Sukriti
Nag, MD, PhD,[3] David A Bennett, MD[3]; Aron S Buchman, MD[3]

Corresponding Author:

Jingjing Yang, jingjing.yang@emory.edu

Affiliation Information for All Authors: 1. Center for Computational and Quantitative
Genetics, Department of Human Genetics, Emory University School of Medicine, 615 Michael
St, Atlanta, GA, 30322, USA; 2. Department of Biostatistics, Yale University School of Public
Health, New Haven, CT, USA; 3. Rush Alzheimer's Disease Center, Rush University Medicine
Center, Chicago, IL, 60612, USA.

ABSTRACT

Background and Objectives: Develop imputation models using clinical measures to infer
Alzheimer's disease neuropathologic changes (AD-NC) in living adults to identify adults at risk
for Alzheimer's disease (AD).

Methods: We used clinical and postmortem data of two prospective cohort studies — Memory
and Aging Project (MAP) and Religious Orders Study (ROS). We used generalized linear
regression models with Elastic-Net penalty to train imputation models of AD-NC traits (β-
Amyloid, tau tangles, global AD pathology, and NIA-Reagan), in MAP decedents using clinical
measures collected at last visit as predictors. ROS cohort was used as an independent
validation and test data. We validated these models in ROS decedents and applied the models
to baseline clinical data of ROS participants to infer baseline AD-NC traits. Baseline clinical
data were collected an average of 8 years before last follow-up. We used Cox proportional
hazard models to test if inferred baseline AD-NC traits predicted incident AD dementia (ADD).
In addition, two-sample t-tests were used to examine if inferred baseline AD-NC traits
predicted adults with high risk of pathologic AD profiled at death.

Results: By applying imputation models to clinical measures collected at last visit in ROS to
validate the imputation models, we obtained predicted R^2 as 0.188 for β-Amyloid, 0.316 for tau
tangles, and 0.262 for global AD pathology. Prediction area under the receiver operating
characteristic curve (AUC) for the dichotomous NIA-Reagan was 0.765. All four inferred AD-
NC traits at last visit strongly discriminated postmortem NIA-Reagan status (p-values $< 10^{-28}$).
The inferred baseline levels of all four AD-NC traits predicted ADD, with higher accuracies for
predicting ADD in Year 3 (AUC ranging in 0.861 – 0.919) versus Year 5 (AUC 0.842 – 0.896),
and the highest accuracy was obtained using inferred NIA-Reagan status. The inferred
baseline levels of all four AD-NC traits significantly discriminate individuals with postmortem
pathologic AD (all p-values $< 1.5 \times 10^{-7}$).

Conclusions: Inferred baseline levels of AD-NC traits derived from clinical measures
discriminate adults at risk for ADD and pathologic AD profiled at death. Further studies are
needed to determine if repeated measures of inferred AD-NC traits can be used to monitor the
accumulation of AD-NC traits during the prolonged course of AD.

Keywords & Subject Ares: Alzheimer's disease (AD), Cohort studies, Computational
imputation model, Alzheimer's disease neurologic change (AD-NC)

INTRODUCTION

Alzheimer's disease dementia (ADD) is a chronic progressive disorder that develops over years before manifesting impaired cognition. In its earliest asymptomatic stage, Alzheimer's disease neuropathologic changes (AD-NC) accumulate in adults with normal cognition[1], and higher levels of AD-NC during early stages are associated with an increased risk of ADD.[2-6] Yet, ADD is commonly caused by mixed-brain pathologies, so clinical risk models identifying adults at risk for clinical ADD do not inform on which AD-NC traits underlie the predicted risk of ADD. Thus, developing a model that can infer the level of AD-NC traits in living adults with normal cognition can identify adults with specific high levels of AD-NC traits being at risk of ADD, which could facilitate early targeted treatments of specific AD-NC traits.[7-14]

Currently, indices of AD-NC can only be obtained after death via autopsy. Yet, AD-NC accumulate over years before cognitive impairment manifests[15]. Recent efforts to identify AD-NC before impaired cognition have focused on using brain imaging or CSF AD-NC biomarkers[16]. Yet, these biomarkers are not widely available due to their costs, invasiveness, and difficultly to deploy at scale. This explains in part why current aging research is focused on identifying blood biomarkers for AD-NC traits that can be used in general population.[17-21] A computational model that can infer AD-NC traits based on clinical measures in living adults is an understudied alternative that if successful could be deployed widely and might also be combined with fluid biomarkers to improve identification of adults at risk of ADD.

Machine learning techniques such as penalized regression have been employed to impute missing data or estimate data that are difficult to measure directly in many areas of biomedical research[22]. Prior efforts to impute AD-NC traits have used a combination of brain imaging and clinical measures to infer AD-NC traits.[23] However, brain imaging is expensive and not widely available. In contrast, a validated imputation model can be conveniently applied to infer AD-NC traits in any living adults with the same clinical measures used in the imputation model.

To develop imputation models for AD-NC traits, we used clinical and postmortem data from two harmonized, independent cohort studies — Rush Memory and Aging Project (MAP) and the Religious Orders Study (ROS).[24] First, we trained imputation models to infer AD-NC traits using clinical measures at the last visit in MAP decedents and validated model performance in a second independent cohort, ROS decedents. Then we applied the imputation models to clinical measures collected in ROS participants at study entry to infer baseline levels of AD-NC traits. The baseline is on average 8 years before the last follow up in MAP and ROS. We examined if inferred baseline AD-NC traits predicted future ADD and pathologic AD profiled at death.

METHODS

We use the STOBE cohortreporting guidelines.[25]

Participants

Participants were community-dwelling older adults enrolled without known clinical dementia in one of two ongoing cohort studies of chronic conditions of aging, MAP (n=1179, ~500 autopsied) and ROS (n=1103, ~600 autopsied). Both cohorts employed a harmonized data collection battery administered by the same research assistants facilitating joint analyses. We

95   only analyzed samples that had at least two follow-ups. At study entry, MAP and ROS have
96   1742 samples that had no cognitive impairment (NCI) and 540 samples that had mild cognitive
97   impairment (MCI) (**Table S1**). The duration of follow-up for participants ranged from 2 to 26
98   years, with an average of 8 years and standard deviation 5.42 (**Fig S1; Tables S1-S2**).

Assessment of Alzheimer's Disease Neuropathologic Traits (AD-NC)

After death, the brain was removed and hemisected following standard procedure, as
previously described[24]. Tissue blocks were dissected from predetermined regions and used for
making the postmortem diagnosis of pathologic AD. Structured autopsy collected indices of
neurodegenerative pathologies that were collected blinded to all prior clinical and cognitive
data. This study focuses on studying three continuous measures summarizing the burden of
AD-NC in five cortical regions — β-amyloid, tau tangles, and global AD pathology (i.e., a
summary of neuritic and diffuse plaques), and one dichotomous NIA-Reagan status (**Table
S2**). Measures of β-amyloid and tau tangles were square-root transformed to improve their
asymptotic normality as previously reported[26, 27]. NIA-Reagan status was also taken as
diagnosis of pathologic AD based on modified AD NIA-Reagan criteria[28].

Assessment of Cognition and Cognitive Status

A structured exam was administered annually. The neuropsychological battery included 19
tests that assessed five domains of cognitive ability. Raw test scores were standardized per
test using baseline means and standard deviations (SDs) of both cohorts; the resulting Z-
scores were then averaged across 17 of these cognitive tests to derive a summary composite
global cognition score and scores for five cognitive abilities including: episodic memory,
semantic memory, working memory, visuospatial abilities, and perceptual speed. A
neuropsychologist and a neurologist with expertise in dementia reviewed annual cognitive
testing and classified cognitive status of participants as NCI, MCI, ADD, or other dementias
according to established criterion of clinical diagnosis of Alzheimer's disease[29].

Other Clinical Covariates

Five groups of clinical variables (a total of 57) were considered as predictors for developing
imputation models of AD-NC traits (**Tables S2-S5**) — i) *Common Alzheimer's disease risk
factors* including age, sex, education, Mini Mental State Exam (MMSE) score[30], and *APOE* E4
allele; ii) *Health measures* such as blood pressure, depression, and cardiovascular diseases;
iii) *Medication usage* such as lipid lowering medications, and antidepressants; iv) *Variables
uniquely profiled by MAP/ROS* such as composite global cognition scores based on 17
cognitive tests, self-reported physical activity and social network size[24]; v) *Motor and sleep
metrics* such as dexterity, hand strength, gait function, four parkinsonian signs[31], and four self-
reported variables about sleep quality and duration[32].

These 57 clinical variables were selected by excluding variables with a high proportion of
missing values (>20%) and those that were highly correlated with other selected clinical
variables (correlation>0.95). Missing values of longitudinal variables were imputed from the
corresponding values at participants' nearest visits. A total of 345 individuals with missing
cross-sectional variables were excluded, leaving n=1179 MAP and n=1103 ROS participants

for the analyses in this study, including both living participants and decedents. A heat map
showing the correlations of the clinical variables examined in this study is shown in **Fig S2**.

Analytic Approach

Training and validating imputation models to infer AD-NC

Training imputation models in MAP. We trained an imputation model per AD-NC trait using
clinical variables obtained at the last visit before death as predictors, employing a generalized
linear regression model with Elastic-Net penalty (GLM-EN) (**Fig 1A**). A total of 57 clinical
variables were considered as predictors for imputation models of AD-NC traits (**Tables S2-S5;**
Fig S2). By using the GLM-EN method, variable selection was implemented and potential
collinearity among clinical variables was accounted for during model training. Ten-fold cross
validation was used during model training to select Elastic-Net penalty parameters to ensure
best imputation accuracy.

Validating imputation model in ROS. Next, we validated the performance of these imputation
models by inferring AD-NC traits using clinical variables measured at the last visit proximate to
death in the 2nd independent test cohort, ROS decedents (**Fig 1A**). Prediction R^2, the squared
correlation between inferred and measured values, was used for assessing imputation
accuracy for continuous AD-NC traits. Prediction accuracy for the dichotomous NIA-Reagan
was evaluated by the predicted area under curve (AUC) values of receiver operating
characteristic curve (ROC)[33].

Applying imputation models to infer AD-NC at study baseline

Once imputation models are trained and validated, they can be applied to any living adult with
the same clinical variables measured. Thus, we applied the imputation models to the clinical
data obtained from the ROS cohort at study entry (baseline) with an average of eight years of
follow-up to infer the levels of the four AD-NC traits at baseline (**Fig 1B**). We examined if these
inferred AD-NC traits could be employed as a biomarker of AD-NC through two sets of
analyses. We examined if inferred AD-NC traits at study baseline: a) predict adults at risk for
ADD, b) predict adults at risk for pathologic AD measured at death.

Inferred AD-NC traits at study entry and incident ADD.

We fitted Cox proportional hazard models using covariates of age, sex, education, and a single
inferred baseline AD-NC trait in MAP cohort for predicting ADD, and evaluated the prediction
performance in year 3 and year 5 in ROS cohort (**Fig S3**). Our fitted Cox proportional hazard
risk prediction models[34-36] also accounted for the competing risk of death. The annual cognitive
status diagnosis and the follow up year were used to identify the first occurrence of ADD. For
each participant, the year of enrollment is considered as baseline (time 0), the year of first
diagnosis of ADD is considered as the time when the event occurs (incident ADD), and the last
visit of participants without the considered event during all follow-ups is considered as the right
censored time for living participants or the time of death for dead participants without ADD.
Sample size distributions with respect to cognitive status at baseline and the cognition event
types are shown in **Tables S1**. All Cox models were trained using data from MAP participants
and tested with data from ROS participants[37]. Both living and deceased ROS participants were
used to evaluate the prediction performance by Cox models.

We first considered all individuals including adults without dementia at baseline, and then
included only individuals with NCI at baseline. We calculated AUC to evaluate the prediction
accuracy of developing ADD in year 3 and year 5. Since Cox models provides a continuous
risk score for developing ADD, by selecting a risk score threshold corresponding the ~80%
specificity (the proportion of correctly predicted non-ADD in test ROS samples, i.e., 1 – false
positive fraction), we could compare the overall classification, sensitivity (the proportion of true
positive predictions in all test cases, i.e., true positive fraction), and accuracy (the proportion of
true discrimination of ADD in all test samples). Samples with risk scores greater than the
selected threshold were predicted as positives (i.e., ADD) or otherwise negatives (i.e., no
ADD). The sensitivity and specificity corresponding to a selected risk score threshold reflect
risk model performance at one point in the ROC curves.

Additionally, we also sequentially added the other three inferred baseline AD-NC traits into the
Cox model with covariates of age, sex, education, and inferred baseline β-amyloid. The
prediction performance of the models with additional AD-NC traits was also evaluated for
developing incident ADD in year 3 and year 5 in ROS cohort.

Inferred AD-NC traits at study entry and pathologic AD.

We evaluated the discrimination of these inferred baseline levels of AD-NC traits for pathologic
AD at the time of death, i.e., postmortem NIA-Reagan status (**Fig 1B**). These evaluations only
use ROS decedents who also had postmortem NIA-Reagan status profiled. Boxplots and two-
sample t-tests were used to evaluate the discrimination of inferred baseline AD-NC traits and
postmortem NIA-Reagan status. Scatter plots and ROC curve were used to evaluate the
consistence between inferred baseline AD-NC traits and the corresponding profiled
postmortem AD-NC traits in ROS samples.

Research Ethics and Informed Consent
Both MAP and ROS studies were approved by an Institutional Review Board of Rush
University Medical Center. All participants agreed to annual clinical evaluations and autopsy at
the time of death. Written informed consent was obtained from all study participants as was an
Anatomical Gift Act for organ donation.
210

Data Access and Availability Statement

All data analyzed in this study are de-identified and available to any qualified investigator by
submitting a request through the Rush Alzheimer's Disease Center Research Resource
Sharing Hub, https://www.radc.rush.edu, which has descriptions of the studies and available
data.

RESULTS

Training imputation models of AD-NC traits in MAP

Imputation models were trained for AD-NC traits by GLM-EN method by taking 57 clinical
variables measured at last visit before death in MAP decedents as predictors. The estimated
coefficients (i.e., effect sizes) of strongest predictors of AD-NC traits are shown in **Fig 2**,

including *APOE* E4, age, and composite cognition score. **Fig 2** illustrates that the imputation
models isolated different combinations of most predictive clinical measures to infer the levels
of each of the four AD-NC traits.

Validating imputation models of AD-NC traits in ROS

Inferred levels of AD-NC traits at last visit of ROS decedents can be generated by using the
corresponding trained imputation model per AD-NC trait and measured clinical variables at last
visit before death. Scatter plots illustrate the comparison between the inferred levels of
continuous AD-NC traits and the corresponding indices measured at autopsy in ROS
decedents (**Fig S4, A-C).** ROC plot illustrates the consistency between the inferred NIA-
Reagan status at last visit versus the profiled status by autopsy (**Fig S4D**). The prediction R^2
was 0.188 for β-Amyloid, 0.316 for tau tangles, and 0.262 for global AD pathology. AUC for the
dichotomous NIA-Reagan was 0.765 (**Table S6)**. As shown by the boxplots (**Fig S4**) of inferred
AD-NC traits at last visit with respect to pathologic AD status (i.e., NIA-Reagan status profiled
by autopsy), inferred AD-NC traits at last visit in ROS decedents discriminated profiled
pathologic AD at autopsy, with two-sample test p-values $< 10^{-28}$ for all four inferred AD-NC
traits. These results validated the inference accuracy of these trained imputation models of
AD-NC traits.

Association of inferred baseline AD-NC traits and ADD

To infer baseline levels of four AD-NC traits at study entry, we applied the validated imputation
models to clinical data collected at baseline in all available ROS participants, an average of 8
years before their last visit (n=1103) (**Fig 1B**). Cox models were fitted by using covariates of
age, sex, education, and each single inferred baseline AD-NC trait, to predict the risk of
developing ADD (**Fig S3**). Coefficient estimates of the inferred baseline AD-NC traits in the
Cox models were provided along with the corresponding p-values in **Table S7**. All inferred
baseline AD-NC traits had significant predictive coefficients with p-values $< 10^{-30}$ for
predicting the development of ADD from adults with NCI or MCI, and p-values $< 10^{-5}$ for
predicting the development of ADD from adults with only NCI. The inferred baseline NIA-
Reagan was found with the largest coefficient effect size with the most significant p-values,
compared to other three inferred baseline AD-NC traits.

The inferred baseline levels of all four AD-NC traits predicted ADD (**Fig 3, Upper Row**). For
all four AD-NC traits, model performance was higher in Year 3 (AUC ranging in 0.861 – 0.919)
versus Year 5 (AUC 0.842 – 0.896). Similar results were observed when we restricted the
analyses to the prediction of ADD in individuals with NCI at baseline (**Fig 3, Lower Row**). Of
the four AD-NC traits, inferred baseline NIA-Reagan had the highest predictivity (Year 3, AUC
0.919; Year 5, AUC 0.896) for ADD. By selecting a risk score threshold corresponding the
~80% specificity, we calculated sensitivity and accuracy based on the prediction results by all
Cox models (**Table 1**). Inferred baseline NIA-Reagan also had the highest accuracy rates
(80%) and sensitivity for predicting ADD in year 3 (0.911) and year 5 (0.829) (**Table 1**) as
compared to the other AD-NC traits.

In further analyses, we examined if adding terms for multiple inferred AD-NC traits in a single
Cox model would improve the prediction accuracy for incident ADD. As shown in **Fig S5**

(Upper Row), we observed slightly improved AUC for prediction of incident ADD in adults with
NCI or MCI, by sequentially adding inferred baseline β-amyloid, tau tangles, global AD
pathology, and NIA-Reagan. Yet, including covariates of all four AD-NC traits together did not
perform any better than using the inferred NIA-Reagan alone (**Last Column of Fig3 versus**
Fig S5). Sequentially adding inferred baseline β-amyloid, tau tangles, global AD pathology did
not increase the prediction of incident ADD in adults with only NCI (**Fig S5, Lower Row**). Still,
adding inferred baseline NIA-Reagan would improve the prediction of incident ADD in adults
with only NCI, but still not perform any better than using the inferred NIA-Reagan alone (**Last**
Column of Fig3 versus Fig S5).

Association of inferred baseline AD-NC traits with postmortem pathologic AD

By comparing inferred AD-NC traits in ROS decedents to their corresponding postmortem
measurements, we observed significant concordance as shown in scatter and ROC plots in
Fig 4. Also, inferred baseline AD-NC traits obtained in ROS decedents who had an autopsy at
the time of death, discriminated individuals with a postmortem diagnosis of pathologic AD (i.e.,
NIA-Reagan status profiled by autopsy), with significant p-values $< 1.5 \times 10^{-7}$ by two-sample t-
tests (**Fig 5**). The postmortem pathologic AD and AD-NC traits were profiled at death an
average of 8 years after baseline.

DISCUSSION

We developed and validated imputation models of four AD-NC traits based on clinical measures
in living adults obtained at their last visit about one year before death. To demonstrate the utility
of inferred AD-NC traits, we applied these models to clinical measures obtained in living adults
at their initial clinical assessment at study entry about eight years before death. Inferred baseline
AD-NC traits showed good accuracy in predicting adults at risk for developing clinical ADD 3
and 5 years during follow-up and also discriminated adults at risk for pathologic AD profiled at
the time of death. These data suggest that imputation models to infer AD-NC traits based on
clinical measures may have potential to provide a low cost, non-invasive classifier for clinical
Alzheimer disease dementia. A classifier based on clinical measures could be deployed in
diverse populations and areas with limited medical infrastructure. Further work will be needed to
determine if repeated measures of inferred AD-NC traits might be used as a reliable proxy for
monitoring the accumulation of AD-NC traits during the prolonged course of Alzheimer's disease.
This approach might be extended to infer the presence of other AD related dementia (ADRD)
pathologies that commonly accumulate in aging brains.

Tools detecting preclinical Alzheimer's disease, i.e., adults with normal cognition at risk for future
ADD, can benefit from early treatments that are crucial to prevent symptomatic ADD. To fill this
gap, current research efforts have focused on identifying AD biomarkers that correlate with
postmortem indices of AD-NC traits.[7-14, 38] Yet, imaging or CSF biomarkers are not widely
available due to their costs, invasiveness, and difficultly to deploy on a large scale. Aging
research is currently focused on identifying blood biomarkers to fill this gap.[18-21] Developing an
effective AD classifier based on clinical measures alone is an understudied alternative.

The concept of fitting imputation models has been used in diverse disciplines to infer data that
are not easily available.[39-44] The imputation model learns the predictive information of
postmortem AD-NC traits like tau tangles from clinical indices at last visit before decedents

undergoing autopsy. The imputation model works by mathematically "explaining" the variation
of observed AD-NC traits in autopsied adults by their equivalence in clinical indices. Efforts to
impute AD-NC traits using clinical and brain imaging indices have been reported, but brain
imaging is costly and not widely available.[45, 46] Recent work, that compared tau and amyloid
PET brain imaging to AD indices measured at autopsy, suggest that current imaging may not
reliably detect the early stages of AD pathology.[47, 48]

Clinical measures prior to death and postmortem AD-NC traits are needed to develop
imputation models for AD-NC traits. Obtaining a large range of clinical measures within a year
prior of death in large numbers of older adults as well as a structured autopsy is difficult. Thus,
two large cohorts like MAP or ROS with the same clinical and postmortem indices of AD-NC
traits are rare, and may explain in part the paucity of previous studies that have tried to
develop imputation models for AD-NC traits based on clinical measurements alone. Thus, the
current study provides novel data demonstrating the feasibility of developing imputation
models to infer AD-NC traits from clinical measures obtained at the last visit about one year
prior to death.

Once an imputation model has been validated, it can be applied to infer AD-NC traits in any
older adult with similar clinical measures.[49] To demonstrate the effectiveness of the imputation
models that were developed, we applied these models to baseline clinical measures obtained
at study entry with an average of 8 years before last visit, in an independent test cohort. As
would be expected for an effective AD classifier, adults without dementia at study entry who
had higher inferred baseline levels of AD-NC traits had a higher risk of subsequent ADD. Cox
proportional hazard regression models using covariates of inferred baseline AD-NC traits,
along with age at visit, sex, and education, showed good accuracy for predicting ADD three
and five years after baseline. Inferred baseline AD-NC traits also predicted adults with a high
risk of pathologic AD profiled on average eight years later at death. That is, while the
imputation models were developed based on clinical measures at last visit prior to death,
inferred levels of AD-NC traits based on clinical data obtained almost a decade before death
showed good predictions of clinical ADD during the course of the study as well as postmortem
measurements of AD-NC traits. These promising initial results will require further work to
validate the accuracy of repeated measures of inferred AD-NC traits in living adults before they
can be employed to monitor the accumulation of AD-NC pathologic traits or the efficacy of
clinical drug trials.

This study has several limitations. First, participants were predominantly Americans of
European descent and have higher than average levels of education, so our findings will need
to be replicated in more diverse populations. Second, this study only inferred AD-NC traits.
Since most older adults show mixed-brain pathologies, further work is needed to extend
imputation models to accommodate the presence of combinations of multiple AD/ADRD
pathologies. Third, brain imaging, as well as serum or fluid biomarkers were not examined and
might enhance predictions of the inferred AD-NC traits. Further validation using brain imaging
or fluid AD-NC biomarkers are needed. Last, the current study used diverse clinical predictors,
many that are not available outside the research setting. Further work is needed to identify a
parsimonious set of clinical predictors that are widely available to enhance the use of this
approach in diverse populations and geographic locations that have limited medical personnel
and resources. Despite these limitations, this study is best conceptualized as an important first

step highlighting the potential of developing inferred AD-NC traits based on clinical measures
in living adults.

The study has several strengths that lend confidence for the current findings. All subjects were
recruited from the community, underwent an annual detailed clinical evaluation, and were
without dementia based on their clinical assessment at study entry. Large numbers of men and
women underwent annual assessments, and follow-up rates were very high (~90%) with a
medium of 8 years follow up. Uniform and structured procedures were employed for both the
collection of clinical measures and postmortem AD-NC traits. An important strength of the
current study design is that we evaluated model performance in a second cohort, independent
of the training cohort, that employed similar staff and data collection procedures[50].

363
364
365

Acknowledgments

We are deeply indebted to all participants who contributed their data and agreed to autopsy at
the time of their death. We thank the Rush Alzheimer's Disease Center staff for their efforts.

Study Funding: This work was supported by the National Institute of Health R35GM138313,
P30AG10161, P30AG72975, K01AG054700, R01AG15819, R01AG17917, R01AG56352;
R01AG79133, the Illinois Department of Public Health; and the Robert C. Borwell Endowment
Fund. The funding organizations had no role in the design or conduct of the study; collection,
management, analysis, or interpretation of the data; or preparation, review, or approval of the
manuscript.

Disclosures: There are no disclosures for any of the authors.
378
379

380 **Supplemental:** One supplementary data file with 7 supplement tables and 5 supplement
381 figures are provided
382

Table 1. Prediction accuracy (with 95% confidence interval) and sensitivity with respect to selected risk score thresholds that ensure ~80% specificity by Cox models using a single inferred AD-NC trait. Values in this table are reflecting the Cox risk model prediction performance at one point in the ROC curves as shown in **Fig 3**, with corresponding risk score thresholds. Samples with predicted risk scores greater than the selected threshold were considered as Predicted Positives (incident ADD), otherwise Predicted Negatives (not developing ADD).

		β-Amyloid		Tau tangles		Global AD Pathology		NIA-Reagan	
		Y3	Y5	Y3	Y5	Y3	Y5	Y3	Y5
NCI/MCI -> ADD	Accuracy[a] (95% CI)	0.798 (0.773, 0.822)	0.788 (0.763, 0.813)	0.802 (0.777, 0.825)	0.795 (0.771, 0.818)	0.801 (0.775, 0.823)	0.796 (0.771, 0.819)	0.809 (0.784, 0.831)	0.803 (0.778, 0.826)
	Sensitivity[b]	0.772	0.699	0.822	0.756	0.797	0.764	0.911	0.829
	Specificity[c]	0.800	0.800	0.800	0.800	0.800	0.800	0.800	0.800
NCI -> ADD	Accuracy[a] (95% CI)	0.796 (0.768, 0.823)	0.792 (0.763, 0.7819)	0.794 (0.765, 0.821)	0.791 (0.761, 0.818)	0.795 (0.767, 0.823)	0.793 (0.764, 0.820)	0.804 (0.775, 0.830)	0.805 (0.777, 0.832)
	Sensitivity[b]	0.650	0.634	0.550	0.609	0.600	0.658	0.950	0.902
	Specificity[c]	0.800	0.800	0.800	0.800	0.800	0.800	0.800	0.800

a. Accuracy= (# True Positive Predictions + # True Negative Predictions) / (# of test samples)

b. Sensitivity = (# True Positive Predictions) / (# Positives in test samples) = True Positive Fraction

c. Specificity = (# True Negative Predictions) / (# Negatives in test samples) = 1−False Positive Fraction

Main Figures

Fig 1. Study design used to develop and validate imputation models that infer the burden of AD-NC traits based on clinical measures in living adults. This figure provides an overview contrasting the complementary components of this study. **A.** We trained imputation models for AD-NC traits using clinical data from the last visit before death in MAP decedents that underwent autopsy. Then we validated these models in a 2nd independent sample of ROS decedents. **B.** Then to show the utility of how these imputation models could be used in any adults with the requisite clinical measures, we applied the validated imputation models to clinical data obtained at the initial clinical assessment of ROS participants to infer AD-NC traits at the ROS baseline visit about a decade before the last follow-up assessment.

Fig 2. **Different combinations of clinical measures obtained at the last visit before death are used by imputation models to infer each of the four AD-NC traits.** GLM-EN method was used to train imputation models for each of the four AD-NC traits. Effect sizes (beta) of the standardized predictors in the prediction model with |beta| > 0.01 are plotted in the figure to highlight the differences and common and AD-NC trait specific measures in the four imputation models used to infer AD-NC traits.

Fig 3. Inferred AD-NC traits based on clinical measures at study entry for all ROS participants predict Alzheimer's Disease Dementia (ADD). *We applied validated imputation models to clinical measures collected at study entry in ROS participants to infer 4 AD-NC pathologic traits. Row 1 shows that prediction of ADD for the inferred burden of each of the four AD-NC pathologic traits in adults without dementia at study entry. Row 2 shows results for inferred burden of each of the four AD-NC pathologic traits in adults with NCI at baseline. Inferred NIA-Reagan was found with best predictivity for the incidence of ADD in year 3 and year 5.*

Fig 4. Inferred baseline AD-NC traits using clinical measures at the time of study entry predict the pathologic AD profiled at autopsy in ROS decedents. *Scatter plots for continuous AD pathologic traits and ROC curve for binary NIA-Reagan show that the inferred AD-NC traits based on clinical measures at study entry were concordant with their corresponding measures profiled at the time of autopsy about a decade later.*

Fig 5. Boxplots of inferred AD-NC traits at study baseline versus postmortem NIA-Reagan status profiled at autopsy in ROS decedents with autopsy. *Two-sample t-test p-values are 1.5×10^{-10} for β-amyloid (A), 7.3×10^{-11} for tau tangles (B), 7.1×10^{-10} for global AD pathology (C), and 1.2×10^{-7} for NIA-Reagan (D).*

1 **Appendix. Authors**

Name	Location	Contribution
Jingjing Yang, PhD	Center for Computational and Quantitative Genetics, Department of Human Genetics, Emory University School of Medicine, Atlanta, GA	Drafting/revision of the manuscript for content, including medical writing for content; study concept or design; analysis or interpretation of data
Xizhu Liu	Department of Biostatistics, Yale University School of Public Health, New Haven, CT	Drafting/revision of the manuscript for content, including medical writing for content; analysis or interpretation of data
Shahram Oveisgharan, MD	Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago, IL	Drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data; analysis or interpretation of data
Andrea R. Zammit, PhD	Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago, IL	Drafting/revision of the manuscript for content, including medical writing for content; analysis or interpretation of data
Sukriti Nag, MD, PhD	Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago, IL	Drafting/revision of the manuscript for content, including medical writing for content; analysis or interpretation of data
David A. Bennett, MD	Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago, IL	Drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data; analysis or interpretation of data
Aron S. Buchman, MD	Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago, IL	Drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data; study concept or design; analysis or interpretation of data

References

1.   Hyman BT, Phelps CH, Beach TG, et al. National Institute on Aging–Alzheimer's Association guidelines for the neuropathologic assessment of Alzheimer's disease. Alzheimer's and Dementia 2012;8:1-13.

2.   Ballard C, Gauthier S, Corbett A, Brayne C, Aarsland D, Jones E. Alzheimer's disease. Lancet 2011;377:1019-1031.

3.   Jack CR, Jr., Albert M, Knopman DS, et al. Introduction to revised criteria for the diagnosis of Alzheimer's disease: National Institute on Aging and the Alzheimer's Association workgroup. Alzheimers Dement 2011;7:256-262.

4.   Wilson RS, Wang T, Yu L, Bennett DA, Boyle PA. Normative Cognitive Decline in Old Age. Ann Neurol 2020;87:816-829.

5.   Boyle PA, Yu L, Leurgans SE, et al. Attributable risk of Alzheimer's dementia attributed to age-related neuropathologies. Ann Neurol 2019;85:114-124.

6.   Yu L, Wang T, Wilson RS, et al. Common age-related neuropathologies and yearly variability in cognition. Annals of clinical and translational neurology 2019;6:2140-2149.

7.   Lan MJ, Ogden RT, Kumar D, et al. Utility of Molecular and Structural Brain Imaging to Predict Progression from Mild Cognitive Impairment to Dementia. J Alzheimers Dis 2017;60:939-947.

8.   Lu M, Pontecorvo MJ, Devous MD, Sr., et al. Aggregated Tau Measured by Visual Interpretation of Flortaucipir Positron Emission Tomography and the Associated Risk of Clinical Progression of Mild Cognitive Impairment and Alzheimer Disease: Results From 2 Phase III Clinical Trials. JAMA neurology 2021;78:445-453.

9.   Skillbäck T, Kornhuber J, Blennow K, Zetterberg H, Lewczuk P. Erlangen Score Predicts Cognitive and Neuroimaging Progression in Mild Cognitive Impairment Stage of Alzheimer's Disease. J Alzheimers Dis 2019;69:551-559.

10.   Janelidze S, Mattsson N, Palmqvist S, et al. Plasma P-tau181 in Alzheimer's disease: relationship to other biomarkers, differential diagnosis, neuropathology and longitudinal progression to Alzheimer's dementia. Nat Med 2020;26:379-386.

11.   Ansari A, Maffioletti E, Milanesi E, et al. miR-146a and miR-181a are involved in the progression of mild cognitive impairment to Alzheimer's disease. Neurobiol Aging 2019;82:102-109.

12.   Faura J, Bustamante A, Penalba A, et al. CCL23: A Chemokine Associated with Progression from Mild Cognitive Impairment to Alzheimer's Disease. J Alzheimers Dis 2020;73:1585-1595.

13.   Silbert LC, Howieson DB, Dodge H, Kaye JA. Cognitive impairment risk: white matter hyperintensity progression matters. Neurology 2009;73:120-125.

14.   Greene SJ, Killiany RJ. Subregions of the inferior parietal lobule are affected in the progression to Alzheimer's disease. Neurobiol Aging 2010;31:1304-1311.

15.   Sperling RA, Aisen PS, Beckett LA, et al. Toward defining the preclinical stages of Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. Alzheimers Dement 2011;7:280-292.

16.   Ezzati A, Harvey DJ, Habeck C, et al. Predicting Amyloid-beta Levels in Amnestic Mild Cognitive Impairment Using Machine Learning Techniques. J Alzheimers Dis 2020;73:1211-1219.

17.   Jack CR, Jr., Holtzman DM. Biomarker modeling of Alzheimer's disease. Neuron 2013;80:1347-1358.

18.   Chételat G, Arbizu J, Barthel H, et al. Amyloid-PET and (18)F-FDG-PET in the diagnostic investigation of Alzheimer's disease and other dementias. Lancet Neurol 2020;19:951-962.

19.   Isaacs JD, Boenink M. Biomarkers for dementia: too soon for routine clinical use. Lancet Neurol 2020;19:884-885.

20.   Bălașa AF, Chircov C, Grumezescu AM. Body Fluid Biomarkers for Alzheimer's Disease-An Up-To-Date Overview. Biomedicines 2020;8.

21.   Darst BF, Lu Q, Johnson SC, Engelman CD. Integrated analysis of genomics, longitudinal metabolomics, and Alzheimer's risk factors among 1,111 cohort participants. Genet Epidemiol 2019;43:657-674.

22.   Hastie T, Tibshirani R, Friedman J. The elements of statistical learning: prediction, inference and data mining. Springer-Verlag, New York 2009.

23.   Braak H, Thal DR, Ghebremedhin E, Del Tredici K. Stages of the pathologic process in Alzheimer disease: age categories from 1 to 100 years. J Neuropathol Exp Neurol 2011;70:960-969.

24.   Bennett DA, Buchman AS, Boyle PA, Barnes LL, Wilson RS, Schneider JA. Religious Orders Study and Rush Memory and Aging Project. J Alzheimers Dis 2018;64:S161-S189.

25.   von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: guidelines for reporting observational studies. Int J Surg 2014;12:1495-1499.

26.   Bennett DA, Schneider JA, Wilson RS, Bienias JL, Arnold SE. Neurofibrillary tangles mediate the association of amyloid load with clinical Alzheimer disease and level of cognitive function. Arch Neurol 2004;61:378-384.

27.   Bennett DA, Wilson RS, Schneider JA, et al. Apolipoprotein E epsilon4 allele, AD pathology, and the clinical expression of Alzheimer's disease. Neurology 2003;60:246-252.

28.   Hyman BT, Phelps CH, Beach TG, et al. National Institute on Aging-Alzheimer's Association guidelines for the neuropathologic assessment of Alzheimer's disease. Alzheimers Dement 2012;8:1-13.

29.   McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM. Clinical diagnosis of Alzheimer's disease: report of the NINCDS-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. Neurology 1984;34:939-944.

30.   Tombaugh TN, McIntyre NJ. The mini-mental state examination: a comprehensive review. J Am Geriatr Soc 1992;40:922-935.

31.   Buchman AS, Yu L, Oveisgharan S, et al. Cortical proteins may provide motor resilience in older adults. Scientific Reports 2021;11:11311.

32.   Park M, Buchman AS, Lim AS, Leurgans SE, Bennett DA. Sleep complaints and incident disability in a community-based cohort study of older persons. Am J Geriatr Psychiatry 2014;22:718-726.

33.   Linden A. Measuring diagnostic and predictive accuracy in disease management: an introduction to receiver operating characteristic (ROC) analysis. J Eval Clin Pract 2006;12:132-139.

34.   Cox DR. Regression models and life-tables. Journal of the Royal Statistical Society: Series B (Methodological) 1972;34:187-202.

35.   Gerds TA, Scheike TH, Andersen PK. Absolute risk regression for competing risks:
interpretation, link functions, and prediction. Stat Med 2012;31:3921-3930.
36.   Ozenne B, Sørensen AL, Scheike T, Torp-Pedersen C, Gerds TA. riskRegression:
predicting the risk of an event using Cox regression models. The R Journal 2017;9:440-460.
37.   Hastie T, Tibshirani R, Friedman JS. The Elements of Statistical Learning, 2 ed:
Springer-Verlag New York, 2009.
38.   Frölich L, Peters O, Lewczuk P, et al. Incremental value of biomarker combinations to
predict progression of mild cognitive impairment to Alzheimer's dementia. Alzheimers Res Ther
2017;9:84.
39.   Feng X, Chen L, Wang Z, Li SC. I-Impute: a self-consistent method to impute single cell
RNA sequencing data. BMC genomics 2020;21:618.
40.   Li WV, Li JJ. An accurate and robust imputation method scImpute for single-cell RNA-
seq data. Nat Commun 2018;9:997.
41.   Gamazon ER, Wheeler HE, Shah KP, et al. A gene-based association method for
mapping traits using reference transcriptome data. Nature genetics 2015;47:1091-1098.
42.   Nguyen M, He T, An L, Alexander DC, Feng J, Yeo BTT. Predicting Alzheimer's disease
progression using deep recurrent neural networks. Neuroimage 2020;222:117203.
43.   Nagpal S, Meng X, Epstein MP, et al. TIGAR: An Improved Bayesian Tool for
Transcriptomic Data Imputation Enhances Gene Mapping of Complex Traits. American journal
of human genetics 2019;105:258-266.
44.   Luningham JM, Chen J, Tang S, et al. Bayesian Genome-wide TWAS Method to
Leverage both cis- and trans-eQTL Information through Summary Statistics. American journal
of human genetics 2020;107:714-726.
45.   Tosun D, Chen YF, Yu P, et al. Amyloid status imputed from a multimodal classifier
including structural MRI distinguishes progressors from nonprogressors in a mild Alzheimer's
disease clinical trial cohort. Alzheimers Dement 2016;12:977-986.
46.   Tosun D, Joshi S, Weiner MW. Neuroimaging predictors of brain amyloidosis in mild
cognitive impairment. Ann Neurol 2013;74:188-198.
47.   Ghirelli A, Tosakulwong N, Weigand SD, et al. Sensitivity-Specificity of Tau and Amyloid
β Positron Emission Tomography in Frontotemporal Lobar Degeneration. Ann Neurol
2020;88:1009-1022.
48.   Soleimani-Meigooni DN, Iaccarino L, La Joie R, et al. 18F-flortaucipir PET to autopsy
comparisons in Alzheimer's disease and other neurodegenerative diseases. Brain
2020;143:3477-3494.
49.   Luningham JM, Chen J, Tang S, et al. Bayesian Genome-wide TWAS Method to
Leverage both cis- and trans-eQTL Information through Summary Statistics. American journal
of human genetics 2020.
50.   Licher S, Yilmaz P, Leening MJG, et al. External validation of four dementia prediction
models for use in the general community-dwelling population: a comparative analysis from the
Rotterdam Study. Eur J Epidemiol 2018;33:645-655.
135