Inferring Alzheimer's disease pathologic traits from clinical measures in living adults

Author(s):

Jingjing **Yang**, PhD[1]*; Xizhu **Liu**, BS[2]; Shahram **Oveisgharan**, MD[3]; Andrea R. **Zammit**, PhD[3]; Sukriti **Nag**, MD, PhD,[3] David A **Bennett**, MD[3]; Aron S **Buchman**, MD[3]

Affiliation Information for All Authors:
1. Center for Computational and Quantitative Genetics, Department of Human Genetics, Emory University School of Medicine, 615 Michael St, Atlanta, GA, 30322, USA;
2. Department of Biostatistics, Yale University School of Public Health, 60 College St, New Haven, CT, 06510, USA;
3. Rush Alzheimer's Disease Center, Rush University Medicine Center, 1620 W Harrison St, Chicago, IL, 60612, USA.

Corresponding Author:

* Jingjing Yang

Address: Department of Human Genetics, Emory University School of Medicine, 615 Michael Street, Suite 305K, Atlanta, GA, 30322, USA

Email: jingjing.yang@emory.edu

Phone: 404-727-3481

ABSTRACT

Background: Alzheimer's disease neuropathologic changes (AD-NC) are important for

identify people with high risk for AD dementia (ADD) and subtyping ADD.

Objective: Develop imputation models based on clinical measures to infer AD-NC.

Methods: We used penalized generalized linear regression to train imputation models

for four AD-NC traits (amyloid-β, tangles, global AD pathology, and pathologic AD) in

Rush Memory and Aging Project decedents, using clinical measures at the last visit

prior to death as predictors. We validated these models by inferring AD-NC traits with

clinical measures at the last visit prior to death for independent Religious Orders Study

(ROS) decedents. We inferred baseline AD-NC traits for all ROS participants at study

entry, and then tested if inferred AD-NC traits at study entry predicted incident ADD and

postmortem pathologic AD.

Results: Inferred AD-NC traits at the last visit prior to death were related to postmortem

measures with R^2=(0.188,0.316,0.262) respectively for amyloid-β, tangles, and global

AD pathology, and prediction Area Under the receiver operating characteristic Curve

(AUC) 0.765 for pathologic AD. Inferred baseline levels of all four AD-NC traits

predicted ADD. The strongest prediction was obtained by the inferred baseline

probabilities of pathologic AD with AUC=(0.919,0.896) for predicting the development of

ADD in 3 and 5 years from baseline. The inferred baseline levels of all four AD-NC traits

significantly discriminated pathologic AD profiled eight years later with p-

values$<1.4 \times 10^{-10}$.

Conclusion: Inferred AD-NC traits based on clinical measures may provide effective

AD biomarkers that can estimate the burden of AD-NC traits in aging adults.

KEYWORDS: Alzheimer's disease; Computational modeling; Alzheimer's disease

neurologic change; Dementia; Pathology

INTRODUCTION

The accumulation of Alzheimer's disease neuropathologic changes (AD-NC),

such as amyloid-β and intracellular neurofibrillary tangles, underlying Alzheimer's

disease dementia (ADD) has been observed even during the initial stages of ADD when

cognition is normal [1]. Higher levels of AD-NC during the early stages of ADD have

been shown to be associated with an increased risk of ADD [2-6]. Separate therapies

have been developed for targeting the peptide amyloid-β in extracellular amyloid

plaques and the protein tau in intracellular neurofibrillary tangles [7-9]. Yet, conventional

prediction models identifying adults at risk for clinical ADD do not inform on which AD-

NC traits underlie the risk of ADD. Thus, procedures that can accurately infer elevated

levels of AD-NC traits in living adults with normal cognition have the potential to

facilitate early targeted treatments [10-17].

Direct measures of brain AD-NC traits can only be obtained at autopsy. Recent

efforts to quantify AD-NC during life have focused on identifying biomarkers of AD-NC

traits by using brain imaging or fluid AD biomarkers [18-20]. Recent work comparing tau

and amyloid positron emission tomography (PET) brain imaging measures to indices

measured at autopsy, suggests that current imaging may not reliably detect the early

stages of AD pathology [21, 22]. Particularly, brain imaging and CSF biomarkers are not

widely available due to their costs, invasiveness, and difficultly to deploy at scale.

Studies focusing on serum biomarkers have advanced rapidly in recent years, yet the

field has not converged on specific biomarkers that can be employed in the general

population [23-27].

Rapid advances in machine learning methods such as penalized generalized

linear regression [28] have been employed to impute missing data or infer data that are

difficult to be measured directly in biomedical research fields [29-34]. Similarly, the

penalized generalized linear regression method could be deployed to develop

imputation models based on clinical measures in older adults to infer levels of AD-NC

traits. Such imputation models learn the predictive information of postmortem AD-NC

traits like tangles from clinical measures obtained prior to death in decedents

undergoing autopsy. The imputation model works by mathematically "explaining" the

variation of observed AD-NC traits measured in decedents by their equivalence of

weighted linear combinations of predictive clinical measures. Once imputation models

developed for AD-NC traits are validated in an independent cohort, they can be applied

to any older adults with similar clinical measures to infer AD-NC traits.

Comprehensive clinical and postmortem data are necessary to develop and

validate imputation models for AD-NC traits. This multi-stage study leveraged clinical

and postmortem data from two harmonized, independent, longitudinal prospective

cohort studies —— Rush Memory and Aging Project (MAP) and the Religious Orders

Study (ROS) [35]. First, we trained imputation models for four AD-NC traits (amyloid-β,

tangles, global AD pathology, and pathologic AD diagnosis) by applying the penalized

generalized linear regression method to clinical measures obtained at the last visit

before death and postmortem AD-NC indices measured at autopsy in MAP decedents.

Second, we validated the imputation models in a second independent cohort (ROS)

which collected the same clinical and postmortem AD-NC traits. Third, we applied the

imputation models to clinical measures collected in adults without dementia at study

entry to infer baseline levels of AD-NC traits that were on average about eight years

before death for decedents (or before last visit for living participants). We demonstrated

the efficacy of these interfered baseline AD-NC traits as effective AD biomarkers, by

showing their predictivity of future clinical ADD and discrimination of postmortem

pathologic AD.

MATERIALS AND METHODS

Participants

Participants were community-dwelling older adults enrolled without known dementia and

with at least two annual visits in one of two ongoing longitudinal prospective cohort

studies of chronic conditions of aging — MAP (n=1179 with ~500 autopsied) and ROS

(n=1103 with ~600 autopsied). Both cohorts employed a harmonized data collection

battery administered by the same research assistants facilitating joint analyses. For this

study, we included adults without clinical evidence of dementia at enrollment with at

least two annual follow-up cognitive assessments. At study entry, 1742 adults had no

cognitive impairment (NCI) and 540 adults had mild cognitive impairment (MCI) (**Table**

S1). The duration of annual follow-up for participants ranged from 2 to 26 years, with an

average follow-up of 8 years (SD, 5.42 years) [**Fig S1; Tables S1-S2**].

Assessment of AD-NC Traits

After the death of ROS/MAP participants, their brains were removed and

hemisected following the standard procedure, as previously described [35]. Tissue

blocks were dissected from predetermined regions and used for postmortem diagnosis

of pathologic AD. Structured autopsy collected indices of AD/ADRD pathologies that

were collected blinded to all prior clinical and cognitive data. This study focuses on the

development of imputation models to infer the following four AD-NC traits that were

measured in autopsied decedents (**Table S2**).

Amyloid-β was labeled with an N-terminus–directed monoclonal antibody (10D5;

Elan, Dublin, Ireland; 1:1,000). Immunohistochemistry was performed as previously

described using diaminobenzidine as the reporter, with 2.5% nickel sulfate to enhance

immunoreaction product contrast.

PHFtau (Tangles) was labeled with an antibody specific for phosphorylated tau

(AT8; Innogenetics, San Ramon, CA; 1:1,000). Amyloid-β load and tangles were

quantified in 8 brain regions (anterior cingulate cortex, superior frontal cortex, mid

frontal cortex, inferior temporal cortex, hippocampus, entorhinal cortex, angular

gyrus/supramarginal gyrus, and calcarine cortex). Overall amyloid-β load was calculated

through averaging mean percent area of amyloid-β deposition per region, across

multiple brain regions. Tangles densities were derived by averaging tangles densities

across corresponding brain regions. Measures of amyloid-β and tangles were further

square-root transformed to improve their asymptotic normality as previously reported

[36, 37].

Global AD Pathology*: A modified Bielschowsky silver stain was used to

visualize neuritic plaques, diffuse plaques, and neurofibrillary tangles in five cortical

areas (hippocampus, entorhinal, midfrontal, middle temporal, and inferior parietal).

Neuritic and diffuse plaques, and neurofibrillary tangles were counted in the region that

appeared to have the maximum density of each pathology as previously described. A

standardized score was created for each neuropathology in each region by dividing the

raw count by the standard deviation of the mean for the same neuropathology in the

same region. This standardization procedure puts the pathologic indices on a relatively

common scale. A summary global AD pathology score was made based on the average

of the greatest density of neuritic plaques, diffuse plaques, and neurofibrillary tangles in

one mm^2 [38, 39]

Pathologic Diagnosis of AD: The National Institute on Aging-Reagan criteria

were used with intermediate and high likelihood cases indicating a pathologic diagnosis

of AD, which is a binary indicator with value 1 denoting the present of pathologic AD

and 0 denoting the absent of pathologic AD [40].

Assessment of Composite Cognition Score and Cognitive Status

A structured cognitive assessment was administered annually. The

neuropsychological battery included 19 tests that assessed five cognitive abilities

(episodic memory, semantic memory, working memory, visuospatial ability/perceptual

orientation, and perceptual speed). Raw test scores were standardized for each test

using baseline means and standard deviations (SDs) of both cohorts; the resulting Z-

scores were then averaged across these cognitive tests to derive a single summary

composite cognition score as described in prior publications [35, 41].

Cognitive diagnoses were made in a three-step process. Cognitive testing was

scored by a computer program and the results were reviewed by a neuropsychologist to

diagnose cognitive impairment. Then participants were evaluated by a physician who

used available cognitive and clinical data to classify cognitive status at each annual

visit. Dementia required meaningful decline in cognitive function with impairment in

multiple areas of cognition, and AD required dementia and progressive loss of episodic

memory. Individuals with cognitive impairment who did not meet dementia criteria were

diagnosed with mild cognitive impairment (MCI). Individuals without dementia or MCI

were classified as having no cognitive impairment (NCI). Clinical diagnosis of cognitive

status was based on published criteria [42-44]. Participants with dementia due to

primary cause other than AD are excluded in this study. At the time of death, select

clinical data from the entire study were reviewed by a neurologist, blinded to

postmortem data, to render a final cognitive status diagnosis[35].

Clinical Covariates

Diverse clinical measures were used to develop imputation models for four AD-

NC traits. **Table S3** shows the complete list and groupings of the 57 clinical measures

examined in this study, including both cross-sectional and longitudinal variables. Cross-

sectional clinical measures were collected only once for each participant, such as sex,

education, and *APOE* genotype. Longitudinal clinical measures were collected each

time during participants' annual visits. Baseline clinical characteristics are provided in

Tables S2, S4, and S5 and a heat map (**Fig S2**) is included to show the inter-

correlations of the clinical variables examined in this study. The measures analyzed in

this study were selected after excluding cross-sectional variables with a high proportion

of missing values and those that were highly correlated with other selected clinical

measures (correlation>0.95). Samples with missing values in the selected cross-

sectional clinical measures were excluded. Missing values of longitudinal clinical

measures were assumed to be missed at random and were imputed from the measured

values at the nearest visit of the same participant, by using the "*fill(.direction = "up")*"

function from R library "*tidyr*". That is, if a participant had a missing value at last visit for

a longitudinal variable, the missing value would be imputed as the measured value of

this variable in this participant's nearest previous visit with collected measurement. The

percentage of missing data (i.e., missing rate) in the longitudinal clinical measurements

at last visit and study baseline were presented in **Fig, S3**. The number of samples that

were actually used in the analyses are presented in **Tables S6**, and **S7**.

Analytic Approach

A multi-stage analytic approach was employed to develop, validate, and

demonstrate the effectiveness of the inferred the levels of four AD-NC traits at study

entry as potential AD biomarkers.

Developing imputation models to infer AD-NC traits

Stage 1. Training imputation models.

We trained an imputation model for each of the four AD-NC traits using 57

clinical variables obtained in MAP participants at the last visit before death as

predictors, by using the generalized linear regression model with Elastic-Net penalty

(GLM-EN) [45] (**Fig 1A**). Only MAP decedents with autopsy were used for developing

imputation models, because profiled AD-NC traits were required. By using the GLM-EN

method, variable selection was implemented and potential collinearity among clinical

variables was accounted for during model training. Since the Elastic-Net penalty is a

linear combination of L1 (i.e. LASSO) [46] and L2 (i.e., Ridge) [47] penalties on the

coefficients of clinical variables, variable selection are handled by the L1 penalty (i.e.,

penalizing the L1 norm of the coefficient vector) while potential collinearity is accounted

for by the L2 penalty (i.e., penalizing the L2 norm of the coefficient vector). Ten-fold

cross validation was used during model training to select Elastic-Net penalty parameters

(i.e., the proportions of L1 and L2 penalty) to ensure optimal imputation accuracy.

Stage 2. Validating imputation models in a second independent cohort.

Next, we validated the performance of the imputation models developed to infer

four AD-NC traits using clinical variables measured at the last visit proximate to death in

a second independent cohort with ROS decedents (**Fig 1A**). Prediction R^2, the squared

correlation between inferred and measured values, was used for assessing imputation

accuracy for continuous AD-NC traits. Prediction accuracy for the dichotomous

pathologic AD diagnosis was evaluated by the predicted area under curve (AUC) values

of receiver operating characteristic curve (ROC) [48].

Stage 3. Infer AD-NC traits at study entry and test their effectiveness as AD biomarkers

Once imputation models are trained and validated, they can be applied to any

living adult with the same clinical variables measured. So, to illustrate the use of these

imputation models, we applied these validated models to the clinical data obtained from

ROS participants at study entry (i.e., baseline that was on average of eight years before

death or last follow-up visit) to infer baseline AD-NC traits (**Fig 1B**). Scatter plots and

ROC curves were used to evaluate the consistency between inferred AD-NC traits at

baseline and the corresponding measured postmortem AD-NC traits profiled at autopsy

in ROS decedents. Then we examined the effectiveness of inferred AD-NC traits as

potential AD biomarkers through two complementary analyses, one for evaluating the

predictivity for incident ADD by Cox proportional hazard model (Stage 3A) and the other

one for evaluating the discrimination of postmortem pathologic AD (Stage 3B).

Stage 3A. We fitted Cox proportional hazard models using covariates of age,

sex, education, and a single inferred baseline AD-NC trait in MAP cohort for predicting

incident ADD. Then we evaluated the performance of the Cox models for predicting

incident ADD in year 3 and year 5 from baseline in ROS cohort (**Fig S4**). Our fitted Cox

proportional hazard risk prediction models [49-51] also accounted for the competing risk

of death. The annual cognitive status diagnosis and the follow up year were used to

identify the first occurrence of ADD. For each participant, the year of enrollment is

considered as baseline (time 0), the year of first diagnosis of ADD is considered as the

time when the event occurs (incident ADD), and the last visit of participants without the

considered event during all follow-ups is considered as the right censored time for living

participants or the time of death for dead participants without ADD. Sample size

distributions with respect to cognitive status at baseline and the cognition event types

are shown in **Tables S1**. All Cox models were trained using data from MAP participants

(both living and deceased) and tested with data from ROS participants (both living and

deceased).

For fitting and testing Cox models for predicting incident ADD, we first used all

individuals without dementia at baseline, and then fitted and tested another set of Cox

models by using only individuals with NCI at baseline. For each set of Cox models fitted

by using MAP participants, we calculated model accuracy (AUC) for predicting incident

ADD in year 3 and year 5 from baseline.

Since Cox models provide a continuous risk score for incident ADD, by selecting

a risk score threshold corresponding to ~80% specificity (the proportion of correctly

predicted non-ADD in test ROS participant's, i.e., 1 – false positive fraction), we could

calculate sensitivity (the proportion of true positive predictions in all test cases, i.e., true

positive fraction), and the overall classification accuracy (the proportion of true

discrimination of ADD in all test samples). Samples with risk scores greater than the

selected threshold were considered to develop ADD and less than the threshold as not

developing ADD in a specific year. That is, given the known ADD status of participants

in year 3 and year 5 from baseline, we can compare the predicted risk of incident ADD

to the actual incident ADD status to calculate the overall classification accuracy,

specificity, and sensitivity. Also, the sensitivity and specificity corresponding to a

selected risk score threshold reflect risk model performance at one point in the ROC

plots.

We also sequentially added the other three inferred baseline AD-NC traits into

the Cox model, in addition of covariates of age, sex, education, and inferred baseline

amyloid-β, and examined the prediction performance of these models.

Stage 3B. We examined the discrimination of inferred baseline levels of AD-NC

traits with respect to postmortem pathologic AD diagnosis (**Fig 1B**). Boxplots and two-

sample t-tests were used to evaluate the discrimination of postmortem pathologic AD by

inferred baseline AD-NC traits. Only ROS decedents with profiled pathologic AD

diagnoses were use in this discrimination analysis.

RESULTS

Developed Imputation Models for Inferring AD-NC Traits

Imputation models were trained to infer AD-NC traits by applying the GLM-EN

method using clinical measures in MAP decedents at their last visit before death as

predictors. Selected predictive clinical measures with standardized effect sizes

|beta| >0.01 estimated by the imputation models are shown in **Fig 2.** Composite

cognition score and *APOE* E4 allele were the strongest predictors that were selected for

all four AD-NC traits. Yet, **Fig 2** also illustrates that varied non-cognitive clinical

measures including motor function such as motor gait and dexterity, health conditions

such as anxiety and hypertension, and medications such as lipid lowering and anti-

inflammatory medications were also selected. The effect size for motor gait was nearly

as strong as APOE for tangles, amyloid-beta, and global AD pathology.

These estimated effect sizes of selected clinical measure predictors in the

imputation models are used as weights to construct weighted sum with clinical

measurements to infer AD-NC trait levels. So, while all four AD-NC traits are related and

may share composite cognition score and *APOE E4* allele as important predictors, the

different sets of selected predictors by these four imputation models highlight that

mathematically different combinations of clinical measures with different estimated

effect sizes are necessary to best capture unique features of these inter-related AD-NC

traits.

Validation of Imputation Models for Inferring AD-NC Traits

To validate the imputation models developed in MAP decedents, we applied the imputation models to clinical measures obtained at the last visit prior to death for decedents in a second independent ROS cohort to infer their levels of four AD-NC traits. Scatter plots illustrate the correlations between the inferred levels of the three continuous AD-NC traits and their corresponding indices measured at autopsy (**Fig S5, A-C).** The prediction R^2 was 0.188 for amyloid-β, 0.316 for tangles, and 0.262 for global AD pathology (**Table S8**). An ROC plot (with AUC=0.765) illustrates the consistency between the inferred probabilities of pathologic AD based on clinical measures obtained at last visit before death versus the profiled pathologic AD status by autopsy (**Fig S5D; Table S8)**. As would be expected for effective AD biomarkers, all four inferred AD-NC traits discriminated profiled pathologic AD at autopsy, with two-sample test P values < 10^{-28} (Box plots in **Fig S5**; **Table S8**). Together these results in a second independent cohort validated the accuracy of the imputation models developed in MAP for all four AD-NC traits.

Inferred Baseline AD-NC Traits Predicted Incident ADD

The imputation models were developed and validated using clinical measures at the last visit prior to death in MAP and ROS decedents. In further analyses we examined the effectiveness of inferred baseline AD-NC traits at study entry as AD biomarkers. To infer baseline levels of four AD-NC traits at study entry, we applied the validated imputation models to clinical data collected at baseline in all ROS participants (n=1103; both living and decedents), an average of 8 years before death for decedents

(or last follow-up for living participants) (**Fig 1B**). Scatter plots of the three continuous

AD-NC traits and an ROC plot of the binary pathologic AD illustrate the correlation

between the inferred baseline versus the profiled AD-NC traits by autopsy (**Fig S6)**.

Although the correlations were lower than the inferred AD-NC traits at last visit before

death (**Fig S5**), the following analyses with Cox models still demonstrated the

predictivity of the inferred baseline AD-NC traits for predicting incident ADD.

We employed separate Cox models that considered covariates age, sex,

education, and each one of the four inferred baseline AD-NC traits to examine the

predictivity for incident ADD (**Fig 3**). Coefficient estimates of the inferred baseline AD-

NC traits in these Cox models were provided along with the corresponding p-values in

Table S9. All inferred baseline AD-NC traits were strongly associated with incident

ADD, with p-values $< 10^{-30}$ in Cox models predicting incident ADD for adults without

dementia at baseline, and p-values $< 10^{-5}$ in Cox models for predicting incident ADD

from adults with NCI at baseline.

For all four AD-NC traits, model performance was higher in Year 3 (AUC ranging

in 0.861 – 0.919) versus Year 5 (AUC ranging in 0.842 – 0.896) from baseline (**Fig 3,**

Upper Row), for predicting incident ADD for adults without dementia at baseline. Of the

four AD-NC traits, inferred baseline probabilities of pathologic AD had the highest

predictivity (AUC 0.919 in Year 3; AUC 0.896 in Year 5) for incident ADD from adults

without dementia at baseline. Similar results were observed when we restricted the

analyses to the prediction of incident ADD in individuals with NCI at baseline (**Fig 3,**

Lower Row).

By selecting a risk score threshold corresponding the ~80% specificity, we

calculated sensitivity and accuracy based on the prediction results for all of the Cox

models (**Table 1**). Inferred baseline probabilities of pathologic AD also had the highest

accuracy rates (80%) and sensitivity for predicting incident ADD in Year 3 (0.911) and

Year 5 (0.829) from baseline (**Table 1**), compared to the other AD-NC traits.

In further analyses, we examined if adding covariates of additional inferred

baseline AD-NC traits in a single Cox model would improve the prediction accuracy for

incident ADD. As shown in **Fig S7 (Upper Row)**, we observed slightly improved AUC

for prediction of incident ADD in adults without dementia, when we sequentially added

each of the four AD-NC traits. Yet, modeling all four AD-NC traits together did not yield

better results than using inferred probabilities of pathologic AD alone (**Last Column of**

Fig3 versus Fig S7).

Sequentially adding the inferred baseline levels of three continuous AD-NC traits

(amyloid-β, tangles, and global AD pathology) in a single Cox model did not improve the

prediction of incident ADD for adults with baseline NCI (**Fig S7, Lower Row**). Although

adding the inferred baseline levels of the probabilities of pathologic AD improved the

prediction accuracy of incident ADD for adults with baseline NCI, but the prediction

accuracy was comparable as using the inferred baseline levels of the probabilities of

pathologic AD alone (**Last Column of Fig3 versus Fig S7**).

Inferred Baseline AD-NC Traits Discriminated Postmortem Pathologic AD

By examining if the inferred baseline AD-NC traits would discriminate

postmortem pathologic AD diagnosis, we presented boxplots of the inferred baseline

AD-NC traits of ROS decedents with respect to their postmortem pathologic AD

diagnosis by autopsy in **Fig 4**. By two-sample t-tests, we showed that all four inferred

baseline AD-NC traits discriminated individuals with postmortem pathologic AD

diagnosis by autopsy, with significant p-values $< 1.4 \times 10^{-10}$.

DISCUSSION

This study applied the machine learning GLM-EN methods to clinical

measurements and postmortem indices of four AD-NC traits (amyloid-β, tangles, global

AD pathology, and pathologic AD) obtained from the same older adults to develop

imputation models that could be used to infer levels of four AD-NC traits based on

clinical measurements. We validated these imputation models of AD-NC traits in a

second independent cohort of older adults that collected similar clinical and postmortem

indices. We applied the validated imputation models to clinical measures obtained at

study entry to infer baseline AD-NC traits in adults about an average of eight years

before death (decedents) or their last follow-up visit (living participants).

Adults without dementia at study entry who had higher baseline levels of inferred

AD-NC traits had a higher risk of developing incident ADD during follow-up years and

they also had a higher risk of having postmortem pathologic AD. These data suggest

that inferred levels of AD-NC traits based on clinical measures may provide a low cost,

non-invasive effective AD biomarker that can estimate the burden of AD-NC traits

during the chronic course of Alzheimer's disease. Inferred AD-NC traits may provide a

way to monitor the clinical course of the accumulation of AD-NC traits underlying

Alzheimer's disease, improve the homogeneity of clinical trials and catalyze early

targeted treatments to prevent the development of ADD in aging adults. Further studies

to validate longitudinal inferred AD-NC traits will be needed.

Novelty of this Study

Currently AD-NC traits in brain can only be measured at autopsy. Recent work has

focused on identifying effective AD biomarkers that can be used to assess levels of AD-

NC traits in living adults, especially during early stages of AD when cognition is still normal.

There are many prior studies that have examined the associations of clinical measures,

such as *APOE E4* allele, age, and cognitive measures, with future cognitive status or with

different AD/ADRD indices measured at autopsy including the postmortem diagnosis of

pathologic AD. Yet, it is important to note that the aim of these prior studies was not to

infer levels of the different AD-NC traits examined in this study, nor to test their

effectiveness as AD biomarkers [52-56].

Brain imaging studies of AD-NC traits have tried to employ serial imaging or CSF

fluid biomarkers as proxies to obtain measures of AD-NC traits in brains to assess the

accumulation of amyloid-β and tangles in early stages of Alzheimer's disease [18-20].

Recent work that compared tau and amyloid PET brain imaging to AD indices measured

at autopsy, suggests that current imaging may not reliably detect the early stages of AD

pathology [21, 22]. Yet, the expense and limited availability of brain imaging and the

invasiveness of obtaining CSF biomarkers make them difficult to be deployed at scale for

the general population. This study fills this gap by employing machine learning methods

that could be used to mathematically infer an AD-NC trait like "tangles" and estimate its

burden at any time point prior to death in any adult with the requisite clinical measures.

Obtaining structured autopsy and diverse clinical measures during annual follow-

ups, especially within a year prior of death, in large numbers of older adults is difficult.

Thus, it is novel to have two large cohorts like MAP or ROS with the same clinical and

postmortem indices of AD-NC traits that can be leveraged to develop imputation models

in one cohort and validate these imputation models in a second independent sample of

older adults. The rarity of these resources may explain in part the paucity of previous

studies trying to infer AD-NC traits based on clinical measurements alone.

Another novel feature of this study is that we provide evidence that inferred

baseline AD-NC traits on average 8 years before death for decedents (or before last visit

for living participants) may be used as effective AD biomarkers as they predicted incident

ADD and discriminated postmortem pathologic AD. That is, this study provides novel data

demonstrating the feasibility and effectiveness of developing imputation models to infer

AD-NC traits from clinical measures alone.

Implications and Future Directions

This study is best conceptualized as an important first step highlighting that new

machine learning analytic techniques can be used to infer AD-NC traits based on

clinical measures collected in older adults. Further studies are still needed to determine

if repeated inferred levels of AD-NC traits inform on trajectories of the accumulation of

these different AD-NC traits. Currently, the temporal course of accumulation of these

different AD-NC traits and the onset of their associations with impaired cognition are

unknown. Such data are crucial to determine if inferred AD-NC traits could be used to

assess the ongoing clinical course of Alzheimer's disease, and to assess the efficacy of

guiding treatments targeting specific AD-NC traits. For example, separate therapies

have been under study for targeting the peptide amyloid-β in extracellular amyloid

plaques and the protein tau in intracellular neurofibrillary tangles [7-9].

The analytic approach implemented in this study might also be extended to infer

the presence of other pathologies that are hard to measure in living adults and untangle

the effects of mixed-brain pathologies underlying late-life cognitive impairment and

dementia. Since many older adults with ADD show mixed-brain pathologies [57], further

work will be needed to develop analytic approaches that can infer and account for the

presence of different combinations of varied AD/ADRD pathologies.

Brain imaging as well as serum or fluid biomarkers were not examined in this

study, which could be used to validate the inferred levels of AD-NC traits by using our

developed imputation models. Additionally, the brain imaging and fluid biomarkers might

be included as additional predictors to enhance the imputation accuracy of AD-NC

traits, which could be crucial for untangling the relative contributions of mixed-brain

pathologies driving ADD in aging adults.

Limitations and Strengths of this Study

This study still has several limitations. First, participants were predominantly

Americans of European descent and have higher than average levels of education, so

our findings will need to be replicated in more diverse populations. Second, the current

study used diverse clinical predictors, many of that might not be available outside the

research setting such as the composite cognition score based on 19 cognitive tests

designed for ROS/MAP studies. Further work is needed to identify a parsimonious set of

clinical predictors that are more widely available to enhance the use of this approach in

diverse populations and geographic locations. Despite these limitations, this study is

best conceptualized as an important first step highlighting the potential of using machine

learning methods to infer AD-NC traits or other AD/ADRD pathologies based on clinical

measures that can be collected via remote phenotyping or electronic health records.

Nonetheless, this study has several strengths that lend confidence for the current

findings. All subjects were recruited from the community, underwent an annual detailed

clinical evaluation, and were without dementia based on their clinical assessment at

study entry. Large numbers of men and women underwent annual assessments, and

follow-up rates were very high (~90%) with an average of 8 years follow up. Uniform

and structured procedures were employed for the collection of clinical measures and

postmortem AD-NC traits in both MAP and ROS cohorts. An important strength of the

current study design is that we developed imputation models in MAP cohorts, and then

evaluated model performance in the second independent ROS cohort that employed

similar staff and data collection procedures [58].

ACKNOWLEGEMENTS

We are deeply indebted to all participants who contributed their data and agreed to

autopsy at the time of their death. We thank the Rush Alzheimer's Disease Center staff

for their efforts.

FUNDING

This work was supported by the National Institute of Health R35GM138313,

P30AG10161, P30AG72975, K01AG054700, R01AG15819, R01AG17917,

R01AG56352; R01AG79133, the Illinois Department of Public Health; and the Robert C.

Borwell Endowment Fund. The funding organizations had no role in the design or

conduct of the study; collection, management, analysis, or interpretation of the data; or

preparation, review, or approval of the manuscript.

CONFLICT OF INTEREST

Jingjing Yang is an Editorial Board Member of this journal, but was not involved in the

peer-review process nor had access to any information regarding its peer-review. There

are no other disclosures nor conflict of interest for any of the authors.

CONSENT STATEMENT

Both MAP and ROS studies were approved by an Institutional Review Board of Rush

University Medical Center. All participants agreed to annual clinical evaluations and

autopsy at the time of death. Written informed consent was obtained from all study

participants as was an Anatomical Gift Act for organ donation.

DATA AVAILABILITY

All data analyzed in this study are de-identified and available to any qualified

investigator by submitting a request through the Rush Alzheimer's Disease Center

Research Resource Sharing Hub, https://www.radc.rush.edu, which has descriptions of

the studies and available data.

REFERENCES

[1]   Hyman BT, Phelps CH, Beach TG, Bigio EH, Cairns NJ, Carrillo MC, Dickson DW, Duyckaerts C, Frosch MP, Masliah E, Mirra SS, Nelson PT, Schneider JA, Thal DR, Thies B, Trojanowski JQ, Vinters HV, Montine TJ (2012) National Institute on Aging–Alzheimer's Association guidelines for the neuropathologic assessment of Alzheimer's disease. *Alzheimer's and Dementia* **8**, 1-13.

[2]   Ballard C, Gauthier S, Corbett A, Brayne C, Aarsland D, Jones E (2011) Alzheimer's disease. *Lancet* **377**, 1019-1031.

[3]   Jack CR, Jr., Albert M, Knopman DS, McKhann GM, Sperling RA, Carrillo MC, Thies B, Phelps CH (2011) Introduction to revised criteria for the diagnosis of Alzheimer's disease: National Institute on Aging and the Alzheimer's Association workgroup. *Alzheimers Dement* **7**, 256-262.

[4]   Wilson RS, Wang T, Yu L, Bennett DA, Boyle PA (2020) Normative Cognitive Decline in Old Age. *Ann Neurol* **87**, 816-829.

[5]   Boyle PA, Yu L, Leurgans SE, Wilson RS, Brookmeyer R, Schneider JA, Bennett DA (2019) Attributable risk of Alzheimer's dementia attributed to age-related neuropathologies. *Ann Neurol* **85**, 114-124.

[6]   Yu L, Wang T, Wilson RS, Leurgans S, Schneider JA, Bennett DA, Boyle PA (2019) Common age-related neuropathologies and yearly variability in cognition. *Ann Clin Transl Neurol* **6**, 2140-2149.

[7]   Zhang Y, Chen H, Li R, Sterling K, Song W (2023) Amyloid beta-based therapy for Alzheimer's disease: challenges, successes and future. *Signal Transduct Target Ther* **8**, 248.

[8]   Gotz J, Ittner A, Ittner LM (2012) Tau-targeted treatment strategies in Alzheimer's disease. *Br J Pharmacol* **165**, 1246-1259.

[9]   Lee HE, Lim D, Lee JY, Lim SM, Pae AN (2019) Recent tau-targeted clinical strategies for the treatment of Alzheimer's disease. *Future Med Chem* **11**, 1845-1848.

[10]   Lan MJ, Ogden RT, Kumar D, Stern Y, Parsey RV, Pelton GH, Rubin-Falcone H, Pradhaban G, Zanderigo F, Miller JM, Mann JJ, Devanand DP (2017) Utility of Molecular and Structural Brain Imaging to Predict Progression from Mild Cognitive Impairment to Dementia. *J Alzheimers Dis* **60**, 939-947.

[11]   Lu M, Pontecorvo MJ, Devous MD, Sr., Arora AK, Galante N, McGeehan A, Devadanam C, Salloway SP, Doraiswamy PM, Curtis C, Truocchio SP, Flitter M, Locascio T, Devine M, Zimmer JA, Fleisher AS, Mintun MA (2021) Aggregated Tau Measured by Visual Interpretation of Flortaucipir Positron Emission Tomography and the Associated Risk of Clinical Progression of Mild Cognitive Impairment and Alzheimer Disease: Results From 2 Phase III Clinical Trials. *JAMA Neurol* **78**, 445-453.

[12]   Skillbäck T, Kornhuber J, Blennow K, Zetterberg H, Lewczuk P (2019) Erlangen Score Predicts Cognitive and Neuroimaging Progression in Mild Cognitive Impairment Stage of Alzheimer's Disease. *J Alzheimers Dis* **69**, 551-559.

[13]   Janelidze S, Mattsson N, Palmqvist S, Smith R, Beach TG, Serrano GE, Chai X, Proctor NK, Eichenlaub U, Zetterberg H, Blennow K, Reiman EM, Stomrud E, Dage JL, Hansson O (2020) Plasma P-tau181 in Alzheimer's disease: relationship to other biomarkers,

differential diagnosis, neuropathology and longitudinal progression to Alzheimer's
dementia. *Nat Med* **26**, 379-386.

[14] Ansari A, Maffioletti E, Milanesi E, Marizzoni M, Frisoni GB, Blin O, Richardson JC, Bordet
R, Forloni G, Gennarelli M, Bocchio-Chiavetto L (2019) miR-146a and miR-181a are
involved in the progression of mild cognitive impairment to Alzheimer's disease.
Neurobiol Aging **82**, 102-109.

[15] Faura J, Bustamante A, Penalba A, Giralt D, Simats A, Martínez-Sáez E, Alcolea D, Fortea
J, Lleó A, Teunissen CE, van der Flier WM, Ibañez L, Harari O, Cruchaga C, Hernández-
Guillamón M, Delgado P, Montaner J (2020) CCL23: A Chemokine Associated with
Progression from Mild Cognitive Impairment to Alzheimer's Disease. *J Alzheimers Dis* **73**,
1585-1595.

[16] Silbert LC, Howieson DB, Dodge H, Kaye JA (2009) Cognitive impairment risk: white
matter hyperintensity progression matters. *Neurology* **73**, 120-125.

[17] Greene SJ, Killiany RJ (2010) Subregions of the inferior parietal lobule are affected in the
progression to Alzheimer's disease. *Neurobiol Aging* **31**, 1304-1311.

[18] Braak H, Thal DR, Ghebremedhin E, Del Tredici K (2011) Stages of the pathologic process
in Alzheimer disease: age categories from 1 to 100 years. *J Neuropathol Exp Neurol* **70**,
960-969.

[19] Pettigrew C, Soldan A, Wang J, Wang MC, Greenberg B, Albert M, Moghekar A, Team BR
(2022) Longitudinal CSF Alzheimer's disease biomarker changes from middle age to late
adulthood. *Alzheimers Dement (Amst)* **14**, e12374.

[20] Johnson KA, Fox NC, Sperling RA, Klunk WE (2012) Brain imaging in Alzheimer disease.
Cold Spring Harb Perspect Med **2**, a006213.

[21] Ghirelli A, Tosakulwong N, Weigand SD, Clark HM, Ali F, Botha H, Duffy JR, Utianski RL,
Buciuc M, Murray ME, Labuzan SA, Spychalla AJ, Pham NTT, Schwarz CG, Senjem ML,
Machulda MM, Baker M, Rademakers R, Filippi M, Jack CR, Jr., Lowe VJ, Parisi JE,
Dickson DW, Josephs KA, Whitwell JL (2020) Sensitivity-Specificity of Tau and Amyloid β
Positron Emission Tomography in Frontotemporal Lobar Degeneration. *Ann Neurol* **88**,
1009-1022.

[22] Soleimani-Meigooni DN, Iaccarino L, La Joie R, Baker S, Bourakova V, Boxer AL, Edwards
L, Eser R, Gorno-Tempini ML, Jagust WJ, Janabi M, Kramer JH, Lesman-Segev OH,
Mellinger T, Miller BL, Pham J, Rosen HJ, Spina S, Seeley WW, Strom A, Grinberg LT,
Rabinovici GD (2020) 18F-flortaucipir PET to autopsy comparisons in Alzheimer's disease
and other neurodegenerative diseases. *Brain* **143**, 3477-3494.

[23] Jack CR, Jr., Holtzman DM (2013) Biomarker modeling of Alzheimer's disease. *Neuron*
80, 1347-1358.

[24] Darst BF, Lu Q, Johnson SC, Engelman CD (2019) Integrated analysis of genomics,
longitudinal metabolomics, and Alzheimer's risk factors among 1,111 cohort
participants. *Genet Epidemiol* **43**, 657-674.

[25] Chételat G, Arbizu J, Barthel H, Garibotto V, Law I, Morbelli S, van de Giessen E, Agosta
F, Barkhof F, Brooks DJ, Carrillo MC, Dubois B, Fjell AM, Frisoni GB, Hansson O, Herholz
K, Hutton BF, Jack CR, Jr., Lammertsma AA, Landau SM, Minoshima S, Nobili F, Nordberg
A, Ossenkoppele R, Oyen WJG, Perani D, Rabinovici GD, Scheltens P, Villemagne VL,

597    Zetterberg H, Drzezga A (2020) Amyloid-PET and (18)F-FDG-PET in the diagnostic
598    investigation of Alzheimer's disease and other dementias. *Lancet Neurol* **19**, 951-962.
599 [26] Isaacs JD, Boenink M (2020) Biomarkers for dementia: too soon for routine clinical use.
600    *Lancet Neurol* **19**, 884-885.
601 [27] Bălașa AF, Chircov C, Grumezescu AM (2020) Body Fluid Biomarkers for Alzheimer's
602    Disease-An Up-To-Date Overview. *Biomedicines* **8**, 421.
603 [28] Hastie T, Tibshirani R, Friedman J (2009) The elements of statistical learning: prediction,
604    inference and data mining. *Springer-Verlag, New York*.
605 [29] Feng X, Chen L, Wang Z, Li SC (2020) I-Impute: a self-consistent method to impute single
606    cell RNA sequencing data. *BMC Genomics* **21**, 618.
607 [30] Li WV, Li JJ (2018) An accurate and robust imputation method scImpute for single-cell
608    RNA-seq data. *Nat Commun* **9**, 997.
609 [31] Gamazon ER, Wheeler HE, Shah KP, Mozaffari SV, Aquino-Michaels K, Carroll RJ, Eyler
610    AE, Denny JC, Nicolae DL, Cox NJ, Im HK (2015) A gene-based association method for
611    mapping traits using reference transcriptome data. *Nat Genet* **47**, 1091-1098.
612 [32] Nguyen M, He T, An L, Alexander DC, Feng J, Yeo BTT (2020) Predicting Alzheimer's
613    disease progression using deep recurrent neural networks. *Neuroimage* **222**, 117203.
614 [33] Nagpal S, Meng X, Epstein MP, Tsoi LC, Patrick M, Gibson G, De Jager PL, Bennett DA,
615    Wingo AP, Wingo TS, Yang J (2019) TIGAR: An Improved Bayesian Tool for
616    Transcriptomic Data Imputation Enhances Gene Mapping of Complex Traits. *Am J Hum*
617    *Genet* **105**, 258-266.
618 [34] Luningham JM, Chen J, Tang S, De Jager PL, Bennett DA, Buchman AS, Yang J (2020)
619    Bayesian Genome-wide TWAS Method to Leverage both cis- and trans-eQTL Information
620    through Summary Statistics. *Am J Hum Genet* **107**, 714-726.
621 [35] Bennett DA, Buchman AS, Boyle PA, Barnes LL, Wilson RS, Schneider JA (2018) Religious
622    Orders Study and Rush Memory and Aging Project. *J Alzheimers Dis* **64**, S161-S189.
623 [36] Bennett DA, Schneider JA, Wilson RS, Bienias JL, Arnold SE (2004) Neurofibrillary tangles
624    mediate the association of amyloid load with clinical Alzheimer disease and level of
625    cognitive function. *Arch Neurol* **61**, 378-384.
626 [37] Bennett DA, Wilson RS, Schneider JA, Evans DA, Aggarwal NT, Arnold SE, Cochran EJ,
627    Berry-Kravis E, Bienias JL (2003) Apolipoprotein E epsilon4 allele, AD pathology, and the
628    clinical expression of Alzheimer's disease. *Neurology* **60**, 246-252.
629 [38] Bennett DA, Wilson RS, Schneider JA, Evans DA, Mendes De Leon CF, Arnold SE, Barnes
630    LL, Bienias JL (2003) Education modifies the relation of AD pathology to level of
631    cognitive function in older persons. *Neurology* **60**, 1909-1915.
632 [39] Bennett DA, Wilson RS, Boyle PA, Buchman AS, Schneider JA (2012) Relation of
633    neuropathology to cognition in persons without cognitive impairment. *Ann Neurol* **72**,
634    599-609.
635 [40] (1997) Consensus recommendations for the postmortem diagnosis of Alzheimer's
636    disease. The National Institute on Aging, and Reagan Institute Working Group on
637    Diagnostic Criteria for the Neuropathological Assessment of Alzheimer's Disease.
638    *Neurobiol Aging* **18**, S1-2.

639   [41]   Wilson RS, Boyle PA, Yu L, Barnes LL, Sytsma J, Buchman AS, Bennett DA, Schneider JA
640   (2015) Temporal course and pathologic basis of unawareness of memory loss in
641   dementia. *Neurology* **85**, 984-991.
642   [42]   Boyle PA, Wilson RS, Aggarwal NT, Tang Y, Bennett DA (2006) Mild cognitive
643   impairment: risk of Alzheimer disease and rate of cognitive decline. *Neurology* **67**, 441-
644   445.
645   [43]   McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack CR, Jr., Kawas CH, Klunk WE,
646   Koroshetz WJ, Manly JJ, Mayeux R, Mohs RC, Morris JC, Rossor MN, Scheltens P, Carillo
647   MC, Thies B, Weintraub S, Phelps CH (2011) The diagnosis of dementia due to
648   Alzheimer's disease: Recommendations from the National Institute on Aging and the
649   Alzheimer's Association workgroup. *Alzheimers Dement*.
650   [44]   Bennett DA, Schneider JA, Arvanitakis Z, Kelly JF, Aggarwal NT, Shah RC, Wilson RS
651   (2006) Neuropathology of older persons without cognitive impairment from two
652   community-based studies. *Neurology* **66**, 1837-1844.
653   [45]   Zou H, Hastie T (2005) Regularization and variable selection via the elastic net. *Journal of*
654   *the Royal Statistical Society: Series B (Statistical Methodology)* **67**, 301-320.
655   [46]   Tibshirani R (1996) Regression Shrinkage and Selection via the Lasso. *Journal of the*
656   *Royal Statistical Society. Series B (Methodological)* **58**, 267-288.
657   [47]   Hoerl AE, Kennard RW (2000) Ridge Regression: Biased Estimation for Nonorthogonal
658   Problems. *Technometrics* **42**, 80-86.
659   [48]   Linden A (2006) Measuring diagnostic and predictive accuracy in disease management:
660   an introduction to receiver operating characteristic (ROC) analysis. *J Eval Clin Pract* **12**,
661   132-139.
662   [49]   Cox DR (1972) Regression models and life-tables. *Journal of the Royal Statistical Society:*
663   *Series B (Methodological)* **34**, 187-202.
664   [50]   Gerds TA, Scheike TH, Andersen PK (2012) Absolute risk regression for competing risks:
665   interpretation, link functions, and prediction. *Stat Med* **31**, 3921-3930.
666   [51]   Ozenne B, Sørensen AL, Scheike T, Torp-Pedersen C, Gerds TA (2017) riskRegression:
667   predicting the risk of an event using Cox regression models. *The R Journal* **9**, 440-460.
668   [52]   Powell MR, Smith GE, Knopman DS, Parisi JE, Boeve BF, Petersen RC, Ivnik RJ (2006)
669   Cognitive measures predict pathologic Alzheimer disease. *Arch Neurol* **63**, 865-868.
670   [53]   Dickerson BC, Sperling RA, Hyman BT, Albert MS, Blacker D (2007) Clinical prediction of
671   Alzheimer disease dementia across the spectrum of mild cognitive impairment. *Arch*
672   *Gen Psychiatry* **64**, 1443-1450.
673   [54]   Petersen RC, Smith GE, Ivnik RJ, Tangalos EG, Schaid DJ, Thibodeau SN, Kokmen E,
674   Waring SC, Kurland LT (1995) Apolipoprotein E status as a predictor of the development
675   of Alzheimer's disease in memory-impaired individuals. *JAMA* **273**, 1274-1278.
676   [55]   Bondi MW, Salmon DP, Galasko D, Thomas RG, Thal LJ (1999) Neuropsychological
677   function and apolipoprotein E genotype in the preclinical detection of Alzheimer's
678   disease. *Psychol Aging* **14**, 295-303.
679   [56]   Phongpreecha T, Cholerton B, Bukhari S, Chang AL, De Francesco D, Thuraiappah M,
680   Godrich D, Perna A, Becker MG, Ravindra NG, Espinosa C, Kim Y, Berson E, Mataraso S,
681   Sha SJ, Fox EJ, Montine KS, Baker LD, Craft S, White L, Poston KL, Beecham G,

Aghaeepour N, Montine TJ (2023) Prediction of neuropathologic lesions from clinical
data. *Alzheimers Dement* **19**, 3005-3018.
[57]	Schneider JA, Arvanitakis Z, Bang W, Bennett DA (2007) Mixed brain pathologies account
for most dementia cases in community-dwelling older persons. *Neurology* **69**, 2197-
2204.
[58]	Licher S, Yilmaz P, Leening MJG, Wolters FJ, Vernooij MW, Stephan BCM, Ikram MK,
Ikram MA (2018) External validation of four dementia prediction models for use in the
general community-dwelling population: a comparative analysis from the Rotterdam
Study. *Eur J Epidemiol* **33**, 645-655.
691

Figure Legends

Fig 1. Overall study design to develop and validate imputation models that infer
AD-NC traits based on clinical measures in older adults. A multi-stage analytic
approach was employed to develop, validate, and demonstrate the effectiveness of
inferred levels of four AD-NC traits derived from clinical measures as AD biomarkers. **A.**
We trained imputation models for four AD-NC traits using clinical data obtained at the last
visit before death in MAP decedents that underwent autopsy (**Fig 2**). Then we validated
these models in an independent cohort study (ROS) that collected the same clinical and
postmortem measures. **B**. We tested the effectiveness as AD biomarkers for the inferred
levels of four AD-NC traits at baseline, which were obtained by applying the validated
imputation models to clinical measures obtained at study entry. We examined if the
inferred baseline AD-NC traits predicted incident ADD (**Fig 3**) and discriminated adults at
risk for postmortem (on average 8 years after baseline) pathologic AD in ROS cohort (**Fig**
4).

Fig 2. **Machine learning methods were used to select different combinations of**
clinical measures to infer each of the four AD-NC traits. GLM-EN method was used
to train an imputation model for each of the four AD-NC traits. Standardized effect sizes
(beta) of selected predictive predictors with $|\text{beta}| > 0.01$ for each of the four imputation
models were plotted. The inferred values of each of the inferred AD-NC traits are
determined by the weighted averages of the corresponding selected predictors, with
weights given by estimated standard effect sizes. Although all four AD-NC traits are
inter-related and share cognition and *APOE E4* allele as important predictors, different
sets of selected predictors by their imputation models highlight that different
combinations of clinical measures with different effect sizes are necessary for inferring
the unique features of these inter-related AD-NC traits.
717
Fig 3. Inferred baseline AD-NC traits predicted incident Alzheimer's Disease
Dementia (ADD). We used Cox proportional hazard models to examine the predictivity
of each of the inferred baseline AD-NC traits along with age, sex, and education
covariates for incident ADD in 3 and 5 years after study entry. Top four panels show the
prediction accuracies (ROC plots) with each the four inferred baseline AD-NC traits in
adults without dementia (NCI+MCI) at study entry. Bottom four panels show prediction
accuracies with each of the four AD-NC traits in adults with NCI at study entry. As
expected for an effective AD biomarker, each of the inferred baseline AD-NC traits
predicted ADD.

Fig 4. Inferred AD-NC traits at study baseline discriminated postmortem pathologic
AD profiled at autopsy. Pathologic AD here is the binary postmortem NIA-Reagan status
profiled at autopsy, with value 1 representing pathologic AD (teal boxplots) and 0
representing no pathologic AD (red boxplots). Two-sample t-test p-values are 1.4×10^{-10}
for amyloid-β (A), 1.2×10^{-11} for tangles (B), 9.8×10^{-12} for global AD pathology (C), and
1.6×10^{-10} for pathologic AD (D).
733

Table 1. Prediction accuracy (with 95% confidence interval) and sensitivity with respect to selected risk score thresholds that ensure ~80% specificity by Cox models using a single inferred AD-NC trait. Values in this table are reflecting the Cox risk model prediction performance at one point in the ROC curves as shown in **Fig 3**, with corresponding risk score thresholds. Samples with predicted risk scores greater than the selected threshold were considered as Predicted Positives (incident ADD), otherwise Predicted Negatives (not developing ADD).

		Amyloid-β		Tangles		Global AD Pathology		Pathologic AD	
		Y3	*Y5*	*Y3*	*Y5*	*Y3*	*Y5*	*Y3*	*Y5*
NCI/MCI -> ADD	*Accuracy[a] (95% CI)*	*0.798 (0.773, 0.822)*	*0.788 (0.763, 0.813)*	*0.802 (0.777, 0.825)*	*0.795 (0.771, 0.818)*	*0.801 (0.775, 0.823)*	*0.796 (0.771, 0.819)*	*0.809 (0.784, 0.831)*	*0.803 (0.778, 0.826)*
	Sensitivity[b]	*0.772*	*0.699*	*0.822*	*0.756*	*0.797*	*0.764*	*0.911*	*0.829*
	Specificity[c]	*0.800*	*0.800*	*0.800*	*0.800*	*0.800*	*0.800*	*0.800*	*0.800*
NCI -> ADD	*Accuracy[a] (95% CI)*	*0.796 (0.768, 0.823)*	*0.792 (0.763, 0.7819)*	*0.794 (0.765, 0.821)*	*0.791 (0.761, 0.818)*	*0.795 (0.767, 0.823)*	*0.793 (0.764, 0.820)*	*0.804 (0.775, 0.830)*	*0.805 (0.777, 0.832)*
	Sensitivity[b]	*0.650*	*0.634*	*0.550*	*0.609*	*0.600*	*0.658*	*0.950*	*0.902*
	Specificity[c]	*0.800*	*0.800*	*0.800*	*0.800*	*0.800*	*0.800*	*0.800*	*0.800*

a. Accuracy= (# True Positive Predictions + # True Negative Predictions) / (# of test samples)

b. Sensitivity = (# True Positive Predictions) / (# Positives in test samples) = True Positive Fraction

c. Specificity = (# True Negative Predictions) / (# Negatives in test samples) = 1−False Positive Fraction

Fig 1. Overall study design to develop and validate imputation models that infer AD-NC traits based on clinical measures in older adults. A multi-stage analytic approach was employed to develop, validate, and demonstrate the effectiveness of inferred levels of four AD-NC traits derived from clinical measures as AD biomarkers. **A.** We trained imputation models for four AD-NC traits using clinical data obtained at the last visit before death in MAP decedents that underwent autopsy (**Fig 2**). Then we validated these models in an independent cohort study (ROS) that collected the same clinical and postmortem measures. **B.** We tested the effectiveness as AD biomarkers for the inferred levels of four AD-NC traits at baseline, which were obtained by applying the validated imputation models to clinical measures obtained at study entry. We examined if the inferred baseline AD-NC traits predicted incident ADD (**Fig 3**) and discriminated adults at risk for postmortem (on average 8 years after baseline) pathologic AD in ROS cohort (**Fig 4**).

Fig 2. **Machine learning methods were used to select different combinations of clinical measures to infer each of the four AD-NC traits.** GLM-EN method was used to train an imputation model for each of the four AD-NC traits. Standardized effect sizes (beta) of selected predictive predictors with |beta| > 0.01 for each of the four imputation models were plotted. The inferred values of each of the inferred AD-NC traits are determined by the weighted averages of the corresponding selected predictors, with weights given by estimated standard effect sizes. Although all four AD-NC traits are inter-related and share cognition and *APOE E4* allele as important predictors, different sets of selected predictors by their imputation models highlight that different combinations of clinical measures with different effect sizes are necessary for inferring the unique features of these inter-related AD-NC traits.

Fig 3. Inferred baseline AD-NC traits predicted incident Alzheimer's Disease Dementia (ADD). We used Cox proportional hazard models to examine the predictivity of each of the inferred baseline AD-NC traits along with age, sex, and education covariates for incident ADD in 3 and 5 years after study entry. Top four panels show the prediction accuracies (ROC plots) with each the four inferred baseline AD-NC traits in adults without dementia (NCI+MCI) at study entry. Bottom four panels show prediction accuracies with each of the four AD-NC traits in adults with NCI at study entry. As expected for an effective AD biomarker, each of the inferred baseline AD-NC traits predicted ADD.

Fig 4. Inferred AD-NC traits at study baseline discriminated postmortem pathologic AD profiled at autopsy. Pathologic AD here is the binary postmortem NIA-Reagan status profiled at autopsy, with value 1 representing pathologic AD (teal boxplots) and 0 representing no pathologic AD (red boxplots). Two-sample t-test p-values are 1.4×10^{-10} for amyloid-β (A), 1.2×10^{-11} for tangles (B), 9.8×10^{-12} for global AD pathology (C), and 1.6×10^{-10} for pathologic AD (D).